

Validation of the SCOFF Questionnaire for Eating Disorders in a Multiethnic General Population Sample

Francesca Solmi, PhD^{1*}
Stephani L. Hatch, PhD²
Matthew Hotopf, PhD²
Janet Treasure, PhD³
Nadia Micali, PhD¹

ABSTRACT

Introduction: This study aimed to validate the SCOFF, an eating disorders (ED) screening questionnaire, in a multiethnic general population sample of adults.

Method: A two-stage design was employed using the South East London Community Health Study phases I and II data. A total of 1,669 participants were screened using the SCOFF in SELCoHI, and 145 were administered an ED clinical interview in SELCoHII. We explored the diagnostic validity of the questionnaire restricting to the 145 individuals with the clinical questionnaire.

Results: Sensitivity and specificity of the SCOFF were 53.7 and 93.5%, respectively.

Conclusion: The SCOFF showed good levels of specificity but low sensitivity, resulting in a high percentage of false negatives. Given the low sensitivity found in our sample the SCOFF is likely to be a suboptimal measure for the identification of ED in the community. © 2014 The Authors International Journal of Eating Disorders Published by Wiley Periodicals, Inc.

Keywords: SCOFF; eating disorders; diagnostic validity

(*Int J Eat Disord* 2015; 48:312-316)

Introduction

Eating disorders (ED) and other specified feeding or eating disorders (OSFED) [the umbrella definition for subthreshold ED in the Diagnostic and Statistical Manual of mental disorders 5th ed (DSM-5)] have a lifetime prevalence of up to 9%¹ and are associated with considerable physical and psychiatric comorbidity.² However, ED are often undetected in the general population resulting in small proportions of individuals receiving treatment.³

The introduction of the SCOFF (an acronym describing five key screening questions for ED, which can be recalled through the mnemonic 'Sick,

Control, One stone, Fat, Food')⁴ as a screening tool for ED in clinical settings has opened a window of opportunity for extending its use to routine screenings in the general population. The SCOFF has been validated in a number of primary-care based studies; in the UK⁴⁻⁷ and internationally,⁸⁻¹³ both in written and oral delivery;¹⁴ and compared to other instruments.¹⁵ Early studies have employed clinical ED cases and controls,^{4,5,7} used mainly female populations^{5,7,15-17} and few have included individuals older than 40 years of age.^{6,16,18} Overall, studies using clinical populations have yielded higher values of sensitivity (Se) and specificity (Sp)^{5,6,18} than community studies, the latter usually finding higher Sp than Se.¹¹⁻¹³

However, most validation studies in the community have relied on young^{8,11} and homogeneous populations^{6,12} (i.e., females, limited ethnic representation), limiting the scope for the generalizability of these results. Therefore, in order to investigate the suitability of the SCOFF as both a screening tool at the community level and in general population surveys, we aimed to pilot a validation of the instrument in a multiethnic population-based sample of adults aged 16-90.

Method

Sample, Measures

This study employed data from the South East London Community Health Study (SELCoH) I and II, a two-phase

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Accepted 12 November 2014

Additional Supporting Information may be found in the online version of this article.

*Correspondence to: Francesca Solmi, Ph.D.; Behavioural and Brain Science Unit, Institute of Child Health, 30 Guildford St, London WC1N 1EH, United Kingdom
E-mail: francesca.solmi@ucl.ac.uk

¹ Behavioural and Brain Sciences Unit, Institute of Child Health, University College London, London, United Kingdom

² Psychological Medicine, Institute of Psychiatry, King's College London, London, United Kingdom

³ Eating Disorders Unit, Institute of Psychiatry, King's College London, London, United Kingdom

Published online 12 December 2014 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/eat.22373

© 2014 The Authors International Journal of Eating Disorders Published by Wiley Periodicals, Inc.

general population survey of 1,669 individuals aged 16+ living in the London (UK) boroughs of Lambeth and Southwark. More details on the rationale, sampling, representativeness, and assessment of participants in SELCoHI is provided elsewhere.¹⁹ In SELCoHI, 1,669 participants completed the SCOFF questionnaire and underwent objective anthropometric measurements to calculate body mass index (BMI). In SELCoHII, participants who had given their consent to be re-contacted in SELCoHII and did not need an interpreter ($N = 1,560$) were eligible for inclusion in a clinical assessment of ED using the ED section of the Structured Clinical Interview for DSM-IV Axis I Disorders nonpatient edition (SCID-I-NP),²⁰ if they: (i) had screened positive at the SCOFF in SELCoHI ($N = 158$); or (ii) had screened negative and had not screened positive for other mental health conditions ($N = 599$). SCOFF positive participants were gender matched with a randomly selected sample of eligible screen negatives.

Participants were asked for ED symptoms occurred at the time of and since the SELCoHI assessment when answering SCID-I questions. Despite referring to DSM-IV diagnosis, the SCID-I also contains a section on binge eating disorder (BED), which means that all ED diagnoses were explored. Moreover, interviewers did not apply the 'skip-rules' of the SCID-I in order to avoid underestimating the prevalence of diagnoses²¹ and gathered information on type, frequency, and duration of ED behaviors in order to be able to subsequently derive DSM5 diagnoses. SELCoHI data was collected between June 2008 and December 2010 and SELCoHII data between August 2011 and March 2013. Consent was collected prior to participation to the study.

Analyses

Sample characteristics were described using cross tabulations and chi-square tests. Sensitivity, specificity, and positive and negative predictive values (PPV, NPV) were calculated on the sample interviewed in SELCoHII ($N = 145$) using sampling weights accounting for: (i) proportion of SCOFF positive and negative participants interviewed with the SCID over the whole sample who completed the SCOFF who agreed to take part to SELCoHII (N total: 1,560; screen positive: 158; screen negative: 1402); and (ii) proportion of ED diagnoses amongst screen positives and negatives, as previously recommended in two-phase epidemiological studies.²² These weights account for the real prevalence of the condition in the population when circumstances do not allow maintaining adequate sampling ratios, but do not account for characteristics associated with participation at follow-up (e.g., age, gender, and ethnicity). We weighted our data based on the sample of the 1,560 participants who agreed to be followed up, on the *a priori* knowledge that we could have not assessed the real ED status of the remaining 109 participants with the

SCID-I and that limiting analyses to participants eligible for inclusion (i.e., without mental health comorbidities, common in individuals with ED) could have underestimated the number of false negatives and, thus overestimated the sensitivity. As a sensitivity analysis, we calculated sensitivity and specificity using the whole initial sample ($N = 1,669$) and the results did not change qualitatively (Se: 52.8; Sp: 93.7). Analyses were run in Stata12.

Results

Sample

Of the 322 participants who were eligible to take part to the ED module (158 screen positive and 164 screen negatives), 89 (56.3%) participants who had screened positive on the SCOFF and 88 (53.6%) of those who screened negative were lost to follow up. As seen in Supporting Information Table S1, no systematic differences existed between participants who took part in the study and those who were lost to follow-up with respect to age, marital status, ethnicity, education, and age. Among screen positive participants, however, more obese (67%) and underweight (100.0%) participants were lost to follow-up.

A total of 145 participants [76 (46.3%) SCOFF negative, 69 (43.7%) SCOFF positive] were assessed using the SCID-I interview. Of these, 31 (21.4%) received a threshold or subthreshold ED diagnosis. The majority of participants were female (75%), of White ethnicity (43%), between the ages of 25 and 34 years (28%), with a normal BMI (54%) and had at least a General Certificate of Secondary Education (GCSE) qualification (90%). No participant was underweight (**Table 1**).

Diagnostic Validity

DSM5 diagnosis was correctly predicted by the SCOFF for a total of 101 (69.7%; $N = 73$: no ED; $N = 28$: ED) participants; 3 (2.0%) participants were misclassified by the SCOFF as not having an ED and 41 (28.3%) as having an ED. Two of the three false negative participants had a diagnosis of binge eating disorder (BED) and one of OSFED presenting with excessive exercise (not in table). All false negatives were women, of White ethnicity, between the ages of 35 and 54 and ~60% were obese (Supporting Information **Table S2**).

Based on the established cutoff of ≥ 2 positive answers, the weighted sensitivity, specificity, PPV and NPV of the SCOFF were 53.7 (95% CI: 36.2–71.2), 93.5 (95% CI: 88.9–98.0), 40.6 (95% CI: 28.9–53.1), and 96.1 (95% CI: 88.9–99.2), respectively.

TABLE 1. Socio-demographic characteristics of the interviewed sample in SELCoHII

Socio-Demographic Characteristics	N (%)
Total	145 (100)
Gender	
Male	36 (24.8)
Female	109 (75.2)
Ethnicity	
White	83 (57.2)
Black	42 (29)
Asian	4 (2.8)
Other	16 (11)
Education	
No qualification	14 (9.7)
GCSE/A-level	70 (48.3)
Degree level or above	61 (42)
BMI	
Underweight	0 (0)
Normal weight	75 (54.3)
Overweight	31 (22.5)
Obese	32 (23.2)
Age	
16–24	38 (26.2)
25–34	41 (28.3)
35–44	23 (15.9)
45–54	26 (17.9)
55–64	9 (6.2)
65+	8 (5.5)

Discussion

This study aimed to assess the validity of the SCOFF as a screening tool for ED in a multiethnic general population sample of adults in London (UK).

In line with previous general population studies, we found that specificity of the instrument was higher than its sensitivity, and that the latter was lower than what previously found in some studies,^{11,12} but not others.¹³ All of these studies employed a younger population than ours. PPV was low, which is common for low prevalence conditions, but NPV was high. Low sensitivity suggests that high proportions of individuals with an ED are not identified by the SCOFF.

Several factors could account for this finding. It is possible that, in the absence of follow-up questions (i.e., such as probing questions contained in the SCID) the ego-syntonicity of ED could lead to negative answers. Moreover, the focus on the fat/thin dichotomy in assessing body dissatisfaction could introduce gender biases not accounting for different cognitions in men²³ or in individuals who are overweight or obese. Although the former were not represented amongst the participants who were false negatives in our sample, two of the three of false negatives were obese. More research using larger mixed gender sample is warranted to test the validity of the SCOFF in men. The limited numbers of questions contained in the SCOFF could also mean that important behaviors (e.g., laxative use, exces-

TABLE 2. Summary of weighted diagnostic validity measures for the SCOFF questionnaire

Diagnostic Measures	Values (95%CI)
Sensitivity	53.7 (36.2–71.2)
Specificity	93.5 (88.9–98.0)
Positive predictive value	40.6 (28.9–53.1)
Negative predicted value	96.1 (88.9–99.2)

sive exercise) central to the diagnosis of bulimia nervosa or OSFED such as purging disorder, are missed, as in the case of the participant presenting with excessive exercise. Similarly, although we could not provide evidence of this from our sample, it can be speculated that without a measure of BMI, cases of AN where the individual is stable on a low, unhealthy BMI, could be missed as question 3 of the SCOFF only enquires about recent weight loss. Finally, the SCOFF was developed when BED was not yet a recognized diagnosis and it is possible that its questions are limited in identifying the condition, which might be more prevalent in older adults, as was the case amongst false negatives in our sample. This could explain the lower sensitivity we found compared to other community studies employing younger participants.

This study has several strengths. It employed a large representative and ethnically diverse general population sample, suggesting that findings are generalizable to similarly diverse populations. Although the validation was conducted on a subsample of individuals (less than 10% of the full study population), sampling weights were employed to account for differential sampling across screen positives and negatives to ensure that estimates of sensitivity and specificity reflected the prevalence of ED in the whole sample. However, some limitations should also be accounted for. The SCID interview was conducted 2–3 years following the administration of the SCOFF; recall bias and regression to the mean could thus have occurred to some extent. Since ED are chronic conditions and no incident cases (i.e., new ED onsets between the administration of the SCOFF and the SCID) were found in interviews, we suggest that the diagnoses were not greatly under- or over-estimated and that false negatives could not be attributed to ED cases with onset occurring after the administration of the SCOFF. It is possible, however, that different recall patterns could have occurred. We tried to minimize this by asking about present and past ED behaviors in general and then for their duration in order to identify overlaps with the time of the SCOFF interview. Interviewers were also blinded as to the screening status of participants in order to avoid observer bias. Substantial losses to follow-up occurred and whilst the sample of screen

negative seems to be representative of the overall sample (Supporting Information Table S1), 4 (100%) underweight participants amongst the screen positives lost to follow-up could index missed ED cases (especially AN) (Supporting Information Table S1). Moreover, it appears that losses to follow-up in the screen positive group could have occurred with respect to participants of Asian and other ethnic backgrounds (Supporting Information Table S1). This could introduce some degree of selection bias, which our sampling weights could not account for. However, given the small proportion and the lower weight assigned to screen positive participants interviewed compared to screen negatives, this is unlikely to bias the overall sensitivity and specificity estimates.^{11,12} Only participants with no mental health comorbidities were eligible to be assessed in SELCoHII and were therefore interviewed. Although we weighted Se and Sp to represent the whole sample that agreed to be followed up, we could be overestimating or underestimating Se. On the one hand, as ED are comorbid with a number of psychiatric conditions²⁴ and more false negatives could have occurred and been missed in the noneligible sample. On the other hand, underestimation could also have occurred, as the false negatives ratio was based on three individuals only and, although weights were applied, uncertainty around the estimate (reflected in the wide 95% CI) exists. Future studies should aim at conducting both interviews simultaneously and on the whole sample to improve accuracy of findings.

Recent studies found high levels of psychiatric comorbidity in individuals who screened positive on the SCOFF.^{25–27} Given its high specificity, this is likely to reflect the high levels of psychopathology in individuals with ED. However, although good at ruling out an ED (i.e., high specificity) the SCOFF should be used with caution as an ED screening tool, as the low sensitivity in our sample indicates that a substantial number of individuals with ED might be missed. This is particularly relevant to population-based samples and suggestions of using the SCOFF as a screening tool in the community.¹³ Based on our findings, more research is needed to assess whether rephrasing some of its questions could improve the diagnostic validity of the SCOFF without compromising its brevity. Moreover, research exploring the diagnostic validity of the SCOFF across different population subgroups (e.g., overweight and obese, males, ethnic minorities) is warranted to improve the measure.

SLH and MH receive salary support from the National Institute for Health Research (NIHR) Mental Health Biomedical Research Centre at South London and Maudsley

NHS Foundation Trust and Institute of Psychiatry, King's College London. This research was supported by the Biomedical Research Nucleus data management and informatics facility at South London and Maudsley NHS Foundation Trust, which is funded by the National Institute for Health Research (NIHR) Mental Health Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London and a joint infrastructure grant from Guy's and St Thomas' Charity and the Maudsley Charity. This research was also funded by a National Institute of Health Research (NIHR) clinician scientist award to Dr N Micali and by a grant received by the British Academy. These funders had no involvement in study design, data collection, analysis or the decision to submit for publication. The authors have no financial involvement (including employment, fees, share ownership) or affiliation with any organisation whose financial interests may be affected by material in the manuscript, or which might potentially bias it. This publication is the work of the authors and Nadia Micali will serve as guarantors for the contents of this paper. The authors are grateful for the support received the Open Access Funding Team at the University College London Library Services towards publishing this manuscript as an open access publication.

References

1. Smink FRE, van Hoeken D, Hoek HW. Epidemiology of eating disorders: Incidence, prevalence and mortality rates. *Curr Psychiatry Rep* 2012;14:406–414.
2. Hudson JI, Hiripi E, Pope, Jr., HG, Kessler RC. The prevalence and correlates of eating disorders in the national comorbidity survey replication. *Biol Psychiatry* 2007;61:348–358.
3. Mond JM, Hay PJ, Rodgers B, Owen C. Health service utilization for eating disorders: findings from a community-based study. *Int J Eat Disord* 2007;40:399–408.
4. Morgan JF, Reid F, Lacey JH. The SCOFF questionnaire: A new screening tool for eating disorders. *West J Med* 2000;172:164–165.
5. Morgan JF, Reid F, Lacey JH. The SCOFF questionnaire: Assessment of a new screening tool for eating disorders. *BMJ* 1999;319:1467–1468.
6. Luck AJ. The SCOFF questionnaire and clinical interview for eating disorders in general practice: comparative study. *BMJ* 2002;325:755–756.
7. Hill LS, Reid F, Morgan JF, Lacey JH. SCOFF, the development of an eating disorder screening questionnaire. *Int J Eat Disord* 2010;43:344–351.
8. Muro-Sans P, Amador-Campos JA, Morgan JF. The SCOFF-c: Psychometric properties of the Catalan version in a Spanish adolescent sample. *J Psychosom Res* 2008;64:81–86.
9. Garcia FD, Grigioni S, Chelali S, Meyrignac G, Thibaut F, Dechelotte P. Validation of the French version of SCOFF questionnaire for screening of eating disorders among adults. *World J Biol Psychiatry Inform Healthcare Stockholm* 2010;11:888–893.
10. Pannocchia L, Fiorino M, Giannini M, Vanderlinden J. A psychometric exploration of an Italian translation of the SCOFF questionnaire. *Eur Eat Disord Rev* 2011;19:371–373.
11. Leung SF, Lee KL, Lee SM, Leung SC, Hung WS, Lee WL, et al. Psychometric properties of the SCOFF questionnaire (Chinese version) for screening eating disorders in Hong Kong secondary school students: A cross-sectional study. *Int J Nurs Stud* 2009;46:239–247.

12. Lähteenmäki S, Aalto-Setälä T, Suokas JT, Saarni SE, Perälä J, Saarni SI, et al. Validation of the Finnish version of the SCOFF questionnaire among young adults aged 20 to 35 years. *BMC Psychiatry* 2009;9:5.
13. Parker SC, Lyons J, Bonner J. Eating disorders in graduate students: Exploring the SCOFF questionnaire as a simple screening tool. *J Am Coll Health* Routledge 2005;54:103–107.
14. Perry L, Morgan J, Reid F, Brunton J, O'Brien A, Luck A, et al. Screening for symptoms of eating disorders: Reliability of the SCOFF screening tool with written compared to oral delivery. *Int J Eat Disord* 2002;32:466–472.
15. Mond JM, Myers TC, Crosby RD, Hay PJ, Rodgers B, Morgan JF, et al. Screening for eating disorders in primary care: EDE-Q versus SCOFF. *Behav Res Ther* 2008;46:612–622.
16. Cotton M-A, Ball C, Robinson P. Four simple questions can help screen for eating disorders. *J Gen Intern Med* 2003;18:53–56.
17. Siervo M, Boschi V, Papa A, Bellini O, Falconi C. Application of the SCOFF, Eating Attitude Test 26 (EAT 26) and Eating Inventory (TFEQ) Questionnaires in young women seeking diet-therapy. *Eat Weight Disord* 2005;10:76–82.
18. Garcia-Campayo J, Sanz-Carrillo C, Ibañez JA, Lou S, Solano V, Alda M. Validation of the Spanish version of the SCOFF questionnaire for the screening of eating disorders in primary care. *J Psychosom Res* 2005;59:51–55.
19. Hatch SL, Frissa S, Verdecchia M, Stewart R, Fear NT, Reichenberg A, et al. Identifying socio-demographic and socioeconomic determinants of health inequalities in a diverse London community: The South East London Community Health (SELCoH) study. *BMC Public Health* 2011;11:861.
20. First MB, Spitzer RL, Miriam G, Williams JB. Structured Clinical Interview for DSM-IV-TR Axis I Disorders, Research Version, Non-patient Edition (SCID-I/NP). Biometrics Research NYSPI, editor. New York, 2002.
21. Swanson SA, Crow SJ, Le Grange D, Swendsen J, Merikangas KR. Prevalence and correlates of eating disorders in adolescents. Results from the national comorbidity survey replication adolescent supplement. *Arch Gen Psychiatry* 2011;68:714–723.
22. Dunn G, Pickles A, Tansella M, Vazquez-Barquero L. Two-phase epidemiological surveys in psychiatric research. *Br J Psychiatry* 1999;174:95–100.
23. Strother E, Lemberg R, Stanford SC, Turberville D. Eating disorders in men: Underdiagnosed, undertreated, and misunderstood. *Eat Disord* Routledge 2012;20:346–355.
24. O'Brien KM, Vincent NK. Psychiatric comorbidity in anorexia and bulimia nervosa: Nature, prevalence, and causal relationships. *Clin Psychol Rev* 2003;23:57–74.
25. Herpertz-Dahlmann B, Wille N, Hölling H, Vloet TD, Ravens-Sieberer U. Disordered eating behaviour and attitudes, associated psychopathology and health-related quality of life: Results of the BELLA study. *Eur Child Adolesc Psychiatry* 2008;17 Suppl 1:82–91.
26. McBride O, McManus S, Thompson J, Palmer RL, Brugha T. Profiling disordered eating patterns and body mass index (BMI) in the English general population. *Soc Psychiatry Psychiatr Epidemiol* 2012;48:783–793.
27. Solmi F, Hatch SL, Hotopf M, Treasure J, Micali N. Prevalence and correlates of disordered eating in a general population sample: The South East London Community Health (SELCoH) study. *Soc Psychiatry Psychiatr Epidemiol* 2014; 49:1335–1346.