



Published in final edited form as:

Cancer Prev Res (Phila). 2014 September ; 7(9): 886–895. doi:10.1158/1940-6207.CAPR-14-0058.

Plasma Tocopherols and Risk of Prostate Cancer in the Selenium and Vitamin E Cancer Prevention Trial (SELECT)

Demetrius Albanes¹, Cathee Till², Eric A. Klein³, Phyllis J. Goodman², Alison M. Mondul¹, Stephanie J. Weinstein¹, Philip R. Taylor¹, Howard L. Parnes⁴, J. Michael Gaziano⁵, Xiaoling Song⁶, Neil E. Fleshner⁷, Powel H. Brown⁸, Frank L. Meyskens Jr.⁹, and Ian M. Thompson¹⁰

¹Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, MD

²SWOG Statistical Center, Fred Hutchinson Cancer Research Center, Seattle, WA

³Department of Urology, Glickman Urological and Kidney Institute, Cleveland Clinic, Cleveland, OH

⁴Division of Cancer Prevention, National Cancer Institute, Bethesda, MD

⁵Massachusetts Veteran's Epidemiology, Research and Information Center, VA Boston Healthcare System, Boston, MA

⁶Biomarker Laboratory, Fred Hutchinson Cancer Research Center, Seattle, WA

⁷Princess Margaret Hospital, University of Toronto, Toronto, Ontario, Canada

⁸The University of Texas MD Anderson Cancer Center, Houston, TX

⁹University of California at Irvine, Department of Medicine, Orange, CA

¹⁰Cancer Therapy and Research Center, University of Texas Health Science Center, San Antonio, TX

Abstract

The Selenium and Vitamin E Cancer Prevention Trial (SELECT) showed higher prostate cancer incidence in men supplemented with high-dose α -tocopherol. We therefore examined whether pre-supplementation plasma α -tocopherol or γ -tocopherol was associated with overall or high-grade prostate cancer. A stratified case-cohort sample that included 1,746 incident prostate cancer cases diagnosed through June, 2009 and a subcohort of 3,211 men was derived from the SELECT trial of 35,533 men. Plasma was collected at entry in 2001–2004, and median follow-up was 5.5 years (range, 0 – 7.9 years). Incidence of prostate cancer as a function of plasma α -tocopherol, γ -tocopherol, and supplementation with α -tocopherol or selenomethionine was estimated by the hazard ratio (HR). Plasma γ -tocopherol was not associated with prostate cancer. Men with higher α -tocopherol concentrations appeared to have risk similar to that of men with lower concentrations [overall HR for fifth (Q5) vs. first quintile (Q1), 1.21 (95% confidence interval (CI), 0.88–1.66, *P*-

Corresponding Author: Demetrius Albanes, Division of Cancer Epidemiology and Genetics, National Cancer Institute, NIH, 9609 Medical Center Drive, Room 6E342, Bethesda, MD, 20892-9704. Phone: 240-276-7212; Fax: 240-276-7837; daa@nih.gov.

Conflicts of interest: No conflicts to disclose.

trend=0.24; in the trial placebo arm, Q5 HR, 0.85, 95% CI, 0.44–1.62, *P*-trend=0.66]. We found a strong positive plasma α -tocopherol association among men receiving the trial selenomethionine supplement [Q5 HR, 2.04, 95% CI, 1.29–3.22; *P*-trend=0.005]. A positive plasma α -tocopherol-prostate cancer association also appeared limited to high-grade disease (Gleason grade 7–10, overall Q5 HR, 1.59, 95% CI, 1.13–2.24, *P*-trend=0.001; among men receiving selenomethionine, HR, 2.12, 95% CI, 1.32–3.40; *P*-trend=0.0002). Our findings indicate that higher plasma α -tocopherol concentrations may interact with selenomethionine supplements to increase high-grade prostate cancer risk, suggesting a biological interaction between α -tocopherol and selenium itself or selenomethionine.

Keywords

α -tocopherol; γ -tocopherol; vitamin E; selenium; methionine; chemoprevention; randomized trial; prostate cancer; case-cohort study

Introduction

How vitamin E influences prostate carcinogenesis and cancer risk is not fully understood. Laboratory studies point to anti-prostate-tumorigenic properties for most of the eight vitamin E compounds (including four tocopherols and four tocotrienols), but findings from large controlled trials have been contradictory. Whereas the Alpha-Tocopherol, Beta-Carotene Cancer Prevention (ATBC) Study found a 32% and 41% reduction in prostate cancer incidence and mortality, respectively, in male smokers taking α -tocopheryl-acetate 50 IU daily (1), the Selenium and Vitamin E Cancer Prevention Trial (SELECT) showed 17% higher prostate cancer incidence rates among men supplemented with 400 IU daily compared to those receiving placebo (2,3), and the contemporaneous Physicians' Health Study (PHS) II found no effect using a 400 IU dose every other day (4). At the same time, observational studies indicate that higher vitamin E status (e.g., plasma α -tocopherol) is associated with reduced risk of prostate cancer, particularly among smokers, and possibly only for aggressive disease (5–13). It is therefore possible that physiological vitamin E status and high-dose vitamin E supplementation have independent, synergistic or antagonistic effects on prostate carcinogenesis.

To address this timely issue, we examined pre-randomization plasma concentrations of two key vitamin E compounds for humans, α -tocopherol and γ -tocopherol, in relation to risk of developing prostate cancer, overall and within each of the four intervention arms, in a case-cohort study of 4,754 men within SELECT. The plasma tocopherol-risk association was also evaluated according to biologically relevant factors including Gleason grade, time from randomization to prostate cancer diagnosis, and smoking status. The impact of the trial's α -tocopherol and selenomethionine supplementation on prostate cancer risk was also examined within categories of the pre-randomization plasma concentrations.

Materials and Methods

Study population

SELECT was a randomized, double-blind, placebo-controlled trial of daily supplementation of selenium (200 µg elemental selenium as *L*-selenomethionine), vitamin E (400 IU of *all-rac*- α -tocopheryl acetate), both, or placebo (14). African-American men \geq 50 years and non-African-American men \geq 55 years (total $n = 35,533$) were recruited from 427 sites in the United States, Canada, and Puerto Rico between August, 2001 and June, 2004. Men with a prostate-specific antigen level >4 ng/ml or a digital rectal examination suspicious for prostate cancer were ineligible. Study supplementation continued through October, 2008 (2), after which approximately 18,000 men were enrolled into the observational cohort follow-up study (15). Sociodemographic and risk factor information was collected at baseline including age, race, diet, vitamin supplement use, smoking status, medical history, and family history of cancer (16,17). Participants were monitored twice annually for blood pressure, weight, smoking status, and new medical events. Information regarding the standard-of-care monitoring and biopsy cores for SELECT has been reported previously (2,16). The study was approved by the local institutional review boards of each study site, and all participants provided written informed consent (Clinicaltrials.gov Identifier: NCT00006392).

Incident prostate cancer cases were self-reported and confirmed by central review. Medical records were abstracted for diagnostic method and clinical stage, and prostate tissue with corresponding pathology reports were sent for central pathology review, diagnostic confirmation, and Gleason score determination (2). A total of 1,856 prostate cancer cases serving as the primary endpoints were identified through June, 2009, including 1,746 with plasma available for tocopherol, phospholipid (18,19), and other analyses. A sub-cohort, representative of the SELECT participants, was then created *a priori* as the comparison group for all biomarker studies of prostate cancer and other endpoints. Men were sampled from nine race/age categories of, 1) African-American vs. not, and, 2) ages < 55 (African-Americans only), 55–59, 60–64, 65–70, and ≥ 70 years for both African-Americans and others. For each case, men were selected for the sub-cohort at random from each race/age category using a ratio of 1 case to 3 controls for the African-American categories, and 1 to 1.5 for others. The higher proportion of controls for African-Americans was aimed at boosting study power for that subgroup. Annually beginning in 2005, the number of prostate cancer cases in each of the nine race/age strata was determined, and sub-cohort members were randomly selected within each stratum, based on the sampling ratio. The sub-cohort was comprised of 3,211 individuals (including some of the cases) for this analysis, with a total combined prostate cancer-sub-cohort sample of 4,754 men whose follow-up time ranged from 0 to 7.9 years (median, 5.5 years). Supplementary Table S1 shows the age and race groups of the sub-cohort.

Plasma tocopherols

At randomization, participants provided a blood sample (EDTA), collected at least 3 hours after a meal, processed for plasma, buffy coat, and red blood cells, and stored at -70°C (available for 84% of the SELECT participants) (17). All cases and a representative fraction

of the cohort were assayed annually to minimize batch and storage time effects. Plasma α -tocopherol and γ -tocopherol concentrations were determined by reverse-phase high-performance liquid chromatography as described (20). Cases and sub-cohort participant plasma samples randomly were ordered within individual batches along with 381 blinded pooled quality control (QC) duplicate and triplicate samples. The weighted mean coefficients of variation (CV) were 3.1% and 4.5% for α -tocopherol and γ -tocopherol, respectively. Tocopherol concentrations did not differ based on year of plasma collection, with median α -tocopherol values of 12.9, 12.7, 12.5, 12.3, and 12.6 mg/L for 2005, 2006, 2007, 2008, and 2009, respectively, and 1.7, 1.7, 1.8, 1.7, and 1.9 mg/L, respectively, for γ -tocopherol. Total cholesterol was measured using a Roche Cobas Mira Plus Chemistry Analyzer and Roche Multi-Analyte Serum Calibrator. Samples were run in duplicate and every 10 study samples were bracketed by QC duplicate samples (BioRad Liquid Assayed Multiqual 1, 2, 3) with a CV of 2.5%.

Statistical analysis

Spearman rank order correlation coefficients were determined within the sub-cohort. Proportional hazards modelling was used to test the association between plasma tocopherols and the risk of total and high-grade (Gleason 7) prostate cancer. Because the sampling scheme used in creating the sub-cohort was stratified, all analyses were stratified by the nine race-age groups. The proportional hazards assumptions were applied to each race-age group, and the resulting summary parameters are a weighted mean across the strata, with sampling weights derived as the inverse of the probability of being selected for the cohort within each stratum (21,22). A second type of weighting was used in constructing the pseudo-likelihood function. Non-cases within the sub-cohort are weighted equally, from baseline until censor date. Cases outside the sub-cohort are weighted only at the time of diagnosis. Cases within the selected cohort carry a weight of 1, but are considered non-cases up until the time of diagnosis. A SAS macro that computes the weighted estimates and the robust covariance matrix was used (<http://lib.stat.cmu.edu/general/robphreg>). Proportional hazards assumptions were tested and met using the methods of Lin et al. (23).

The primary analysis examines the plasma tocopherol associations overall, within the four intervention arms of the trial, and by any selenomethionine and any α -tocopherol supplementation. Secondary analyses of key a priori hypotheses included examination of the plasma tocopherol associations within the four intervention arms and any selenomethionine and any α -tocopherol by disease grade (Gleason 6 vs. 7–10), smoking status (never vs. current and former), and follow-up time (< 3 years post randomization vs. 3 years). In addition, the effects of the intervention supplementation vs. placebo within tertiles of tocopherol concentrations were examined (tertiles rather than quintiles being used in order to maintain adequate sample sizes). Quintiles of α -tocopherol and γ -tocopherol based on the entire sub-cohort sample were modelled as indicator variables, with the lowest used as the referent category. Multivariable models were adjusted for baseline PSA, family history of prostate cancer, body mass index (continuous), diabetes, cholesterol (continuous), smoking status (current, former, never), and mutually adjusted for the other tocopherol. Results are also age- and race-adjusted as a result of all models being stratified by age/race groups before being weighted and combined to generate summary statistics. Tests for linear trend

were obtained by treating the quantile ordering as a continuous variable. All subgroup associations were examined using subgroup-specific sampling weights. Statistical interaction was evaluated by comparing models with and without a cross-product interaction term using the log-likelihood ratio test. Statistical analyses were performed by the SELECT/SWOG Statistical Center using SAS software version 9.2 (SAS Institute, Inc., Cary, North Carolina). All statistical tests were two-sided.

Results

Baseline characteristics of case-cohort participants

Median follow-up was 3.4 years and 6.2 years for prostate cancer cases and sub-cohort non-cases, respectively. Cases were more likely to be college graduates, non-smokers, have a family history of prostate cancer and higher PSA, and less likely to be diabetic (Table 1). Men with higher baseline plasma α -tocopherol were more likely to be Caucasian, college graduates, former smokers, had lower energy and fat intake, higher calcium and vitamin E intake (diet only), and lower plasma γ -tocopherol and higher cholesterol concentrations (Supplementary Table S2). Men with higher plasma γ -tocopherol had lower α -tocopherol but higher cholesterol and were more likely to be African-American, diabetic, and to have a higher BMI, and less likely to have a college degree, a history of benign prostatic hyperplasia, or to smoke. These men also had higher energy and fat intake, and lower calcium, vitamin D, and vitamin E intake (diet only), (Supplementary Table S2). Baseline plasma α -tocopherol and γ -tocopherol among the sub-cohort non-cases (means, 13.6 and 2.0 mg/L, respectively) were consistent with nationally representative values for men in the U.S. (i.e., 11.6 and 2.1 mg/L, respectively); (24), and the two tocopherols were inversely correlated in our study sample ($r = -0.21$, $P < .0001$).

Plasma tocopherol-prostate cancer association

Pre-randomization concentrations of plasma α -tocopherol were not significantly associated with risk of prostate cancer overall or in men receiving the trial placebo or α -tocopherol supplement alone. Men with higher baseline α -tocopherol who received the trial selenomethionine supplement alone or in combination with the α -tocopherol supplement, however, had substantially increased prostate cancer risk (all men receiving selenomethionine, hazard ratio (HR) for the fifth (Q5) vs. first quintile (Q1), 2.04 (95% CI 1.29–3.22), P -trend = 0.005). HR estimates from a joint effects model with men in the first plasma α -tocopherol quintile who received the trial placebo as the referent category yielded similar results overall (Supplementary Table S3).

Plasma γ -tocopherol was not associated with prostate cancer overall or within the trial intervention arms, although concentrations appeared inversely related to risk in the placebo arm and possibly positively associated with risk among men receiving α -tocopherol alone, although neither test for trend was statistically significant (Table 2). The primary findings for plasma α -tocopherol and γ -tocopherol did not differ by race/ethnicity (data not shown).

Secondary analyses revealed that a positive association between plasma α -tocopherol and prostate cancer was evident only for high-grade disease (Gleason grade 7–10, Q5 HR, 1.59,

95% CI, 1.13–2.24; P -trend = 0.001), particularly among men who received the selenomethionine supplement (alone or with α -tocopherol) (Table 3). The positive α -tocopherol risk association and interaction with selenomethionine was similar among never and current/former smokers, but was only observed within the first three years of the trial. The γ -tocopherol–prostate cancer relation did not differ by Gleason grade, smoking status or follow-up time, although the interaction with follow-up time was statistically significant (P = 0.009) (Supplementary Table S4).

The previously reported higher prostate cancer incidence in men receiving the SELECT α -tocopherol supplement, alone or with selenomethionine (versus placebo) (2,3), appeared restricted to those in the second and third tertiles of baseline plasma α -tocopherol (HR 1.11 and 1.25, respectively; Table 4), with an HR for any α -tocopherol vs. placebo of 1.15 (95% CI 0.91–1.47) for the two tertiles combined. Risk was also elevated for men receiving selenomethionine if they were in the highest plasma α -tocopherol tertile. All 95% CIs included 1.0, however, reflecting diminished power for testing the intervention effects in these subgroups, and trend tests were not significant. Median (range) on-study plasma α -tocopherol concentrations 6 months into the trial in the first, second and third tertiles of baseline α -tocopherol were 9.6 (5.7–15.0), 11.3 (7.4–16.2), and 15.0 (9.0–22.2) mg/L among men not supplemented with α -tocopherol, and 15.2 (7.3–33.2), 18.0 (10.6–28.6), and 17.9 (9.8–55.4) mg/L among men receiving α -tocopherol. With respect to plasma γ -tocopherol, only men in the highest baseline tertile appeared to experience increased prostate cancer risk in response to the α -tocopherol supplementation (versus placebo), although these findings also were not statistically significant (Table 4).

Discussion

Within the SELECT trial that recently showed that supplementation with 400 IU α -tocopheryl acetate daily resulted in higher prostate cancer incidence (3), we found that men with the highest pre-randomization plasma α -tocopherol concentrations were twice as likely to be diagnosed with prostate cancer if they received the trial selenomethionine supplement. Plasma α -tocopherol was unrelated to risk among men in the α -tocopherol only and placebo trial arms. In contrast to recent data from transgenic rat for adenocarcinoma of prostate (TRAP) model experiments (25), there was no prostate cancer–plasma γ -tocopherol association overall, although a modest inverse association was suggested in the placebo arm.

Higher plasma concentrations of α -tocopherol result from supplementation and increased dietary intake, higher cholesterol and triglyceride levels, and some genetic variants related to vitamin E transport and metabolism (24,26,27). Experimental and other human data show a wide range of effects relevant to prostate tumorigenesis for vitamin E compounds, including inhibition of membrane-associated lipid peroxidation, inflammation, angiogenesis, and cell proliferation, enhanced apoptosis, and reduced circulating androgen levels (27–33). Protective associations have been observed between higher serum α -tocopherol and prostate cancer risk in several (5–13), but not all (34–40), studies, and one large controlled trial found a one-third reduction in prostate cancer incidence in men taking a relatively low, 50 I.U. dose of α -tocopheryl acetate daily (1).

How elevated plasma α -tocopherol concentrations resulted in higher prostate cancer risk among men supplemented with selenium is unknown and, to our knowledge, has not been previously observed. The fact that selenomethionine was the compound administered in SELECT leaves open the possibility that either elemental selenium or the amino acid moiety (or both) was responsible for the observed harmful interaction with plasma α -tocopherol. Selenium and vitamin E have previously been shown to enhance each other's antioxidant functions in various model systems (41,42), yet our findings suggest the possibility of an alternative biological interaction when both micronutrients are elevated. It is noteworthy in this regard that metabolites of methionine as well as the related one-carbon metabolic pathway have been associated with prostate cancer recurrence. In one study, prostate cancer patients whose malignancies recurred within 2 years were more likely to have higher methionine, cysteine and sarcosine metabolite concentrations at the time of prostatectomy than were patients who remained in remission for at least 5 years (43). The fact that higher pre-randomization plasma α -tocopherol was primarily related to increased risk of high-grade prostate cancer and within the first three years of SELECT is consistent with these clinical observations regarding increased tumor progression and recurrence; i.e., selenomethionine in the setting of higher α -tocopherol concentrations may have promoted the growth of subclinical cancers (44). Also relevant in this regard are the higher overall prostate cancer risk in the selenomethionine arms in the highest plasma α -tocopherol tertile (Table 4), and the somewhat increased incidence of high-grade prostate cancers in men receiving selenomethionine in SELECT (Table 3 in reference (3), HR 1.21 ($P = 0.11$) for selenomethionine alone, and HR 1.23 ($P = 0.08$) for selenomethionine plus α -tocopherol, compared to the placebo arm). How increased plasma α -tocopherol under conditions of methionine or selenium supplementation might lead to higher prostate cancer risk requires further study.

Given the increased prostate cancer incidence in SELECT participants randomized to a high, commonly used dose of α -tocopherol, it is possible that men with higher pre-randomization plasma levels were taking α -tocopherol supplements prior to enrolment and, as a result, remained at elevated risk for the first three years of the trial. Pre-study "vitamin E" supplement use (i.e., use of specific tocopherols or formulations was not queried) was comparable to patterns for U.S. men [i.e., 29% in SELECT vs. 27%; (45)], but was in fact substantially higher among men in the two highest plasma α -tocopherol quintiles (46%) compared to men in the two lowest quintiles (14%). Such a continuing effect in the earliest trial years is consistent with data reported from the ATBC Study showing a 2–3 year post-trial lag for return from lower to baseline prostate cancer risk (46). We found no material association between plasma PSA and α -tocopherol ($r = -0.004$) or γ -tocopherol ($r = -0.04$), and no intervention arm differences in PSA or DRE testing in SELECT were found (3). Therefore, increased risk of high-grade disease during the first 3 years of intervention is not likely due to detection bias.

The present findings also indicate that the impact of the daily 400 IU α -tocopherol supplement on prostate cancer in SELECT may have been modified by pre-randomization α -tocopherol status, with elevated risk suggested primarily among men with higher plasma concentrations. This could reflect a stronger adverse impact of high-dose α -tocopherol

among men who had higher baseline biochemical status who may have experienced a heightened response to supplementation. Men in the α -tocopherol arms with higher baseline plasma α -tocopherol (e.g., >11 mg/L) did achieve 18–24% higher follow-up concentrations as compared with men of lower baseline status (i.e., medians of 18 mg/L and 15 mg/L, respectively, and means of 20.0 mg/L and 16.1 mg/L), suggesting the possibility of an α -tocopherol threshold above which a deleterious impact on the development of prostate cancer may occur, and adding to the growing debate concerning the risks and benefits of high-dose vitamin supplementation. Our finding of excess risk of high-grade, Gleason 7–10 cancers is of particular concern in this regard.

Strengths of the present study include the ability to examine pre-randomization vitamin E biochemical status (i.e., plasma α -tocopherol and γ -tocopherol) within the setting of a large controlled trial of α -tocopherol and selenium supplementation, and to detect strong interactions that may have biological significance. With the exception of being over-sampled for African-Americans, the sub-cohort was representative of the SELECT population and likely unbiased. Laboratory assays of the plasma tocopherols demonstrated high quality control, and all prostate cancer diagnoses were reviewed at a central pathology laboratory. We were, however, limited by the relatively short follow-up, most of which occurred during the trial supplementation period, and by the requirement for PSA and DRE results to be negative for prostate cancer at baseline, resulting in an event rate that reflected men at average risk. The latter eligibility criterion led to the preponderance of early, small incident tumors with very little advanced disease being diagnosed. Our study was also somewhat underpowered for evaluating interactions between the trial intervention arms and plasma tocopherol concentrations (47).

Ambiguous human data for the role of vitamin E in prostate cancer risk and prevention have resulted in uncertainty concerning the potential benefit of dietary supplementation with this essential nutrient. High-dose α -tocopherol supplementation was shown in SELECT to increase prostate cancer incidence, and the present study suggests that men with higher pre-randomization plasma α -tocopherol status experienced that adverse effect disproportionately, possibly as a result of pre-trial self-supplementation or an accentuated response to the trial supplementation. This strengthens the argument that attaining supra-physiological concentrations of vitamin E has deleterious consequences, and suggests that some persons having lower vitamin E “set-points” may not be susceptible to those effects. Whether (and how) higher α -tocopherol status interacts with selenomethionine supplementation to increase prostate cancer risk, and high-grade disease in particular, requires further investigation.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding: This work was supported in part by Public Health Service Cooperative Agreement grant 7U10CA037429-29 awarded by the National Cancer Institute in cooperation with the National Center for Complementary and Alternative Medicine National Institutes of Health, National Institutes of Health, Department

of Health and Human Services (C. Till, E. Klein, P. Goodman, F. Meyskens, Jr., I. Thompson), and by the Intramural Program of the U.S. Public Health Service, National Institutes of Health, National Cancer Institute (D. Albanes, A. Mondul, S. Weinstein, P. Taylor). Study agents and packaging were provided by Perrigo Company (Allegan, Michigan), Sabinsa Corporation (Piscataway, New Jersey), Tishcon Corporation (Westbury, New York), and DSM Nutritional Products, Inc. (Parsippany, New Jersey).

Abbreviations

| | |
|---------------|---|
| ATBC | Alpha-Tocopherol, Beta-Carotene Cancer Prevention |
| BMI | body mass index |
| CI | confidence interval |
| CV | coefficient of variation |
| HR | hazard ratio |
| PHS | Physicians' Health Study |
| PSA | prostate specific antigen |
| QC | quality control |
| SELECT | Selenium and Vitamin E Cancer Prevention Trial |

Reference List

1. Heinonen OP, Albanes D, Virtamo J, Taylor PR, Huttunen JK, Hartman AM, et al. Prostate cancer and supplementation with alpha-tocopherol and beta-carotene: incidence and mortality in a controlled trial. *J Natl Cancer Inst.* 1998; 90:440–6. [PubMed: 9521168]
2. Lippman SM, Klein EA, Goodman PJ, Lucia MS, Thompson IM, Ford LG, et al. Effect of selenium and vitamin E on risk of prostate cancer and other cancers: the Selenium and Vitamin E Cancer Prevention Trial (SELECT). *JAMA.* 2009; 301:39–51. [PubMed: 19066370]
3. Klein EA, Thompson IM, Tangen CM, Crowley JJ, Lucia M, Goodman PJ, et al. The Selenium and Vitamin E Cancer Prevention Trial (SELECT). Vitamin E and the Risk of Prostate Cancer. *JAMA.* 2011; 306:1549–56. [PubMed: 21990298]
4. Gaziano JM, Glynn RJ, Christen WG, Kurth T, Belanger C, MacFadyen J, et al. Vitamins E and C in the prevention of prostate and total cancer in men: the Physicians' Health Study II randomized controlled trial. *JAMA.* 2009; 301:52–62. [PubMed: 19066368]
5. Eichholzer M, Stahelin HB, Gey KF, Ludin E, Bernasconi F. Prediction of male cancer mortality by plasma levels of interacting vitamins: 17-year follow-up of the prospective Basel study. *Int J Cancer.* 1996; 66:145–50. [PubMed: 8603802]
6. Goodman GE, Schaffer S, Omenn GS, Chen C, King I. The association between lung and prostate cancer risk, and serum micronutrients: results and lessons learned from beta-carotene and retinol efficacy trial. *Cancer Epidemiol Biomarkers Prev.* 2003; 12:518–26. [PubMed: 12814997]
7. Weinstein SJ, Wright ME, Pietinen P, King I, Tan C, Taylor PR, et al. Serum alpha-tocopherol and gamma-tocopherol in relation to prostate cancer risk in a prospective study. *J Natl Cancer Inst.* 2005; 97:396–9. [PubMed: 15741576]
8. Weinstein SJ, Wright ME, Lawson KA, Snyder K, Mannisto S, Taylor PR, et al. Serum and dietary vitamin E in relation to prostate cancer risk. *Cancer Epidemiol Biomarkers Prev.* 2007; 16:1253–9. [PubMed: 17548693]
9. Watters JL, Gail MH, Weinstein SJ, Virtamo J, Albanes D. Associations between alpha-tocopherol, beta-carotene, and retinol and prostate cancer survival. *Cancer Res.* 2009; 69:3833–41. [PubMed: 19383902]

10. Cheng T, Barnett MJ, Kristal AE, Ambrosone CB, King IB, Thornquist MD, et al. Genetic variation in myeloperoxidase modifies the association of serum alpha-tocopherol with aggressive prostate cancer among current smokers. *J Nutr.* 2011; 141:1731–7. [PubMed: 21795425]
11. Chan JM, Stampfer MJ, Ma J, Rimm EB, Willett WC, Giovannucci EL. Supplemental vitamin E intake and prostate cancer risk in a large cohort of men in the United States. *Cancer Epidemiol Biomarkers Prev.* 1999; 8:893–9. [PubMed: 10548318]
12. Gann PH, Ma J, Giovannucci E, Willett W, Sacks FM, Hennekens CH, et al. Lower prostate cancer risk in men with elevated plasma lycopene levels: results of a prospective analysis. *Cancer Res.* 1999; 59:1225–30. [PubMed: 10096552]
13. Weinstein SJ, Peters U, Ahn J, Friesen MD, Riboli E, Hayes RB, et al. Serum alpha-tocopherol and gamma-tocopherol concentrations and prostate cancer risk in the PLCO Screening Trial: a nested case-control study. *PLoS One.* 2012; 7:e40204. [PubMed: 22792240]
14. Lippman SM, Goodman PJ, Klein EA, Parnes HL, Thompson IM Jr, Kristal AR, et al. Designing the Selenium and Vitamin E Cancer Prevention Trial (SELECT). *J Natl Cancer Inst.* 2005; 97:94–102. [PubMed: 15657339]
15. Goodman PJ, Hartline JA, Tangen CM, Crowley JJ, Minasian LM, Klein EA, et al. Moving a randomized clinical trial into an observational cohort. *Clinical Trials.* 2013; 10:131–42. [PubMed: 23064404]
16. Lippman SM, Goodman PJ, Klein EA, Parnes HL, Thompson IM Jr, Kristal AR, et al. Designing the Selenium and Vitamin E Cancer Prevention Trial (SELECT). *J Natl Cancer Inst.* 2005; 97:94–102. [PubMed: 15657339]
17. Hoque A, Albanes D, Lippman SM, Spitz MR, Taylor PR, Klein EA, et al. Molecular epidemiologic studies within the Selenium and Vitamin E Cancer Prevention Trial (SELECT). *Cancer Causes Control.* 2001; 12:627–33. [PubMed: 11552710]
18. Brasky TM, Till C, White E, Neuhaus ML, Song X, Goodman P, et al. Serum phospholipid fatty acids and prostate cancer risk: results from the prostate cancer prevention trial. *Am J Epidemiol.* 2011; 173:1429–39. [PubMed: 21518693]
19. Stavrovskaya IG, Bird SS, Marur VR, Sniatynski MJ, Baranov SV, Greenberg HK, et al. Dietary macronutrients modulate the fatty acyl composition of rat liver mitochondrial cardiolipins. *J Lipid Res.* 2013; 54:2623–35. [PubMed: 23690505]
20. Kristal AR, Till C, Platz EA, Song X, King IB, Neuhaus ML, et al. Serum lycopene concentration and prostate cancer risk: results from the Prostate Cancer Prevention Trial. *Cancer Epidemiol Biomarkers Prev.* 2011; 20:638–46. [PubMed: 21335507]
21. Barlow WE. Robust variance estimation for the case-cohort design. *Biometrics.* 1994; 50:1064–72. [PubMed: 7786988]
22. Barlow WE, Ichikawa L, Rosner D, Izumi S. Analysis of case-cohort designs. *J Clin Epidemiol.* 1999; 52:1165–72. [PubMed: 10580779]
23. Lin DY, Wei LJ, Ying Z. Checking the cox model with cumulative sums of Martingale-based residuals. *Biometrika.* 1993; 80:557–72.
24. Ford ES, Schleicher RL, Mokdad AH, Ajani UA, Liu S. Distribution of serum concentrations of alpha-tocopherol and gamma-tocopherol in the US population. *Am J Clin Nutr.* 2006; 84:375–83. [PubMed: 16895886]
25. Takahashi S, Takeshita K, Seeni A, Sugiura S, Tang M, Sato SY, et al. Suppression of prostate cancer in a transgenic rat model via gamma-tocopherol activation of caspase signaling. *Prostate.* 2009; 69:644–51. [PubMed: 19143023]
26. Major JM, Yu K, Wheeler W, Zhang H, Cornelis MC, Wright ME, et al. Genome-wide association study identifies common variants associated with circulating vitamin E levels. *Hum Mol Genet.* 2011; 20:3876–83. [PubMed: 21729881]
27. Ricciarelli R, Zingg JM, Azzì A. Vitamin E: protective role of a Janus molecule. *FASEB J.* 2001; 15:2314–25. [PubMed: 11689457]
28. Fleshner NE. Vitamin E and prostate cancer. *Urol Clin North Am.* 2002; 29:107–13. ix. [PubMed: 12109337]

29. Hartman TJ, Dorgan JF, Woodson K, Virtamo J, Tangrea JA, Heinonen OP, et al. Effects of long-term alpha-tocopherol supplementation on serum hormones in older men. *Prostate*. 2001; 46:33–8. [PubMed: 11170129]
30. Food and Nutrition Board, Institute of Medicine. Dietary reference intakes for vitamin C, vitamin E, selenium, and carotenoids. Washington, DC: National Academy Press; 2000.
31. Venkateswaran V, Klotz LH. Diet and prostate cancer: mechanisms of action and implications for chemoprevention. *Nat Rev Urol*. 2010; 7:442–53. [PubMed: 20647991]
32. Basu A, Grossie B, Bennett M, Mills N, Imrhan V. Alpha-tocopheryl succinate (alpha-TOS) modulates human prostate LNCaP xenograft growth and gene expression in BALB/c nude mice fed two levels of dietary soybean oil. *Eur J Nutr*. 2007; 46:34–43. [PubMed: 17180484]
33. Schindler R, Mentlein R. Flavonoids and vitamin E reduce the release of the angiogenic peptide vascular endothelial growth factor from human tumor cells. *J Nutr*. 2006; 136:1477–82. [PubMed: 16702307]
34. Key TJ, Appleby PN, Allen NE, Travis RC, Roddam AW, Jenab M, et al. Plasma carotenoids, retinol, and tocopherols and the risk of prostate cancer in the European Prospective Investigation into Cancer and Nutrition study. *Am J Clin Nutr*. 2007; 86:672–81. [PubMed: 17823432]
35. Gill JK, Franke AA, Steven MJ, Cooney RV, Wilkens LR, Le ML, et al. Association of selenium, tocopherols, carotenoids, retinol, and 15-isoprostane F(2t) in serum or urine with prostate cancer risk: the multiethnic cohort. *Cancer Causes Control*. 2009; 20:1161–71. [PubMed: 19212706]
36. Huang HY, Alberg AJ, Norkus EP, Hoffman SC, Comstock GW, Helzlsouer KJ. Prospective study of antioxidant micronutrients in the blood and the risk of developing prostate cancer. *Am J Epidemiol*. 2003; 157:335–44. [PubMed: 12578804]
37. Helzlsouer KJ, Huang HY, Alberg AJ, Hoffman S, Burke A, Norkus EP, et al. Association between alpha-tocopherol, gamma-tocopherol, selenium, and subsequent prostate cancer. *J Natl Cancer Inst*. 2000; 92:2018–23. [PubMed: 11121464]
38. Nomura AM, Stemmermann GN, Lee J, Craft NE. Serum micronutrients and prostate cancer in Japanese Americans in Hawaii. *Cancer Epidemiol Biomarkers Prev*. 1997; 6:487–91. [PubMed: 9232334]
39. Knekt P, Aromaa A, Maatela J, Aaran RK, Nikkari T, Hakama M, et al. Serum vitamin E and risk of cancer among Finnish men during a 10-year follow-up. *Am J Epidemiol*. 1988; 127:28–41. [PubMed: 3276161]
40. Hayes RB, Bogdanovic JF, Schroeder FH, De Bruijn A, Raatgever JW, Van der Maas PJ, et al. Serum retinol and prostate cancer. *Cancer*. 1988; 62:2021–6. [PubMed: 3167814]
41. Zu K, Ip C. Synergy between selenium and vitamin E in apoptosis induction is associated with activation of distinctive initiator caspases in human prostate cancer cells. *Cancer Res*. 2003; 63:6988–95. [PubMed: 14583501]
42. Venkateswaran V, Fleshner NE, Klotz LH. Synergistic effect of vitamin E and selenium in human prostate cancer cell lines. *Prostate Cancer Prostatic Dis*. 2004; 7:54–6. [PubMed: 14745439]
43. Stabler S, Koyama T, Zhao Z, Martinez-Ferrer M, Allen RH, Luka Z, et al. Serum methionine metabolites are risk factors for metastatic prostate cancer progression. *PLoS One*. 2011; 6:e22486. [PubMed: 21853037]
44. Stawiarska-Pieta B, Bielec B, Birkner K, Birkner E. The influence of vitamin E and methionine on the activity of enzymes and the morphological picture of liver of rats intoxicated with sodium fluoride. *Food Chem Toxicol*. 2012; 50:972–8. [PubMed: 22266362]
45. Bailey RL, Gahche JJ, Lentino CV, Dwyer JT, Engel JS, Thomas PR, et al. Dietary supplement use in the United States, 2003–2006. *J Nutr*. 2011; 141:261–6. [PubMed: 21178089]
46. Virtamo J, Pietinen P, Huttunen JK, Korhonen P, Malila N, Virtanen MJ, et al. Incidence of cancer and mortality following alpha-tocopherol and beta-carotene supplementation: a postintervention follow-up. *JAMA*. 2003; 290:476–85. [PubMed: 12876090]
47. Gaziano J, Sesso HD, Christen WG, Bubes V, Smith JP, MacFadyen J, et al. Multivitamins in the prevention of cancer in men: The Physicians' Health Study II randomized controlled trial. *JAMA*. 2012:1–10.

Table 1

Baseline demographic and health-related characteristics for prostate cancer cases and sub-cohort members, SELECT case-cohort study

| | Prostate Cancer Cases n=1,746 (%) | Sub-Cohort Non-Cases n=3,008 (%) |
|--|--|---|
| Age, y, mean (SD) | 63.5 (6.1) | 63.3 (6.6) |
| Race/ethnicity, n (%) | | |
| African-American | 251 (14.4) | 735 (24.4) |
| Hispanic | 58 (3.3) | 139 (4.6) |
| Caucasian | 1406 (80.5) | 2083 (69.2) |
| Other | 31 (1.8) | 51 (1.7) |
| College graduate, n (%) | 948 (54.3) | 1428 (47.5) |
| History of benign prostatic hyperplasia, n (%) | | |
| Yes | 292 (16.7) | 470 (15.6) |
| Missing/unknown | 142 (8.1) | 190 (6.3) |
| Family history of prostate cancer, n (%) | 506 (29.0) | 423 (14.1) |
| History of diabetes, n (%) | 127 (7.3) | 386 (12.8) |
| Height, cm, mean (SD) | 176.8 (7.4) | 176.8 (7.5) |
| Body mass index, kg/m ² , mean (SD) | 28.5 (4.4) | 28.8 (4.6) |
| Smoking status, n (%) | | |
| Lifetime non-smoker | 829 (47.5) | 1250 (41.6) |
| Former smoker | 821 (47.0) | 1497 (49.8) |
| Current smoker | 96 (5.5) | 261 (8.7) |
| Trial supplementation arm, n (%) | | |
| α -Tocopherol only | 479 (27.4) | 765 (25.4) |
| Selenomethionine only | 437 (25.0) | 749 (24.9) |
| α -Tocopherol and selenomethionine | 424 (24.3) | 753 (25.0) |
| Placebo | 406 (23.3) | 741 (24.6) |
| Dietary intake/day, mean (SD) | | |
| Energy, kcal | 2352 (1081) | 2312 (1185) |
| Total fat, g | 94.2 (52.1) | 93.6 (56.4) |
| Calcium, mg | 1059 (597) | 1012 (604) |
| Vitamin D (calciferol), μ g | 7.1 (5.0) | 6.8 (5.0) |
| Vitamin E, IU | 18.1 (12) | 17.9 (14.4) |
| Supplemental vitamin E use prior to trial, n (%) | 501 (28.7) | 860 (28.6) |
| Plasma biochemical measures, mean (SD) | | |
| Alpha-tocopherol, mg/L | 14.0 (5.3) | 13.6 (5.6) |
| Gamma-tocopherol, mg/L | 1.9 (1.1) | 2.0 (1.2) |
| AT:GT molar ratio | 12.5 (17.0) | 11.6 (16.0) |
| Cholesterol, mg/dL | 203 (37) | 200 (38) |
| Prostate-specific antigen (μ g/mL), n (%) | | |
| <1.00 | 130 (7.4) | 1330 (44.2) |
| 1.00–1.99 | 464 (26.6) | 1039 (34.5) |

| | Prostate Cancer Cases n=1,746 (%) | Sub-Cohort Non-Cases n=3,008 (%) |
|---|--|---|
| 2.00–2.99 | 587 (33.6) | 420 (14.0) |
| >=3.00 | 564 (32.3) | 219 (7.3) |
| Prostate-specific antigen (µg/mL), median (IQR) | 2.4 (1.7–3.2) | 1.1 (0.6–1.8) |

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Association between quintiles of plasma α -tocopherol, γ -tocopherol, and prostate cancer risk overall and by intervention arm, SELECT case-cohort study

Table 2

| | Plasma tocopherol quintiles | | | | | <i>P</i> _{trend} |
|---|-----------------------------|------------------|------------------|------------------|------------------|---------------------------|
| | 1 | 2 | 3 | 4 | 5 | |
| α-Tocopherol | 9.83 | > 9.83 – 11.71 | > 11.71 – 13.70 | > 13.70 – 16.80 | > 16.80 | |
| Median, mg/L | 8.64 | 10.75 | 12.65 | 15.01 | 20.09 | |
| Cases/sub-cohort, n ^a | 299/681 | 343/656 | 361/629 | 375/616 | 362/624 | |
| Overall HR ^b (95% CI) | 1.00 | 1.07 (0.82–1.40) | 1.06 (0.81–1.41) | 1.12 (0.83–1.49) | 1.21 (0.88–1.66) | 0.24 |
| HR ^b (95% CI) within intervention arms | | | | | | |
| Placebo | 1.00 | 0.97 (0.56–1.68) | 0.91 (0.49–1.68) | 0.96 (0.53–1.74) | 0.85 (0.44–1.62) | 0.66 |
| α -Tocopherol only | 1.00 | 0.60 (0.35–1.03) | 0.72 (0.43–1.21) | 0.67 (0.38–1.19) | 0.59 (0.33–1.07) | 0.24 |
| Selenomethionine only | 1.00 | 1.21 (0.72–2.03) | 1.42 (0.79–2.55) | 1.69 (0.96–2.99) | 1.83 (0.95–3.54) | 0.04 |
| α -Tocopherol and selenomethionine | 1.00 | 1.72 (1.03–2.89) | 1.34 (0.78–2.30) | 1.53 (0.86–2.73) | 2.05 (1.15–3.67) | 0.06 |
| Any selenomethionine | 1.00 | 1.47 (1.02–2.13) | 1.44 (0.97–2.14) | 1.62 (1.07–2.43) | 2.04 (1.29–3.22) | 0.005 |
| Any α -tocopherol | 1.00 | 1.00 (0.69–1.46) | 0.96 (0.66–1.39) | 0.99 (0.66–1.49) | 1.12 (0.74–1.70) | 0.64 |
| γ-Tocopherol | 0.96 | > 0.96 – 1.47 | > 1.47 – 2.01 | > 2.01 – 2.76 | > 2.76 | |
| Median, mg/L | 0.66 | 1.24 | 1.73 | 2.35 | 3.41 | |
| Cases/sub-cohort, n ^a | 355/633 | 361/628 | 360/628 | 363/629 | 301/688 | |
| Overall HR ^b (95% CI) | 1.00 | 1.02 (0.78–1.33) | 1.11 (0.84–1.45) | 1.13 (0.84–1.50) | 0.93 (0.69–1.24) | 0.90 |
| HR (95% CI) within intervention arms | | | | | | |
| Placebo | 1.00 | 0.84 (0.50–1.43) | 0.98 (0.56–1.72) | 0.71 (0.39–1.30) | 0.68 (0.37–1.25) | 0.17 |
| α -Tocopherol only | 1.00 | 1.24 (0.74–2.07) | 1.52 (0.91–2.53) | 1.32 (0.75–2.31) | 1.52 (0.88–2.60) | 0.16 |
| Selenomethionine only | 1.00 | 0.90 (0.52–1.55) | 0.97 (0.56–1.68) | 1.38 (0.79–2.39) | 0.75 (0.42–1.36) | 0.82 |
| α -Tocopherol and selenomethionine | 1.00 | 0.96 (0.56–1.62) | 0.94 (0.55–1.60) | 1.05 (0.62–1.76) | 0.77 (0.44–1.36) | 0.58 |
| Any selenomethionine | 1.00 | 0.93 (0.63–1.37) | 0.94 (0.63–1.39) | 1.20 (0.80–1.79) | 0.77 (0.50–1.16) | 0.64 |
| Any α -tocopherol | 1.00 | 1.15 (0.79–1.65) | 1.22 (0.85–1.77) | 1.21 (0.82–1.79) | 1.14 (0.77–1.69) | 0.48 |

^aThe sub-cohort contains both cases and non-cases, with overlap between case and sub-cohort numbers.

^bHazard ratios (HR) based on proportional hazard regression, adjusted for PSA (continuous), family history of prostate cancer, BMI (continuous), diabetes at baseline, cholesterol (continuous), and smoking status (current smoker, former smoker, never smoker) and mutually adjusted for γ -tocopherol or α -tocopherol (continuous). Results are also age- and race-adjusted as a result of all models being stratified by age/race groups before being weighted and combined to generate summary statistics. Overall HR also adjusted for intervention arm. CI, confidence interval. The first plasma tocopherol quintiles serve as

Referent categories. Interaction between treatment arm and tocopherol quintiles was performed using a linear trend for quintiles in a 3 degrees-of-freedom chi-square test, with the *P*-value for α -tocopherol being 0.20 and for γ -tocopherol being 0.27.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3

Association between quintiles of plasma α -tocopherol and prostate cancer risk stratified by Gleason grade, smoking status, follow-up time, and by trial supplementation, SELECT case-cohort study

| Histologic grade | Plasma α -tocopherol quintiles | | | | | P_{trend} |
|---|---------------------------------------|------------------|------------------|------------------|------------------|--------------------|
| | 1 | 2 | 3 | 4 | 5 | |
| Gleason 2-6 | | | | | | |
| Cases/sub-cohort, n ^a | 171/681 | 201/656 | 209/629 | 209/616 | 203/624 | |
| Overall HR ^b (95% CI) | 1.00 | 1.14 (0.89-1.45) | 1.13 (0.88-1.44) | 1.12 (0.88-1.44) | 1.06 (0.82-1.36) | 0.80 |
| Placebo | 1.00 | 1.04 (0.64-1.69) | 1.16 (0.71-1.89) | 0.94 (0.57-1.53) | 1.04 (0.63-1.73) | 0.93 |
| α -Tocopherol only | 1.00 | 0.83 (0.52-1.32) | 0.95 (0.60-1.52) | 0.82 (0.51-1.32) | 0.56 (0.34-0.93) | 0.04 |
| Selenomethionine only | 1.00 | 1.12 (0.69-1.81) | 1.33 (0.82-2.16) | 1.22 (0.75-1.99) | 1.10 (0.67-1.79) | 0.65 |
| α -Tocopherol and selenomethionine | 1.00 | 1.72 (1.04-2.85) | 0.94 (0.56-1.58) | 1.62 (0.98-2.68) | 1.75 (1.06-2.91) | 0.08 |
| Any selenomethionine | 1.00 | 1.37 (0.96-1.94) | 1.15 (0.81-1.63) | 1.41 (0.99-2.00) | 1.39 (0.98-1.98) | 0.10 |
| Any α -tocopherol | 1.00 | 1.17 (0.83-1.64) | 0.96 (0.68-1.35) | 1.12 (0.80-1.59) | 0.98 (0.69-1.40) | 0.82 |
| Gleason 7-10 | | | | | | |
| Cases/sub-cohort, n ^a | 72/681 | 89/656 | 103/629 | 117/616 | 117/624 | |
| Overall HR ^b (95% CI) | 1.00 | 1.15 (0.81-1.64) | 1.45 (1.03-2.04) | 1.58 (1.13-2.21) | 1.59 (1.13-2.24) | 0.001 |
| Placebo | 1.00 | 1.64 (0.75-3.60) | 2.50 (1.18-5.30) | 1.81 (0.85-3.86) | 1.73 (0.78-3.82) | 0.28 |
| α -Tocopherol only | 1.00 | 0.89 (0.47-1.69) | 0.89 (0.47-1.68) | 1.24 (0.67-2.32) | 0.93 (0.48-1.80) | 0.77 |
| Selenomethionine only | 1.00 | 1.13 (0.52-2.45) | 2.10 (1.01-4.36) | 2.41 (1.19-4.88) | 2.58 (1.27-5.26) | 0.001 |
| α -Tocopherol and selenomethionine | 1.00 | 1.27 (0.65-2.50) | 1.24 (0.64-2.41) | 1.51 (0.78-2.92) | 1.75 (0.92-3.32) | 0.08 |
| Any selenomethionine | 1.00 | 1.21 (0.73-2.01) | 1.62 (1.00-2.64) | 1.92 (1.18-3.10) | 2.12 (1.32-3.40) | 0.0002 |
| Any α -tocopherol | 1.00 | 1.08 (0.68-1.71) | 1.06 (0.67-1.68) | 1.35 (0.86-2.13) | 1.29 (0.81-2.04) | 0.17 |
| Smoking status | | | | | | |
| Never smokers | | | | | | |
| Cases/sub-cohort, n ^a | 146/289 | 163/276 | 168/268 | 172/264 | 177/259 | |
| Overall HR ^b (95% CI) | 1.00 | 0.96 (0.72-1.30) | 1.05 (0.78-1.41) | 1.03 (0.76-1.40) | 1.10 (0.81-1.49) | 0.44 |
| Placebo | 1.00 | 1.00 (0.54-1.85) | 1.21 (0.65-2.24) | 0.81 (0.42-1.57) | 0.99 (0.51-1.92) | 0.78 |
| α -Tocopherol only | 1.00 | 0.66 (0.37-1.19) | 0.66 (0.37-1.19) | 0.67 (0.37-1.21) | 0.58 (0.32-1.05) | 0.11 |

| | Plasma α -tocopherol quintiles | | | | | P_{trend} |
|---|---------------------------------------|------------------|------------------|------------------|------------------|-------------|
| | 1 | 2 | 3 | 4 | 5 | |
| Selenomethionine only | 1.00 | 1.29 (0.71–2.35) | 1.64 (0.90–2.97) | 1.47 (0.80–2.69) | 1.84 (1.00–3.39) | 0.05 |
| α -Tocopherol and selenomethionine | 1.00 | 1.19 (0.65–2.18) | 0.89 (0.49–1.63) | 1.08 (0.58–2.01) | 1.66 (0.90–3.06) | 0.18 |
| Any selenomethionine | 1.00 | 1.19 (0.78–1.82) | 1.16 (0.76–1.77) | 1.35 (0.88–2.08) | 1.70 (1.11–2.61) | 0.01 |
| Any α -tocopherol | 1.00 | 0.90 (0.59–1.36) | 0.79 (0.52–1.20) | 0.84 (0.55–1.29) | 0.91 (0.59–1.39) | 0.62 |
| Current & former smokers | | | | | | |
| Cases/sub-cohort, n ^a | 146/402 | 187/569 | 193/563 | 202/552 | 186/564 | |
| Overall HR ^b (95% CI) | 1.00 | 1.40 (1.06–1.84) | 1.35 (1.03–1.78) | 1.43 (1.09–1.88) | 1.26 (0.95–1.67) | 0.18 |
| Placebo | 1.00 | 1.18 (0.67–2.10) | 1.18 (0.67–2.06) | 1.14 (0.65–1.99) | 1.01 (0.57–1.80) | 0.93 |
| α -Tocopherol only | 1.00 | 1.05 (0.62–1.77) | 1.27 (0.75–2.16) | 0.99 (0.58–1.67) | 0.85 (0.48–1.50) | 0.51 |
| Selenomethionine only | 1.00 | 1.17 (0.63–2.16) | 1.48 (0.82–2.66) | 1.64 (0.91–2.93) | 1.52 (0.84–2.75) | 0.09 |
| α -Tocopherol and selenomethionine | 1.00 | 1.68 (0.98–2.89) | 1.11 (0.64–1.92) | 1.86 (1.09–3.16) | 1.53 (0.88–2.64) | 0.15 |
| Any selenomethionine | 1.00 | 1.33 (0.87–2.03) | 1.47 (0.98–2.21) | 1.60 (1.06–2.41) | 1.44 (0.95–2.20) | 0.06 |
| Any α -tocopherol | 1.00 | 1.33 (0.92–1.93) | 1.16 (0.80–1.70) | 1.32 (0.90–1.92) | 1.11 (0.75–1.63) | 0.72 |
| Follow-up time to diagnosis | | | | | | |
| < 3 years | | | | | | |
| Cases/sub-cohort, n ^a | 113/681 | 124/656 | 158/629 | 149/616 | 154/624 | |
| Overall HR ^b (95% CI) | 1.00 | 1.08 (0.80–1.45) | 1.41 (1.06–1.87) | 1.29 (0.97–1.72) | 1.30 (0.98–1.74) | 0.03 |
| Placebo | 1.00 | 0.96 (0.55–1.69) | 1.27 (0.74–2.19) | 0.74 (0.41–1.34) | 0.88 (0.48–1.59) | 0.42 |
| α -Tocopherol only | 1.00 | 0.89 (0.51–1.55) | 1.05 (0.61–1.80) | 1.22 (0.71–2.08) | 0.73 (0.41–1.27) | 0.61 |
| Selenomethionine only | 1.00 | 1.14 (0.59–2.19) | 2.40 (1.31–4.37) | 1.94 (1.06–3.54) | 2.09 (1.15–3.79) | 0.003 |
| α -Tocopherol and selenomethionine | 1.00 | 1.31 (0.73–2.36) | 1.24 (0.69–2.25) | 1.67 (0.93–2.99) | 2.04 (1.16–3.61) | 0.01 |
| Any selenomethionine | 1.00 | 1.27 (0.82–1.96) | 1.78 (1.17–2.70) | 1.78 (1.17–2.71) | 2.08 (1.38–3.12) | 0.0001 |
| Any α -tocopherol | 1.00 | 1.09 (0.73–1.64) | 1.14 (0.76–1.70) | 1.39 (0.93–2.06) | 1.22 (0.81–1.83) | 0.17 |
| 3 years | | | | | | |
| Cases/sub-cohort, n ^a | 186/631 | 219/614 | 203/594 | 226/575 | 208/577 | |
| Overall HR ^b (95% CI) | 1.00 | 1.10 (0.87–1.40) | 1.00 (0.79–1.28) | 1.08 (0.85–1.38) | 1.01 (0.78–1.29) | 0.94 |
| Placebo | 1.00 | 1.18 (0.70–1.98) | 1.33 (0.79–2.25) | 1.15 (0.69–1.91) | 1.04 (0.60–1.81) | 0.99 |
| α -Tocopherol only | 1.00 | 0.78 (0.49–1.23) | 0.79 (0.49–1.26) | 0.68 (0.42–1.10) | 0.58 (0.35–0.96) | 0.03 |
| Selenomethionine only | 1.00 | 1.05 (0.65–1.70) | 1.00 (0.61–1.65) | 1.29 (0.80–2.09) | 1.14 (0.69–1.86) | 0.40 |

| | Plasma α -tocopherol quintiles | | | | | P_{trend} |
|---|---------------------------------------|------------------|------------------|------------------|------------------|--------------------|
| | 1 | 2 | 3 | 4 | 5 | |
| α -Tocopherol and selenomethionine | 1.00 | 1.56 (0.95–2.55) | 0.94 (0.57–1.53) | 1.37 (0.84–2.25) | 1.36 (0.82–2.24) | 0.44 |
| Any selenomethionine | 1.00 | 1.27 (0.90–1.79) | 0.99 (0.70–1.40) | 1.33 (0.94–1.88) | 1.28 (0.90–1.82) | 0.19 |
| Any α -tocopherol | 1.00 | 1.09 (0.78–1.52) | 0.87(0.62–1.21) | 0.95 (0.67–1.35) | 0.90 (0.63–1.27) | 0.36 |

^aThe sub-cohort contains both cases and non-cases, with overlap between case and sub-cohort numbers.

^bHazard ratios (HR) and 95% confidence intervals (CI) based on proportional hazard regression, mutually adjusted for γ -tocopherol or α -tocopherol (continuous), with PSA, family history of prostate cancer, BMI, diabetes, cholesterol, and smoking status included in the models. Results are also age- and race-adjusted as a result of all models being stratified by age/race groups before being weighted and combined to generate summary statistics. The first plasma tocopherol quintiles serve as Referent categories. Interaction P -values for plasma α -tocopherol x smoking status are 0.95, 0.75, and 0.94 for the four intervention arm, any selenium, and any vitamin E strata (respectively). Interaction P -values for plasma α -tocopherol x follow-up time to diagnosis are 0.21, 0.35, and 0.30 for the four intervention arm, any selenium, and any vitamin E strata (respectively).

Supplementation effects on prostate cancer hazard ratios within tertiles of plasma α -tocopherol and γ -tocopherol, SELECT case-cohort study

Table 4

| | Plasma tocopherol tertiles | | | <i>P</i> _{trend} |
|---|----------------------------|------------------|------------------|---------------------------|
| | 1 | 2 | 3 | |
| α-Tocopherol | | | | |
| Median, mg/L | 11.05 | > 11.05 – 14.56 | > 14.56 | |
| Cases/sub-cohort, <i>n</i> ^a | 9.50 | 12.65 | 17.61 | |
| | 529/1108 | 602/1059 | 609/1039 | |
| Intervention HR ^b (95% CI) | | | | |
| α -Tocopherol vs. Placebo | 1.03 (0.68–1.58) | 1.24 (0.84–1.84) | 1.10 (0.76–1.59) | 0.54 |
| Selenomethionine vs. Placebo | 0.84 (0.55–1.27) | 1.06 (0.72–1.57) | 1.26 (0.85–1.88) | 0.31 |
| α -Tocopherol and selenomethionine vs. Placebo | 0.97 (0.65–1.45) | 0.92 (0.62–1.37) | 1.34 (0.90–2.01) | 0.39 |
| Any selenomethionine vs. Placebo | 0.90 (0.63–1.29) | 1.05 (0.74–1.49) | 1.38 (0.98–1.95) | |
| Any α -tocopherol vs. Placebo | 1.03 (0.72–1.48) | 1.11 (0.78–1.57) | 1.25 (0.90–1.74) | |
| γ-Tocopherol | | | | |
| Median, mg/L | 1.32 | > 1.32 – 2.23 | > 2.23 | |
| Cases/sub-cohort, <i>n</i> ^a | 0.86 | 1.73 | 2.96 | |
| | 597/1051 | 613/1031 | 530/1124 | |
| Intervention HR ^b (95% CI) | | | | |
| α -Tocopherol vs. Placebo | 0.98 (0.66–1.45) | 1.14 (0.77–1.68) | 1.37 (0.93–2.03) | 0.42 |
| Selenomethionine vs. Placebo | 1.08 (0.73–1.58) | 1.03 (0.70–1.52) | 1.22 (0.80–1.86) | 0.74 |
| α -Tocopherol and selenomethionine vs. Placebo | 1.19 (0.80–1.78) | 1.02 (0.69–1.50) | 1.22 (0.81–1.83) | 0.48 |
| Any selenomethionine vs. Placebo | 1.17 (0.84–1.64) | 1.02 (0.73–1.43) | 1.25 (0.86–1.82) | |
| Any α -tocopherol vs. Placebo | 1.09 (0.77–1.54) | 1.03 (0.73–1.44) | 1.29 (0.90–1.83) | |

^aThe sub-cohort contains both cases and non-cases, with overlap between case and sub-cohort numbers.

^bHazard ratios (HR) and 95% confidence intervals (CI) based on proportional hazard regression, stratified by tocopherol tertile and adjusted for PSA (continuous), family history of prostate cancer, BMI (continuous), diabetes at baseline, cholesterol (continuous), and smoking status (current smoker, former smoker, never smoker) and mutually adjusted for γ -tocopherol or α -tocopherol (continuous). Results are also age- and race-adjusted as a result of all models being stratified by age/race groups before being weighted and combined to generate summary statistics. For each HR, the referent category is placebo arm within that category of plasma tocopherol concentration.