

Two new species of the *Clubiona corticalis*-group from Yunnan Province, China (Araneae, Clubionidae)

Pan-Long Wu¹, Guo Zheng², Feng Zhang¹

1 *The Key Laboratory of Invertebrate Systematics and Application, College of Life Sciences, Hebei University, Baoding, Hebei 071002, P. R. China* **2** *College of Life Science and Chemistry, Shenyang Normal University, Shenyang, Liaoning 110034, P. R. China*

Corresponding author: Feng Zhang (dudu06042001@163.com)

Academic editor: Y. Marusik | Received 9 January 2015 | Accepted 19 March 2015 | Published 16 April 2015

<http://zoobank.org/F1480639-1CC1-4399-AFFB-99B6B626CFE4>

Citation: Wu P-L, Zheng G, Zhang F (2015) Two new species of the *Clubiona corticalis*-group from Yunnan Province, China (Araneae, Clubionidae). ZooKeys 496: 15–25. doi: 10.3897/zookeys.496.9223

Abstract

The present paper describes two new *Clubiona corticalis*-group species collected from Xishuangbanna, Yunnan Province of China: *Clubiona submoralis* **sp. n.** (♀♂) and *C. pollicaris* **sp. n.** (♀♂).

Keywords

Sac spiders, *Paraclubiona*, *Atalia*, taxonomy, South-East Asia

Introduction

Clubiona Latreille, 1804, the largest genus of the Clubionidae, currently includes 468 species (Platnick 2014) widely distributed around the world (except South America). Because of its high species diversity, several revisions have been published by Simon (1932) for the French species, Lohmander (1944) for Swedish species, Wiehle (1965) for German species, Wunderlich (2011) for European species, Edwards (1958) for the North American species, Dondale and Redner (1982) for Canadian and Alaskan species, Mikhailov (1990, 1991, 1995, 2002, 2012) for Palaeartic species, and Deeleman-Reinhold (2001) for southeast Asian species.

Table 1. A list of *Clubiona corticalis*-group species in China.

	Species name	Known sex	Distribution
1	<i>C. altissimoides</i> Liu et al., 2007	♂♀	Yunnan
2	<i>C. applanata</i> Liu et al., 2007	♂♀	Yunnan
3	<i>C. brachyptera</i> Zhu, Ren & Chen, 2012	♂♀	Hainan
4	<i>C. cordata</i> Zhang & Zhu, 2009	♂♀	Sichuan, Xizang
5	<i>C. cylindrata</i> Liu et al., 2007	♂♀	Yunnan
6	<i>C. didentata</i> Zhang & Yin, 1998	♂	Yunnan
7	<i>C. kurosawai</i> Ono, 1986	♂♀	Taiwan
8	<i>C. lamina</i> Zhang, Zhu & Song, 2007	♂	Yunnan
9	<i>C. lyriformis</i> Song & Zhu, 1991	♀	Hubei
10	<i>C. moralis</i> Song & Zhu, 1991	♂♀	Hubei
11	<i>C. parallela</i> Hu & Li, 1987	♂♀	Xizang
12	<i>C. pyrifer</i> Schenkel, 1936	♂♀	Gansu
13	<i>C. qiyunensis</i> Xu, Yang & Song, 2003	♂♀	Fujian, Anhui
14	<i>C. taiwanica</i> Ono, 1994	♂♀	Yunnan, Taiwan
15	<i>C. tengchong</i> Zhang, Zhu & Song, 2007	♂	Yunnan
16	<i>C. yaginumai</i> Hayashi, 1989	♂♀	Taiwan
17	<i>C. submoralis</i> sp. n.	♂♀	Yunnan
18	<i>C. pollicaris</i> sp. n.	♂♀	Yunnan

Clubiona corticalis-group was first recognized by Simon (1932). *Atalia* Thorell, 1887 (type species *A. concinna* Thorell, 1887, belongs to the *corticalis*-group) and subgenus *Paraclubiona* Lohmander, 1944 (with type species *C. corticalis*) are currently considered as junior synonyms of *Clubiona*. Species of this group can be recognized by the following characters: inflated tegulum; long filiform, spiniform, or short embolus; simple and weakly developed retrolateral tibial apophysis; the anterior position of copulatory openings (Mikhailov 1995).

Currently, the *corticalis*-group includes 48 species mainly distributed in Eurasia and Australia (Mikhailov 1995, Deeleman-Reinhold 2001, Liu et al. 2007). Among these, at least 18 species have been recorded from China (see Table 1).

While examining *Clubiona* specimens collected from Xishuangbanna Prefecture, Yunnan Province of China, we found two new species belonging to the *corticalis*-group, which are described in this paper.

Material and methods

All specimens studied are stored in 75% ethanol and deposited in the Institute of Zoology, Chinese Academy of Sciences in Beijing (IZCAS). All specimens were examined under a Tech XTL-II stereomicroscope. The photos, drawings and measurements were prepared using a Leica M205A stereomicroscope equipped with a DFC450 CCD camera and a drawing tube. Carapace length was measured from the anterior mar-

gin to the posterior margin of the carapace medially. The eyes were measured as the maximum diameter of the lens in dorsal or frontal view. The measurements of legs are shown as total length (femur, patella, tibia, metatarsus, tarsus). The epigyne was cleared in a solution of potassium hydroxide (KOH) and transferred to 75% ethanol for taking photos, drawing and measuring. All measurements are in millimeters.

The following abbreviations are used: ALE, anterior lateral eyes; AME, anterior median eyes; B, bursae; C, conductor; CO, copulatory openings; E, embolus; FD, fertilization ducts; MOA, median ocular area; PLE, posterior lateral eyes; PME, posterior median eyes; PPA, prolateral patellar apophysis; RFA, retrolateral femoral apophysis; RPA, retrolateral patellar apophysis; RTA, retrolateral tibial apophysis; S, spermathecae.

Taxonomy

Clubiona submoralis sp. n.

<http://zoobank.org/3299A436-1170-4959-9BE3-95190B84F461>

Figs 1–12

Type material. Holotype ♂, CHINA, Yunnan Province, Xishuangbanna Prefecture, Mengla County, Menglun Town, XTBG (Xishuangbanna Tropical Botanical Garden), Yunnan Rubber Plantation (575 m; 21°54.46'N, 101°15.98'E), 21 July 2007, Guo Zheng leg. Paratypes: 9 ♂, 10 ♀, same data as holotype.


Diagnosis. Among the Chinese species of the *corticalis*-group, this new species resembles *C. moralis*, but differs by: a wider and triangular embolus (filiform and coiled in *C. moralis*), presence of only one round, black marking on tegulum (several fan-shaped markings in *C. moralis*); fertilization ducts opening in the medio-ventral side of the spermathecae (anteriorly in *C. moralis*); and the spermathecae almost as large as the bursae (the bursae twice longer than the spermathecae in *C. moralis*) (Figs 3–12).

Etymology. The species name refers to a close resemblance between the new species and *C. moralis*.

Description. Male. Total length 4.36–4.86. Holotype (Fig. 1): body 4.73 long; carapace 2.14 long, 1.96 wide; abdomen 2.49 long, 1.30 wide. Carapace brown. Median furrow longitudinal. In dorsal view, anterior eye row recurved, posterior eye row recurved. Eye sizes and interdistances: AME 0.10, ALE 0.11, PME 0.11, PLE 0.10; AME–AME 0.07, AME–ALE 0.06, PME–PME 0.27, PME–PLE 0.15. MOA 0.31 long, front 0.30 wide, back 0.48 wide. Chelicerae dark brown, promargin with six teeth, retromargin with three teeth. Endites brown, longer than wide. Labium dark brown, longer than wide. Sternum 1.19 long, 0.80 wide. Abdomen oval, brown, with conspicuous anterior tufts of hairs, dorsum with fine brown hairs. Legs yellow, both tibia I and II with two pairs of ventral spines, both metatarsi I and II with one pair of ventral spines. Measurements of legs: I 5.50 (1.55, 0.75, 1.57, 1.07, 0.56), II 6.02 (1.65, 0.80, 1.80, 1.21, 0.56), III 4.73 (1.26, 0.61, 1.13, 1.30, 0.43), IV 6.73 (1.78, 0.71, 1.65, 2.01, 0.58).



Figures 1–7. *Clubiona submoralis* sp. n. **1** male habitus, dorsal view **2** female habitus, dorsal view **3** epigyne, ventral view **4** vulva **5** left male palp, ventral view **6** same, dorsal view, showing tibial apophysis **7** same, retrolateral view. Scale bars: 1 mm (**1–2**); 0.2 mm (**3–7**).



Figures 8–12. *Clubiona submoralis* sp. n. **8** epigyne, ventral view **9** vulva **10** left male palp, ventral view **11** tibial apophysis, dorsal view **12** left male palp, retrolateral view. Scale bars: 0.25 mm (**8–12**).

Palp (Figs 5–7, 10–12). Femur unmodified; patella swollen, almost globular, twice wider than tibia, and 1.5 wider than femur, with short and rounded pro- and retro-lateral apophyses, retrolateral side with short modified spines near the base; tibia with small retro-ventral membranous apophysis. Cymbium longer than tegulum. Tegulum inflated, with a round, black marking medially; embolus short, wide, almost triangular

in ventral view; conductor membranous, folded in the middle position, and almost threefold longer than wide.

Female. Total length 4.45–4.92. One paratype (Fig. 2) measured, body 4.47 long; carapace 1.87 long, 1.43 wide; abdomen 2.49 long, 1.39 wide. Eye sizes and interdistances: AME 0.10, ALE 0.11, PME 0.09, PLE 0.07; AME–AME 0.08, AME–ALE 0.04, PME–PME 0.24, PME–PLE 0.14. MOA 0.22 long, front 0.23 wide, back 0.41 wide. Sternum 0.96 long, 0.70 wide. Measurements of legs: I 4.53 (1.31, 0.66, 1.21, 0.84, 0.51), II 5.14 (1.48, 0.74, 1.41, 0.97, 0.54), III 4.22 (1.25, 0.57, 0.90, 1.13, 0.37), IV 6.14 (1.66, 0.70, 1.40, 1.89, 0.49). Coloration lighter than in male. Other characters as in male.

Epigyne (Figs 3–4, 8–9). Copulatory openings located anteriorly; in ventral view, the anterior part of copulatory ducts well visible and extending posteriorly, then connecting to bursae; spermathecae located anterior to bursae, both of them almost spherical, and with the same size as bursae.

Distribution. China (Yunnan).

***Clubiona pollicaris* sp. n.**

<http://zoobank.org/A815DA83-1925-4B07-BB84-D2DC0D5DA3C8>

Figs 13–19, 23–27

Type material. Holotype ♂, CHINA, Yunnan Province, Xishuangbanna Prefecture, Mengla County, Menglun Nature Reserve (710 m; 21°57.70'N, 101°11.89'E), 7 August 2007, Guo Zheng leg. Paratypes: 6 ♂, 6 ♀, same data as holotype.


Diagnosis. The new species differs from all other *Clubiona* species by a very long retrolateral femoral apophysis (almost as long as femur) and resembles *C. qiyunensis* (Figs 20–22; also see Wu and Zhang (2014): 211, f. 13–23), but differs by: a smaller and inconspicuous embolus; the much longer femoral apophysis (in *C. qiyunensis*, femoral apophysis is short, less than diameter of femur); a thumb-shaped prolateral patellar apophysis; anteriorly situated spermathecae; and rectangular bursae (Figs 15–19, 23–27).

Etymology. The species name is derived from the Latin word “*pollicaris*”, and refers to the prolateral patellar apophysis, which in ventral view is thumb-shaped.


Description. Male. Total length 3.20–3.49. Holotype (Fig. 13) body 3.27 long; carapace 1.51 long, 1.45 wide; abdomen 1.61 long, 0.90 wide. Carapace brown. Median furrow longitudinal. Both anterior and posterior eye row recurved in dorsal view. Eye sizes and interdistances: AME 0.06, ALE 0.10, PME 0.09, PLE 0.08; AME–AME 0.06, AME–ALE 0.03, PME–PME 0.19, PME–PLE 0.08. MOA 0.21 long, front 0.19 wide, back 0.37 wide. Chelicerae brown, promargin with five teeth, retromargin with four teeth. Both endites and labium brown, longer than wide. Sternum 0.83 long, 0.56 wide. Abdomen oval, brown, with conspicuous anterior tufts of hairs, dorsum with fine yellow hairs. Legs brown, both tibia I and II with two pairs of ventral spines, both metatarsi I and II with one pair of ventral spines. Measurements of legs: I 3.09 (0.84, 0.47, 0.83, 0.62, 0.33), II 3.46 (0.90,



Figures 13–16. *Clubiona pollicaris* sp. n. **13** male habitus, dorsal view **14** female habitus, dorsal view **15** left male palp, ventral view **16** same, dorsal view, showing patellar apophysis. Scale bars: 1 mm (13–14); 0.2 mm (15, 16).



Figures 17–22. 17–19 *Clubiona pollicaris* sp. n. 17 epigyne, ventral view 18 vulva 19 left male palp, retrolateral view; 20–22 *Clubiona qiyunensis* Xu, Yang & Song, 2003. 20 epigyne, ventral view 21 vulva 22 left male palp, retrolateral view. Scale bars: 0.1 mm (17, 18) 0.2 mm (19–22).



Figures 23–27. *Clubiona pollicaris* sp. n. **23** epigyne, ventral view **24** vulva **25** left male palp, ventral view **26** patellar apophysis, dorsal view **27** left male palp, retrolateral view. Scale bars: 0.25 mm (**23–27**).

0.50, 1.02, 0.69, 0.35), III 2.92 (0.84, 0.43, 0.63, 0.76, 0.26), IV 4.11 (1.18, 0.50, 0.96, 1.15, 0.32).

Palp (Figs 15–16, 19, 25–27). Femur modified, with long and thin retrolateral apophysis originating from median part, apophysis longer than tibia and subequal in length to femur; patella twice longer and 1.5 wider than tibia, with a round apophysis and thumb-shaped prolateral apophysis; tibia without apophyses; cymbium shorter than tegulum; tegulum inflated; sperm duct obvious, almost straight in retrolateral view; embolus short, originating from the apical tegulum prolaterally, directed almost horizontally.

Female. Total length 3.12–3.92. One paratype (Fig. 14) measured: body 3.90 long, carapace 1.52 long, 1.13 wide; abdomen 2.09 long, 1.18 wide. Eye sizes and interdistances: AME 0.07, ALE 0.09, PME 0.08, PLE 0.07; AME–AME 0.08, AME–ALE 0.05, PME–PME 0.20, PME–PLE 0.12. MOA 0.21 long, front 0.20 wide, back

0.35 wide. Sternum 0.85 long, 0.57 wide. Measurements of legs: I 2.69 (0.84, 0.38, 0.64, 0.51, 0.32), II 3.09 (0.90, 0.46, 0.79, 0.60, 0.34), III 2.63 (0.80, 0.35, 0.55, 0.67, 0.26), IV 3.92 (1.02, 0.50, 0.93, 1.10, 0.37). Coloration slightly lighter than in male; other characters as in male.

Epigyne (Figs 17–18, 23–24). Copulatory openings small, located almost in the centre of the epigynal plate; in ventral view copulatory ducts inconspicuous; spermathecae long, tubular and sinuous; bursae large, membranous, almost rectangular.

Distribution. China (Yunnan).

Comments. Retrolateral femoral apophysis is known only in three species of Clubionidae: *Clubiona pollicaris*, *C. qiyunensis* and *C. brachyptera*.

Acknowledgements

We thank Dr. Shuqiang Li (Institute of Zoology, Chinese Academy of Sciences, Beijing) for providing valuable specimens. We are grateful to Dr. K.G. Mikhailov and Dr. R. Bosmans for reviewing the manuscript. Prof. Victor Fet kindly helped reviewing the English of the manuscript. This work was supported by the National Natural Science Foundation of China (No. 31372154, 31172121, 31372224).

References

- Deeleman-Reinhold CL (2001) Forest spiders of South East Asia: with a revision of the sac and ground spiders (Araneae: Clubionidae, Corinnidae, Liocranidae, Gnaphosidae, Prodidomidae and Trochanteriidae). Brill, Leiden, 591 pp.
- Dondale CD, Redner JH (1982) The insects and arachnids of Canada, Part 9. The sac spiders of Canada and Alaska, Araneae: Clubionidae and Anyphaenidae. Research Branch Agriculture Canada Publication 1724: 1–194.
- Edwards RJ (1958) The spider subfamily Clubioninae of the United States, Canada and Alaska (Araneae: Clubionidae). Bulletin of the Museum of Comparative Zoology, Harvard 118(6): 365–436.
- Hayashi T (1989) Four new species of the genus *Clubiona* (Araneae: Clubionidae) from Japan. In: Nishikawa Y, Ono H (Eds) Arachnological Papers Presented to Takeo Yaginuma on the Occasion of his Retirement. Osaka Arachnologists' Group, Osaka, 103–110.
- Hu JL, Li AH (1987) The spiders collected from the fields and the forests of Xizang Autonomous Region, China. (II). Agricultural Insects, Spiders, Plant Diseases and Weeds of Xizang 2: 247–353.
- Huang WJ, Chen SH (2012) The Spider Fauna of Taiwan. Araneae. Miturgidae, Anyphaenidae, Clubionidae. National Taiwan Normal University, Taipei, 130 pp.
- Liu P, Yan HM, Griswold C, Ubick D (2007) Three new species of the genus *Clubiona* from China (Araneae: Clubionidae). Zootaxa 1456: 63–68.
- Lohmander H (1944) Vorläufige Spinnennotizen. Arkiv för Zoologi 35(A, 16): 1–21.

- Mikhailov KG (1990) The spider genus *Clubiona* Latreille 1804 in the Soviet Far East, 1 (Arachnida, Aranei, Clubionidae). *Korean Arachnology* 5(2): 139–175.
- Mikhailov KG (1991) The spider genus *Clubiona* Latreille 1804 in the Soviet Far East, 2 (Arachnida, Aranei, Clubionidae). *Korean Arachnology* 6(2): 207–235.
- Mikhailov KG (1995) Erection of infrageneric groupings within the spider genus *Clubiona* Latreille, 1804 (Aranei Clubionidae): a typological approach. *Arthropoda Selecta* 4(2): 33–48.
- Mikhailov KG (2002) The spider genus *Clubiona* Latreille, 1804 (Aranei: Clubionidae) in the fauna of the former USSR: 2003 update. *Arthropoda Selecta* 11: 283–317.
- Mikhailov KG (2012) Reassessment of the spider genus *Clubiona* (Aranei, Clubionidae). *Vestnik Zoologii* 46(2): 177–180. doi: 10.2478/v10058-012-0015-z
- Ono H (1994) Spiders of the genus *Clubiona* from Taiwan (Araneae: Clubionidae). *Acta Arachnologica*, Tokyo 43: 71–85. doi: 10.2476/asjaa.43.71
- Ono H (1986) A new spider of the group of *Clubiona corticalis* (Araneae, Clubionidae) found in Japan. In: *Entomological Papers Presented to Yoshihiko Kurosawa on the Occasion of his Retirement*. Tokyo, 19–25.
- Platnick NI (2014) The World Spider Catalog, Version 15.5. American Museum of Natural History. <http://www.wsc.nmbe.ch/family/17/Clubionidae> [accessed 30 December 2014]
- Schenkel E (1936) Schwedisch-chinesische wissenschaftliche Expedition nach den nordwestlichen Provinzen Chinas, unter Leitung von Dr Sven Hedin und Prof. Sü Ping-chang. Araneae gesammelt vom schwedischen Arzt der Exped. *Arkiv för Zoologi* 29(A1): 1–314.
- Simon E (1932) Les arachnides de France. Tome VI. Synopsis générale et catalogue des espèces françaises de l'ordre des Araneae; 4e partie. Paris, 6: 773–978.
- Song DX, Zhu MS, Gao SS, Guan JD (1991) Six species of clubionid spiders (Araneae: Clubionidae) from China. *Journal of Xinjiang University* 8: 66–72.
- Wiehle H (1965) Die *Clubiona*-Arten Deutschlands, ihre natürliche Gruppierung und die Einheitlichkeit im Bau ihrer Vulva (Arach., Araneae). *Senckenbergiana Biologica* 46: 471–505.
- Wu PL, Zhang F (2014) A new species of the spider genus *Clubiona* from China, with description of the male of *Clubiona qiynensis* (Araneae: Clubionidae). *Acta Zoologica Academiae Scientiarum Hungaricae* 60(3): 207–215.
- Wunderlich J (2011) Extant and fossil spiders (Araneae). *Beträge zur Araneologie* 6: 1–640.
- Xu YJ, Yang JY, Song DX (2003) Two new species of the genus *Clubiona* from Anhui, China (Araneae: Clubionidae). *Journal of Hebei University (Natural Science Edition)* 23: 411–413.
- Zhang F, Zhu MS (2009) Three new species of the genus *Clubiona* from Xizang and Sichuan, China (Araneae, Clubionidae). *Acta Zootaxonomica Sinica* 34: 725–729.
- Zhang F, Zhu MS, Song DX (2007) Three new species of the genus *Clubiona* from Yunnan Province, China (Araneae, Clubionidae). *Journal of Hebei University, Natural Science Edition* 27: 407–411.
- Zhang YJ, Yin CM (1998) Six new species of the spiders of genus *Clubiona* from China (Araneae: Clubionidae). *Acta Zootaxonomica Sinica* 23: 9–17.
- Zhu BR, Ren HQ, Chen J (2012) A new species of the genus *Clubiona* Latreille (Araneae, Clubionidae) from Hainan Island, China. *Zootaxa* 3167: 53–56.