



Published in final edited form as:

*Biochim Biophys Acta*. 2014 September ; 1840(9): 2813–2828. doi:10.1016/j.bbagen.2014.04.016.

## The cytochrome P450 genes of channel catfish: their involvement in disease defense responses as revealed by meta-analysis of RNA-Seq datasets

Jiaren Zhang<sup>a</sup>, Jun Yao<sup>a</sup>, Ruijia Wang<sup>a</sup>, Yu Zhang<sup>a</sup>, Shikai Liu<sup>a</sup>, Luyang Sun<sup>a</sup>, Yanliang Jiang<sup>a</sup>, Jianbin Feng<sup>a</sup>, Nannan Liu<sup>b</sup>, David Nelson<sup>c</sup>, Geoff Waldbieser<sup>d</sup>, and Zhanjiang Liu<sup>a,\*</sup>

<sup>a</sup>The Fish Molecular Genetics and Biotechnology Laboratory, School of Fisheries, Aquaculture and Aquatic Sciences and Program of Cell and Molecular Biosciences, Aquatic Genomics Unit, Auburn University, Auburn, AL 36849, USA

<sup>b</sup>Department of Entomology and Plant Pathology, Auburn University, Auburn, AL 36849, USA

<sup>c</sup>Department of Microbiology, Immunology and Biochemistry, University of Tennessee, Memphis TN 38163, USA

<sup>d</sup>USDA, ARS, Catfish Genetics Research Unit, 141 Experiment Station Road, Stoneville, Mississippi 38776, USA

### Abstract

**Background**—Cytochrome P450s (*CYP*s) encode one of the most diverse enzyme superfamily in nature. They catalyze oxidative reactions of endogenous molecules and exogenous chemicals.

**Methods**—We identified *CYP*s genes through *in silico* analysis using EST, RNA-Seq and genome databases of channel catfish. Phylogenetic analyses and conserved syntenic analyses were conducted to determine their identities and orthologies. Meta-analysis of RNA-Seq databases was conducted to analyze expression profile of *CYP* genes following bacterial infection.

**Results**—A full set of 61 *CYP* genes were identified and characterized in channel catfish. Phylogenetic tree and conserved synteny provided strong evidence of their identities and orthology. Lineage-specific gene duplication was evident in a number of clans in channel catfish. *CYP46A1* is missing in the catfish genome as observed with syntenic analysis and RT-PCR analysis. Thirty *CYP*s were found up- or down-regulated in liver, while seven and eight *CYP*s were observed regulated in intestine and gill following bacterial infection.

**Conclusion**—We systematically identified and characterized a full set of 61 *CYP* genes in channel catfish and studied their expression profiles after bacterial infection. Strikingly large numbers of *CYP* genes appear to be involved in the bacterial defense processes.

**General significance**—This work provides an example to systematically study *CYP* genes in non-model species. Moreover, it provides a basis for further toxicological and physiological studies in channel catfish.

\*Corresponding author: 203 Swingle Hall, School of Fisheries, Aquaculture and Aquatic Sciences, Auburn University, Auburn, AL 36849, USA; Tel.: 1-334-844-4054; fax: 1-334-844-9208; liuzhan@auburn.edu.

## Keywords

Cytochrome P450; *CYP*; catfish; genome; immunity; bacterial infection

---

## 1. Introduction

Cytochrome P450s (*CYP*s), named for their characteristic spectral property of Soret absorption peak at 450 nm when binding with carbon monoxide, constitute a widespread and highly diverse heme-thiolate enzyme superfamily (1-4). These enzymes catalyze the oxidative reactions, which are not only involved in synthesizing or metabolizing of endogenous molecules, such as arachidonic acid, eicosanoid, retinoic acid, steroid, vitamin D3, bile acid, biogenic amine, prostaglandin and cholesterol, but also engaged in the detoxification processes of exogenous substrates, including pharmaceuticals, and foreign chemicals (5-10). Moreover, it has been demonstrated that the expression level of many *CYP*s is regulated during the host defense responses in liver, as well as in extrahepatic tissues such as kidney and brain in mice or rats (11-21).

With the application of the next-generation sequencing, rapid progress has been made for expanding *CYP* family from model species to non-model species. At present, thousands of *CYP*s have been identified from all domains of life including Animalia, Plantae, Fungi, Protista, Archaea, Bacteria and even virus (22-33). All these *CYP*s are classified into clans, families and subfamilies based on sequence similarities (40% amino acid sequence identity rule for membership in a family and 55% amino acid sequence identity rule for membership in a subfamily), phylogenetic relationships and syntenic relationships (23, 34-36). According to these rules, vertebrate *CYP*s could be clustered into 19 families within 10 clans-including *CYP* clan 2 (include *CYP1*, *CYP2*, *CYP17* and *CYP21* family), *CYP* clan 3 (include *CYP3* and *CYP5* family), *CYP* clan 4 (include *CYP4* family), *CYP* clan 7 (include *CYP7*, *CYP8* and *CYP39* family), *CYP* clan 19 (include *CYP19* family), *CYP* clan 20 (include *CYP20* family), *CYP* clan 26 (include *CYP16* and *CYP26* family), *CYP* clan 46 (include *CYP46* family), *CYP* clan 51 (include *CYP51* family) and mitochondrial clan (include *CYP11*, *CYP24* and *CYP27* family) (22, 23, 35). Generally, families among *CYP5-51* are involved in endogenous metabolism, whereas *CYP1-3* families and several subfamilies of *CYP4* play an important role in detoxification processes.

Currently, all 19 vertebrate *CYP* families have been identified in teleost fish, including rainbow trout (*Oncorhynchus mykiss*) (37-44), Japanese pufferfish (*Fugu rubripes*) (45, 46), zebrafish (*Danio rario*) (47-56), Atlantic salmon (*Salmo salar*) (57-60), European seabass (*Dicentrarchus labrax*) (61-63), largemouth bass (*Micropterus salmoides*) (64), medaka (*Oryzias latipes*) (65-71), common carp (*Cyprinus carpio*) (72-74), mummichog (*Fundulus heteroclitus*) (75-78), three-spined stickleback (*Gasterosteus aculeatus*) (70), gilthead seabream (*Sparus aurata*) (79), fathead minnow (*Pimephales promelas*) (80, 81), half-smooth tongue sole (*Cynoglossus semilaevis*) (82, 83). By far, with the accomplishment of whole genome sequence (84, 85), *CYP* genes in Japanese pufferfish (45) and zebrafish (55) have been analyzed at the whole genome level, revealing 54 *CYP*s (update to 61 *CYP*s, Table 3) and 94 *CYP*s are presented in the Japanese pufferfish and zebrafish, respectively.

Several channel catfish *CYP*s have been characterized previously including *CYP1B1*, *CYP2X1*, *CYP11A1*, *CYP17A1*, *CYP19A1* and *CYP19A2* (86-89). In recent years, following the development of genomic resources of channel catfish (90), particularly the ESTs (91, 92), transcriptome sequences generated by RNA-Seq (93-95) and the draft whole genome sequence, the systematic analysis of *CYP* genes in channel catfish genome becomes feasible. Channel catfish (*Ictalurus punctatus*) is the leading aquaculture species in the United States. Its sustainable production is threatened by the widespread ESC and columnaris disease outbreaks caused by the two pathogens: *E. ictaluri* and *F. columnare*. Here we report the identification of a full set of *CYP* genes, their phylogenetic analysis and syntenic analysis in the channel catfish, and their involvement in response to bacterial infections with *E. ictaluri* and *F. columnare*.

## 2. Materials and Methods

### 2.1. *CYP* homologous genes collection and database mining

In order to identify the complete set of *CYP* genes in channel catfish, we collected all the fish *CYP*s (*Anguilla japonica*, *Carassius auratus*, *Carassius carassius*, *Cyprinus carpio*, *Cynoglossus semilaevis*, *Danio rario*, *Dicentrarchus labrax*, *Fugu rubripes*, *Fundulus heteroclitus*, *Gasterosteus aculeatus*, *Gobiocypris rarus*, *Micropterus salmoides*, *Oncorhynchus mykiss*, *Oreochromis niloticus*, *Oryzias latipes*, *Pimephales promelas*, *Salmo salar*, *Sparus aurata*, *Squalus acanthias*, *Stenotomus chrysops*) from the database of NCBI (<http://www.ncbi.nlm.nih.gov>), Ensembl (<http://www.ensembl.org>) and *CYP* homepage (<http://drnelson.uthsc.edu/CytochromeP450.html>) (22). In addition, other *CYP* homologous genes from human, mouse, chicken and frog (*Xenopus laevis* and *Xenopus tropicalis*) were also collected.

More than 1000 collected *CYP*s were first used as query sequences to search against channel catfish EST and RNA-Seq databases. In order to pull all potential *CYP* genes, the cutoff value was set at the level of  $1e-5$  such that conserved transcripts are captured initially for further analysis. The retrieved sequences were then translated using ORF Finder (<http://www.ncbi.nlm.nih.gov/gorf/gorf.html>). Further, the predicted ORFs were verified by BLASTP against NCBI non-redundant (Nr) protein sequence database. All identified channel catfish *CYP* transcripts and all other query sequences were utilized to search channel catfish draft genome sequence using TBLASTN program. The following steps are the same with above with the exception of applying FGENESH from SoftBerry (<http://linux1.softberry.com/berry.phtml>) to predict exons and amino acid sequences using genomic sequences.

### 2.2. Phylogenetic and syntenic analyses

In order to identify channel catfish *CYP*s, all the amino acids from channel catfish and other species were used to conduct phylogenetic analyses. We constructed separate phylogenetic trees for different *CYP* clans since the number of sequences under study was too many to fit a page. Sequence alignment was performed using MUSCLE (Multiple Sequence Comparison by Log-Expectation) (96). JTT (Jones-Taylor-Thornton) and gamma distributed

rate with invariant sites (G+I) model was proposed by ProtTest3 (97). Maximum likelihood phylogenetic trees were built with MEGA 5.10 with 1000 bootstrap replications (98).

Shared synteny was searched by examining the conserved co-localization of neighboring genes on scaffold (unpublished data) of channel catfish and zebrafish. Generally, in order to obtain the location information of these channel catfish *CYP*s on scaffolds, all identified *CYP* transcripts together with query sequences from other species were used as queries to blast against channel catfish draft genome sequence. Neighboring genes of the channel catfish *CYP*s were predicted by FGENESH (<http://linux1.softberry.com/berry.phtml?topic=fgenesh&group=programs&subgroup=gfind>). Neighboring genes of zebrafish were identified from Zv9 database in ENSEMBL.

### 2.3. Degenerate primers design and Touchdown RT-PCR for *CYP46*

*CYP46* sequences from human, mouse, rat, pig, cow, frog, Japanese pufferfish, green spotted pufferfish, and zebrafish were first aligned using software MUSCLE. The degenerate primers were designed by CODEHOP (Consensus-Degenerate Hybrid Oligonucleotide Primers) (99) using the alignment file generated as the input file. Parameters were selected as maximum core degeneracy: 64, target clamp temperature: 60°C, Genetic code: standard, and Codon usage: *Ictalurus punctatus*. Two sets of degenerate primers were chosen to conduct the reaction with the estimated size of production around 400 bp and 200 bp, respectively (Table 1).

One healthy channel catfish (Marion strain) was sacrificed in this study. Following manufacturer's protocol, total RNA (RNeasy Mini Kit, Qiagen, USA) was extracted from brain. The extracted RNA was quantified using UV-spectrophotometer and then reverse-transcribed using iScript™ cDNA Synthesis Kit (Bio-Rad, USA). Degenerated primers were applied to amplify brain cDNA. PCR amplification was carried out using KAPA HiFi PCR kit (Kapa Biosystems, USA). The 25 µl PCR reaction mixture contained 5.0 µl of 5×KAPA HiFi Fidelity Buffer, 1 µl of KAPA dNTP Mix (10 mM), 0.5 µl of KAPA HiFi DNA Polymerase, 1 µl (10 pmol/µl) of each primers, 1 µl of template, and 15.5 µl PCR-grade water. Touchdown PCR was carried out on a Bio-Rad PCR system using the cycling conditions: denaturation, 95°C/4 min, 10 cycles of 95°C /30s, 65°C /30s (each cycle decrease 1°C), and 72°C/30s, 20 cycles of 95°C/30s, 55°C/30s and 72°C /30s, then, 72°C 10 min. PCR products were resolved on 1.5% agarose gel.

### 2.4. Meta-analyses of RNA-Seq datasets

Illumina-based RNA-Seq reads were retrieved from three RNA-Seq datasets: liver samples from catfish experimentally challenged with *E. ictaluri* (3 day and 14 day post infection) (SRA accession number SRP028159) (93), intestine samples from catfish challenged with *E. ictaluri* (3 h, 24 h and 3 day post infection) (SRA accession number SRP009069) (94) and gill samples from catfish challenged with *F. columnare* (4 h, 24 h and 48 h post infection) (SRA accession number SRP012586) (95). Trimmed high quality RNA-seq reads were mapped onto the deduced channel catfish *CYP*s reference assembly using CLC Genomics Workbench software (version 5.5.2; CLC bio, Aarhus, Denmark). Mapping parameters were set as 95% of the reads in perfect alignment and 2 mismatches. After the total mapped

reads number for each transcript was determined, normalization was conducted in order to determine RPKM (Reads Per Kilobase of exon per Million mapped reads). The proportions-based Kal's test was performed to identify the differently expressed genes comparing with control sample and fold changes were calculated. Transcripts with absolute fold change value  $\geq 2$ ,  $p$ -value  $\leq 0.05$  and total read number  $\geq 5$  were included in the analyses as differently expressed genes.

### 3. Results

#### 3.1. Identification of *CYP* genes in channel catfish

*CYP* genes collected from various species (as detailed in Materials and Methods) were used as queries to conduct BLAST searches against the channel catfish transcriptome and genome databases. All sequences with significant hits were assembled into unique sequences of 61 *CYP* genes in channel catfish (Table 2). Among all these genes, 40 sequences were identified in both databases with full-length or nearly full-length *CYP*s, while the remaining 21 genes were identified in the genome database with partial sequences of transcripts in the transcriptome database. All these sequences have been deposited to GenBank with their accession numbers being summarized in Table 2.

Based on the rules described previously for *CYP* genes nomenclature (35), the 61 *CYP* genes can be classified into 9 *CYP* clans (Table 2) including *CYP* clans 2, 3, 4, 7, 19, 20, 26, 51 and mitochondrial clan. These represented all vertebrate *CYP* clans but clan 46 (*CYP*46) which is absent in channel catfish.

The 61 channel catfish *CYP*s fall into 16 families (Table 2) including *CYP*1, *CYP*2, *CYP*3, *CYP*4, *CYP*5, *CYP*7, *CYP*8, *CYP*11, *CYP*17, *CYP*19, *CYP*20, *CYP*21, *CYP*24, *CYP*26, *CYP*27 and *CYP*51 families. Three families, *CYP*16, *CYP*39 and *CYP*46, of the vertebrate *CYP*s are missing from channel catfish. Of the 16 *CYP* families in channel catfish, *CYP*2, *CYP*3, *CYP*4 and *CYP*27 are the four biggest families contributed more than half (32 *CYP*s) of the *CYP* genes in channel catfish.

#### 3.2. Location of *CYP*s on linkage groups

To better understand the genome distribution of channel catfish *CYP*s, we examined their locations on channel catfish linkage map (101). Of the 61 channel catfish *CYP*s, genomic locations of 58 *CYP*s can be determined, while genomic locations for three *CYP*s: *CYP*1A1, *CYP*11C1, and *CYP*27A3, cannot yet be determined. The 58 channel catfish *CYP*s were located on 21 linkage groups of the channel catfish genome (Table 2). Among which, linkage group 28 had the largest number of *CYP*s including five family of eight different genes, followed by LG17 and LG20 with five *CYP*s each.

#### 3.3. Specific *CYP*s and comparative studies

A comparison of *CYP* gene numbers in various species is shown in Table 3. Mammals (human and mouse) had eight specific subfamilies: *CYP*2A, *CYP*2B, *CYP*2E, *CYP*2S, *CYP*2T, *CYP*4A, *CYP*4X and *CYP*4Z; chickens had one specific subfamily, *CYP*2H that is absent from other taxa analyzed. Amphibians have eight specific subfamilies: *CYP*2Q,

CYP2AM, CYP2AN, CYP2AP, CYP2AQ, CYP2AR, CYP2AS and *CYP2AT*. A large number of 18 *CYP* subfamilies appeared to be teleost-specific: CYP2K, CYP2M, CYP2N, CYP2P, CYP2V, CYP2X, CYP2Y, CYP2Z, CYP2AA, CYP2AD, CYP2AE, CYP3B, CYP3C, CYP3D, CYP7C, CYP7D, CYP11C, and *CYP16A* (Table 3). It is noteworthy that these teleost-specific *CYPs* belong to five families, and the vast majority (11) of these belongs to *CYP2* family. Of the 18 teleost-specific subfamilies, 10 were identified from channel catfish (Table 3).

Within the 37 *CYP* subfamilies identified in channel catfish, 13 (CYP2K, CYP2X, CYP2Y, CYP2AD, CYP3A, CYP4B, CYP4F, CYP4T, CYP8A, CYP8B, CYP17A, CYP19A and *CYP27A*) possess multiple copies of *CYP* genes while 24 possess a single copy for each *CYP* gene (Table 3). Among these subfamilies, *CYP27A* subfamily obtained the largest number of *CYP* gene copies of seven (Table 3).

Compared to zebrafish, channel catfish genome appeared to have lost several subfamilies including *CYP2N*, *CYP2P*, *CYP2V*, *CYP2AE*, *CYP39A*, and *CYP46A*. In addition, channel catfish had much fewer copies for *CYP2K*, *CYP2AA*. Zebrafish had 11 copies of *CYP2K* while channel catfish had only 2 copies; similarly, zebrafish had 12 copies of *CYP2AA*, while channel catfish had just a single copy of *CYP2AA* (Table 3).

### 3.4. Phylogenetic analysis and lineage-specific gene duplication

A total of 61 channel catfish *CYP* genes have been phylogenetically analyzed. First, the 61 genes were categorized into Clans by initial phylogenetic analysis. Each clade (Clan) then was subsequently analyzed separately due to the enormous size of the phylogenetic tree. As shown in Figs 1-9, phylogenetic analyses provided clear evidence for the identities of most channel catfish *CYP* genes. Among these, CYP1A1, CYP1B1, CYP1C1, CYP2K8, CYP2K17, CYP2M1, CYP2R1, CYP2U1, CYP2AD6, CYP4V7, CYP4V8, CYP5A1, CYP7A1, CYP7C1, CYP7D1, CYP11A1, CYP11C1, CYP17A1, CYP17A2, CYP19A1, CYP19A2, CYP20A1, CYP21A1, CYP24A1, CYP26A1, CYP26B1, CYP26C1, CYP27A2, CYP27A3, CYP27B1, CYP27C1 and *CYP51A1* were placed into their corresponding clades containing other teleost equivalents, respectively, with strong bootstrap support.

Lineage-specific gene duplication was evident in a number of clans (Clan 2, Clan 3, Clan 4, Clan 7 and mitochondrial clan). For instance, within Clan 2, two *CYP2Y* genes (*CYP2Y7* and *CYP2Y8*), four *CYP2X* genes (*CYP2X1*, *CYP2X20*, *CYP2X21* and *CYP2X22*), one *CYP2AA* gene (*CYP2AA14*), and three *CYP2AD* genes (*CYP2AD8*, *CYP2AD9* and *CYP2AD10*) were clustered together; similarly, four *CYP3A* genes (*CYP3A125*, *CYP3A126*, *CYP3A127* and *CYP3A128*), one *CYP3C* gene (*CYP3C5*), two *CYP4F* genes (*CYP4F60* and *CYP4F66*), two *CYP4T* genes (*CYP4T16* and *CYP4T17*), two *CYP8A* genes (*CYP8A1a* and *CYP8A1b*), three copies of *CYP8B* genes (*CYP8B5*, *CYP8B6* and *CYP8B7*) and five *CYP27A* genes (*CYP27A14*, *CYP27A15*, *CYP27A16*, *CYP27A17* and *CYP27A18*) clustered together to form their own clade adjacent to zebrafish counterparts in teleost branch, respectively, suggesting channel catfish lineage-specific gene duplication. Similar lineage-specific gene duplication may have also occurred as many of zebrafish genes were clustered together themselves as well (Figs 1-4, Fig9).

### 3.5. Syntenic analysis and tandem duplication of channel catfish CYPs

Though phylogenetic relationships provide strong support for the identities of most *CYP* genes, syntenic analyses were required to provide additional evidence for orthologies or otherwise the paralogies. Syntenic analyses for a group of *CYP* genes were conducted, including subfamily 2K, subfamily 2X, subfamily 2Y, subfamily 2AA, subfamily 2AD, subfamily 3A, subfamily 3C, subfamily 4F, subfamily 4T, subfamily 8A, subfamily 27A, subfamily 27B and subfamily 27C.

Positions of these channel catfish *CYP*s and their neighbor genes were identified from the draft genome scaffolds. And the genes were also identified from the zebrafish genome. As shown in Figs. 10-14, conserved syntenic blocks were characterized in these subfamilies between channel catfish and zebrafish counterparts. On the one hand, the conserved syntenies provide the strongest evidence for the orthologies, and on the other hand, a number of *CYP* genes were found to be located together in tandem, apparently as a result of tandem gene duplications. For instance, channel catfish *CYP2AA14* shared the conserved neighbor genes with a cluster of eight zebrafish *CYP2AA* genes on chromosome 23; channel catfish *CYP3C5* shared the conserved neighboring genes with three *CYP3C* genes on chromosome 3; channel catfish *CYP27B1* shared the conserved neighboring genes with zebrafish *CYP27B1* on chromosome 11; and channel catfish *CYP27C1* shared the conserved neighboring genes with zebrafish *CYP27C1* on chromosome 6. Tandem *CYP* genes were found within the conserved syntenies, suggesting both lineage-specific gene duplication in channel catfish and zebrafish, and the paralogies among the channel catfish genes and among the zebrafish genes within the conserved syntenic blocks. For instance, four channel catfish *CYP2AD* genes (*CYP2AD6*, *CYP2AD8*, *CYP2AD9* and *CYP2AD10*) were present in tandem in the channel catfish genome within the conserved syntenic region in zebrafish genome that included 11 *CYP2* genes arranged in tandem arrays (*CYP2J*, five *CYP2Ps*, one *CYP2V*, three *CYP2ADs* and one *CYP2N*); Similarly, two channel catfish *CYP2K* genes (*CYP2K8* and *CYP2K17*) corresponded six *CYP2K* genes, all present in tandem arrays in both species; three channel catfish *CYP2X* genes (*CYP2X1*, *CYP2X20* and *CYP2X21*), two channel catfish *CYP2Y* genes (*CYP2Y7* and *CYP2Y8*), four channel catfish *CYP3A* genes (*CYP3A125*, *CYP3A126*, *CYP3A127* and *CYP3A128*), two channel catfish *CYP4F* genes (*CYP4F60* and *CYP4F66*), two channel catfish *CYP4T* genes (*CYP4T16* and *CYP4T17*) and three channel catfish *CYP27* genes (*CYP27A14*, *CYP27A15* and *CYP27A16*) were all present in tandem in the channel catfish genome, corresponding the conserved syntenies of zebrafish genomic regions containing four *CYP2X* genes, two *CYP2Y* genes, *CYP3A65*, *CYP4F3*, *CYP4T8* and four *CYP27A* genes of zebrafish, respectively. In addition, two *CYP8A* genes were identified in the channel catfish genome that had only one counterpart in the zebrafish genome, the *CYP8A1* gene. We annotated the one with identical neighboring genes in channel catfish as *CYP8A1a* gene, and its paralog as *CYP8A1b* in channel catfish because this paralog shared only *KCNB1* neighboring gene with zebrafish *CYP8A1* gene, following the nomenclature of duplicated genes in zebrafish.

### 3.6. Meta-analyses of CYP expression after bacterial infection using RNA-Seq datasets

Three RNA-Seq datasets, RNA-Seq data from the liver of catfish infected with *E. ictaluri* (93), RNA-Seq data from the intestine of catfish infected with *E. ictaluri* (94), and RNA-Seq

data from the gill of catfish infected with *F. columnare* (95), were used in this study. These datasets were analyzed to determine expression profile of all *CYP* genes after bacterial infection, using the cut-off of 2-fold change, p-value < 0.05, and reads number per gene. There were a total of 36 channel catfish *CYP*s showed significant differential expression for at least one time point post infection in the three tissues compared with control (Table 4).

Several channel catfish *CYP*s exhibited drastic induction or suppression after bacterial infection. As shown in table 4, 3 days after *E. ictaluri* infection in the liver, drastic up-regulation was found in *CYP8A1b*, *CYP8A1a* and *CYP1C1*. *CYP8A1b* was up-regulated over 30-fold; *CYP8A1a* was up-regulated over 27-fold; while *CYP1C1* was up-regulated 17-fold. In contrast, dramatic down-regulation was observed with *CYP2X22*, *CYP4V7*, *CYP2X21*, *CYP2X20* and *CYP7A1* with 52-fold, 32-fold, 31-fold, 14-fold and 11-fold reduction in expression after bacterial infection, respectively.

The involvement of *CYP* genes in the disease response appeared to be at early stages after infection. As shown in Table 4, numerous *CYP* genes exhibited differential expression 3 days after infection, but at the time point of 14 days after *E. ictaluri* infection in the liver, only two *CYP* genes, *CYP4F60* and *CYP8B7*, were highly up-regulated or down-regulated, with eight-fold up and six-fold down for *CYP4F60* and *CYP8B7*, respectively.

Many *CYP* genes were regulated by bacterial infection in the liver, but only few were regulated in the intestine and the gill (Table 4). Compared with the liver where 30 *CYP* genes were significantly regulated by bacterial infection compared with control, only 7 *CYP* genes were regulated in the intestine, and 8 *CYP* genes were regulated in the gill. Not only the number of *CYP* genes under regulation was much fewer in the intestine and gill, the level of induction and suppression was also much less dramatic. The most highly down-regulated *CYP* gene in the intestine is *CYP2X1* (down 5.6X), and the most highly down-regulated *CYP* gene in the gill is *CYP26A1* (down 5.9 X). Apparently, much of this difference was caused by the expression patterns of *CYP* genes, which are naturally expressed most abundantly in the liver (Table 4). Nonetheless, the use of the high throughput next generation sequencing RNA-Seq allowed detection of *CYP* gene expression in tissues other than the liver. A total of seven *CYP* genes were detected in the intestine: *CYP2K17*, *CYP2R1*, *CYP2X1*, *CYP2AA14*, *CYP4T16*, *CYP26A1* and *CYP51A1*; and a total of eight *CYP* genes were detected in the gill: *CYP2X20*, *CYP2AD6*, *CYP4T16*, *CYP17A1*, *CYP19A1*, *CYP26A1*, *CYP27A17*, and *CYP51A1*. Three of these genes, *CYP4T16*, *CYP26A1*, and *CYP51A1* were expressed in all three tested tissues.

#### 4. Discussion

*CYP* genes play key roles in many crucial biological processes including oxidative transformation of xenobiotics and metabolism of endogenous substrates. They belong to one of the most widespread and diverse gene families that consist more than 18,500 members (24) among various species. In spite of their importance, only a few *CYP* genes were characterized from channel catfish (86-89). Systematic analysis of channel catfish *CYP* genes has been lacking. In this study, we identified 61 catfish *CYP* genes, which may represent the vast majority, if not all, *CYP* genes in the channel catfish genome. This



assessment is based on the resources we used for the identification of these genes: several hundred thousands of ESTs (101-106), RNA-Seq (93-95), and the draft genome sequences that are yet not published, but represent over 200× genome coverage (unpublished data). This repertoire of genomic resources allowed thorough identification of *CYPs* in channel catfish. However, it is possible that additional *CYP* genes are yet to be discovered. At any rate, we believe that the vast majority of channel catfish *CYPs* in channel catfish have been discovered in this study.

Phylogenetic and syntenic analyses allowed annotation of these genes, and revealed that many of the *CYP* genes had gone through lineage-specific gene duplications, leading to the presence of a large number of paralogues within a species. Meta-analysis was conducted for the analysis of *CYP* gene expression. To our surprise, a large number of *CYP* genes, 36 in total, were up- or down-regulated after bacterial infection, suggesting their involvement in disease responses. The transcripts of all 61 P450 genes in the channel catfish not only were detected from the liver, where they are believed to be expressed, a significant number of *CYP* genes were expressed in tissues other than the liver, 7 in the intestine and 8 in the gill.

#### 4.1. *CYP* families 1-4

The significance of *CYP* genes in detoxification of a variety of environmental pollutants, food additives, organic compounds and even drugs in aquatic species has been well established in previous studies (32, 55). This feature is particularly important for channel catfish that inhabits at the bottom of water column. Among all the *CYPs*, the genes in families of *CYP1*, *CYP2*, *CYP3* and *CYP4* were reported to be highly involved in metabolism of xenobiotics, drugs and fatty acids (55). Identification of *CYPs* in these four families is critical for understanding of the detoxification mechanism in channel catfish.

In the present study, we identified 30 *CYP* genes belong to *CYP* families 1, 2, 3 and 4 in channel catfish (Table 2). In contrast to those *CYPs* primarily involved in endogenous metabolism, *CYP* genes among families 1-4 exhibited a larger degree of divergence across species (Table 3). This is strikingly significant in some fish-specific subfamilies belong to *CYP2* family and *CYP3* family. For example, gene number differences were observed in *CYP2K*, *CYP2N*, *CYP2P*, *CYP2X*, *CYP2AA*, *CYP2AD*, *CYP3A* and *CYP3C* subfamilies among species (Table 3). This observation indicated that these *CYP* subfamilies exhibited high level of divergence could be result of individual gene duplication or gene loss, or remnants of the genome duplication (55). Among these, *CYP2N* and *CYP2P* genes were commonly found in teleost species (45, 55, 64, 108, 109) but absent in catfish. In killifish, these two genes have been shown to catalyze benzphetamine N-demethylation and metabolize arachidonic acid (108, 109). In catfish, however, their function could be replaced by other *CYP* genes during evolution. For example, *CYP2X1* in channel catfish have been proved to possess benzphetamine demethylase activity (110). In addition, only subfamilies *CYP1A*, *CYP1B*, *CYP2U* and *CYP2R* appear to be evolutionarily conserved across species. Of these four subfamilies, *CYP2R1* and *CYP2U1* catalyze modifications on the endogenous substrates vitamin D and arachidonic acid (101-112), while *CYP1A* and *CYP1B* are induced by a variety of drugs or contaminants (32, 113-115).

Phylogenetic analysis and syntenic analysis provided strong support for the identity of the majority of *CYPs* in channel catfish. However, for subfamily *CYP2Y*, *CYP2X*, *CYP2AA*, *CYP2AD*, *CYP3A*, *CYP3C*, *CYP4F* and *CYP4T*, phylogenetic analysis did not yield informative conclusion concerning their identities because members in these subfamilies formed their own clades in the phylogenetic tree, respectively. Consistent with phylogeny, syntenic analyses showed that genes in these *CYP* subfamilies existed as tandem duplication arrays which shared synteny with corresponding gene clusters in zebrafish, indicating that members of these subfamilies could be derived from recent lineage-specific gene duplication events. As such, these tandem arranged *CYP* genes are paralogous one another within a conserved syntenic block. The high level of lineage-specific multiplication of these *CYP* genes may suggest that the involved organisms were under evolutionary selection for the rapid expansion of such *CYP* genes, perhaps in the face of heavy environmental pollution.

#### 4.2. *CYP* families 5-51

*CYP* genes in families 5-51, including *CYP5A1*, *CYP7A1*, *CYP7C1*, *CYP7D1*, *CYP11A1*, *CYP11C1*, *CYP17A1*, *CYP17A2*, *CYP19A1*, *CYP19A2*, *CYP20A1*, *CYP21A1*, *CYP24A1*, *CYP26A1*, *CYP26B1*, *CYP26C1*, *CYP27A2*, *CYP27A3*, *CYP27B1*, *CYP27C1* and *CYP51A1* are mainly involved in the metabolism of endogenous substrates (32, 36). In contrast to *CYP* gene families 1-4, all *CYP* subfamilies in families 5-51 have single copy gene in channel catfish with the exception of *CYP8A*, *CYP8B*, *CYP17A*, *CYP19A* and *CYP27A*, in which 2, 3, 2, 2 and 7 copies are presented in each of them. Phylogenetic analysis provides clear evidence for majority of *CYP* genes in these families, where they exhibit 1:1 correspondence with teleost counterparts, indicating conservation of enzyme activities and physiological functions.

Syntenic analysis of *CYP8A* genes indicated that there were two conserved syntenic blocks in channel catfish in comparison with just one in zebrafish. One of them contains five conserved genes and the other one possesses only two conserved genes in the block. We simply named the first one as *CYP8A1a* and the second one as *CYP8A1b*, following the nomenclature rule of zebrafish duplicated genes. Apparently, segmental duplication in the catfish genome accounted for the observed additional copy of the *CYP8A1* gene. It is likely that several genes in the conserved syntenic block of the catfish genome were lost after segmental duplication.

For *CYP27A* subfamily, syntenic analyses were only available for *CYP27A14*, *CYP27A15* and *CYP27A16*. These three genes showed a conserved syntenic block with zebrafish *CYP27A* cluster. Syntenic analysis of *CYP8B* genes, *CYP27A17* gene and *CYP27A18* gene were not available at this stage because missing information of their neighboring genes. Gap filling in the genomic sequence is required to provide a better resource for syntenic analysis in the future.

*CYP46A1* gene in mammals is a cholesterol 24-hydroxylase enzyme and only present in the brain. It has been widely identified among species including teleost fish and plays an essential role in the majority of cholesterol turnover in vertebrate central nervous system (116). Though the function of *CYP46* gene has not been studied in teleost, mutation of *CYP46A1* gene could lead to serious neurodegenerative disease including Multiple Sclerosis,

Alzheimer and Huntington Diseases in human (117). Four *CYP46A* genes have been identified in zebrafish (55). However, in channel catfish, no *CYP46* homolog was found in any databases (EST, RNA-seq and genome sequence) in this study. Syntenic analyses with zebrafish indicated a potential gene loss in the conserved block between *ak7a* gene and *SPTLC2* gene (Fig 15). We had also tried to amplify this gene in the catfish brain tissue using degenerate primers, in case that the gene may have existed, but had not been found. We were unable to amplify the *CYP46* gene transcripts from the brain (data not shown). Although the negative PCR results still does not exclude the possibility of *CYP46* presence in channel catfish, it is highly possible that channel catfish has lost the *CYP46* gene set in its genome. Further analyses are required to validate this speculation. In addition to *CYP46*, the catfish appeared not to possess *CYP16* and *CYP39*. *CYP16* was also lost in the zebrafish genome (55), and *CYP39* was lost in the fugu genome (45).

### 4.3. *CYP* gene expression in disease defense response

RNAseq-based expression analysis has become a robust method to assess transcriptional profile to different challenge experiment (118). As described in our recent studies (93-95), using the 100 bp paired-end reads, we successfully captured comprehensive transcriptome from catfish intestine and gill after *E. ictaluri* and *Columnare* infection, respectively. The expression patterns of differentially expressed genes from these two studies were validated by quantitative real-time RT-PCR with average correlation coefficient around 0.9 ( $p < 0.001$ ). In the present work, these two datasets from RNA-Seq of the intestine and gill after *E. ictaluri* and *Columnare* infection, along with dataset from RNA-Seq of the liver after ESC challenge, were utilized to analyze expression profiles of *CYP* genes in channel catfish after infection.

The expression patterns of *CYP* genes following *E. ictaluri* infection in the liver exhibited drastic differences as compared to the control. Up-regulation was observed 3 days after infection for a large number of *CYPs* including *CYP1C1*, *CYP2AD8*, *CYP3A128*, *CYP3C5*, *CYP4F60*, *CYP4F66*, *CYP4T16*, *CYP7D1*, *CYP8A1a*, *CYP8A1b*, *CYP8B5*, *CYP8B6*, *CYP11A1*, *CYP21A1*, *CYP27A2* and *CYP51A1*. This is the first report involving induction of this large number of *CYP* genes after infection. In a study conducted by Sewer et al. (14), mRNAs of all three of the *CYP4A* subfamily members (*CYP4A1*, *CYP4A2* and *CYP4A3*) were found to be induced 2- to 6-fold in the F344 rat livers after LPS administration. Later (15), the same group observed that this induction was not unique to LPS, *CYP4A* mRNA expression levels were also induced by irritants such as  $\text{SiO}_2$  and  $\text{BaSO}_4$ . Though the mechanism behind this is not clear, it is possible that these *CYPs* are involved in the synthesis of mediators of inflammation cascade. Although the detailed mechanism for the involvement of such a large number of *CYP* genes in defense responses after infection is unknown at present, the disease induced expression suggested that they are important mediators in defense. Additional research is warranted to explore how the *CYP* genes are involved.

Down-regulation of *CYP* genes were observed 3 days after infection in liver with *CYP2M1*, *CYP2R1*, *CYP2X1*, *CYP2X20*, *CYP2X21*, *CYP2X22*, *CYP2Y8*, *CYP4V7*, *CYP7A1*, *CYP8B7*, *CYP24A1*, *CYP26A1*, *CYP27A3* and *CYP27A14*. Similar observation has been

reported by Chaluvadi et al., who demonstrated that infection of mice with enteropathogenic bacterium *C. rodentium* could cause selective down-regulation of hepatic cytochrome P450 mRNA and protein levels (119). Cui et al. also found that *CYP4F4* and *CYP4F5* were suppressed at early stage but induced after 24h of LPS treatment, and reached the highest levels at 2 weeks post-injury (18). In addition, Renton and Nicholson reported that *CYPs* were down-regulated after LPS treatment in the brain of rats (16). In all these cases, it is hypothesized that the pattern of down-regulation *CYPs* is a pathophysiological consequence of inflammatory process, and regulated by inflammatory mediators, such as cytokines. Similar study has been reported by Morgan (11), who found that down-regulation in enzyme activities of *CYP1A1*, *CYP2C11*, *CYP2C12*, *CYP2E1* and *CYP3A2* were triggered by inflammation mediators including cytokines IL-1, IL-6, or tumor necrosis factor (TNF) in liver of rats. Though the mechanisms under this down-regulation of *CYP* genes have not been clearly established, several hypotheses have been proposed by Morgan (12): firstly, the down-regulation of *CYP* genes may be associated with oxidative damage in liver; secondly, *CYP* genes down-regulation during an inflammatory response could be related to their function in metabolism of arachidonic acid; Lastly, the reason for down-regulation of *CYP* enzymes in the liver could be related to their ability to form nitric oxide. Further work is needed to elucidate the role of *CYPs* in the inflammation response, and particularly after the bacterial infection.

## 5. Conclusion

In summary, we systematically identified and characterized a repertoire of 61 *CYP* genes in channel catfish and studied their expression profile after bacterial infection. Strikingly large numbers of *CYP* genes appear to be involved in the bacterial defense processes.

## Acknowledgements

We thank Alabama Supercomputer Center for providing the computer capacity for the bioinformatic analysis. This project was supported by Agriculture and Food Research Initiative Competitive Grant no. 2009-2009-35205-05101, 2010-65205-20356 and 2012-67015-19410 from the USDA National Institute of Food and Agriculture (NIFA). D. Nelson contribution supported by NIH U41-HG003345.

## Reference

1. Omura T, Sato R. The carbon monoxide-binding pigment of liver microsomes I. evidence for its hemoprotein nature. *The Journal of biological chemistry*. 1964; 239:2370–2378. [PubMed: 14209971]
2. Werck-reichhart D, Feyereisen R. Protein family review Cytochromes P450 : a success story. *Genome biology*. 2000; 1:3003.1–3003.9.
3. Bernhardt R. Cytochromes P450 as versatile biocatalysts. *Journal of biotechnology*. 2006; 124:128–145. [PubMed: 16516322]
4. Hrycay EG, Bandiera SM. The monooxygenase, peroxidase, and peroxygenase properties of cytochrome P450. *Archives of biochemistry and biophysics*. 2012; 522:71–89. [PubMed: 22266245]
5. Nehert DW, Gonzalez F. P450 Genes: Structure, Evolution, and Regulation. *Biochemistry*. 1987; 56:945–993.
6. Degtyarenko KN, Archakov AI. Molecular evolution of P450 superfamily and P450-containing monooxygenase systems. *FEBS letters*. 1993; 332:1–8. [PubMed: 8405421]

7. Denisov IG, Makris TM, Sligar SG, Schlichting I. Structure and chemistry of cytochrome P450. *Chemical reviews*. 2005; 105:2253–2277. [PubMed: 15941214]
8. Hannemann F, Bichet A, Ewen KM, Bernhardt R. Cytochrome P450 systems--biological variations of electron transport chains. *Biochimica et biophysica acta*. 2007; 1770:330–344. [PubMed: 16978787]
9. Nebert DW, Russell DW. Clinical importance of the cytochromes P450. *Lancet*. 2002; 360:1155–1162. [PubMed: 12387968]
10. Nebert DW, Wikvall K, Miller WL. Human cytochromes P450 in health and disease. *Philosophical Transactions of The Royal Society B*. 2013; 368 20120431.
11. Morgan ET. Regulation of cytochromes p450 during inflammation and infection. *Drug Metabolism Reviews*. 1997; 29:1129–1188. [PubMed: 9421688]
12. Morgan ET. Regulation of cytochrome p450 by inflammatory mediators: why and how? *Drug metabolism and disposition*. 2001; 29:207–212. [PubMed: 11181485]
13. Renton KW. Cytochrome P450 regulation and drug biotransformation during inflammation and infection. *Current Drug Metabolism*. 2004; 5:235–243. [PubMed: 15180493]
14. Sewer MB, Koop DR, Morgan ET. Endotoxemia in rats is associated with induction of the P450 4A subfamily and suppression of several other forms of cytochrome P450. *Drug Metabolism and Disposition*. 1996; 4:401–407. [PubMed: 8801054]
15. Sewer MB, Koop DR, Morgan ET. Differential inductive and suppressive effects of endotoxin and particulate irritants on hepatic and renal cytochrome P-450 expression. *The Journal of pharmacology and experimental therapeutics*. 1997; 280:1445–1454. [PubMed: 9067334]
16. Renton KW, Nicholson TE. Hepatic and central nervous system cytochrome P450 are down-regulated during lipopolysaccharide-evoked localized inflammation in brain. *The Journal of pharmacology and experimental therapeutics*. 2000; 294:524–530. [PubMed: 10900228]
17. Monshouer M, Witkamp RF, Nuijmeijer SM, Van Amsterdam JG, Van Miert a S. Suppression of cytochrome P450- and UDP glucuronosyl transferase-dependent enzyme activities by proinflammatory cytokines and possible role of nitric oxide in primary cultures of pig hepatocytes. *Toxicology and applied pharmacology*. 1996; 137:237–244. [PubMed: 8661349]
18. Cui X, Kalsotra a, Robida a M, Matzilevich D, Moore a N, Boehme CL, Morgan ET, Dash PK, Strobel HW. Expression of cytochromes P450 4F4 and 4F5 in infection and injury models of inflammation. *Biochimica et biophysica acta*. 2003; 1619:325–331. [PubMed: 12573492]
19. Richardson TA, Sherman M, Antonovic L, Kardar SS, Henry W, Kalman D, Morgan ET. Hepatic and renal cytochrome P450 gene regulation during *Citrobacter rodentium* infection in wildtype and Toll-like Receptor 4 mutant mice. *Drug Metabolism and Disposition*. 2007; 34:354–360. [PubMed: 16339354]
20. Richardson TA, Morgan ET. Hepatic cytochrome p450 gene regulation during endotoxin induced inflammation in nuclear receptor knockout mice. *The journal of pharmacology and experimental therapeutics*. 2005; 314:703–709. [PubMed: 15860574]
21. Satarug S, Lang MA, Yongvanit P, Lang MA. Induction of cytochrome P450 2A6 expression in humans by the carcinogenic parasite infection, *opisthorchiasis viverrini*. *Cancer Epidemiol Biomarkers Prevention*. 1996; 5:795–800.
22. Nelson DR. The cytochrome p450 homepage. *Human genomics*. 2009; 4:59–65. [PubMed: 19951895]
23. Nelson DR. Progress in tracing the evolutionary paths of cytochrome P450. *Biochimica et biophysica acta*. 2011; 1814:14–18. [PubMed: 20736090]
24. Nelson DR. A world of cytochrome P450s. *Philosophical Transactions of The Royal Society B*. 2013; 368 20120430.
25. Nelson DR, Zeldin DC, Hoffman SMG, Maltais LJ, Wain HM, Nebert DW. Comparison of cytochrome P450 (CYP) genes from the mouse and human genomes, including nomenclature recommendations for genes, pseudogenes and alternative-splice variants. *Pharmacogenetics*. 2003; 450:1–18.
26. Bak S, Beisson F, Bishop G, Hamberger B, Höfer R, Paquette S, Werck-Reichhart D. Cytochromes p450. *The Arabidopsis book / American Society of Plant Biologists*. 2011; 9:e0144. [PubMed: 22303269]

27. Baldwin WS, Marko PB, Nelson DR. The cytochrome P450 (CYP) gene superfamily in *Daphnia pulex*. *BMC genomics*. 2009; 10:169. [PubMed: 19383150]
28. Guo H, Bao Z, Du H, Zhang L, Wang S, Sun L, Mou X, Hu X. Identification of Cytochrome P450 (CYP) genes in Zhikong scallop (*Chlamys farreri*). *Journal of Ocean University of China*. 2012; 12:97–102.
29. Ai J, Zhu Y, Duan J, Yu Q, Zhang G, Wan F, Xiang Z. Genome-wide analysis of cytochrome P450 monooxygenase genes in the silkworm, *Bombyx mori*. *Gene*. 2011; 480:42–50. [PubMed: 21440608]
30. Diaz GJ, Murcia HW, Cepeda SM. Cytochrome P450 enzymes involved in the metabolism of aflatoxin B1 in chickens and quail. *Poultry science*. 2010; 89:2461–2469.
31. Nelson DR, Ming R, Alam M, Schuler M. a. Comparison of Cytochrome P450 Genes from Six Plant Genomes. *Tropical Plant Biology*. 2008; 1:216–235.
32. Uno T, Ishizuka M, Itakura T. Cytochrome P450 (CYP) in fish. *Environmental toxicology and pharmacology*. 2012; 34:1–13. [PubMed: 22418068]
33. Lee DS, Nioche P, Hamberg M, Raman CS. Structural insights into the evolutionary paths of oxylipin biosynthetic enzymes. *Nature*. 2008; 455:363–368. [PubMed: 18716621]
34. Nebert DW, Adesnik M, Coon MJ, Estabrook RW, Gonzalez FJ, Guengerich FP, Gunsalus IC, Johnson EF, Kemper B, Levin W, et al. The P450 gene superfamily: recommended nomenclature. *DNA*. 1987; 6:1–11. [PubMed: 3829886]
35. Nelson DR. Cytochrome P450 nomenclature, *Methods in molecular biology* (Clifton, N.J.). 2004; 320:1–10.
36. Nelson DR, Goldstone JV, Stegeman JJ. The cytochrome P450 genesis locus : the origin and evolution of animal cytochrome P450s. *Philosophical Transactions of the Royal Society*. 2013; 368 20120474.
37. Sakai N, Tanaka M, Adachi S, Millerh WL, Nagahama Y. Rainbow trout cytochrome P-450c17 (17 alpha-hydroxylase/17, 20-lyase). cDNA cloning, enzymatic properties and temporal pattern of ovarian P450c17 mRNA expression during oogenesis. *FEBS letters*. 1992; 301:60–64. [PubMed: 1451787]
38. Buhler DR, Yang YH, Dreher TW, Miranda CL, Wang JL. Cloning and sequencing of the major rainbow trout constitutive cytochrome P450 (CYP2K1): identification of a new cytochrome P450 gene subfamily and its expression in mature rainbow trout liver and trunk kidney. *Archives of biochemistry and biophysics*. 1994; 312:45–51. [PubMed: 8031145]
39. Activity A, Yang Y, Wang J, Miranda CL, Buhler DR. CYP2M1 : cloning, sequencing, and expression of a new cytochrome P450 from rainbow trout liver with fatty acid (omega-6)-hydroxylation activity. *Archives of biochemistry and biophysics*. 1998; 352:271–280. [PubMed: 9587416]
40. Carvan M, Ponomareva L, Solis W, Matlib R, Puga A, Nebert D. Trout CYP1A3 Gene: Recognition of Fish DNA Motifs by Mouse Regulatory Proteins. *Marine biotechnology (New York, N.Y.)*. 1999; 1:155–166.
41. Råbergh CM, Vrolijk NH, Lipsky MM, Chen TT. Differential expression of two CYP1A genes in rainbow trout (*Oncorhynchus mykiss*). *Toxicology and applied pharmacology*. 2000; 165:195–205. [PubMed: 10860869]
42. Kusakabe M, Kobayashi T, Todo T, Mark Lokman P, Nagahama Y, Young G. Molecular cloning and expression during spermatogenesis of a cDNA encoding testicular 11beta-hydroxylase (P45011beta) in rainbow trout (*Oncorhynchus mykiss*). *Molecular reproduction and development*. 2002; 62:456–469. [PubMed: 12112578]
43. Lee SJ, Buhler DR. Cloning, tissue distribution, and functional studies of a new cytochrome P450 3A subfamily member, CYP3A45, from rainbow trout (*Oncorhynchus mykiss*) intestinal ceca. *Archives of Biochemistry and Biophysics*. 2003; 412:77–89. [PubMed: 12646270]
44. Gomez CF, Constantine L, Moen M, Vaz A, Wang W, Huggett DB. Ibuprofen metabolism in the liver and gill of rainbow trout, *Oncorhynchus mykiss*. *Bulletin of environmental contamination and toxicology*. 2011; 86:247–51. [PubMed: 21301807]
45. Nelson DR. Comparison of P450s from human and fugu: 420 million years of vertebrate P450 evolution. *Archives of biochemistry and biophysics*. 2003; 409:18–24. [PubMed: 12464240]

46. Kim JH, Raisuddin S, Ki JS, Lee JS, Han KN. Molecular cloning and beta-naphthoflavone-induced expression of a cytochrome P450 1A (CYP1A) gene from an anadromous river pufferfish, *Takifugu obscurus*. *Marine pollution bulletin*. 2008; 57:433–440. [PubMed: 18304588]
47. Chiang EF, Yan YL, Guiguen Y, Postlethwait J, Chung B, Two Cyp19 (P450 aromatase) genes on duplicated zebrafish chromosomes are expressed in ovary or brain. *Molecular biology and evolution*. 2001; 18:542–550. [PubMed: 11264405]
48. Goto-Kazeto R, Kight KE, Zohar Y, Place AR, Trant JM. Localization and expression of aromatase mRNA in adult zebrafish. *General and comparative endocrinology*. 2004; 139:72–84. [PubMed: 15474538]
49. Wang Y, Ge W. Cloning of zebrafish ovarian P450c17 (CYP17, 17 $\alpha$ -hydroxylase/17, 20-lyase) and characterization of its expression in gonadal and extra-gonadal tissues. *General and Comparative Endocrinology*. 2004; 135:241–249. [PubMed: 14697311]
50. Gu X, Xu F, Wang X, Gao X, Zhao Q. Molecular cloning and expression of a novel CYP26 gene (cyp26d1) during zebrafish early development. *Gene expression patterns*. 2005; 5:733–739. [PubMed: 15979416]
51. Tseng H, Hseu T, Buhler DR, Wang W, Hu C. Constitutive and xenobiotics-induced expression of a novel CYP3A gene from zebrafish larva. *Toxicology and applied pharmacology*. 2005; 205:247–258. [PubMed: 15922010]
52. Wang-Buhler JL, Lee SJ, Chung WG, Stevens JF, Tseng HP, Hseu TH, Hu CH, Westerfield M, Yang YH, Miranda CL, Buhler DR. CYP2K6 from zebrafish (*Danio rerio*): cloning, mapping, developmental/tissue expression, and aflatoxin B1 activation by baculovirus expressed enzyme. *Comparative biochemistry and physiology. Toxicology & pharmacology*. 2005; 140:207–219.
53. Corley-Smith GE, Su H, Wang-Buhler J, Tseng H, Hu C, Hoang T, Chung W, Buhler DR. CYP3C1, the first member of a new cytochrome P450 subfamily found in zebrafish (*Danio rerio*). *Biochemical and biophysical research communications*. 2006; 340:1039–1046. [PubMed: 16414346]
54. Jönsson ME, Orrego R, Woodin BR, Goldstone JV, Stegeman JJ. Basal and 3, 3', 4, 4', 5-pentachlorobiphenyl-induced expression of cytochrome P450 1A, 1B and 1C genes in zebrafish. *Toxicology and applied pharmacology*. 2007; 221:29–41. [PubMed: 17445853]
55. Goldstone JV, McArthur AG, Kubota A, Zanette J, Parente T, Jönsson ME, Nelson DR, Stegeman JJ. Identification and developmental expression of the full complement of Cytochrome P450 genes in Zebrafish. *BMC genomics*. 2010; 11:643. [PubMed: 21087487]
56. Scornaienchi ML, Thornton C, Willett KL, Wilson JY. Cytochrome P450-mediated 17 $\beta$ -estradiol metabolism in zebrafish (*Danio rerio*). *The Journal of endocrinology*. 2010; 206:317–325. [PubMed: 20522564]
57. Arukwe A. Complementary DNA cloning, sequence analysis and differential organ expression of beta-naphthoflavone-inducible cytochrome P4501A in Atlantic salmon (*Salmo salar*). *Comparative biochemistry and physiology. Toxicology & pharmacology*. 2002; 133:613–624.
58. Lyssimachou A, Jenssen BM, Arukwe A. Brain cytochrome P450 aromatase gene isoforms and activity levels in Atlantic salmon after waterborne exposure to nominal environmental concentrations of the pharmaceutical ethynylestradiol and antifoulant tributyltin. *Toxicological sciences : an official journal of the Society of Toxicology*. 2006; 91:82–92. [PubMed: 16484284]
59. Vang S-H, Kortner TM, Arukwe A. Steroidogenic acute regulatory (StAR) protein and cholesterol side-chain cleavage (P450scc) as molecular and cellular targets for 17 $\alpha$ -ethynylestradiol in salmon previtellogenic oocytes. *Chemical research in toxicology*. 2007; 20:1811–1819. [PubMed: 18001058]
60. Sanden M, Olsvik PA. Intestinal cellular localization of PCNA protein and CYP1A mRNA in Atlantic salmon *Salmo salar* L. exposed to a model toxicant. *BMC physiology*. 2009; 9:3. [PubMed: 19309504]
61. Stien X, Amichot M, Berge JB, Lafaurie M. Molecular cloning of a CYP1A cDNA from the teleost fish *Dicentrarchus labrax*. *Comparative biochemistry and physiology Toxicology & pharmacology*. 1998; 121:241–248. 1998.

62. Sabourault C, Bergé J, Lafaurie M, Girard JP, Amichot M. Molecular cloning of a phthalate-inducible CYP4 gene (CYP4T2) in kidney from the sea bass, *Dicentrarchus labrax*. Biochemical and biophysical research communications. 1998; 251:213–219. [PubMed: 9790933]
63. Vaccaro E, Salvetti A, Del Carratore R, Nencioni S, Longo V, Gervasi PG. Cloning, tissue expression, and inducibility of CYP 3A79 from sea bass (*Dicentrarchus labrax*). Journal of biochemical and molecular toxicology. 2007; 21:32–40. [PubMed: 17366542]
64. Barber DS, McNally AJ, Garcia-Reyero N, Denslow ND. Exposure to p, p' -DDE or dieldrin during the reproductive season alters hepatic CYP expression in largemouth bass (*Micropterus salmoides*). Aquatic Toxicology. 2007; 81:27–35. [PubMed: 17145087]
65. Melo AC, Ramsdell JS. Sexual dimorphism of brain aromatase activity in medaka: induction of a female phenotype by estradiol. Environmental health perspectives. 2001; 109:257–264. [PubMed: 11333187]
66. Kim I, Kim YJ, Yoon Y, Kawamura S, Lee Y, Lee J. Cloning of cytochrome P450 1A (CYP1A) genes from the hermaphrodite fish *Rivulus marmoratus* and the Japanese medaka *Oryzias latipes*. Marine environmental research. 2004; 58:125–129. [PubMed: 15178024]
67. Kullman SW, Kashiwada S, Hinton DE. Analysis of medaka cytochrome P450 3A homotropic and heterotropic cooperativity. Marine environmental research. 2004; 58:469–473. [PubMed: 15178067]
68. Kashiwada S, Hinton DE, Kullman SW. Functional characterization of medaka CYP3A38 and CYP3A40: kinetics and catalysis by expression in a recombinant baculovirus system. Comparative biochemistry and physiology Toxicology & pharmacology. 2005; 141:338–348.
69. Yokota H, Abe T, Nakai M, Murakami H, Eto C, Yakabe Y. Effects of 4-tert-pentylphenol on the gene expression of P450 11beta-hydroxylase in the gonad of medaka (*Oryzias latipes*). Aquatic toxicology (Amsterdam, Netherlands). 2005; 71:121–132.
70. Goldstone JV, Stegeman JJ. Gene structure of the novel cytochrome P4501D1 genes in stickleback (*Gasterosteus aculeatus*) and medaka (*Oryzias latipes*). Marine environmental research. 2008; 66:19–20. [PubMed: 18400289]
71. Yamauchi R, Ishibashi H, Hirano M, Mori T, Kim J, Arizono K. Effects of synthetic polycyclic musks on estrogen receptor, vitellogenin, pregnane X receptor, and cytochrome P450 3A gene expression in the livers of male medaka (*Oryzias latipes*). Aquatic toxicology (Amsterdam, Netherlands). 2008; 90:261–268.
72. El-kady MA, Mitsuo R, Kaminishi Y, Itakura T. cDNA cloning, sequence analysis and expression of 3-methylcholanthrene-inducible cytochrome P450 1B1 in carp (*Cyprinus carpio*). Environmental Science. 2004; 11:231–240. 2004.
73. Itakura T, El-Kady M, Mitsuo R, Kaminishi Y. Complementary DNA cloning and constitutive expression of cytochrome P450 1C1 in the gills of carp (*Cyprinus carpio*). Environmental Science. 2005; 12:111–120.
74. Barney ML, Patil JG, Gunasekera RM, Carter CG. Distinct cytochrome P450 aromatase isoforms in the common carp (*Cyprinus carpio*): sexual dimorphism and onset of ontogenic expression. General and comparative endocrinology. 2008; 156:499–508. [PubMed: 18430424]
75. Morrison HG, Weil EJ, Karchner SI, Sogin ML, Stegeman JJ. Molecular cloning of CYP1A from the estuarine fish *Fundulus heteroclitus* and phylogenetic analysis of CYP1 genes: update with new sequences. Comparative biochemistry and physiology Part C, Pharmacology, toxicology & endocrinology. 1998; 121:231–240.
76. Hegelund T, Celander MC. Hepatic versus extrahepatic expression of CYP3A30 and CYP3A56 in adult killifish (*Fundulus heteroclitus*). Aquatic Toxicology. 2003; 64:277–291. [PubMed: 12842592]
77. Wang L, Scheffler BE, Willett KL. CYP1C1 messenger RNA expression is inducible by benzo[a]pyrene in *Fundulus heteroclitus* embryos and adults. Toxicological sciences. 2006; 93:331–340. 2006. [PubMed: 16873418]
78. Zanette J, Jenny MJ, Goldstone JV, Woodin BR, Watka L. a, Bainy ACD, Stegeman JJ. New cytochrome P450 1B1, 1C2 and 1D1 genes in the killifish *Fundulus heteroclitus*: Basal expression and response of five killifish CYP1s to the AHR agonist PCB126. Aquatic toxicology (Amsterdam, Netherlands). 2009; 93:234–243.



79. Calò M, Bitto a, Lo Cascio P, Polito F, Lauriano ER, Minutoli L, Altavilla D, Squadrito F. Cytochrome P450 (CYP1A) induction in sea bream (*Sparus Aurata*) gills and liver following exposure to polychlorobiphenyls (PCBs). *Veterinary research communications*. 2009; 33:181–184. [PubMed: 19588260]
80. Halm S, Kwon JY, Rand-Weaver M, Sumpter JP, Pounds N, Hutchinson TH, Tyler CR. Cloning and gene expression of P450 17 $\alpha$ -hydroxylase, 17, 20-lyase cDNA in the gonads and brain of the fathead minnow *Pimephales promelas*. *General and comparative endocrinology*. 2003; 130:256–266. [PubMed: 12606268]
81. Christen V, Caminada D, Arand M, Fent K. Identification of a CYP3A form (CYP3A126) in fathead minnow (*Pimephales promelas*) and characterisation of putative CYP3A enzyme activity. *Analytical and bioanalytical chemistry*. 2010; 396:585–595. [PubMed: 19898817]
82. Deng S, Chen S, Xu J, Liu B. Molecular cloning, characterization and expression analysis of gonadal P450 aromatase in the half-smooth tongue-sole, *Cynoglossus semilaevis*. *Aquaculture*. 2009; 287:211–218.
83. Chen CF, Wen HS, Wang ZP, He F, Zhang JR, Chen XY, Jin GX, Shi B, Shi D, Yang YP, Li JF, Qi BX, Li N. Cloning and expression of P450c17-I (17 $\alpha$ -hydroxylase/17, 20-lyase) in brain and ovary during gonad development in *Cynoglossus semilaevis*. *Fish physiology and biochemistry*. 2010; 36:1001–1012. [PubMed: 20069358]
84. Aparicio S, Chapman J, Stupka E, Putnam N, Chia J-M, Dehal P, Christoffels A, Rash S, Hoon S, Smit A, Gelpke MDS, Roach J, Oh T, Ho IY, Wong M, Detter C, Verhoeve F, Predki P, Tay A, Lucas S, Richardson P, Smith SF, Clark MS, Edwards YJK, Doggett N, Zharkikh A, Tavtigian SV, Pruss D, Barnstead M, Evans C, Baden H, Powell J, Glusman G, Rowen L, Hood L, Tan YH, Elgar G, Hawkins T, Venkatesh B, Rokhsar D, Brenner S. Whole-genome shotgun assembly and analysis of the genome of *Fugu rubripes*. *Science (New York, N.Y.)*. 2002; 297:1301–1310.
85. Howe K, Clark MD, Torroja CF, Torrance J, Berthelot C, Muffato M, Collins JE, Humphray S, McLaren K, Matthews L, McLaren S, Sealy I, Caccamo M, Churcher C, Scott C, Barrett JC, Koch R, Rauch G-J, White S, Chow W, Kilian B, Quintais LT, Guerra-Assunção J. a. Zhou Y, Gu Y, Yen J, Vogel J-H, Eyre T, Redmond S, Banerjee R, et al. The zebrafish reference genome sequence and its relationship to the human genome. *Nature*. 2013 doi: 10.1038.
86. Willett KL, Ganesan S, Patel M, Metzger C, Quiniou S, Waldbieser G, Scheffler B. In vivo and in vitro CYP1B mRNA expression in channel catfish. *Marine environmental research*. 2006; 62(Suppl):332–336.
87. Schlenk D, Furnes B, Zhou X, Debusk BC. Cloning and sequencing of cytochrome P450 2X1 from channel catfish (*Ictalurus punctatus*). *Marine environmental research*. 2002; 54:391–394. [PubMed: 12408592]
88. Trant JM. Isolation and characterization of the cDNA encoding the channel catfish (*Ictalurus punctatus*) form of cytochrome P450arom. *General and comparative endocrinology*. 1994; 95:155–168. [PubMed: 7958747]
89. Kazeto Y, Trant JM. Molecular biology of channel catfish brain cytochrome P450 aromatase (CYP19A2): cloning, preovulatory induction of gene expression, hormonal gene regulation and analysis of promoter region. *Journal of molecular endocrinology*. 2005; 35:571–583. [PubMed: 16326841]
90. Liu Z. Development of genomic resources in support of sequencing, assembly, and annotation of catfish genome. *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics*. 2011; 6:11–17.
91. Li P, Peatman E, Wang S, Feng J, He C, Baoprasertkul P, Xu P, Kucuktas H, Nandi S, Somridhivej B, Serapion J, Simmons M, Turan C, Liu L, Muir W, Dunham R, Brady Y, Grizzle J, Liu Z. Towards the ictalurid catfish transcriptome: generation and analysis of 31,215 catfish ESTs. *BMC Genomics*. 2007; 8:177. [PubMed: 17577415]
92. Wang S, Abernathy J, Waldbieser G, Lindquist E, Richardson P, Lucas S, Wang M, Li P, Thimmapuram J, Liu L, Vullaganti D, Kucuktas H, Murdock C, Small B, Wilson M, Liu H, Jiang Y, Lee Y, Chen F, Lu J, Wang W, Peatman E, Xu P, Somridhivej B, Baoprasertkul P, Quilang J, Sha Z, Bao B, Wang Y, Wang Q, Takano T, Nandi S, Liu S, Wong L, Kaltenboeck L, Quiniou S, Bengten E, Miller N, Trant J, Rokhsar D, Liu Z. Catfish Genome Consortium: Assembly of 500,000 inter-specific catfish expressed sequence tags and large scale gene-associated marker


- development for whole genome association studies. *Genome Biology*. 2010; 11:R8. [PubMed: 20096101]
93. Wang R, Sun L, Bao L, Zhang J, Jiang Y, Yao J, Song L, Feng J, Liu S, Liu Z. Bulk segregant RNA-seq reveals expression and positional candidate genes and allele-specific expression for disease resistance against enteric septicemia of catfish. *BMC Genomics*. 2013; 14:929. [PubMed: 24373586]
  94. Li C, Zhang Y, Wang R, Lu J, Nandi S, Mohanty S, Terhune J, Liu Z, Peatman E. RNA-Seq analysis of mucosal immune responses reveals signatures of intestinal barrier disruption and pathogen entry following *Edwardsiella ictaluri* infection in channel catfish, *Ictalurus punctatus*. *Fish & Shellfish Immunology*. 2012; 32:816–827. [PubMed: 22366064]
  95. Sun F, Peatman E, Li C, Liu S, Jiang Y, Zhou Z, Liu Z. Transcriptomic signatures of attachment, NF- $\kappa$ B suppression and IFN stimulation in the catfish gill following columnaris bacterial infection. *Developmental and Comparative Immunology*. 2012; 38:169–180. [PubMed: 22669032]
  96. Liu S, Zhang Y, Zhou Z, Waldbieser G, Sun F, Lu J, Zhang J, Jiang Y, Zhang H, Wang X, Rajendran KV, Kucuktas H, Peatman E, Liu Z. Efficient assembly and annotation of the transcriptome of catfish by RNA-Seq analysis of a doubled haploid homozygote. *BMC Genomics*. 2012; 13:595. [PubMed: 23127152]
  97. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic acids research*. 2004; 32:1792–1797. [PubMed: 15034147]
  98. Darriba D, Taboada GL, Doallo R, Posada D. ProtTest 3: fast selection of best-fit models of protein evolution. *Bioinformatics (Oxford, England)*. 2011; 27:1164–1165.
  99. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. *Molecular Biology and Evolution*. 2011; 28:2731–2739. [PubMed: 21546353]
  100. Rose TM, Schultz ER, Henikoff JG, Pietrokovski S, McCallum CM, Henikoff S. Consensus-degenerate hybrid oligonucleotide primers for amplification of distantly related sequences. *Nucleic acids research*. 1998; 26:1628–1635. [PubMed: 9512532]
  101. Ninwichian P, Peatman E, Liu H, Kucuktas H, Somridhivej B, Liu S, Li P, Jiang Y, Sha Z, Kaltenboeck L, Abernathy JW, Wang W, Chen F, Lee Y, Wong L, Wang S, Lu J, Liu Z. Second-generation genetic linkage map of catfish and its integration with the BAC-based physical map. G3 (Bethesda, Md). 2012; 2:1233–1241.
  102. Ju Z, Karsi A, Kocabas A, Patterson A, Li P, Cao D, Dunham R, Liu Z. Transcriptome analysis of channel catfish (*Ictalurus punctatus*): genes and expression profile from the brain. *Gene*. 2000; 261:373–382. [PubMed: 11167026]
  103. Cao D, Kocabas A, Ju Z, Karsi A, Li P, Patterson A, Liu Z. Transcriptome of channel catfish (*Ictalurus punctatus*): initial analysis of genes and expression profiles of the head kidney. *Animal genetics*. 2001; 32:169–188. [PubMed: 11531695]
  104. Karsi A, Patterson A, Feng J, Liu Z. Translational machinery of channel catfish: I. A transcriptomic approach to the analysis of 32 40S ribosomal protein genes and their expression. *Gene*. 2002; 291:177–186. [PubMed: 12095691]
  105. Kocabas AM, Li P, Cao D, Karsi A, He C, Patterson A, Ju Z, Dunham R, Liu Z. Expression profile of the channel catfish spleen: analysis of genes involved in immune functions. *Marine biotechnology (New York, N.Y.)*. 2002; 4:526–536.
  106. Li P, Peatman E, Wang S, Feng J, He C, Baoprasertkul P, Xu P, Kucuktas H, Nandi S, Somridhivej B, Serapion J, Simmons M, Turan C, Liu L, Muir W, Dunham R, Brady Y, Grizzle J, Liu Z. Towards the ictalurid catfish transcriptome: generation and analysis of 31,215 catfish ESTs. *BMC genomics*. 2007; 8:177. [PubMed: 17577415]
  107. Wang S, Peatman E, Abernathy J, Waldbieser G, Lindquist E, Richardson P, Lucas S, Wang M, Li P, Thimmapuram J, Liu L, Vullaganti D, Kucuktas H, Murdock C, Small BC, Wilson M, Liu H, Jiang Y, Lee Y, Chen F, Lu J, Wang W, Xu P, Somridhivej B, Baoprasertkul P, Quilang J, Sha Z, Bao B, Wang Y, Wang Q, Takano T, Nandi S, Liu S, Wong L, Kaltenboeck L, Quiniou S, Bengten E, Miller N, Trant J, Rokhsar D, Liu Z. Assembly of 500,000 inter-specific catfish expressed sequence tags and large scale gene-associated marker development for whole genome association studies. *Genome biology*. 2010; 11:R8. [PubMed: 20096101]

108. Oleksiak MF, Wu S, Parker C, Karchner SI, Stegeman JJ, Zeldin DC. Identification, functional characterization, and regulation of a new cytochrome P450 subfamily, the CYP2Ns. *J. Biol. Chem.* 2000; 275:2312–2321. [PubMed: 10644680]
109. Oleksiak MF, Wu S, Parker C, Qu W, Cox R, Zeldin DC, Stegeman JJ. Identification and regulation of a new vertebrate cytochrome P450 subfamily, the CYP2Ps, and functional characterization of CYP2P3, a conserved arachidonic acid epoxygenase/19-hydroxylase. *Arch. Biochem. Biophys.* 2003; 411:223–234. [PubMed: 12623071]
110. Mosadeghi S, Furnes B, Matsuo AY, Schlenk D. Expression and characterization of cytochrome P450 2X1 in channel catfish (*Ictalurus punctatus*). *Biochim. Biophys. Acta.* 2007; 1770:1045–1052. [PubMed: 17459590]
111. Cheng JB, Levine MA, Bell NH, Mangelsdorf DJ, Russell DW. Genetic evidence that the human CYP2R1 enzyme is a key vitamin D 25-hydroxylase. *Proc. Natl. Acad. Sci. USA.* 2004; 101:7711–7715. [PubMed: 15128933]
112. Chuang SS, Helvig C, Taimi M, Ramshaw H. a, Collop AH, Amad M, White J. a, Petkovich M, Jones G, Korczak B. CYP2U1, a novel human thymus- and brain-specific cytochrome P450, catalyzes omega- and (omega-1)-hydroxylation of fatty acids. *The Journal of biological chemistry.* 2004; 279:6305–6314. [PubMed: 14660610]
113. Jung J-H, Kim M, Yim UH, Ha SY, An JG, Won JH, Han GM, Kim NS, Addison RF, Shim WJ. Biomarker responses in pelagic and benthic fish over 1 year following the Hebei Spirit oil spill (Taean, Korea). *Marine pollution bulletin.* 2011; 62:1859–1866. [PubMed: 21658730]
114. Brammell BF, McClain JS, Oris JT, Price DJ, Birge WJ, Elskus A. CYP1A expression in caged rainbow trout discriminates among sites with various degrees of polychlorinated biphenyl contamination. *Archives of environmental contamination and toxicology.* 2010; 58:772–782. [PubMed: 19662324]
115. Bugiak B, Weber LP. Hepatic and vascular mRNA expression in adult zebrafish (*Danio rerio*) following exposure to benzo-a-pyrene and 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin. *Aquatic toxicology (Amsterdam, Netherlands).* 2009; 95:299–306.
116. Russell DW, Halford RW, Ramirez DMO, Shah R, Kotti T. Cholesterol 24-hydroxylase: an enzyme of cholesterol turnover in the brain. *Annual review of biochemistry.* 2009; 78:1017–1040.
117. Leoni V, Caccia C. 24S-hydroxycholesterol in plasma: a marker of cholesterol turnover in neurodegenerative diseases. *Biochimie.* 2013; 95:595–612. [PubMed: 23041502]
118. Oshlack A, Robinson MD, Young MD. From RNA-seq reads to differential expression results. *Genome Biology.* 2010; 11:220. [PubMed: 21176179]
119. Chaluvadi MR, Kinloch RD, Nyagode BA, Richardson TA, Raynor MJ, Sherman M, Antonovic L, Strobel HW, Dillehay DL, Morgan ET. Regulation of Hepatic Cytochrome P450 Expression in Mice with Intestinal or Systemic Infections of *Citrobacter rodentium*. *Drug Metabolism and Disposition.* 2009; 37:366–374.




Figure 1.


Phylogenetic relationship of channel catfish *CYP* genes in clan 2. (A) Phylogenetic relationship of channel catfish *CYP2* subfamily. MUSCLE alignments of all the amino acid sequences were used to generate a phylogenetic tree using Maximum likelihood method. The tree was shown with 10,000 replicates from the bootstrap test and the percentage of bootstrap values were given next to the branches. (B) Phylogenetic relationship of channel catfish *CYP1*, *CYP7* and *CYP21* subfamilies. MUSCLE alignments of all the amino acid sequences were used to generate a phylogenetic tree using Maximum likelihood method. The tree was shown with 10,000 replicates from the bootstrap test and the percentage of bootstrap values were given next to the branches.



**Figure 2.** Phylogenetic relationship of channel catfish *CYP* genes in clan 3. MUSCLE alignments of all the amino acid sequences were used to generate a phylogenetic tree using Maximum likelihood method. The tree was shown with 10,000 replicates from the bootstrap test and the percentage of bootstrap values were given next to the branches.




**Figure 3.** Phylogenetic relationship of channel catfish *CYP* genes in clan 4. MUSCLE alignments of all the amino acid sequences were used to generate a phylogenetic tree using Maximum likelihood method. The tree was shown with 10,000 replicates from the bootstrap test and the percentage of bootstrap values were given next to the branches.



**Figure 4.**


Phylogenetic relationship of channel catfish *CYP* genes in clan 7. MUSCLE alignments of all the amino acid sequences were used to generate a phylogenetic tree using Maximum likelihood method. The tree was shown with 10,000 replicates from the bootstrap test and the percentage of bootstrap values were given next to the branches.






**Figure 5.**

Phylogenetic relationship of channel catfish *CYP* genes in clan 19. MUSCLE alignments of all the amino acid sequences were used to generate a phylogenetic tree using Maximum likelihood method. The tree was shown with 10,000 replicates from the bootstrap test and the percentage of bootstrap values were given next to the branches.




**Figure 6.**

Phylogenetic relationship of channel catfish *CYP* genes in clan 20. MUSCLE alignments of all the amino acid sequences were used to generate a phylogenetic tree using Maximum likelihood method. The tree was shown with 10,000 replicates from the bootstrap test and the percentage of bootstrap values were given next to the branches.




**Figure 7.**


Phylogenetic relationship of channel catfish *CYP* genes in clan 26. MUSCLE alignments of all the amino acid sequences were used to generate a phylogenetic tree using Maximum likelihood method. The tree was shown with 10,000 replicates from the bootstrap test and the percentage of bootstrap values were given next to the branches.




**Figure 8.** Phylogenetic relationship of channel catfish *CYP* genes in clan 51. MUSCLE alignments of all the amino acid sequences were used to generate a phylogenetic tree using Maximum likelihood method. The tree was shown with 10,000 replicates from the bootstrap test and the percentage of bootstrap values were given next to the branches.




**Figure 9.** Phylogenetic relationship of channel catfish *CYP* genes in mitochondria clan. MUSCLE alignments of all the amino acid sequences were used to generate a phylogenetic tree using Maximum likelihood method. The tree was shown with 10,000 replicates from the bootstrap test and the percentage of bootstrap values were given next to the branches.



**Figure 10.**  
Syntenic analysis of subfamilies *CYP2AA*, *CYP2AD*, *CYP2K*, *CYP2X* and *CYP2Y*




**Figure 11.**  
Syntenic analysis of subfamilies *CYP3A* and *CYP3C*




**Figure 12.**  
Syntenic analysis of subfamilies *CYP4F* and *CYP4T*






**Figure 13.**  
Syntenic analysis of subfamily *CYP8A*



**Figure 14.**  
Syntenic analysis of subfamilies *CYP27A*, *CYP27B* and *CYP27C*.



**Figure 15.**  
Syntenic analysis of *CYP46*.

**Table 1**Degenerate primers used for *CYP46A1* amplification

primer name	forward primer	reverse primer
primer 1	agagggcccagacggarrtngayga	cggggtcccaggagaanggrwarta
primer 2	gaccttctcatcgccggncangarac	agccgcaggggtccttnamacytg

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Table 2**  
**Cytochrome P450 genes of channel catfish**

Catfish CYPs	Transcripts or genome	Sequence integrity	Accession number of NCBI	Linkage group	Reference
CYP1A1	both	complete	JT412024.1	not available	this study
CYP1B1	both	complete	AAAY90143.1	6	[92]
CYP1C1	both	complete	JT408041.1	16	this study
CYP2K8	both	complete	JT406642.1	3	this study
CYP2K17	both	complete	JT415766.1	3	this study
CYP2M1	both	complete	JT414187.1	22	this study
CYP2R1	both	complete	JT416283.1	21	this study
CYP2U1	both	complete	JT416818.1	21	this study
CYP2X1	both	complete	AAG30296.1	9	[93]
CYP2X20	genome	partial	KF531904	9	this study
CYP2X21	genome	complete	KF531901	9	this study
CYP2X22	genome	partial	KF531905	14	this study
CYP2Y7	genome	partial	KF531905	3	this study
CYP2Y8	genome	partial	KF531907	3	this study
CYP2AA14	both	complete	JT411919.1	11	this study
CYP2AD6	genome	partial	KF531902	28	this study
CYP2AD8	genome	partial	KF531903	28	this study
CYP2AD9	both	complete	JT418663.1	28	this study
CYP2AD10	genome	partial	KF531908	28	this study
CYP3A125	genome	partial	KF531909	20	this study
CYP3A126	both	complete	JT418825.1	20	this study
CYP3A127	genome	partial	KF531910	20	this study
CYP3A128	genome	partial	KF531911	20	this study
CYP3C5	both	complete	JT408739.1	28	this study
CYP4F60	both	complete	JT479785.1	5	this study
CYP4F66	genome	Partial	KF531912	5	this study
CYP4T16	genome	partial	KF531913	19	this study
CYP4T17	both	partial	JT320237.1	19	this study
CYP4V7	both	complete	JT408109.1	9	this study
CYP4V8	both	complete	JT414161.1	29	this study
CYP5A1	both	complete	JT316268.1	4	this study
CYP7A1	both	complete	JT408390.1	1	this study
CYP7C1	both	complete	JT411121.1	11	this study
CYP7D1	both	complete	JT415644.1	18	this study
CYP8A1a	both	complete	JT411628.1	3	this study
CYP8A1b	both	partial	JT470032.1	7	this study

<b>Catfish CYPs</b>	<b>Transcripts or genome</b>	<b>Sequence integrity</b>	<b>Accession number of NCBI</b>	<b>Linkage group</b>	<b>Reference</b>
<i>CYP8B5</i>	both	complete	JT323293.1	28	this study
<i>CYP8B6</i>	genome	partial	KF531914	8	this study
<i>CYP8B7</i>	genome	partial	KF531915	20	this study
<i>CYP11A1</i>	both	complete	NP_001187241.1	14	unpublished
<i>CYP11C1</i>	both	complete	JT399700.1	not available	this study
<i>CYP17A1</i>	both	complete	NP_001187242.1	6	unpublished
<i>CYP17A2</i>	both	partial	JT364637.1	11	this study
<i>CYP19A1</i>	both	complete	Q92111.1	6	[94]
<i>CYP19A2</i>	both	complete	AF417239.1	18	[95]
<i>CYP20A1</i>	both	complete	JT411689.1	13	this study
<i>CYP21A1</i>	both	complete	JT407837.1	1	this study
<i>CYP24A1</i>	genome	partial	KF531916	7	this study
<i>CYP26A1</i>	both	complete	JT413111.1	5	this study
<i>CYP26B1</i>	both	complete	JT416269.1	23	this study
<i>CYP26C1</i>	both	complete	JT411012.1	7	this study
<i>CYP27A2</i>	both	complete	JT244274.1	13	this study
<i>CYP27A3</i>	genome	partial	KF531917	not available	this study
<i>CYP27A14</i>	both	complete	JT406399.1	17	this study
<i>CYP27A15</i>	genome	partial	KF531918	17	this study
<i>CYP27A16</i>	genome	partial	KF531919	17	this study
<i>CYP27A17</i>	genome	partial	KF531920	17	this study
<i>CYP27A18</i>	genome	partial	KF531921	17	this study
<i>CYP27B1</i>	both	complete	JT483829.1	28	this study
<i>CYP27C1</i>	both	complete	JT408218.1	13	this study
<i>CYP51A1</i>	both	complete	JT418451.1	28	this study

**Table 3**

Gene number variation of *CYPs* among human, chicken, frogs, Japanese pufferfish, zebrafish and catfish in diverse subfamilies. Shading indicates lineage-specific *CYP* subfamilies.

CYP450 subfamily	gene # Human	gene # Mouse	gene # Chicken	gene # Western clawed frog	gene # African clawed frog	gene # Japanese pufferfish	gene # zebrafish	gene # catfish
1A	2	2	2	1	2	1	1	1
1B	1	1	1	1	1	1	1	1
1C			1	1	1	2	2	1
1D				1	1		1	
2A	3	4						
2B	1	5						
2C	4	15	1					
2D	1	9	1	7	5			
2E	1	1						
2F	1	1				1		
2G		1*						
2H			2					
2I								
2J	1	7	5				1	
2K						3	11	2
2M								1
2N						3	1	
2P						1	6	
2Q				8	5			
2R	1	1	1	1	1	1	1	1
2S	1	1						
2T		1						
2U	1	1	1	1	1	1	1	1
2V							1	
2W	1	1	2					
2X						2	7	4
2Y						2	2	2
2Z						2		
2AA							12	1
2AB		1	5	2	3			
2AC			3	12	6			
2AD						1	3	4
2AE							2	
2AM				6	4			
2AN				5	5			
2AP				1				
2AQ				2	2			

CYP450 subfamily	gene # Human	gene # Mouse	gene # Chicken	gene # Western clawed frog	gene # African clawed frog	gene # Japanese pufferfish	gene # zebrafish	gene # catfish
2AR				1				
2AS				1				
2AT				1	1			
3A	4	8	2	6	4	3	1	4
3B						2		
3C							4	1
3D						1		
4A	2	7						
4B	1	1	1					
4F	6	9	1	3	2	1	1	2
4T				4	10	1	1	2
4V	1	1	1	2	2	1	2	2
4X	1	1						
4Z	1							
5A	1	1	1	1	2	1	1	1
7A	1	1	1	1		1	1	1
7B	1	1	1	1				1
7C						1	1	1
7D							1	1
8A	1	1		1		2	1	2
8B	1	1	1	1	1	2	1	3
11A	1	1	1	1	1	1	2	1
11B	2	2		1	1			
11C						1	1	1
16A						1		
17A	1	1	1	1	1	4	2	2
19A	1	1	1	1	1	2	2	2
20A	1	1	1	1	1	1	1	1
21A	1	1	1	1	1	1	1	1
24A	1	1	1			1	1	1
26A	1	1	1	1	2	1	1	1
26B	1	1	1	1	1	1	1	1
26C	1	1	1	1	1	1	1	1
27A	1	1	1	4	3	5	5	7
27B	1	1		1		1	1	1
27C	1		1	1	2	1	1	1
39A	1	1	1	1	1		1	
46A	1	1	1	6	3	1	4	
51A	1	1	1	1	2	1	1	1
Total	57	100	48	95	80	61	94	61

\* CYP2G subfamily is not mammal specific, because it is also found in lizard (belong to tetrapod).



**Table 4**

Differentially expressed CYPs in channel catfish following *Edwardsiella ictaluri* and *Flavobacterium columnare* infection. Bold values indicate time-points where CYP was significantly changed relative to the control and absolute fold change was larger than two. Dash indicates all values of time-points in the experiment were out of threshold (threshold: p-value < 0.05, reads number per gene 5 and fold change 2).

CYP gene	ESC liver		ESC intestine			Columnaris gill			
	3d	14d	3h	24h	3d	4h	24h	48h	
CYP1C1	<b>16.84</b>	1.81	-	-	-	-	-	-	
CYP2K17	-	-	<b>2.26</b>	1.07	1.27	-	-	-	
CYP2M1	<b>-2.96</b>	1.20	-	-	-	-	-	-	
CYP2R1	<b>-6.80</b>	1.10	1.20	<b>-2.50</b>	-1.93	-	-	-	
CYP2X1	-1.07	<b>-2.31</b>	<b>-5.65</b>	<b>-3.76</b>	<b>-4.66</b>	-	-	-	
CYP2X20	<b>-14.78</b>	-1.85	-	-	-	-1.43	<b>-2.73</b>	<b>-2.57</b>	
CYP2X21	<b>-31.69</b>	<b>-2.01</b>	-	-	-	-	-	-	
CYP2X22	<b>-52.52</b>	1.08	-	-	-	-	-	-	
CYP2Y8	<b>-2.56</b>	-1.00	-	-	-	-	-	-	
CYP2AA14	-	-	-1.26	-1.90	<b>-2.23</b>	-	-	-	
CYP2AD6	-	-	-	-	-	1.83	1.07	<b>2.88</b>	
CYP2AD8	<b>2.32</b>	-1.02	-	-	-	-	-	-	
CYP3A128	<b>3.26</b>	<b>2.03</b>	-	-	-	-	-	-	
CYP3C5	<b>5.93</b>	1.58	-	-	-	-	-	-	
CYP4F60	<b>5.29</b>	<b>7.98</b>	-	-	-	-	-	-	
CYP4F66	<b>2.17</b>	1.71	-	-	-	-	-	-	
CYP4T16	<b>2.41</b>	-1.44	<b>2.75</b>	<b>2.76</b>	<b>2.58</b>	1.65	1.98	<b>3.25</b>	
CYP4V7	<b>-32.48</b>	-1.06	-	-	-	-	-	-	
CYP7A1	<b>-11.72</b>	-1.78	-	-	-	-	-	-	
CYP7D1	<b>10.19</b>	<b>2.09</b>	-	-	-	-	-	-	
CYP8A1a	<b>27.60</b>	<b>2.29</b>	-	-	-	-	-	-	
CYP8A1b	<b>30.23</b>	1.67	-	-	-	-	-	-	
CYP8B5	<b>2.05</b>	1.41	-	-	-	-	-	-	
CYP8B6	<b>2.14</b>	1.39	-	-	-	-	-	-	
CYP8B7	<b>-4.68</b>	<b>-6.30</b>	-	-	-	-	-	-	
CYP11A1	<b>2.72</b>	1.19	-	-	-	-	-	-	
CYP17A1	-	-	-	-	-	-1.25	<b>-3.74</b>	<b>-3.39</b>	
CYP19A1	-	-	-	-	-	1.88	<b>2.06</b>	1.86	
CYP21A1	<b>3.35</b>	-1.04	-	-	-	-	-	-	
CYP24A1	<b>-6.78</b>	<b>-2.15</b>	-	-	-	-	-	-	
CYP26A1	<b>-2.87</b>	-1.09	1.16	<b>-2.67</b>	-1.24	-1.29	<b>-5.92</b>	<b>-2.90</b>	
CYP27A2	<b>2.17</b>	<b>2.01</b>	-	-	-	-	-	-	
CYP27A3	<b>-6.63</b>	1.06	-	-	-	-	-	-	

<i>CYP</i> gene	ESC liver		ESC intestine			Columnaris gill			
	3d	14d	3h	24h	3d	4h	24h	48h	
<i>CYP27A14</i>	<b>-2.59</b>	-1.00	-	-	-	-	-	-	
<i>CYP27A17</i>	-	-	-	-	-	<b>-3.93</b>	-1.08	-1.31	
<i>CYP51A1</i>	<b>7.28</b>	<b>2.96</b>	-1.21	<b>-2.33</b>	<b>-2.27</b>	1.22	<b>-2.18</b>	<b>-2.13</b>	

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript