

The Arabidopsis RNA-Binding Protein AtRGGA Regulates Tolerance to Salt and Drought Stress^{1[OPEN]}

Alfredo Ambrosone^{2,3}, Giorgia Batelli², Roberta Nurcato, Vincenzo Aurilia, Paola Punzo, Dhinoth Kumar Bangarusamy⁴, Ida Ruberti, Massimiliano Sassi⁵, Antonietta Leone, Antonello Costa, and Stefania Grillo*

National Research Council of Italy, Institute of Biosciences and Bioresources, Research Division Portici, 80055 Portici (Naples), Italy (A.A., G.B., R.N., P.P., A.C., S.G.); National Research Council of Italy, Institute for Mediterranean Agriculture and Forest Systems, 80056 Ercolano (Naples), Italy (V.A.); Bio-science Core Laboratories, King Abdullah University of Science and Technology, Thuwal 23955–6900, Kingdom of Saudi Arabia (D.K.B.); National Research Council of Italy, Institute of Molecular Biology and Pathology, 00185 Rome, Italy (I.R., M.S.); and Department of Pharmacy, University of Salerno, 84084 Fisciano (Salerno), Italy (A.L.)

Salt and drought stress severely reduce plant growth and crop productivity worldwide. The identification of genes underlying stress response and tolerance is the subject of intense research in plant biology. Through microarray analyses, we previously identified in potato (*Solanum tuberosum*) *StRGGA*, coding for an Arginine Glycine Glycine (RGG) box-containing RNA-binding protein, whose expression was specifically induced in potato cell cultures gradually exposed to osmotic stress. Here, we show that the Arabidopsis (*Arabidopsis thaliana*) ortholog, *AtRGGA*, is a functional RNA-binding protein required for a proper response to osmotic stress. *AtRGGA* gene expression was up-regulated in seedlings after long-term exposure to abscisic acid (ABA) and polyethylene glycol, while treatments with NaCl resulted in *AtRGGA* down-regulation. *AtRGGA* promoter analysis showed activity in several tissues, including stomata, the organs controlling transpiration. Fusion of *AtRGGA* with yellow fluorescent protein indicated that *AtRGGA* is localized in the cytoplasm and the cytoplasmic perinuclear region. In addition, the *rgga* knockout mutant was hypersensitive to ABA in root growth and survival tests and to salt stress during germination and at the vegetative stage. *AtRGGA*-overexpressing plants showed higher tolerance to ABA and salt stress on plates and in soil, accumulating lower levels of proline when exposed to drought stress. Finally, a global analysis of gene expression revealed extensive alterations in the transcriptome under salt stress, including several genes such as *ASCORBATE PEROXIDASE2*, *GLUTATHIONE S-TRANSFERASE TAU9*, and several *SMALL AUXIN UPREGULATED RNA*-like genes showing opposite expression behavior in transgenic and knockout plants. Taken together, our results reveal an important role of *AtRGGA* in the mechanisms of plant response and adaptation to stress.

Abiotic stresses such as salinity and drought account for extensive reductions in yields of agricultural crops. While salt stress has an ionic component specifically brought about by Na⁺ toxicity, both drought and salinity challenge plants by imposing osmotic stress, caused by a reduction in soil water potential (Maggio et al., 2006). As

a result of osmotic stress, a complex response aimed at limiting cellular damages and rescuing a new homeostasis is elicited in plants, which includes a coordination of biochemical and physiological changes, including stomata closure, cell growth alterations, photosynthesis inhibition, flowering time and root architecture modification, and inhibition of seed germination (Zhu, 2002). A key role in the regulation of these processes is played by the hormone abscisic acid (ABA). A major breakthrough in our understanding of the osmotic stress responses has come with the recent identification of the ABA PYRABACTIN RESISTANCE (PYR)/PYRABACTIN RESISTANCE-LIKE1 (PYR1)/REGULATORY COMPONENT OF ABA RECEPTOR (RCAR) receptors and the elucidation of their mechanism of action in ABA-mediated signaling cascades (Fujii et al., 2009; Ma et al., 2009; Park et al., 2009).

Osmotic stress induces an increase in ABA concentration, perceived by the PYR/PYL/RCAR receptors. When ABA is ligated, a conformational change is induced in members of the PYR/PYL/RCAR family of receptors, which become able to bind and inhibit type 2C protein phosphatases, thus releasing SUCROSE NON-FERMENTING1-RELATED PROTEIN KINASE2.2 (SnRK2.2), SnRK2.3, and

¹ This work was supported by the Italian Ministry of University and Research (Fondo per gli Investimenti della Ricerca di Base Project PlantSTRESS and GenoPOM-PRO project no. PON02_00395_3082360).

² These authors contributed equally to the article.

³ Present address: National Research Council of Italy, Istituto di Cibernetica E. Caianiello, Via Campi Flegrei 34, 80078 Pozzuoli, Italy.

⁴ Present address: SciGenom Laboratories, Cochin, Kerala 682037, India.

⁵ Present address: Laboratoire de Reproduction et Développement des Plantes, École Normale Supérieure de Lyon 46, Allée d'Italie, 69364 Lyon cedex 07, France.

* Address correspondence to grillo@unina.it.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantphysiol.org) is: Stefania Grillo (grillo@unina.it).

^[OPEN] Articles can be viewed without a subscription.

www.plantphysiol.org/cgi/doi/10.1104/pp.114.255802

SnRK2.6 kinases from inhibition. In turn, ABA-activated SnRK2s phosphorylate ABA-responsive element-binding transcription factors such as ABF2 to induce the up-regulation of ABA-responsive genes and, in guard cells, plasma membrane-located ion channels such as SLOW ANION CHANNEL1 and POTASSIUM CHANNEL IN *ARABIDOPSIS THALIANA*1 potassium channel to promote and maintain stomata closure (Geiger et al., 2009, 2010; Sato et al., 2009; Hubbard et al., 2010; Klingler et al., 2010). ABA-independent pathways, which may involve ABA-unresponsive members of the SnRK2 family (Fujii and Zhu, 2012), also participate in osmotic stress responses and seem to interact and converge with ABA-mediated pathways (Ishitani et al., 1997).

While outstanding progress has been made in the elucidation of perception and signaling cascades resulting in stress-induced modifications of gene expression and channel activation, RNA regulatory mechanisms such as synthesis, processing, transport, translation, storage, stability, and degradation of RNA molecules are emerging as key processes participating in the modulation of cellular responses to stress (Ambrosone et al., 2012; Nakaminami et al., 2012). The importance of mRNA stability mechanisms is known in the case of the Na⁺/H⁺ antiporter SALT OVERLY SENSITIVE1 (SOS1) required for tolerance to salt stress. SOS1 mRNA is highly unstable in control conditions, but within 10 min after the imposition of salt stress, SOS1 transcript is stabilized in a process mediated by reactive oxygen species (Shi et al., 2003; Chung et al., 2008). A strong impact of stress on translation efficiency has been shown in the case of hypoxia, where Branco-Price et al. (2008) demonstrated that a mechanism of selective mRNA translation without reduction of transcription coordinated metabolic adjustments to oxygen deprivation. Recently, it was shown that the RNA-binding protein (RBP) OLIGOURIDYLATE BINDING PROTEIN1 participates in the selective mRNA translation mechanism during hypoxia by sequestering mRNAs in stress granules. Upon reoxygenation, stress granules dissolve and mRNAs return to actively translating polysomes (Sorenson and Bailey-Serres, 2014).

The regulation of RNA metabolism directly or indirectly involves RBPs, which are distinguished based on the presence and organization of several different functional motifs and domains, with the RNA recognition motif (RRM) and K homology domain being the most common in plants (Lorković, 2009). Other domains and motifs include the Tudor SN domain, Arg repeats, glycine-rich domain (GR), zinc finger domain (Burd and Dreyfuss, 1994; Albà and Pagès, 1998; Lorković and Barta, 2002), Arg/Gly motif, and cold shock domain (Nakaminami et al., 2012; Ambrosone et al., 2013). Several RBPs have recently been shown to be involved in plant development and stress responses. Tudor SN (TSN) proteins are RBPs involved in RNA stability control upon salt stress. Double mutants *tsn1/tsn2* showed a drastic reduction in germination, growth, survival, and fitness under high-salinity stress (dit Frey et al., 2010). RBPs were also shown to be involved in response to heat and cold stress. REGULATOR OF C-REPEAT/DEHYDRATION-

RESPONSIVE ELEMENT BINDING FACTOR GENE EXPRESSION3 is a K homology domain-containing RBP that was shown to be a negative regulator of heat stress response by repressing the expression of several heat stress factors, such as *HSA1a*, *HSA1b*, and *HSA1d* (Guan et al., 2013). The zinc finger-containing glycine-rich RBP AtRZ-1A is induced by cold, and, when overexpressed, increases freezing tolerance in Arabidopsis (*Arabidopsis thaliana*; Kim et al., 2005). Manipulation of the expression of a GR- and RRM domain-containing protein, AtGRP7, impacts stress tolerance under high salinity, drought, or cold stress. Overexpression of AtGRP7 increased freezing tolerance but also caused delayed germination and seedling growth under salt or dehydration stress (Kim et al., 2008).


In an effort to identify genes important for adaptation to osmotic stress, we isolated *StRGGA*, a gene encoding a putative RBP, whose expression was induced in potato (*Solanum tuberosum*) cell cultures gradually adapted to high concentrations of polyethylene glycol (PEG), while no change in *StRGGA* expression was observed when cells were shocked with PEG (Ambrosone et al., 2011). Here, we present the characterization of the putative *RGGA* ortholog in Arabidopsis. *AtRGGA* is expressed in several tissues, including stomata, and transcript abundance is increased in cells and plants exposed to PEG and ABA. *AtRGGA* encodes a cytosolic protein capable of binding RNA in vitro. Transgenic plants overexpressing *AtRGGA* are more tolerant to ABA, drought, and salt stress, whereas *rgga* mutant plants are more sensitive to ABA and osmotic stresses. Manipulation of *AtRGGA* expression has a severe impact on whole-gene expression, indicating that *AtRGGA* has an important functional role in planta.

RESULTS

AtRGGA Gene Expression in Response to Stress and ABA Treatments

In a previous study, we isolated *StRGGA* (GenBank accession no. FM209282), whose gene expression is specifically induced in culture cells of potato gradually adapted to high concentrations of PEG. Exposure of potato cells to abrupt osmotic stress did not elicit a change in *StRGGA* transcript abundance (Ambrosone et al., 2011; Fig. 1A; Supplemental Table S1). The deduced protein sequence of *StRGGA* shares 63% sequence homology with the protein encoded by the locus *At4g16830* of Arabidopsis (Supplemental Fig. S1), which, therefore, is hypothesized to be the Arabidopsis ortholog (*AtRGGA*). To investigate whether *AtRGGA* was also induced by stress treatments in Arabidopsis, we analyzed gene expression in cells and seedlings exposed to NaCl and osmotic (PEG) stress. ABA treatments were also included to assess a possible involvement of the hormone in the regulation of *AtRGGA* transcript abundance. In MM2D cells (Menges and Murray, 2002), NaCl, ABA, and PEG treatments induced a significant up-regulation of *AtRGGA* compared with control untreated cells (Fig. 1B).

Figure 1. Expression analysis of *RGGA* in potato and Arabidopsis. A, Expression of *RGGA* in cells of potato in control conditions and after gradual (PEG-adapted) or abrupt (PEG-shocked) exposure to PEG. B, Gene expression of *RGGA* in Arabidopsis MM2D cells exposed for 24 h to NaCl (150 mM), ABA (50 μ M), or 10% (w/v) PEG. C, *AtRGGA* expression in 14-d-old seedlings of Arabidopsis treated for 24 h with different concentrations of NaCl as indicated. D, *AtRGGA* expression in Arabidopsis seedlings after 48 h of exposure to 35% (w/v) PEG, NaCl (120 mM), or ABA (10 μ M). Gene expression analyses were conducted by quantitative reverse transcription (qRT)-PCR.



Induction seemed to be highest after NaCl treatment, which, however, also caused a reduction in cell viability (Supplemental Fig. S2). In seedlings, 24-h treatments with different concentrations of NaCl caused a slight down-regulation of *AtRGGA* expression (Fig. 1C), while there was an up-regulation in seedlings exposed for 2 d to ABA and PEG (Fig. 1D), indicating that *AtRGGA* transcript abundance is reduced by salt stress in the short term but increased over longer periods of exposure to ABA and osmotic stress.

AtRGGA Binds RNA in Vitro

Protein sequence analysis of AtRGGA showed the presence of a Suppressor of Tom1 (Stm1) domain, found at the N-terminal region of the yeast (*Saccharomyces cerevisiae*) Stm1 nucleic acid-binding protein, and a Hyaluronan-Binding Protein4_Plasminogen Activator Inhibitor-1 mRNA-Binding Protein1 (HABP4_PAIRBP1) domain, found in RBPs, suggesting that AtRGGA could be an RBP (Fig. 2A). To verify this hypothesis, an RNA electromobility shift assay (EMSA) was performed using recombinant, His-tagged AtRGGA (His-RGGA) and total RNA extracted from control as well as salt-stressed whole seedlings. RNA was labeled with biotin and incubated with or without His-RGGA prior to electrophoresis in native conditions. A recombinant version of the PYR1 ABA receptor, His-PYR1, was used as a negative control. As shown in Figure 2B, an RNA mobility shift was specifically observed when RNA was incubated with AtRGGA, indicating that AtRGGA was capable of binding RNA, and the binding was competed by adding an excess of unlabeled RNA, thus showing that AtRGGA is a bona fide RBP. To assess the specificity of AtRGGA binding to RNA, poly(A⁺) and poly(A⁻) RNA fractions were used for RNA EMSA. A band shift after incubation

with His-RGGA was observed when poly(A⁻) RNA was used, indicating that RGGA binds to one or more RNAs contained in the poly(A⁻) RNA fraction.

AtRGGA Promoter Activity in Tissues and Protein Subcellular Localization

To gain insights into the function of *AtRGGA* in plants, we proceeded to analyze its expression pattern and subcellular localization. Transgenic plants expressing the GUS reporter gene driven by the putative promoter (defined as 2 kb upstream of the protein-coding sequence) of *AtRGGA* were stained using 5-bromo-4-chloro-3-indolyl glucuronide to visualize the spatial and temporal patterns of activity of the *AtRGGA* promoter. As shown in Figure 3, GUS activity was visualized both in seedlings and in adult plants in several organs, including leaves, roots, inflorescences, and siliques. Interestingly, within leaves, a strong staining of stomata was observed (Fig. 3D), indicating the expression of *AtRGGA* in guard cells. In reproductive organs, GUS activity was visualized in pollen grains (Fig. 3, G and H) and tubes of germinating pollen (Fig. 3H) as well as in funiculi attaching seeds to siliques (Fig. 3, M and N). Results gathered by GUS staining assays were generally consistent with publicly available expression data showing the presence of *AtRGGA* transcript in all analyzed tissues (Supplemental Fig. S3).

To analyze AtRGGA protein subcellular localization, we generated transgenic plants overexpressing a yellow fluorescent protein (YFP)-RGGA fusion protein. Young seedlings of YFP-RGGA were observed by confocal laser scanning microscopy following a short incubation in propidium iodide to counterstain cell walls. As shown in Figure 4A, in cells of the root apex, where vacuoles are less developed, a clear YFP signal was observed in the




Figure 2. A, Schematic representation of Arabidopsis *AtRGGA* protein domain organization. Gray boxes indicate the locations of the Stm1 N-terminal domain (Stm1; InterPro no. IPR019084) and the hyaluronan/mRNA-binding domain (HABP4_PA11_RBP1; InterPro no. IPR006861). B, EMSA of Arabidopsis RNA incubated with recombinant *AtRGGA* (His-RGGA). RNA was extracted from NaCl-treated (Salt Stress RNA) or untreated (Control RNA) plants and labeled with biotin. Unlabeled RNA (160-fold) was used as a competitor. Recombinant PYR1 (His-PYR1) served as a negative control. C, EMSA of Arabidopsis total, poly(A⁺), and poly(A⁻) RNA incubated without or with recombinant *AtRGGA* (His-RGGA). The brackets indicate labeled RNA, and the arrows indicate RGGA-bound RNA.

cytoplasm. Upon prolonged incubation in propidium iodide solution, minor staining of nuclear DNA could be achieved. In this case, we observed exclusion of YFP-RGGA from the nuclei in cells of the root elongation zone, while strong YFP fluorescence was visualized in the perinuclear region of the cytoplasm (Fig. 4B). In leaf tissues, YFP signal was especially strong in stomata (Fig. 4C). Similar results were obtained using transgenic plants expressing a C-terminal fusion of *AtRGGA* with YFP (Fig. 4D). Together, the protein subcellular localization studies indicate that *AtRGGA* localizes in the cytoplasm and the perinuclear region.


Functional Analysis of *AtRGGA*

To characterize the role of *AtRGGA* in plant responses to salt and drought stress, a transfer DNA (T-DNA)

insertion mutant in which *AtRGGA* gene expression is abolished (SALK_143514; *rgga*; Fig. 5A; Supplemental Fig. S4) was obtained from the Arabidopsis Biological Resource Center, and transgenic plants overexpressing a FLAG-RGGA fusion protein (Fig. 5B; Supplemental Fig. S4) were generated. In control conditions, *rgga* plants appeared to have larger rosettes compared with wild-type Columbia-0 (Col-0) plants and showed delayed flowering in a long-day (16 h of light/8 h of darkness) regime (Supplemental Fig. S4).

Phenotype analyses in the presence of stress were conducted at different developmental stages. At the germination stage, *rgga* displayed a higher sensitivity to NaCl compared with wild-type Col-0. In particular, while an average of 83% of Col-0 seeds presented fully expanded cotyledons after 7 d of exposure to NaCl (120 mM), only 64% of *rgga* seeds were germinated (Fig. 5C). In contrast, seeds of overexpressing plants (35S::FLAG-RGGA) did not

Figure 3. *AtRGGGA* promoter activity in tissues of *Arabidopsis*. GUS staining was performed in vegetative and reproductive tissues of transgenic *Arabidopsis* plants expressing the GUS reporter gene under the control of the *AtRGGGA* promoter. Five-day-old seedling (A), root (B), leaves (C), inflorescences (E), and siliques (L–N) were stained. Closeup views of stomata (D), anther (F), stigma (G), ovary (H), and ovule (I) are also shown.



show any significant differences in their ability to germinate in salt stress medium compared with the wild type or controls transformed with the empty vector (Fig. 5C).

Survival tests instead showed differences in the ability to withstand salt stress conditions of both mutant and overexpressing plants compared with controls. After 7 d of exposure to high-salt conditions, both the knockout mutant and overexpressing plants showed significantly different survival percentages from the wild type. More than 60% of seedlings of three different overexpressing lines did not display any signs of necrosis or bleaching, compared with 36% of Col-0 seedlings, while *rgga* seedlings were more sensitive to salt stress, with only 7% of plants surviving long-term exposure to NaCl (180 mM; Fig. 5D). Differences in sensitivity to salt stress also could be observed visually (Fig. 6A).

Closure of stomata when exposed to a dry environment is an important ABA-dependent mechanism contributing to plant stress tolerance, whose efficiency can be inferred by measuring the decline in fresh weight of detached leaves in a time course (Raschke, 1970; Verslues et al., 2006). Therefore, water-loss measurements during the course of 3 h were performed on both *rgga* mutant

and 35S::FLAG-RGGA plants. As shown in Figure 6B, wild-type Col-0 and *rgga* had lost a similar amount of water (62% and 61% of fresh weight, respectively), while leaves detached from overexpressing plants retained a higher amount of water, having lost 51% of their initial fresh weight, thus suggesting that an increased expression of *AtRGGGA* results in a more efficient closure of stomata in drought stress conditions.

To further assess the role of RGGGA in ABA-dependent mechanisms of response to environmental stresses, root growth and survival tests in the presence of ABA were performed. Root growth experiments on plates showed a hypersensitivity of *rgga* to the presence of ABA in the medium, while overexpressing plants did not display significant differences as compared with Col-0 (Fig. 6, C and D). In terms of survival of ABA exposure, 10 d of treatment could highlight significant differences between genotypes. 35S::FLAG-RGGA plants showed a higher ability to tolerate the presence of 50 μ M ABA in the medium, with 60% of plants still surviving after 10 d of exposure to the hormone, while the mutant only had about 20% of individuals still surviving and the wild type had about 47%.




Figure 4. *AtRGGA* protein localization in Arabidopsis. Confocal microscopy visualization was performed for transgenic Arabidopsis plants expressing a YFP-*RGGA* or *RGGA*-YFP (D) fusion protein. Propidium iodide staining, YFP fluorescence, and merged images of root apex (A), root elongation zone (B and D), and leaf epidermal cells (C) are shown.

Opposite phenotypes of mutant and transgenic plants in terms of survival of stress conditions were also observed in soil-grown plants. When drought or salt stress was imposed, the *rgga* mutant appeared more sensitive than the wild type, with a more obvious bleaching of leaves. In contrast, overexpressing plants showed less wilting and bleaching symptoms in the presence of salt and drought stress compared with Col-0 (Fig. 7A). Pro content after prolonged exposure to salt and drought stress was measured in the wild type, *rgga*, and 35S::FLAG-*RGGA*. As shown in Figure 7B, levels of Pro in control conditions were similar, with less than 0.2 mmol g^{-1} fresh weight for all the tested genotypes. After 7 d of exposure to drought or salt stress, there was a dramatic increase in Pro content, with levels quintupling in Col-0 and *rgga*. In *AtRGGA*-overexpressing plants, the increase in Pro was less sharp, particularly after drought stress, where plants from three

independent transgenic lines accumulated significantly less Pro (Fig. 7B). In salt stress, however, only one of the three tested lines was significantly different from wild-type or *rgga* levels (Fig. 7B).

Gene Expression Analyses in *rgga* and 35S::FLAG-*RGGA* Plants

To analyze the impact of a modified expression of *AtRGGA* on the Arabidopsis transcriptome, microarray analyses were performed on RNA extracted from *AtRGGA* knockout or overexpressing seedlings grown in control conditions or after exposure to salt stress. The results of microarray analysis were validated using quantitative PCR (qPCR; Supplemental Table S2). In general, an overall perturbation of the transcriptome was observed both in control and stress treatments, with large




Figure 5. Characterization of plants with modified expression of *AtRGGGA*. A, Identification of an *RGGA* knockout mutant. The top shows a representative model of the *At4g16830* locus encoding *RGGA* in Arabidopsis, showing the location of the T-DNA insertion in SALK_143514 (*rgga*). The bottom shows semiquantitative reverse transcription-PCR analysis to confirm that the expression of *At4g16830* is abolished in the *rgga* mutant. β -Actin was amplified as an internal standard. B, Immunoblot using α -FLAG antibody of total proteins extracted from Arabidopsis plants transformed to overexpress the fusion protein FLAG-RGGA (35S::FLAG-RGGA). Different transgenic lines (#10, #15, #18, and #20), along with the wild type (Col-0) and controls transformed with the empty binary plasmid (empty vector), are shown. Ponceau staining of Rubisco small subunit (RbcS) served as a loading control. C, Germination analysis of *AtRGGGA* knockout and transgenic plants in the presence of NaCl (120 mM). Germination was scored in terms of fully expanded cotyledons 7 d after stratification. Data reported are means \pm SD from three independent experiments. The asterisk denotes a significant difference between Col-0 and *rgga* ($P < 0.05$) according to Student's *t* test. D, Survival test of 18-d-old seedlings germinated on germination medium (GM; 4.3 g L⁻¹ MS salts, 30% [w/v] Suc, pH 5.7) and transferred to NaCl (180 mM) medium. Survival was scored daily in terms of absence of necrotic or bleached leaves. Data are means \pm SD of three independent experiments ($n = 30$). Asterisks denote statistically significant differences versus Col-0 assessed by χ^2 test (* $P < 0.05$, ** $P < 0.01$).

numbers of genes showing a variation in gene expression in *rgga* and 35S::FLAG-RGGA (Supplemental Tables S3–S10). An analysis using the Singular Enrichment Analysis tool of agriGO (Du et al., 2010) was performed to identify significantly enriched Gene Ontology (GO) terms in each treatment/genotype. In both mutant and overexpressing plants in control and stress conditions, several GO terms were significantly enriched either in down- and up-regulated genes, with the terms response to stimulus, regulation of cellular process, and biological regulation having among the lowest false discovery rate values in most of the conditions considered (Supplemental Figs. S5 and S6). Genes showing opposite behaviors in the mutant and overexpressing lines in terms of up- or down-regulation as compared with Col-0 are summarized for controls (Table I) and salt stress conditions (Table II). As expected, *AtRGGGA* (*At4g16830*) was present in both conditions as down-regulated in the knockout plants

and up-regulated in overexpressing plants. In control conditions, DEHYDRATION-RESPONSIVE ELEMENT-BINDING PROTEIN2A (DREB2A) and DREB19, key transcription factors involved in drought stress responses, and several heat shock proteins, including *HSP21* and *HSP22*, were down-regulated in *rgga* and had an opposite behavior in 35S::FLAG-RGGA (Table I). After stress treatment, several genes coding for SAUR-like auxin-responsive proteins were up-regulated in *rgga* and down-regulated in 35S::FLAG-RGGA (Table II). Among the genes down-regulated in *rgga* and up-regulated in 35S::FLAG-RGGA, we identified the reactive oxygen species scavengers *APX2*, a cytosolic member of the L-ascorbate peroxidase gene family, and *GSTU9*, together with transcription factors *WRKY41* and *WRKY50*, the latter previously shown to be involved in defense responses and the former recently shown to be involved in the modulation of *ABSCISIC ACID INSENSITIVE3*




Figure 6. Phenotypes of *RGGA* knockout and overexpressing plants. A, Phenotypes of Col-0, *rgga*, and 35S::FLAG-*RGGA* plants grown on GM for 14 d and exposed for 7 d to NaCl (180 or 200 mM). B, Water loss of leaves detached from Col-0, *rgga*, and 35S::FLAG-*RGGA* plants. Data are presented as percentages of initial weight lost at different time points (1, 2, and 3 h). Each point consists of average values \pm SD ($n = 5$ for each line). Data relative to 35S::FLAG-*RGGA* represent means of three independent transgenic lines. C, Quantification of primary root length of 14-d-old seedlings germinated for 4 d on GM and transferred to control GM medium or medium containing 20 μ M ABA. Values are means \pm SD ($n = 25$). The asterisk indicates a statistically significant difference assessed by Student's *t* test ($P < 0.001$). D, Photograph of seedlings grown as described in C. E, Survival test of 18-d-old seedlings germinated on GM and transferred to ABA (50 μ M) medium. Survival was scored daily in terms of absence of necrotic or bleached leaves. Data are means \pm SD of three independent experiments ($n = 30$). Asterisks denote statistically significant differences versus Col-0 assessed by χ^2 test (* $P < 0.05$, ** $P < 0.01$).


transcript abundance and ABA sensitivity at the seed and early seedling stages (Gao et al., 2011; Ding et al., 2014).

DISCUSSION

In a previous report, we identified *StRGGA*, a potato gene encoding a putative RBP whose expression was

specifically induced in cultured cells adapted to high concentrations of PEG (Ambrosone et al., 2011; Fig. 1). Here, we have provided evidence that the putative Arabidopsis ortholog, *AtRGGA*, is involved in tolerance to drought and salt stress. *AtRGGA* is up-regulated both in cells and plants upon long-term exposure to PEG and ABA (Fig. 1). *AtRGGA* is expressed in several

Figure 7. Phenotypes of *AtRGGA* knockout and overexpressing plants. A, Representative 4-week-old plants of Col-0, *rgga*, and 35S::FLAG-RGGA genotypes grown in control conditions or after 7-d NaCl (300 mM) or drought treatment. B, Pro concentrations in Col-0, *rgga*, and 35S::FLAG-RGGA plants treated as described in A. Asterisks indicate statistically significant differences assessed by Student's *t* test ($P < 0.01$).



Arabidopsis tissues, including tissues that perceive or respond to osmotic stress conditions, such as roots and stomata, and tissues that undergo extensive dehydration processes, such as pollen (Fig. 3). Finally, the *rgga* knockout mutant displays a seed germination hypersensitive to NaCl and is less tolerant compared with the wild type to salt stress both at the seedling stage and as adult plants (Figs. 5–7). Conversely, transgenic plants overexpressing *AtRGGA* appear to be better able to withstand salt and drought stress and lose water at a slower rate in detached leaf assays as compared with control untransformed plants (Figs. 5 and 6), indicating that a higher expression of *AtRGGA* promotes stomatal closure, a largely ABA-dependent process (Verslues et al., 2006). The hypersensitivity of *rgga* and

the lower sensitivity of 35S::FLAG-RGGA plants to ABA (Fig. 6) further indicate that *AtRGGA* participates in ABA-dependent mechanisms of response to salt and drought stress. When exposed to drought stress, *AtRGGA*-overexpressing plants also accumulate less Pro (Fig. 7), suggesting that the higher tolerance observed is independent of Pro accumulation, a feature shared with several tolerant mutants such as *reduced salt sensitivity1*, *stigma specific protein1 (stig1)*, and *photoautotrophic salt tolerance1* that do not hyperaccumulate this compatible osmolyte under stress conditions (Werner and Finkelstein, 1995; Tsugane et al., 1999; Gao et al., 2006).

Microarray experiments showed that an abolished/increased expression of *AtRGGA* has a profound impact

Table 1. Fold change values obtained by microarray analysis compared with wild-type *Col-0* of genes showing opposite behavior in *rgga* and *35S::FLAG-RGGA* in control conditionsThe locus corresponding to *AtRGGA* is highlighted in boldface.

Probe	Locus	Gene Name	Genotype	
			<i>rgga</i>	<i>35S::FLAG-RGGA</i>
A_84_P20528	<i>BARS1</i>	Baruol synthase	-4.10	2.53
A_84_P12286	<i>AT1G19060</i>	Hypothetical protein	-2.56	3.47
A_84_P835922	<i>UGT88A1</i>	UDP-glucosyltransferase88A1	-2.32	2.20
A_84_P225559	<i>WRKY51</i>	Putative WRKY transcription factor51	-2.09	3.41
A_84_P18337	<i>AOX1B</i>	Alternative oxidase1B	-10.38	2.95
A_84_P18276	<i>AT2G39590</i>	40S ribosomal protein S15a-3	-3.17	7.54
A_84_P806445	<i>AT4G16830</i>	Hyaluronan/mRNA-binding domain-containing protein	-19.86	24.09
A_84_P15686	<i>ATHSP22.0</i>	Heat shock protein22	-5.20	4.74
A_84_P13852	<i>HSP21</i>	Heat shock protein21	-5.66	3.41
A_84_P17108	<i>AT1G71000</i>	Chaperone DnaJ domain-containing protein	-2.71	8.07
A_84_P21479	<i>AT4G16830</i>	Hyaluronan/mRNA binding domain-containing protein	-13.40	13.50
A_84_P822865	<i>DREB2A</i>	Dehydration-responsive element-binding protein2A	-3.60	2.12
A_84_P22181	<i>PMZ</i>	Zinc finger AN1 domain-containing stress-associated protein12	-2.48	2.12
A_84_P10708	<i>AT2G38340</i>	Dehydration-responsive element-binding protein19	-3.23	2.08
A_84_P11439	<i>AT1G52560</i>	HSP20-like chaperone	-2.86	3.09
A_84_P11143	<i>WRKY38</i>	Putative WRKY transcription factor38	-2.50	2.60
A_84_P807598	<i>AT5G22430</i>	Pollen Ole e1 allergen and extensin family protein	2.82	-3.57
A_84_P799181	<i>AT1G05660</i>	Pectin lyase-like protein	2.32	-2.25
A_84_P22316	<i>CYP706A7</i>	Cytochrome P450, family 706, subfamily A, polypeptide 7	2.20	-3.48
A_84_P99106	<i>CLE4</i>	Protein CLAVATA3/Embryo Surrounding Region-related4	4.01	-3.14
A_84_P15932	<i>TIP2;3</i>	Aquaporin TIP2-3	2.33	-9.10
A_84_P20461	<i>AT4G25250</i>	Plant invertase/pectin methylesterase inhibitor domain-containing protein	4.16	-3.20
A_84_P10835	<i>AT3G32030</i>	Terpene cyclase C1 domain-containing protein	3.59	-2.69
A_84_P15498	<i>AT3G02620</i>	Acyl-[acyl-carrier-protein] desaturase	2.69	-3.10
A_84_P11650	<i>AT1G07550</i>	Putative Leu-rich repeat receptor-like Ser/Thr-protein kinase	2.77	-2.11
A_84_P16139	<i>CYP702A1</i>	Cytochrome P450, family 702, subfamily A, polypeptide 1	3.25	-2.79
A_84_P15513	<i>AT3G14540</i>	Terpene cyclase C1 domain-containing protein	3.03	-3.87
A_84_P869688	<i>AGP30</i>	Arabinogalactan protein30	2.69	-4.71
A_84_P14717	<i>EXPA17</i>	Putative expansin A17	3.58	-2.63
A_84_P12815	<i>ATGSTF13</i>	Glutathione S-transferase-like protein	3.47	-2.23
A_84_P13690	<i>AT3G46370</i>	Leu-rich repeat protein kinase-like protein	2.58	-2.12
A_84_P11899	<i>SULTR1;1</i>	Sulfate transporter1.1	3.21	-2.47
A_84_P513890	<i>AT4G11780</i>	Hypothetical protein	2.62	-2.14
A_84_P19188	<i>CYP718</i>	Cytochrome P450, family 718	2.20	-2.38
A_84_P18031	<i>AT1G05660</i>	Pectin lyase-like protein	2.53	-2.74
A_84_P95306	<i>FUT5</i>	Xyloglucan fucosyltransferase	4.46	-2.10
A_84_P759704	<i>AT3G32030</i>	Terpene cyclase C1 domain-containing protein	3.67	-2.07
A_84_P17611	<i>AT4G22460</i>	Bifunctional inhibitor/lipid-transfer protein/seed storage 2S albumin-like protein	2.83	-5.30
A_84_P17872	<i>AT5G60520</i>	Late embryogenesis abundant protein-like protein	2.20	-2.57
A_84_P11794	<i>AT3G45080</i>	Sulfotransferase family protein	2.44	-2.19
A_84_P597193	<i>AT2G48080</i>	Oxidoreductase, 2-Oxoglutarate-Fe(II) oxygenase-like protein	3.83	-2.61
A_84_P13496	<i>AT2G25150</i>	HXXXD-type acyltransferase-like protein	2.14	-4.90
A_84_P18124	<i>AT1G06330</i>	Heavy metal transport/detoxification-like protein	2.99	-2.53
A_84_P22080	<i>AGP30</i>	Arabinogalactan protein30	2.90	-4.48
A_84_P13966	<i>AT5G14650</i>	Polygalacturonase	4.57	-3.00
A_84_P599045	<i>AT3G19320</i>	Leu-rich repeat-containing protein	4.47	-4.51
A_84_P513281	<i>AT5G62330</i>	Hypothetical protein	4.43	-5.10
A_84_P262640	<i>AT5G22430</i>	Pollen Ole e1 allergen and extensin family protein	2.80	-4.44
A_84_P759732	<i>AT3G32030</i>	Terpene cyclase C1 domain-containing protein	3.86	-3.09

on global gene expression, with a misregulation of several thousand genes in control and stress conditions (Tables I and II; Supplemental Tables S3–S10; Supplemental Figs. S5 and S6). Analysis of significantly enriched GO terms showed that terms such as response to stimulus,

regulation of cellular process, and biological regulation were among the ones with lowest false discovery rates, with the exception of genes up-regulated in salt-stressed *AtRGGA*-overexpressing plants, where only the terms response to salt stress, response to stimulus, and response

Table II. Fold change values obtained by microarray analysis compared with wild-type Col-0 of genes showing opposite behavior in *rgga* and *35S::FLAG-RGGA* after salt stress treatment (48 h, 180 mM NaCl)The locus corresponding to *AtRGGA* is highlighted in boldface.

Probe	Locus	Gene Name	Genotype	
			<i>rgga</i>	<i>35S::FLAG-RGGA</i>
A_84_P756195	<i>AT2G04070</i>	Multi Antimicrobial Extrusion Protein efflux family protein	-3.59	3.60
A_84_P768949	<i>AT5G35688</i>	Hypothetical protein	-2.62	2.21
A_84_P11264	<i>GSTU9</i>	GST τ 9	-2.03	2.11
A_84_P762759	<i>AT3G55672</i>	Self-incompatibility S1 family protein	-2.01	4.19
A_84_P89069	<i>AT3G63360</i>	Defensin-like protein11	-2.32	2.74
A_84_P84999	<i>AT3G28580</i>	AAA-type ATPase family protein	-2.42	2.50
A_84_P593216	<i>AT5G60250</i>	C3H4-type zinc finger protein	-2.61	2.17
A_84_P75404	<i>AT5G55150</i>	Hypothetical protein	-3.91	2.19
A_84_P55910	<i>WRKY41</i>	Putative WRKY transcription factor41	-2.07	5.03
A_84_P11156	<i>WRKY50</i>	Putative WRKY transcription factor50	-2.22	3.45
A_84_P580762	<i>AT3G43710</i>	Putative F-box/Kelch-repeat protein	-5.29	4.42
A_84_P806445	<i>AT4G16830</i>	Hyaluronan/mRNA-binding domain-containing protein	-5.34	21.32
A_84_P271800	<i>EPI1</i>	Putative Cys-rich receptor-like protein kinase9	-2.16	2.17
A_84_P23566	<i>AT5G64790</i>	O-Glycosylhydrolase family17 protein	-2.36	2.28
A_84_P606546	<i>AT5G55270</i>	Hypothetical protein	-6.56	3.20
A_84_P833327	<i>AT1G69550</i>	Toll/Interleukin1 Receptor-Nucleotide Binding Site-Leu-rich repeat class disease resistance protein	-2.21	2.24
A_84_P21479	<i>AT4G16830</i>	Hyaluronan/mRNA-binding domain-containing protein	-3.45	10.14
A_84_P23310	<i>AT4G27580</i>	Hypothetical protein	-3.14	3.11
A_84_P20019	<i>AT1G17960</i>	Threonyl-tRNA synthetase	-2.54	2.75
A_84_P11712	<i>AT3G02810</i>	Protein kinase domain-containing protein	-3.58	3.48
A_84_P539380	<i>AT2G35075</i>	Hypothetical protein	-3.75	2.24
A_84_P17261	<i>UGT84B1</i>	UDP-Glc:(indol-3-yl)acetate β -D-glucosyltransferase	-2.19	3.44
A_84_P525660	<i>AT5G52940</i>	Hypothetical protein	-3.96	2.31
A_84_P17379	<i>APX2</i>	L-Ascorbate peroxidase	-2.51	2.55
A_84_P145639	<i>AT5G07610</i>	F-box protein	-7.36	4.13
A_84_P20216	<i>AT3G05950</i>	Germin-like protein subfamily 1, member 7	-2.09	8.36
A_84_P787266	<i>WRKY50</i>	Putative WRKY transcription factor50	-2.45	3.68
A_84_P590126	<i>AT3G22540</i>	Hypothetical protein	3.10	-2.22
A_84_P16241	<i>AT1G52190</i>	Putative peptide transporter	2.87	-2.43
A_84_P22477	<i>MYB40</i>	Myb domain protein40	3.73	-2.85
A_84_P275660	<i>AGP14</i>	Arabinogalactan protein14	2.81	-2.93
A_84_P812392	<i>AT4G33720</i>	Putative pathogenesis-related protein	64.60	-2.05
A_84_P94979	<i>AT5G18060</i>	SAUR-like auxin-responsive protein	2.37	-8.25
A_84_P137439	<i>PAR2</i>	Phytochrome rapidly regulated2 protein	2.09	-6.16
A_84_P20461	<i>AT4G25250</i>	Plant invertase/pectin methylesterase inhibitor domain-containing protein	2.22	-3.57
A_84_P751997	<i>AT1G15630</i>	Hypothetical protein	2.21	-2.88
A_84_P544465	<i>KDR</i>	Basic helix-loop-helix domain-containing protein	2.36	-3.64
A_84_P79415	<i>AT2G18300</i>	Transcription factor basic helix-loop-helix64	2.47	-2.92
A_84_P175621	<i>AGL14</i>	Agamous-like MADS box protein AGL14	2.42	-2.46
A_84_P11420	<i>AT1G64920</i>	UDP-glycosyltransferase-like protein	2.41	-2.28
A_84_P310653	<i>PDF2.5</i>	Defensin-like protein6	2.54	-2.42
A_84_P12525	<i>AT2G47880</i>	Glutaredoxin C13	3.54	-3.34
A_84_P20410	<i>ADS1</i>	Δ -9 acyl-lipid desaturase1	2.46	-2.25
A_84_P732491	<i>AT5G03995</i>	Hypothetical protein	4.90	-2.38
A_84_P14346	<i>SULTR1;3</i>	Sulfate transporter1.3	2.32	-2.49
A_84_P19794	<i>MYB66</i>	Transcription factor WEREWOLF	2.45	-2.86
A_84_P186864	<i>IQD12</i>	IQ-domain12 protein	2.49	-2.85
A_84_P20189	<i>AT3G03830</i>	SAUR-like auxin-responsive protein	2.03	-6.53
A_84_P121222	<i>AT3G21330</i>	Transcription factor basic helix-loop-helix87	3.68	-3.68
A_84_P500468	<i>RALFL27</i>	Protein ralf-like27	3.61	-2.67
A_84_P24090	<i>AT3G45710</i>	Major facilitator protein	2.29	-2.67
A_84_P759704	<i>AT3G32030</i>	Terpene cyclase C1 domain-containing protein	2.17	-3.34
A_84_P16734	<i>SAUR15</i>	SAUR-like auxin-responsive protein	3.08	-2.10
A_84_P166743	<i>AT5G14890</i>	NHL domain-containing protein	2.02	-2.15
A_84_P21135	<i>AT3G03820</i>	SAUR-like auxin-responsive protein	3.45	-2.31

(Table continues on following page.)

Table II. (Continued from previous page.)

Probe	Locus	Gene Name	Genotype	
			<i>rgga</i>	<i>35S::FLAG-RGGA</i>
A_84_P17378	<i>GRI</i>	Stigma-specific Stig1 family protein	2.75	-2.07
A_84_P759796	<i>AT3G46270</i>	Receptor protein kinase-like protein	2.81	-2.07
A_84_P64634	<i>AT4G10910</i>	Hypothetical protein	2.45	-4.97
A_84_P18924	<i>AT1G52130</i>	Man-binding lectin-like protein	2.38	-6.23
A_84_P12394	<i>AT1G14960</i>	Major latex-related protein	2.34	-4.72

to biotic stimulus were significantly enriched, indicating that the tolerance of *RGGA*-overexpressing plants is attributable to proteins involved in protection from biotic and abiotic stress. In *rgga*, the expression of *FLOWERING LOCUS T*, encoding the mobile signal translocated from leaves to the shoot apex to initiate flowering (Abe et al., 2005; Wigge et al., 2005; Jaeger and Wigge, 2007), is severely reduced, possibly accounting for the delayed-flowering phenotype observed (Supplemental Fig. S4). In the mutant, an up-regulation of several members of the *PURPLE ACID PHOSPHATASE (PAP)* family was observed after salt stress treatment. PAPs are metalloenzymes involved in diverse biological processes, such as inorganic phosphate uptake, peroxidation, defense against pathogens, and salt stress, through the alleviation of oxidative damage (Li et al., 2008; Ravichandran et al., 2013). *PAP* transcript up-regulation, therefore, could be an indication of an increased salt stress-caused oxidative damage in *rgga* compared with control plants. Analysis of genes showing an opposite behavior in the mutant versus overexpressing plants as compared with the wild type (Tables I and II) showed that, in control conditions, transcripts of several genes involved in stress responses, such as the heat shock proteins *HSP22* and *HSP21* and the transcription factors *DREB19* and *DREB2A*, are down-regulated in *rgga* and up-regulated in transgenic plants, indicating that higher prestress levels of these genes may partially account for the higher stress tolerance shown by transgenic plants. After salt stress treatment, several members of the *SAUR* gene family were up-regulated in *rgga* and down-regulated in overexpressing plants, including *SAUR15*, *At5g18060*, and *At3g03830*. *SAUR* genes encode small-*M*₁ auxin-responsive proteins whose function is still largely unknown, even though members of the *SAUR19* subfamily, comprising *SAUR19* to *SAUR24*, have been shown recently to have a role in auxin-mediated cell expansion by regulating plasma membrane H⁺-ATPase activity (Spartz et al., 2012, 2014). Transcripts of *SAUR* genes are highly unstable due to the presence in their 3' untranslated region of a conserved downstream element responsible for mRNA instability (Newman et al., 1993; Gil and Green, 1996; Spartz et al., 2012), and, in an early report, *SAUR* mRNAs were shown to be stabilized by the inhibition of protein synthesis (Franco et al., 1990). An interesting hypothesis, therefore, could be that *AtRGGA* might affect RNA stability.

AtRGGA encodes a cytoplasm-localized protein (Fig. 4) with several Gly/Arg motifs that possesses the Stm1

N-terminal and the HABP4_PA1-RBP1 domains characteristic of RNA and nucleic acid-binding proteins (Fig. 2). Stm1 is a yeast G4 quadruplex and purine motif triplex nucleic acid-binding protein of *Saccharomyces cerevisiae* that has been shown to associate with telomeric Y' DNA and ribosomes (Van Dyke et al., 2004). HABP4 binds hyaluronan as well as RNA, while the human PAI-1 mRNA-binding protein binds the type 1 plasminogen activator inhibitor and has been suggested to be involved in the regulation of mRNA stability (Huang et al., 2000; Heaton et al., 2001). As expected from the gene annotation, we show that recombinant *AtRGGA* is capable of efficiently binding RNA in vitro (Fig. 2). The similarity with the Stm1, HABP4, and PAI1 proteins, together with the subcellular localization, suggest that *AtRGGA* might affect posttranscriptional regulation mechanisms such as the control of RNA stability, storage, or translation efficiency rather than the synthesis or nuclear processing of RNAs. In humans and yeast, RGG motif-containing proteins have emerged as key players involved in the posttranscriptional regulation of gene expression, affecting RNA stability as well as RNA translation to protein through interaction with the translation initiation factor eIF4G, which recruits ribosomes to mRNAs (Rajyaguru and Parker, 2012; Walsh and Mohr, 2014). While the possibility that poly(A⁺) RNAs are also bound by *AtRGGA* could not be ruled out, the observed binding to poly(A⁻) (Fig. 2) raises the possibility that *AtRGGA* might interact with one or more RNA components of the ribosomes to modify translation efficiency and/or the stability of ribosome-bound mRNAs. Little evidence suggests that binding to the ribosomes can affect mRNA stability. In bacteria, ribosomes usually protect mRNAs from degradation or, in some instances, promote mRNA decay (Deana and Belasco, 2005). In Arabidopsis, a mutant lacking ribosomal protein S27 shows defects in the ability to degrade selected mRNAs when exposed to genotoxic treatments (Revenkova et al., 1999). More recently, polysome association was correlated to transcript stability during stress conditions in rice (*Oryza sativa*; Park et al., 2012). Further experiments to identify RNA and protein interactors will assess the hypothesis that *AtRGGA* binds to ribosomes and discern whether mRNA stability and/or translation efficiency are affected in stress conditions by the presence of *AtRGGA*.

MATERIALS AND METHODS

Plant Material and Stress Treatments

Potato (*Solanum tuberosum* 'Desirée') cell cultures were generated, maintained, and gradually or abruptly exposed to 20% (w/v) PEG 8000 as described previously (Leone et al., 1994; Ambrosone et al., 2011). Arabidopsis (*Arabidopsis thaliana*) MM2D cells (ecotype Landsberg *erecta*) were maintained as described (Menges and Murray, 2002) in continuous dark conditions. For gene expression analyses, cells were treated for 24 h with NaCl (150 mM), ABA (50 μ M), or 10% (w/v) PEG 8000. Cell viability was assessed by adding fluorescein diacetate (2 μ L of a 5 mg mL⁻¹ stock) to 100 μ L of cell culture. Viability was expressed as the percentage of viable cells in stress versus control conditions.

Arabidopsis plants of the Col-0 ecotype were used throughout this study. A knockout line for *AtRGGGA* (*rgga*; SALK_143514) was obtained from the Nottingham Arabidopsis Stock Centre, and homozygous plants were selected by PCR. Primers were selected using the T-DNA Express primer design tool (<http://signal.salk.edu/tdnaprimers.2.html>; Supplemental Table S1). Transgenic plants overexpressing FLAG-tagged *RGGA* in the Col-0 background were generated by floral dip transformation (Clough and Bent, 1998).

For gene expression studies, 14-d-old seedlings grown on solid GM medium were transferred to plates containing NaCl (50, 100, or 150 mM) and incubated for 24 h or incubated for 7 d on plates containing 35% (w/v) PEG, NaCl (120 mM), or ABA (10 μ M). For germination analyses, seeds were sown in the presence of NaCl (120 mM). Germination was scored in terms of fully expanded cotyledons after 7 d of incubation.

Survival tests were carried out using 18-d-old seedlings germinated on GM plates and transferred to NaCl- or ABA-containing medium (180 mM or 50 μ M, respectively). Survival was scored daily in terms of absence of necrotic or bleached leaves.

To score root growth in the presence of ABA, 4-d-old seedlings grown on GM medium and showing equal primary root length were transferred to GM medium or medium with 20 μ M ABA. Photographs were taken, and root length was scored 10 d after transfer. Stress treatments in soil were performed on 4-week-old plants by water withdrawal for 7 d or by watering with 300 mM NaCl every other day for 7 d. In each irrigation event, the volume of the salt solution was 5% of the pot volume.

For microarray gene expression analyses, 10-d-old seedlings grown on GM plates were transferred to GM or GM + 180 mM NaCl plates for 48 h prior to RNA extraction.

Plasmid Construction

Gateway technology (Life Technologies) was used to obtain binary vectors for promoter and protein localization studies as well as to produce transgenic overexpressing plants. The putative promoter of *AtRGGGA* (corresponding to 2 kb upstream of ATG) was amplified from genomic DNA extracted from Col-0 plant *RGGA* coding sequence amplified from clone U22150 provided by The Arabidopsis Information Resource. PCR amplifications were carried out using Pfu DNA polymerase (Stratagene). Primers used are listed in Supplemental Table S1. The promoter and coding sequences were cloned into pDONR207 using BP clonase to obtain *AtRGGAPromoter::pDONR207* and *AtRGGGA::pDONR207* entry vectors. LR clonase was used for recombination with destination vectors, which were pMDC164 (Curtis and Grossniklaus, 2003) for promoter studies, pEG101 and pEG104 (Earley et al., 2006) for protein localization studies, and pEG202 (Earley et al., 2006) to produce FLAG-tagged overexpressing plants.

To produce His-tagged *AtRGGGA* in *Escherichia coli*, the coding sequence was amplified and cloned between *Sall* and *NotI* restriction sites of pET28a vector. All constructs were sequenced to rule out the presence of mutations introduced by PCR. The His-tagged *AtRGGGA* from pET28a constructs was overexpressed following transformation in *E. coli* BL21 (DE3). Cells were recovered and lysed by 1 mg mL⁻¹ lysozyme after growth at 37°C on Luria-Bertani medium supplemented with 50 μ g mL⁻¹ kanamycin. Induction was carried out by adding 0.5 mM isopropyl- β -D-thiogalactopyranoside to a culture at an optical density measured at 600 nm of 1 and then incubating the culture at 37°C for 4 h. Protein constructs were purified by nickel affinity chromatography as described by the manufacturer (Qiagen). The purified protein was analyzed by SDS-PAGE as described by Laemmli (1970).

Gene Expression Analyses

qPCR gene expression analyses used 1 μ g of DNaseI-treated RNA converted to complementary DNA utilizing SuperScript II Reverse Transcriptase and oligo(dT₂₀) following the manufacturer's instructions (Life Technologies).

The obtained complementary DNA was diluted 1:20, and 4.5 μ L was used for each qRT-PCR, carried out with Platinum SYBR Green qPCR SuperMix and Elongation Factor *EF1 α* as an endogenous control. Primers used (5 μ M) are listed in Supplemental Table S1. qRT-PCR was performed with ABI 7900 HT (Applied Biosystems). At least nine technical replicates per gene per biological replicate were performed. Three biological replicates per genotype were tested, and data were analyzed using RQ Manager followed by analysis with Data Assist (Applied Biosystems) to group together the different replicates and biological replicates and perform statistical analysis.

For microarray gene expression studies, RNA extracted using Trizol (Life Technologies) was quantified and quality checked using the Nanodrop-ND 8000 spectrophotometer (Thermo Fisher Scientific) and the 2100 Bioanalyzer (Agilent Technologies), respectively. Samples with the RNA integrity number of 9 and above were chosen for further analysis. Microarray analysis was performed using the Arabidopsis (V4) Gene Expression Microarray, Design ID: 021169 (Agilent Technologies), containing 43,803 Arabidopsis gene probes and 1,417 Agilent control probes. Total RNA (150 ng) was used to prepare cyanine-3-labeled probe with the help of the low-RNA input linear amplification/labeling kit (Agilent Technologies). The dye incorporation and the copy RNA yield were measured using the Nanodrop-ND 8000 spectrophotometer (Thermo Fisher Scientific). Labeled copy RNA probes (1.65 μ g) were fragmented using fragmentation buffer (Agilent Technologies) and hybridized to the Arabidopsis arrays in the presence of the gene expression hybridization buffer HI-RPM and blocking agent (Agilent Technologies) for 17 h at 65°C with a 10-rpm rotation speed in a hybridization oven (Agilent Technologies). After the 17-h incubation, the arrays were washed using low-stringency wash buffer 1 (Agilent Technologies) at room temperature for 1 min followed by a high-stringency wash using wash buffer 2 (Agilent Technologies) at 37°C. The arrays were air dried and scanned using the high-resolution array scanner (Agilent Technologies) with the appropriate settings for the one-color gene expression arrays. The signal intensities were extracted from the scanned images with the aid of Feature Extraction Software 10.7.1.1 (Agilent Technologies) and subjected to background subtraction and spatial detrending. The outliers and the abnormal features were flagged, and the data were normalized using intraarray percentile shift normalization (threshold of 75 and above) and median-based interarray normalization. GeneSpring GX (Agilent Technologies) was used to calculate the intensity ratios and fold changes. All the genes with *P* values below 0.05 and fold change above 2 were chosen for GO enrichment analysis, which was carried out by Sequentia Italia as described below.

Common sets of differentially expressed genes were identified between the different groups by using Venny (<http://bioinfogp.cnb.csic.es/tools/venny/index.html>; Oliveros, 2007). The identification of enriched GO functional categories was performed by using the Singular Enrichment Analysis tool of agriGO (<http://bioinfo.cau.edu.cn/agriGO/analysis.php>). Arabidopsis was selected as a supported species, and the suggested background was used. A hypergeometric test was used as a statistical method to identify enriched categories, while the other parameters were left as default. When the number of enriched categories was too much for easy understanding, the software REVIGO (<http://revigo.irb.hr>; Supek et al., 2011) was used to obtain reduced lists. The dimension of the output list was set to small, and the database of Arabidopsis GO terms was used.

Immunoblotting

To detect the FLAG-RGGA fusion protein in overexpressing lines, immunoblotting was performed using α -FLAG antibody (Sigma-Aldrich). Total proteins were extracted from 100 mg of tissue using extraction buffer (2 M urea, 0.1 M NaH₂PO₄, and 0.01 M Tris, pH 8) and protease inhibitor cocktail (Sigma-Aldrich). After clarification, protein concentration was estimated using the Bradford reagent assay (Bio-Rad), and 50 μ g of total protein was used for SDS-PAGE followed by transfer on nitrocellulose membranes (Amersham, GE Healthcare). A 1:5,000 dilution of α -FLAG antibody was used for immunoblotting following the manufacturer's instructions.

RNA EMSA

Total RNA (700 ng) extracted from control and salt stress-treated plants was labeled with biotin using the RNA 3' End Biotinylation Kit (Pierce Biotechnology) following the manufacturer's instructions. One microliter of a 1:20 dilution (approximately 30 ng) of the labeled RNA was used for each EMSA reaction. Similar amounts of labeled RNA were also used for poly(A⁺) and poly(A⁻) RNA, which were prepared using the mRNA Isolation Kit (Roche Applied Science). RNA was incubated with 7.5 μ g of His-tagged *AtRGGGA* or His-tagged PYR1. When present, unlabeled RNA was used as a competitor at

560 ng (approximately 160-fold). The RNA EMSA was carried out with the LightShift Chemiluminescent RNA EMSA Kit (Pierce Biotechnology). The binding reaction was analyzed by gel electrophoresis on a native 6% polyacrylamide gel in 0.5× Tris-borate/EDTA buffer and transferred to a nylon membrane.

Pro Content Determination

Pro extraction from Arabidopsis plants grown in soil and subjected to drought stress was carried out as described by Claussen (2005). Briefly, 500 mg of leaves was ground in liquid nitrogen and resuspended in an aqueous solution containing 3% (w/v) 5-sulfosalicylic acid. Suspension was clarified with Ederol round filters (Schleicher & Schull). One milliliter of filtered suspension was diluted with 1 mL of glacial acetic acid and 1 mL of a 2.5% ninhydrin solution (glacial acetic acid:water:orthophosphoric acid, 6:3:1). After incubation at 100°C for 1 h, samples were read at an optical density measured at 546 nm. Three biological replicates were analyzed for each genotype, and each sample was replicated at least three times.

Confocal Imaging

Confocal imaging was carried out with the Leica TCS SP2 system. Propidium iodide staining was carried out as described previously (Sassi et al., 2012). Staining times were as follows: 1 min to counterstain cell walls only, and at least 1 h to achieve minimal nuclear staining.

Supplemental Data

The following supplemental materials are available.

Supplemental Figure S1. Alignment of deduced RGGa protein sequences of potato and Arabidopsis.

Supplemental Figure S2. Cell viability measurements of Arabidopsis MM2D cells.

Supplemental Figure S3. *AtRGGa* gene expression in Arabidopsis tissues and developmental stages.

Supplemental Figure S4. Phenotypes of Arabidopsis plants with modified expression of *AtRGGa*.

Supplemental Figure S5. Enriched GO categories of differentially expressed genes in 35S::FLAG-RGGa plants.

Supplemental Figure S6. Enriched GO categories of differentially expressed genes in *rgga* plants.

Supplemental Table S1. Primers used in this study.

Supplemental Table S2. qPCR validation of microarray results.

Supplemental Table S3. Genes repressed in the *rgga* mutant in salt stress conditions.

Supplemental Table S4. Genes induced in the *rgga* mutant in salt stress conditions.

Supplemental Table S5. Genes induced in the *rgga* mutant in control conditions.

Supplemental Table S6. Genes repressed in the *rgga* mutant in control conditions.

Supplemental Table S7. Genes induced in plants overexpressing RGGa in salt stress conditions.

Supplemental Table S8. Genes repressed in plants overexpressing RGGa in salt stress conditions.

Supplemental Table S9. Genes induced in plants overexpressing RGGa in control conditions.

Supplemental Table S10. Genes repressed in plants overexpressing RGGa in control conditions.

ACKNOWLEDGMENTS

We thank the Nottingham Arabidopsis Stock Centre for providing T-DNA mutant seeds and Gaetano Guarino and Rosario Nocerino for excellent technical assistance.

Received December 17, 2014; accepted March 16, 2015; published March 17, 2015.

LITERATURE CITED

- Abe M, Kobayashi Y, Yamamoto S, Daimon Y, Yamaguchi A, Ikeda Y, Ichinoki H, Notaguchi M, Goto K, Araki T** (2005) FD, a bZIP protein mediating signals from the floral pathway integrator FT at the shoot apex. *Science* **309**: 1052–1056
- Albà MM, Pagès M** (1998) Plant proteins containing the RNA-recognition motif. *Trends Plant Sci* **3**: 15–21
- Ambrosone A, Costa A, Leone A, Grillo S** (2012) Beyond transcription: RNA-binding proteins as emerging regulators of plant response to environmental constraints. *Plant Sci* **182**: 12–18
- Ambrosone A, Costa A, Martinelli R, Massarelli I, De Simone V, Grillo S, Leone A** (2011) Differential gene regulation in potato cells and plants upon abrupt or gradual exposure to water stress. *Acta Physiol Plant* **33**: 1157–1171
- Ambrosone A, Di Giacomo M, Leone A, Grillo MS, Costa A** (2013) Identification of early induced genes upon water deficit in potato cell cultures by cDNA-AFLP. *J Plant Res* **126**: 169–178
- Branco-Price C, Kaiser KA, Jang CJ, Larive CK, Bailey-Serres J** (2008) Selective mRNA translation coordinates energetic and metabolic adjustments to cellular oxygen deprivation and reoxygenation in Arabidopsis thaliana. *Plant J* **56**: 743–755
- Burd CG, Dreyfuss G** (1994) Conserved structures and diversity of functions of RNA-binding proteins. *Science* **265**: 615–621
- Chung JS, Zhu JK, Bressan RA, Hasegawa PM, Shi H** (2008) Reactive oxygen species mediate Na⁺-induced SOS1 mRNA stability in Arabidopsis. *Plant J* **53**: 554–565
- Claussen W** (2005) Proline as a measure of stress in tomato plants. *Plant Sci* **168**: 241–248
- Crough SJ, Bent AF** (1998) Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. *Plant J* **16**: 735–743
- Curtis MD, Grossniklaus U** (2003) A Gateway cloning vector set for high-throughput functional analysis of genes in planta. *Plant Physiol* **133**: 462–469
- Deana A, Belasco JG** (2005) Lost in translation: the influence of ribosomes on bacterial mRNA decay. *Genes Dev* **19**: 2526–2533
- Ding ZJ, Yan JY, Li GX, Wu ZC, Zhang SQ, Zheng SJ** (2014) WRKY41 controls Arabidopsis seed dormancy via direct regulation of ABI3 transcript levels not downstream of ABA. *Plant J* **79**: 810–823
- dit Frey NF, Muller P, Jammes F, Kizis D, Leung J, Perrot-Rechenmann C, Bianchi MW** (2010) The RNA binding protein Tudor-SN is essential for stress tolerance and stabilizes levels of stress-responsive mRNAs encoding secreted proteins in Arabidopsis. *Plant Cell* **22**: 1575–1591
- Du Z, Zhou X, Ling Y, Zhang Z, Su Z** (2010) agriGO: a GO analysis toolkit for the agricultural community. *Nucleic Acids Res* **38**: W64–W70
- Earley KW, Haag JR, Pontes O, Opper K, Juehne T, Song K, Pikaard CS** (2006) Gateway-compatible vectors for plant functional genomics and proteomics. *Plant J* **45**: 616–629
- Franco AR, Gee MA, Guilfoyle TJ** (1990) Induction and superinduction of auxin-responsive mRNAs with auxin and protein synthesis inhibitors. *J Biol Chem* **265**: 15845–15849
- Fujii H, Chinnusamy V, Rodrigues A, Rubio S, Antoni R, Park SY, Cutler SR, Sheen J, Rodriguez PL, Zhu JK** (2009) In vitro reconstitution of an abscisic acid signalling pathway. *Nature* **462**: 660–664
- Fujii H, Zhu JK** (2012) Osmotic stress signaling via protein kinases. *Cell Mol Life Sci* **69**: 3165–3173
- Gao QM, Venugopal S, Navarre D, Kachroo A** (2011) Low oleic acid-derived repression of jasmonic acid-inducible defense responses requires the WRKY50 and WRKY51 proteins. *Plant Physiol* **155**: 464–476
- Gao X, Ren F, Lu YT** (2006) The Arabidopsis mutant stg1 identifies a function for TBP-associated factor 10 in plant osmotic stress adaptation. *Plant Cell Physiol* **47**: 1285–1294
- Geiger D, Scherzer S, Mumm P, Marten I, Ache P, Matschi S, Liese A, Wellmann C, Al-Rasheid KA, Grill E, et al** (2010) Guard cell anion channel SLAC1 is regulated by CDPK protein kinases with distinct Ca²⁺ affinities. *Proc Natl Acad Sci USA* **107**: 8023–8028
- Geiger D, Scherzer S, Mumm P, Stange A, Marten I, Bauer H, Ache P, Matschi S, Liese A, Al-Rasheid KA, et al** (2009) Activity of guard cell

- anion channel SLAC1 is controlled by drought-stress signaling kinase-phosphatase pair. *Proc Natl Acad Sci USA* **106**: 21425–21430
- Gil P, Green PJ** (1996) Multiple regions of the Arabidopsis SAUR-AC1 gene control transcript abundance: the 3' untranslated region functions as an mRNA instability determinant. *EMBO J* **15**: 1678–1686
- Guan Q, Wen C, Zeng H, Zhu J** (2013) A KH domain-containing putative RNA-binding protein is critical for heat stress-responsive gene regulation and thermotolerance in Arabidopsis. *Mol Plant* **6**: 386–395
- Heaton JH, Dlakic WM, Dlakic M, Gelehrter TD** (2001) Identification and cDNA cloning of a novel RNA-binding protein that interacts with the cyclic nucleotide-responsive sequence in the type-1 plasminogen activator inhibitor mRNA. *J Biol Chem* **276**: 3341–3347
- Huang L, Grammatikakis N, Yoneda M, Banerjee SD, Toole BP** (2000) Molecular characterization of a novel intracellular hyaluronan-binding protein. *J Biol Chem* **275**: 29829–29839
- Hubbard KE, Nishimura N, Hitomi K, Getzoff ED, Schroeder JI** (2010) Early abscisic acid signal transduction mechanisms: newly discovered components and newly emerging questions. *Genes Dev* **24**: 1695–1708
- Ishitani M, Xiong L, Stevenson B, Zhu JK** (1997) Genetic analysis of osmotic and cold stress signal transduction in *Arabidopsis*: interactions and convergence of abscisic acid-dependent and abscisic acid-independent pathways. *Plant Cell* **9**: 1935–1949
- Jaeger KE, Wigge PA** (2007) FT protein acts as a long-range signal in Arabidopsis. *Curr Biol* **17**: 1050–1054
- Kim JS, Jung HJ, Lee HJ, Kim KA, Goh CH, Woo Y, Oh SH, Han YS, Kang H** (2008) Glycine-rich RNA-binding protein 7 affects abiotic stress responses by regulating stomata opening and closing in Arabidopsis thaliana. *Plant J* **55**: 455–466
- Kim YO, Kim JS, Kang H** (2005) Cold-inducible zinc finger-containing glycine-rich RNA-binding protein contributes to the enhancement of freezing tolerance in Arabidopsis thaliana. *Plant J* **42**: 890–900
- Klingler JP, Batelli G, Zhu JK** (2010) ABA receptors: the START of a new paradigm in phytohormone signalling. *J Exp Bot* **61**: 3199–3210
- Laemmli UK** (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**: 680–685
- Leone A, Costa A, Tucci M, Grillo S** (1994) Comparative analysis of short- and long-term changes in gene expression caused by low water potential in potato (*Solanum tuberosum*) cell-suspension cultures. *Plant Physiol* **106**: 703–712
- Li WYF, Shao G, Lam HM** (2008) Ectopic expression of GmPAP3 alleviates oxidative damage caused by salinity and osmotic stresses. *New Phytol* **178**: 80–91
- Lorković ZJ** (2009) Role of plant RNA-binding proteins in development, stress response and genome organization. *Trends Plant Sci* **14**: 229–236
- Lorković ZJ, Barta A** (2002) Genome analysis: RNA recognition motif (RRM) and K homology (KH) domain RNA-binding proteins from the flowering plant Arabidopsis thaliana. *Nucleic Acids Res* **30**: 623–635
- Ma Y, Szostkiewicz I, Korte A, Moes D, Yang Y, Christmann A, Grill E** (2009) Regulators of PP2C phosphatase activity function as abscisic acid sensors. *Science* **324**: 1064–1068
- Maggio A, Zhu JK, Hasegawa PM, Bressan RA** (2006) Osmogenetics: Aristotle to Arabidopsis. *Plant Cell* **18**: 1542–1557
- Menges M, Murray JAH** (2002) Synchronous Arabidopsis suspension cultures for analysis of cell-cycle gene activity. *Plant J* **30**: 203–212
- Nakaminami K, Matsui A, Shinozaki K, Seki M** (2012) RNA regulation in plant abiotic stress responses. *Biochim Biophys Acta* **1819**: 149–153
- Newman TC, Ohme-Takagi M, Taylor CB, Green PJ** (1993) DST sequences, highly conserved among plant SAUR genes, target reporter transcripts for rapid decay in tobacco. *Plant Cell* **5**: 701–714
- Oliveros JC** (2007) VENNY: an interactive tool for comparing lists with Venn diagrams. <http://bioinfoqnp.cnb.csic.es/tools/venny/index.html> (October 2013)
- Park SH, Chung PJ, Juntawong P, Bailey-Serres J, Kim YS, Jung H, Bang SW, Kim YK, Do Choi Y, Kim JK** (2012) Posttranscriptional control of photosynthetic mRNA decay under stress conditions requires 3' and 5' untranslated regions and correlates with differential polysome association in rice. *Plant Physiol* **159**: 1111–1124
- Park SY, Fung P, Nishimura N, Jensen DR, Fujii H, Zhao Y, Lumba S, Santiago J, Rodrigues A, Chow TF, et al** (2009) Abscisic acid inhibits type 2C protein phosphatases via the PYR/PYL family of START proteins. *Science* **324**: 1068–1071
- Rajyaguru P, Parker R** (2012) RGG motif proteins: modulators of mRNA functional states. *Cell Cycle* **11**: 2594–2599
- Raschke K** (1970) Stomatal responses to pressure changes and interruptions in the water supply of detached leaves of *Zea mays* L. *Plant Physiol* **45**: 415–423
- Ravichandran S, Stone SL, Benkel B, Prithiviraj B** (2013) Purple Acid Phosphatase5 is required for maintaining basal resistance against *Pseudomonas syringae* in Arabidopsis. *BMC Plant Biol* **13**: 107
- Revenkova E, Masson J, Koncz C, Afsar K, Jakovleva L, Paszkowski J** (1999) Involvement of Arabidopsis thaliana ribosomal protein S27 in mRNA degradation triggered by genotoxic stress. *EMBO J* **18**: 490–499
- Sassi M, Lu Y, Zhang Y, Wang J, Dhonukshe P, Bilou I, Dai M, Li J, Gong X, Jaillais Y, et al** (2012) COP1 mediates the coordination of root and shoot growth by light through modulation of PIN1- and PIN2-dependent auxin transport in Arabidopsis. *Development* **139**: 3402–3412
- Sato A, Sato Y, Fukao Y, Fujiwara M, Umezawa T, Shinozaki K, Hibi T, Taniguchi M, Miyake H, Goto DB, et al** (2009) Threonine at position 306 of the KAT1 potassium channel is essential for channel activity and is a target site for ABA-activated SnRK2/OST1/SnRK2.6 protein kinase. *Biochem J* **424**: 439–448
- Shi H, Lee BH, Wu SJ, Zhu JK** (2003) Overexpression of a plasma membrane Na⁺/H⁺ antiporter gene improves salt tolerance in Arabidopsis thaliana. *Nat Biotechnol* **21**: 81–85
- Sorenson R, Bailey-Serres J** (2014) Selective mRNA sequestration by OLIGOURIDYLATE-BINDING PROTEIN 1 contributes to translational control during hypoxia in Arabidopsis. *Proc Natl Acad Sci USA* **111**: 2373–2378
- Spartz AK, Lee SH, Wenger JP, Gonzalez N, Itoh H, Inzé D, Peer WA, Murphy AS, Overvoorde PJ, Gray WM** (2012) The SAUR19 subfamily of SMALL AUXIN UP RNA genes promote cell expansion. *Plant J* **70**: 978–990
- Spartz AK, Ren H, Park MY, Grandt KN, Lee SH, Murphy AS, Sussman MR, Overvoorde PJ, Gray WM** (2014) SAUR inhibition of PP2C-D phosphatases activates plasma membrane H⁺-ATPases to promote cell expansion in Arabidopsis. *Plant Cell* **26**: 2129–2142
- Supek F, Bošnjak M, Škunca N, Šmuc T** (2011) REVIGO summarizes and visualizes long lists of Gene Ontology terms. *PLoS ONE* **6**: e21800
- Tsugane K, Kobayashi K, Niwa Y, Ohba Y, Wada K, Kobayashi H** (1999) A recessive Arabidopsis mutant that grows photoautotrophically under salt stress shows enhanced active oxygen detoxification. *Plant Cell* **11**: 1195–1206
- Van Dyke MW, Nelson LD, Weilbaecher RG, Mehta DV** (2004) Stm1p, a G4 quadruplex and purine motif triplex nucleic acid-binding protein, interacts with ribosomes and subtelomeric Y' DNA in *Saccharomyces cerevisiae*. *J Biol Chem* **279**: 24323–24333
- Verslues PE, Agarwal M, Katiyar-Agarwal S, Zhu J, Zhu JK** (2006) Methods and concepts in quantifying resistance to drought, salt and freezing, abiotic stresses that affect plant water status. *Plant J* **45**: 523–539
- Walsh D, Mohr I** (2014) Coupling 40S ribosome recruitment to modification of a cap-binding initiation factor by eIF3 subunit e. *Genes Dev* **28**: 835–840
- Werner JE, Finkelstein RR** (1995) Arabidopsis mutants with reduced response to NaCl and osmotic stress. *Physiol Plant* **93**: 659–666
- Wigge PA, Kim MC, Jaeger KE, Busch W, Schmid M, Lohmann JU, Weigel D** (2005) Integration of spatial and temporal information during floral induction in Arabidopsis. *Science* **309**: 1056–1059
- Zhu JK** (2002) Salt and drought stress signal transduction in plants. *Annu Rev Plant Biol* **53**: 247–273