

Continental-scale distributions of dust-associated bacteria and fungi

Albert Barberán^a, Joshua Ladau^b, Jonathan W. Leff^{a,c}, Katherine S. Pollard^b, Holly L. Menninger^d, Robert R. Dunn^{d,e}, and Noah Fierer^{a,c,1}

^aCooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO 80309; ^bGladstone Institutes, University of California, San Francisco, CA 94158; ^cDepartment of Ecology and Evolutionary Biology, University of Colorado, Boulder, CO 80309; ^dDepartment of Biological Sciences, North Carolina State University, Raleigh, NC 27695; and ^eCenter for Macroecology, Evolution and Climate, Natural History Museum of Denmark, University of Copenhagen, Copenhagen DK-2100, Denmark

Edited by William H. Schlesinger, Cary Institute of Ecosystem Studies, Millbrook, NY, and approved March 20, 2015 (received for review October 31, 2014)

It has been known for centuries that microorganisms are ubiquitous in the atmosphere, where they are capable of long-distance dispersal. Likewise, it is well-established that these airborne bacteria and fungi can have myriad effects on human health, as well as the health of plants and livestock. However, we have a limited understanding of how these airborne communities vary across different geographic regions or the factors that structure the geographic patterns of near-surface microbes across large spatial scales. We collected dust samples from the external surfaces of ~1,200 households located across the United States to understand the continental-scale distributions of bacteria and fungi in the near-surface atmosphere. The microbial communities were highly variable in composition across the United States, but the geographic patterns could be explained by climatic and soil variables, with coastal regions of the United States sharing similar airborne microbial communities. Although people living in more urbanized areas were not found to be exposed to distinct outdoor air microbial communities compared with those living in more rural areas, our results do suggest that urbanization leads to homogenization of the airborne microbiota, with more urban communities exhibiting less continental-scale geographic variability than more rural areas. These results provide our first insight into the continental-scale distributions of airborne microbes, which is information that could be used to identify likely associations between microbial exposures in outdoor air and incidences of disease in crops, livestock, and humans.

aerobiology | microbial ecology | microbial dispersal | urbanization | allergens

For nearly 2 centuries, we have known that microbes are ubiquitous in dust and outdoor air (1–3). In the near-surface atmosphere, microbial cells likely account for a significant fraction of aerosolized organic carbon (4), with microbial cell numbers typically ranging from 10^4 to 10^6 cells·m⁻³ over land (5). This means we inhale thousands of microbial cells every hour spent outdoors, and the potential effects of these airborne microbes on human health and the health of plants and animals are well recognized. As just one example, there are currently ~16 million people living in the United States suffering from allergic asthma (6), and there has been considerable attention focused on understanding the airborne and dust-associated microbes influencing asthma and how those microbial triggers vary across space and time (6–9). In addition, the number of virulent fungal infections affecting human populations, wildlife, and plant crops is increasing (10, 11). The effects of those microbes capable of atmospheric transport can even extend to entire ecosystems: Microbes in Saharan dust clouds, for example, have been shown to affect the ecology of alpine lakes in Spain (12) and coral reefs in the Caribbean Sea (13).

Despite the well-recognized importance of airborne microbes and the long history of research on microbial transport through the atmosphere (3), we have only recently been able to describe

the full extent of microbial diversity found in the atmosphere by using molecular approaches to characterize the microbial taxa that are difficult to identify via cultivation-dependent or microscopy-based surveys. Such molecular approaches not only have yielded new insight into the enormous diversity of airborne microbes (14, 15) but also have been instrumental in helping us understand how the composition of the airborne microbial communities varies across time and space (16, 17). However, nearly all of this work has focused on local-scale variability, examining the bacterial or fungal communities found in outdoor air at selected sites. What is missing is a continental-scale understanding of microbial diversity in the near-surface atmosphere and its deposition patterns over land. We do not know whether there are distinct microbial taxa found in the outdoor air from different geographic regions, nor do we know what biotic and abiotic factors may be driving the geographic patterns of dust-associated microbial communities across larger spatial scales. There is some evidence to suggest there are regional differences in exposures to specific bacterial and fungal taxa (18) that may be associated with geographic patterns in allergic disease (19), but the continental-scale biogeographic patterns exhibited by the broad range of microbes that can be found in outdoor air remain undetermined.

Here we used dust samples collected from the external surfaces of ~1,200 homes located across the continental United States to gain our first insight into the continental-scale patterns of bacterial and fungal diversity in the near-surface atmosphere

Significance

We inhale thousands of microbial cells when we breathe in outdoor air, and some of these airborne microbes can serve as pathogens or triggers of allergic disorders. Using settled dust samples from ~1,200 locations, we generated the first atlas, to our knowledge, of airborne bacterial and fungal distributions across the continental United States. We found that airborne microbial communities, such as terrestrial plants and animals, exhibit nonrandom geographic patterns, and we identified the factors that shape the continental-scale distributions of microbial taxa. Furthermore, we found that the airborne microbes found in urban and more rural areas are not distinct in composition, but the dust-associated communities found in more urbanized areas are more homogeneous across the United States.

Author contributions: H.L.M., R.R.D., and N.F. designed research; A.B., J.L., and J.W.L. analyzed data; and A.B., K.S.P., R.R.D., and N.F. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Data deposition: The data reported in this paper have been deposited in the Figshare data repository (figshare.com/articles/1000homes/1270900) and in the National Center for Biotechnology Information database (BioProject ID [PRJNA280363](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA280363)).

¹To whom correspondence should be addressed. Email: noah.fierer@colorado.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1420815112/-DCSupplemental.

this genus include major plant pathogens. Together, these results highlight that many microbial taxa exhibit local- and regional-scale geographic distribution patterns and that we can build range maps of bacterial and fungal taxa comparable to the maps long built by plant and animal ecologists to quantify biogeographic patterns and predict responses to climate change (30).

Our results indicate that the microbial communities found in outdoor air exhibit local and regional-scale geographic patterns, even though airborne microbes are known to disperse many thousands of kilometers (10, 12, 13, 31) and we might expect the complexities of wind patterns in the near-surface atmosphere to obscure the biogeographic distributions of microbes in settled dust. Some of the geographic patterns in airborne microbial communities could be driven by differences in dispersal capabilities. Not all taxa are equally adept at dispersal and survival on the sampled house surfaces, and this variation in the abilities of microbes to disperse to a given location and persist at that location could be an important mechanism generating the biogeographic patterns observed here (32). Likewise, it is important to mention that many local-scale factors that were not effectively captured with our data or statistical models could be responsible for generating some of the observed biogeographic patterns. For example, the types of vegetation found neighboring the homes or other local-scale sources of microbes to the atmosphere (e.g., construction sites, dirt roads) may be influencing the structure of these dust-associated communities at smaller spatial scales.

Effect of Urbanization. Urbanization tends to homogenize plant and animal communities, with more urbanized areas tending to share a large number of plant and animal taxa in common (33). At the same time, urbanization promotes the establishment of nonnative plant and animal species, and thus, cities can often have higher species richness than surrounding rural areas (e.g., refs. 34, 35). We do not know how urbanization affects airborne microbial communities, with some studies suggesting microbial diversity should be lower in more urbanized areas (8) and other studies showing little to no effect of urbanization on microbial diversity (17). This question is important, given that rates of allergenic asthma are higher for the children living in more urbanized areas (reviewed in ref. 9). Although many factors contribute to allergenic disease (19), it has been hypothesized that these geographic differences in allergy rates can be attributed to people living in more urbanized areas being exposed to lower levels of microbial diversity (7, 8). However, at least outdoors, we found no significant effect of urbanization on bacterial or fungal diversity levels ($P > 0.05$, Mann-Whitney test; *SI Appendix*, Fig. S9). Likewise, we found that more urbanized sites did not have microbial communities compositionally distinct from those found in less-urbanized or rural locations ($R^2 < 0.005$ for both bacterial and fungal communities, permutational analysis of variance (PERMANOVA); *SI Appendix*, Fig. S9). We also did not observe any effect of urbanization on bacterial or fungal community composition when we treated urbanization as a quantitative variable (i.e., population density, $R^2 < 0.005$, PERMANOVA), rather than a qualitative one (i.e., urban and rural classification).

Although we observed no significant effects of urbanization on the overall diversity and composition of dust-associated microbial communities, we expected the specific sources of airborne microbes to vary across urban to rural gradients (24). To test this hypothesis, we identified specific bacterial taxa indicative of specific source environments from which dust-associated microbes are most likely to originate (*SI Appendix*, Table S2), including soil, marine waters, feces and skin of humans or other animals, or plants. On the basis of these source assignments, we could then test how urbanization affects the relative importance of specific sources of bacteria to the atmosphere. For example, we would expect fecal bacteria to be more abundant in rural areas close to high densities of livestock (24, 36), and skin-associated microbes to be more common in cities and populated areas. However, we

did not detect any strong or significant effect of urbanization on the proportion of different bacterial indicator taxa (no difference greater than 1% or $P < 0.05$, Mann-Whitney test; Fig. 3). Overall, bacterial communities in the dust samples were primarily derived from bacteria likely to be found on plants and soil, regardless of the degree of urbanization (Fig. 3), a finding in line with other work showing that these sources are the dominant contributors of bacteria to the near-surface atmosphere in most terrestrial environments (13, 15). The imprint of human-associated bacteria was low and primarily a result of skin, rather than stool (with the exception of a single rural sample that was dominated by stool indicator taxa, mainly *Bacteroides* and *Parabacteroides*; Fig. 3). Neither marine waters (and potentially other water bodies including lakes or rivers) nor insects appeared to be important sources of bacteria of the atmosphere (Fig. 3). In the case of the few samples (particularly from urban areas) in which we observed high abundances of bacteria indicative of insects (notably *Rickettsiella*, Entomoplasmatales, *Buchnera*, and *Wolbachia*), these extreme values might be a result of the presence of dead insects (or fragments/depositions of insects) on the outdoor frame of those homes. Although we can demonstrate the importance of plants and soils as sources of microbes in settled dust, our data suggest that people living in more rural areas are not exposed outdoors to bacteria derived from different source environments than those living in more urbanized areas.

However, in line with the patterns commonly observed for plant and animal communities (33), the bacterial and fungal communities found in the dust from more urbanized locations tended to be more homogeneous. In other words, the microbial communities found in cities were not as spatially variable across the continental United States as in those communities found in more rural areas (Fig. 4). People living in more urbanized areas are not likely exposed to different airborne bacteria and fungi outdoors than those living in more rural areas, but urban areas tend to harbor microbial communities that are less spatially




Fig. 3. Proportion of bacterial sequences identified as indicator taxa of potential source environments in urban and rural samples. Chloroplasts were used as indicators of plant influence. Source environments are ordered from left to right and top to bottom according to their median value (medians are shown as horizontal lines). Note that because of the highly skewed distribution, the y axis is squared.




Fig. 4. Relationship between community dissimilarity and spatial distance for (A) bacterial and (B) fungal urban and rural samples. The slopes were significantly different in both cases (bacteria urban = $0.9 \cdot 10^{-5} \pm 1.1 \cdot 10^{-7}$; bacteria rural = $1.7 \cdot 10^{-5} \pm 3.9 \cdot 10^{-7}$; fungi urban = $2.6 \cdot 10^{-5} \pm 1.3 \cdot 10^{-7}$; fungi rural = $4.1 \cdot 10^{-5} \pm 4.4 \cdot 10^{-7}$).

variable than those found in more rural areas. This observed homogenization of dust-associated microbial communities might be a result of a homogenization of environmental conditions and land-use types in urban areas (33) or the homogenization in plant communities or other potential biotic sources of bacteria and fungi to the atmosphere.

Methods

Sample Collection. Outdoor dust samples were collected by volunteers participating in the Wild Life of Our Homes project (homes.yourwildlife.org), a continental-scale citizen science project. We recruited participants from all 50 states and the District of Columbia through our website, social media, and email campaigns over the period from January 2012 to March 2013. Enrolled participants ($n = 1,430$) were provided a written informed consent form approved by the North Carolina State University's Human Research Committee (approval no. 2177), as well as instructions for sampling and a microbe sampling kit containing dual-tipped sterile BBL CultureSwabs. Here we focus on a subset of samples that participants collected from the upper door trim on the outside surface of an exterior door, a sampling location that is found in every home, is unlikely to be cleaned frequently, and serves as a passive collector of outdoor aerosols and dust with little to no direct contact from the home occupants. The amount of time dust has accumulated on these surfaces is unknown and likely variable across homes, so we cannot use this sample set to assess temporal variation in the microbial communities. Participants returned swabs by first-class mail over the period from March 2012 to May 2013, and these swabs were stored in a -20°C freezer until processed. A map of sampling locations and information on the number of samples collected per US state are provided in *SI Appendix, Fig. S1*.

Molecular Analyses. Swabs were prepared for sequencing using the direct PCR approach described previously (37). Swab tips were placed directly into 2-mL 96-well plates (Axygen Inc.), along with the appropriate negative control samples. Plates were processed using the Extract-N-Amp PCR kit (Sigma-Aldrich, Inc.), following a modified version of the manufacturers' instructions. After each well received 250 μL of the Extract-N-Amp Extraction solution, the plate was sealed securely with a 96-round well Impermat Silicon Sealing Mat (Axygen, Inc.) and heated at 90°C for 10 min in a dry bath. Extract-N-Amp Dilution solution was then added to the wells at a 1:1 ratio to the extraction solution and mixed gently by pipetting. The plate was resealed with the mat and stored at 4°C . PCR was conducted in 20- μL triplicate reactions per sample, using 10 μL Extract-N-Amp Ready Mix, 1 μL of the forward and reverse primers, 5 μL PCR-grade water, and 4 μL of the Extract-N-Amp sample solutions from the 96-well plate.

Microbial diversity was assessed using high-throughput sequencing methods to characterize the variation in marker gene sequences. For bacterial and archaeal analyses, we sequenced the V4 hypervariable region of the 16S rRNA gene using the 515-F (GTGCCAGCAGCGCGGTA) and 806-R (GGA-CTACHVGGTWTCTAAT) primer pair (26). For the fungal analyses, we sequenced the first internal transcribed spacer (ITS1) region of the rRNA operon, using the ITS1-F (CTTGTCATTAGAGGAAGTAA) and ITS2 (GCTGCTTCTT-CATCGATGC) primer pair (38). The primers included the appropriate Illumina adapters, with the reverse primers also having an error-correcting 12-bp

barcode unique to each sample to permit multiplexing of samples. Archaeal sequences represented a very small proportion of the recovered sequences (0.13% of the 16S rRNA gene sequences). PCR products from all samples were quantified using the PicoGreen dsDNA assay and pooled together in equimolar concentrations for sequencing on either an Illumina HiSeq or MiSeq instrument. All sequencing runs were completed at the University of Colorado Next Generation Sequencing Facility.

Sequence Processing. The 100-bp sequences were demultiplexed using a custom Python script (<https://github.com/leffj/helper-code-for-uparse>), with quality filtering and phylotype clustering conducted using the UPARSE pipeline (39). For quality filtering, we used a maximum e-value of 0.5 (indicating that on average, a maximum of 0.5 nucleotides were incorrectly assigned in every sequence). Sequences were also dereplicated, and singleton sequences were removed before phylotype determinations. Representative sequences from the phylotypes that were not $\geq 75\%$ similar to sequences contained in either the Greengenes 13.8 database (40) or the UNITE May 2014 database (41) for 16S and ITS rRNA sequences, respectively, were discarded. Raw sequences were then mapped to phylotypes at the 97% similarity threshold. Phylotype taxonomy was determined using the Ribosomal Database Project classifier with a confidence threshold of 0.5 (42), trained on the respective databases for 16S and ITS rRNA sequence. 16S rRNA sequences resulting from direct PCR reagent contaminants (37) were filtered from all samples by removing those classified as *Mycoplasma*, *Pseudomonas*, and *Serratia*, which represented a median 59%, 9%, and 4% of sequences across the controls, respectively. In addition, sequences representing any phylotypes present in $\geq 25\%$ of control samples and those classified as mitochondria or chloroplast were removed (except during analysis of plant presence described here).

To remove potential amplicon sequencing biases, we first removed samples with fewer than 10,000 sequences, and then we normalized the sequence counts using a cumulative-sum scaling, which scales counts by dividing up to a percentile empirically determined (43). Richness and community similarity values after normalization were highly correlated, with values obtained after rarefying to 10,000 sequences per sample (*SI Appendix, Fig. S2*). The total number of samples included in downstream analyses was 1,187 for bacteria and 1,289 for fungi.

Sample Characterization. Latitude and longitude coordinates were derived from location information (address), and the coordinates were used to obtain georeferenced variables for each household. We compiled a set of 31 descriptors: elevation, distance to the coast, climatic variables, plant productivity, soil factors, other organisms' (including humans and livestock) variables, and house characteristics (*SI Appendix, Table S1*). Urban and rural classifications for all sampled locations were obtained from the 2010 US Census Bureau data (www.census.gov). Urban areas are classified as densely developed ($>50,000$ people) residential and commercial territories; 28% of the sampled locations were classified as rural.

Identification of Bacterial Source Indicators. To identify potential taxa indicative of specific source environments (i.e., soil, oceans, human skin, and human/livestock feces), we used Indicator Value analyses (44), as described in ref. 22. We used the proportion of chloroplast sequences in each sample as a proxy for plant presence. To identify those bacteria likely derived from insects, we compiled a list of exclusive symbiont and/or insect pathogens from the literature (45, 46). For the complete list of bacterial indicator taxa, see *SI Appendix, Table S2*.

Statistical Analyses. Microbial community similarity was represented by non-metric multidimensional scaling, using the Bray-Curtis dissimilarity distance metric after Hellinger standardization (47). Bray-Curtis abundance-based dissimilarity matrices were highly correlated with Jaccard incidence-based matrices [$r_M = 0.86$ ($P < 0.001$); $r_M = 0.92$ ($P < 0.001$); Mantel test for bacteria and fungi, respectively]. As some of the descriptors were correlated, we reduced collinearity before our analyses by identifying highly correlated variables by principal component analyses and by examining the variance inflation factor (48). The final set of factors consisted of 18 variables (*SI Appendix, Table S1*). To find the best subset of explanatory variables, we first ran bioenv (49) to maximize the correlation with community dissimilarity. We then performed MRM to estimate the overall explanatory power of the model (50). To test whether community composition was different between urban and rural areas, we used PERMANOVA based on 1,000 permutations (51).

All multivariate statistical analyses were carried out in the R environment (www.r-project.org), using the vegan (vegan.r-forge.r-project.org/), labdsv (ecology.msu.montana.edu/labdsV/), and ecodist (cran.r-project.org/web/)

packages/ecodist/) packages. Nonmetric multidimensional scaling axis scores and the proportional abundances of taxa were mapped by inverse distance weighting interpolation on 100 × 100 grid cells, using the gstat package (<https://r-forge.r-project.org/projects/gstat/>).

Conclusion

We show that the continental-scale distributions of bacteria and fungi in settled dust are correlated with regional environmental factors. These geographic patterns are surprisingly robust, considering the potential for long-distance dispersal of microbes in aerosolized dust particles, the complexity of wind patterns in the near-surface atmosphere, and the difficulties associated with quantifying specific local-scale factors that may influence the types of bacteria and fungi that can be introduced into the atmosphere.

The health effects of this work remain unknown. We do know that bacterial and fungal exposures are distinct across the United States, and some of this variability can be explained by environmental conditions; namely, climatic and soil characteristics.

- Darwin C (1846) An account of the fine dust which often falls on vessels in the Atlantic ocean. *Q J Geol Soc* 2:26–30.
- Pasteur L (1861) Mémoire sur les corpuscules organisés qui existent dans l'atmosphère: Examen de la doctrine des générations spontanées. *Ann Sci Nat* 4e série:5–68.
- Tyndall J (1882) *Essays on the Floating-Matter of the Air* (D. Appleton and Company, New York).
- Wiedinmyer C, et al. (2009) The contribution of biological particles to observed particulate organic carbon at a remote high altitude site. *Atmos Environ* 43(28):4278–4282.
- Burrows SM, Elbert W, Lawrence MG, Pöschl U (2009) Bacteria in the global atmosphere- Part 1: Review and synthesis of literature data for different ecosystems. *Atmos Chem Phys Discuss* 9:9263–9280.
- Denning DW, O'Driscoll BR, Hogaboam CM, Bowyer P, Niven RM (2006) The link between fungi and severe asthma: A summary of the evidence. *Eur Respir J* 27(3):615–626.
- Ege MJ, et al.; GABRIELA Transregio 22 Study Group (2011) Exposure to environmental microorganisms and childhood asthma. *N Engl J Med* 364(8):701–709.
- Hanski I, et al. (2012) Environmental biodiversity, human microbiota, and allergy are interrelated. *Proc Natl Acad Sci USA* 109(21):8334–8339.
- von Mutius E, Verceci D (2010) Farm living: Effects on childhood asthma and allergy. *Nat Rev Immunol* 10(12):861–868.
- Brown JKM, Hovmöller MS (2002) Aerial dispersal of pathogens on the global and continental scales and its impact on plant disease. *Science* 297(5581):537–541.
- Fisher MC, et al. (2012) Emerging fungal threats to animal, plant and ecosystem health. *Nature* 484(7393):186–194.
- Barberán A, Henley J, Fierer N, Casamayor EO (2014) Structure, inter-annual recurrence, and global-scale connectivity of airborne microbial communities. *Sci Total Environ* 487:187–195.
- Prospero JM, Blades E, Mathison G, Naidu R (2005) Interhemispheric transport of viable fungi and bacteria from Africa to the Caribbean with soil dust. *Aerobiologia* 21: 1–19.
- Brodie EL, et al. (2007) Urban aerosols harbor diverse and dynamic bacterial populations. *Proc Natl Acad Sci USA* 104(1):299–304.
- Fröhlich-Nowoiski J, Pickersgill DA, Després VR, Pöschl U (2009) High diversity of fungi in air particulate matter. *Proc Natl Acad Sci USA* 106(31):12814–12819.
- Fierer N, et al. (2008) Short-term temporal variability in airborne bacterial and fungal populations. *Appl Environ Microbiol* 74(1):200–207.
- Bowers RM, McLetchie S, Knight R, Fierer N (2011) Spatial variability in airborne bacterial communities across land-use types and their relationship to the bacterial communities of potential source environments. *ISME J* 5(4):601–612.
- Amend AS, Seifert KA, Samson R, Bruns TD (2010) Indoor fungal composition is geographically patterned and more diverse in temperate zones than in the tropics. *Proc Natl Acad Sci USA* 107(31):13748–13753.
- Salo PM, et al. (2008) Exposure to multiple indoor allergens in US homes and its relationship to asthma. *J Allergy Clin Immunol* 121(3):678–684.
- Tong Y, Lighthart B (1999) Diurnal distribution of total and culturable atmospheric bacteria at a rural site. *Aerosol Sci Technol* 30(2):246–254.
- Shelton BG, Kirkland KH, Flanders WD, Morris GK (2002) Profiles of airborne fungi in buildings and outdoor environments in the United States. *Appl Environ Microbiol* 68(4):1743–1753.
- Dunn RR, Fierer N, Henley JB, Leff JW, Menninger HL (2013) Home life: Factors structuring the bacterial diversity found within and between homes. *PLoS ONE* 8(5): e64133.
- Blanchard DC, Szydek L (1970) Mechanism for the water-to-air transfer and concentration of bacteria. *Science* 170(3958):626–628.
- Bowers RM, et al. (2011) Sources of bacteria in outdoor air across cities in the mid-western United States. *Appl Environ Microbiol* 77(18):6350–6356.
- Lauber CL, Hamady M, Knight R, Fierer N (2009) Pyrosequencing-based assessment of soil pH as a predictor of soil bacterial community structure at the continental scale. *Appl Environ Microbiol* 75(15):5111–5120.

We also know that allergen sensitivities and incidences of asthma, including allergen sensitivities to particular microbial triggers, can vary across the United States (28). Our study was not designed to collect specific allergen sensitivity data for the individuals living in each home, but clearly there are opportunities for future work investigating how the diversity, distribution, and composition of these dust-associated bacteria and fungi may shape incidences of disease across the United States.

ACKNOWLEDGMENTS. We thank the many volunteers who participated in the Wild Life of Our Homes study for collecting home dust samples. Funding for this work was provided by a grant from the A.P. Sloan Microbiology of the Built Environment Program (to N.F. and R.R.D.), with additional support from the US National Science Foundation (DEB0953331 to N.F.). A.B. is supported by a James S. McDonnell Postdoctoral Fellowship. Support was also provided by the National Science Foundation (DMS-1069303 to K.S.P. and J.W.L.), the Gordon and Betty Moore Foundation (3300), the Gladstone Institutes, and a gift from the San Simeon Fund.

- Fierer N, et al. (2012) Cross-biome metagenomic analyses of soil microbial communities and their functional attributes. *Proc Natl Acad Sci USA* 109(52):21390–21395.
- Lee CK, Barbier BA, Bottos EM, McDonald IR, Cary SC (2012) The Inter-Valley Soil Comparative Survey: The ecology of Dry Valley edaphic microbial communities. *ISME J* 6(5):1046–1057.
- Salo PM, et al. (2006) Exposure to *Alternaria alternata* in US homes is associated with asthma symptoms. *J Allergy Clin Immunol* 118(4):892–898.
- Adams RI, Miletto M, Taylor JW, Bruns TD (2013) The diversity and distribution of fungi on residential surfaces. *PLoS ONE* 8(11):e78866.
- MacArthur RH (1972) *Geographical Ecology: Patterns in the Distribution of Species* (Harper & Rowe Publishers, New York).
- Bovallius A, Bucht B, Roffey R, Anäs P (1978) Long-range air transmission of bacteria. *Appl Environ Microbiol* 35(6):1231–1232.
- Peay KG, Schubert MG, Nguyen NH, Bruns TD (2012) Measuring ectomycorrhizal fungal dispersal: Macroecological patterns driven by microscopic propagules. *Mol Ecol* 21(16):4122–4136.
- McKinney ML (2006) Urbanization as a major cause of biotic homogenization. *Biol Conserv* 127(3):247–260.
- Hope D, et al. (2003) Socioeconomics drive urban plant diversity. *Proc Natl Acad Sci USA* 100(15):8788–8792.
- Kühn I, Brandl R, Klotz S (2004) The flora of German cities is naturally species rich. *Evol Ecol Res* 6:749–764.
- Bakutis B, Monstvilienė E, Januskeviciene G (2004) Analyses of airborne contamination with bacteria, endotoxins and dust in livestock barns and poultry houses. *Acta Vet (Brno)* 73(2):283–289.
- Flores GE, Henley JB, Fierer N (2012) A direct PCR approach to accelerate analyses of human-associated microbial communities. *PLoS ONE* 7(9):e44563.
- McGuire KL, et al. (2013) Digging the New York City Skyline: Soil fungal communities in green roofs and city parks. *PLoS ONE* 8(3):e58020.
- Edgar RC (2013) UPARSE: Highly accurate OTU sequences from microbial amplicon reads. *Nat Methods* 10(10):996–998.
- McDonald D, et al. (2012) An improved Greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria and archaea. *ISME J* 6(3):610–618.
- Abarenkov K, et al. (2010) The UNITE database for molecular identification of fungi—recent updates and future perspectives. *New Phytol* 186(2):281–285.
- Wang Q, Garrity GM, Tiedje JM, Cole JR (2007) Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol* 73(16):5261–5267.
- Paulson JN, Stine OC, Bravo HC, Pop M (2013) Differential abundance analysis for microbial marker-gene surveys. *Nat Methods* 10(12):1200–1202.
- Dufrène M, Legendre P (1997) Species assemblages and indicator species: The need for a flexible asymmetrical approach. *Ecol Monogr* 67(3):345–366.
- Feldhaar H (2011) Bacterial symbioses as mediators of ecologically important traits of insect hosts. *Ecol Entomol* 36(5):533–543.
- Engel P, Moran NA (2013) The gut microbiota of insects - diversity in structure and function. *FEMS Microbiol Rev* 37(5):699–735.
- Legendre P, Gallagher E (2001) Ecologically meaningful transformations for ordination of species data. *Oecologia* 129(2):271–280.
- Dormann CF, et al. (2013) Collinearity: A review of methods to deal with it and a simulation study evaluating their performance. *Ecography* 36(1):27–46.
- Clarke K, Ainsworth M (1993) A method of linking multivariate community structure to environmental variables. *Mar Ecol Prog Ser* 92:205–219.
- Lichstein JW (2006) Multiple regression on distance matrices: A multivariate spatial analysis tool. *Plant Ecol* 188(2):117–131.
- Anderson MJ (2001) A new method for non-parametric multivariate analysis of variance. *Austral Ecol* 26(1):32–46.