

TECHNICAL NOTES

Open Access

Dynamic regulation of gene expression using sucrose responsive promoters and RNA interference in *Saccharomyces cerevisiae*

Thomas C Williams, Monica I Espinosa, Lars K Nielsen and Claudia E Vickers*

Abstract

Background: Engineering dynamic, environmentally- and temporally-responsive control of gene expression is one of the principle objectives in the field of synthetic biology. Dynamic regulation is desirable because many engineered functions conflict with endogenous processes which have evolved to facilitate growth and survival, and minimising conflict between growth and production phases can improve product titres in microbial cell factories. There are a limited number of mechanisms that enable dynamic regulation in yeast, and fewer still that are appropriate for application in an industrial setting.

Results: To address this problem we have identified promoters that are repressed during growth on glucose, and activated during growth on sucrose. Catabolite repression and preferential glucose utilisation allows active growth on glucose before switching to production on sucrose. Using sucrose as an activator of gene expression circumvents the need for expensive inducer compounds and enables gene expression to be triggered during growth on a fermentable, high energy-yield carbon source. The ability to fine-tune the timing and population density at which gene expression is activated from the *SUC2* promoter was demonstrated by varying the ratio of glucose to sucrose in the growth medium. Finally, we demonstrated that the system could also be used to repress gene expression (a process also required for many engineering projects). We used the glucose/sucrose system to control a heterologous RNA interference module and dynamically repress the expression of a constitutively regulated GFP gene.

Conclusions: The low noise levels and high dynamic range of the *SUC2* promoter make it a promising option for implementing dynamic regulation in yeast. The capacity to repress gene expression using RNA interference makes the system highly versatile, with great potential for metabolic engineering applications.

Keywords: Yeast, Sucrose, GFP, Metabolic engineering, Synthetic biology, Promoter, *SUC2*, *TEF1*, Diauxic shift

Background

Microorganisms possess a variety of mechanisms for modulating gene expression levels according to changes in their environment. The ability to engineer dynamic regulation is highly desirable in microbial cell factories because many of the most productive genetic modifications used for optimising engineering objectives are also detrimental to cell growth and survival. Engineered pathways compete for the carbon flux, redox potential, and ATP required by the cell for normal growth [1]. Engineered pathways can also result in accumulation of

intermediates, side-products or end-products that are highly toxic to the host cell [2]. If a production strain cannot reach a high population density due to the metabolic burden imposed by an engineered pathway, then product titers are inherently limited. For genetic modifications that impose the most severe limits on growth, it is essential that they be implemented near or after the completion of a growth phase. One of the most promising techniques to optimise cell factory performance is to control the expression/repression of relevant genes using dynamic regulation.

S. cerevisiae (yeast) is an industrial microorganism which has been employed for the production of fuels, chemicals, and pharmaceuticals [3]. There are a limited number of molecular tools available for implementing

* Correspondence: c.vickers@uq.edu.au
Australian Institute for Bioengineering and Nanotechnology (AIBN), The University of Queensland, St. Lucia, QLD 4072, Australia

dynamic regulation in yeast, with most being inadequate for industrial application. For example, the commonly used galactose induction system [4] has a noise level (expression in the absence of inducer) which is unacceptable for some applications such as controlling RNA interference [5]. Furthermore, galactose is prohibitively expensive for use at an industrial scale [6]. Previous efforts to implement dynamic regulation under industrially relevant conditions required the knockout of galactose utilization genes so that a small amount of galactose can be added to fermentations as a gratuitous inducer for gene expression [6]. High concentrations of fermentable carbon sources repress gene expression from galactose promoters (GAL) via a carbon catabolite repression mechanism [7], and galactose utilisation genes are not activated during growth on sucrose [8]. This means that cheap, fermentable sugars such as glucose or sucrose cannot be used during a production phase when the GAL promoters are used to achieve dynamic regulation. To circumvent this limitation, ethanol has been used as a carbon source for fed-batch cultures with the GAL promoters [6]. However, this is also problematic because ethanol is more expensive than commonly used sugar feedstocks, and is in fact a common commercial product of industrial yeast fermentations. Other induction systems such as the doxycycline inducible promoters have the advantage of being completely orthologous to native yeast regulation [9], but are also too expensive to be employed at a large scale. There are many other carbon source regulated promoters in yeast that could potentially be used for dynamic regulation [10], including the *ADH2* promoter that is activated during growth on ethanol, the low-oxygen regulated *DANI* promoter, and the low phosphate activated *PHO5* promoter [11]. These options all rely on the absence of a growth component that enables maximal metabolic flux and gene expression capacity (such as glucose or sucrose). There is therefore a significant unmet need for control systems that enable dynamic regulation and have suitable properties for industrial use of cell factories such as yeast. An ideal promoter would have low noise, high dynamic range, switch-like activation, high absolute expression levels, sustained induction, and high activity on carbon sources that support high metabolic flux.

To expand the toolbox of dynamic regulatory systems in yeast we have explored the use of promoters regulated by glucose de-repression in the presence of sucrose. Sucrose is a preferred feedstock for industrial scale fermentations due to sugarcane being a more environmentally friendly source of sugar than glucose which has been derived from corn [12,13]. Promoters that can regulate high levels of gene expression during a sucrose-fed production phase are therefore highly desirable. We

sought to identify and characterise the spatiotemporal expression dynamics of promoters which are repressed during growth on glucose, and up-regulated during growth on sucrose. We applied these promoters to dynamic over-expression, and also to dynamic repression via a heterologous RNA interference module that can be used to destroy target mRNA according to base pair complementarity of expressed antisense RNA [5].


Results and discussion

Screening for sucrose responsive promoters

The most important properties to consider when assessing mechanisms for dynamic regulation are the dynamic range (induced minus non-induced expression levels) and noise levels (non-induced minus background). Four different promoters were screened with the aim of achieving low expression levels in the presence of glucose, and high expression in the presence of sucrose. Many promoters are known to be de-repressed in the presence of low glucose in yeast [10], little or no published data is available describing how these promoters behave on sucrose as the sole carbon source. It therefore could not be assumed that promoters known to be de-repressed by low glucose would be active on sucrose. For metabolic engineering applications in yeast it is desirable to have a production phase with a cheap carbon source. Promoters which are not only de-repressed by low glucose, but that are activated during growth on the cheap industrial carbon source sucrose are therefore highly desirable for industrial yeast fermentations.

Four promoters were selected based on literature which suggested they may be differentially regulated by sucrose. *SUC2p* is an invertase enzyme that hydrolyses sucrose into glucose and fructose; expression of *SUC2p* is strongly repressed by glucose [10,14]. Transcription of *SUC2* is repressed around 200-fold when glucose levels are high and is activated only after glucose levels fall below 0.1% w/v [15-18]. The *MAL12* gene encodes a disaccharide transporter which is highly induced in the presence of sucrose in non-laboratory strains of *S. cerevisiae* [19,20]. *Glc3p* is a glycogen branching enzyme essential for glycogen accumulation, and the *Gph1p* enzyme breaks down glycogen during stationary phase [21-23]. Both *GLC3* and *GPH1* transcripts have been observed to be up-regulated during growth on sucrose [24].

Yeast strains were generated with promoter-GFP fusion constructs for the promoters *P_{SUC2}*, *P_{MAL12}*, *P_{GLC3}*, or *P_{GPH1}* integrated into the genome at the *URA3* locus in single copy. Each strain was grown in glucose- or sucrose-containing medium with GFP fluorescence measured in the mid exponential phase (OD_{660nm} of ~1.5) after 8 hours of growth (Figure 1).



Each promoter had a significantly higher expression level on sucrose compared to on glucose. The *SUC2* promoter had both the highest dynamic range (12.5-fold induction), and the highest absolute expression level. These properties made the *SUC2* promoter an attractive choice for sucrose-mediated dynamic regulation in *S. cerevisiae*. Although this experiment was successful in identifying the best promoter, it only provided a ‘snapshot’ of expression levels during the exponential growth phase. In order to gain a deeper understanding of noise levels and dynamic range for comparison with other published data it is necessary to understand the spatiotemporal dynamics of gene expression throughout a fermentation.

Fine-tuning temporal gene expression dynamics using different glucose:sucrose ratios

Implementation of sucrose-mediated dynamic regulation would involve use of a growth medium which contains a known ratio of glucose to sucrose. In theory this would result in P_{SUC2} -regulated genes being tightly repressed during a growth phase where glucose is preferentially consumed, then switched ‘on’ during a sucrose-consuming phase. When growth media with different ratios of glucose to sucrose were used to grow a P_{SUC2} -GFP expressing strain, the timing of GFP induction responded according to glucose concentration (Figure 2a-d). A destabilized version of the GFP gene with a half-life of ~20 minutes (γ EGFPCLN2PEST [25]) was used so that any decreases in gene expression level would not be obscured by the high stability (half-life of ~7 hours) of the regular γ EGFP gene.

In addition to the fine-tuning of gene expression, the time-course analysis of *SUC2* promoter activity was necessary to gain more accurate insight into expression levels and dynamic range. As expected, GFP expression from the *SUC2* promoter was strongly repressed during growth on glucose under all conditions. GFP expression levels during the glucose consumption phase were usually indistinguishable from autofluorescence, indicating tight regulation and a lack of undesirable leakiness. Once glucose concentrations dropped below about 5 mM, there was a rapid induction of GFP expression as the *SUC2* promoter was activated. GFP levels from the *SUC2* promoter peaked immediately after the switch from glucose to sucrose at around 4000 au before decreasing to 700–1000 au (Figure 2). This is consistent with the *SUC2* expression level and dynamic range seen in the promoter screen, where populations had been growing on sucrose-only medium for 8 hours and would have progressed past the initial induction phase (Figure 1). As previously reported, the *SUC2* promoter was activated after glucose was depleted even in the absence of sucrose [26]. When sucrose was present in the growth medium (Figure 2a-c) there was a 2 fold higher GFP expression level (~4000 au) (Figure 2a-c) compared to when sucrose was absent (~2000 au) (Figure 2d). As far as we are aware there is no known regulatory mechanism that can account for the increased *SUC2* mediated gene expression in the presence of sucrose compared to in the absence of glucose (during the diauxic shift). It is possible that the phenomenon shown in Figure 2d could simply reflect a greater capacity for gene expression during exponential growth on fermentable carbon sources compared to during the diauxic shift and entry into the stationary phase [27].

It was possible to implement a very fine level of control over the timing and population density at which GFP expression was activated by varying glucose to sucrose ratios in the growth media. For example, with 0.5% glucose and 1.5% sucrose, GFP was fully induced after 5 hours of growth, at an OD_{660nm} of ~2 (Figure 2a). With 1% glucose and 1% sucrose the switch occurred after 9 hours, at an OD_{660nm} of ~6.4 (Figure 2b). When grown in medium containing 1.5% glucose and 0.5% sucrose, GFP expression was activated at 10 hours with a population density of ~7.3 (OD_{660nm}), peaking after 11 hours at OD 8 (Figure 2c). With 2% glucose, and no sucrose in the growth medium, GFP expression was activated after 12 hours at an OD_{660nm} of ~9 (Figure 2d). These variations demonstrate the fine-tuning of gene expression afforded by this fully autonomous and cheap induction system.

For application to metabolic engineering scenarios, it is important that any dynamic regulatory system has a sustained level of induction throughout cultivation.




Figure 2 (See legend on next page.)

(See figure on previous page.)

Figure 2 Tuning the timing of gene expression using different ratios of glucose to sucrose. A strain expressing the destabilized GFP gene driven by the *SUC2* promoter was grown in media containing the indicated concentrations of glucose and sucrose. Extracellular glucose, sucrose, and fructose concentrations were measured using HPLC during the initial growth phase alongside GFP expression levels. Population density (OD_{660nm}) and GFP expression levels were measured up to 56 hours of shake-flask cultivation. All data points and error bars represent the mean and standard deviation from triplicate cultivations.

Measurement of GFP expression levels over 56 hours of shake-flask cultivation showed that expression from the *SUC2* promoter remains at approximately 800–1000 fold higher than during the glucose consumption phase (Figure 2). This is a high expression level and after de-repression has occurred, is comparable to the commonly used *TEF1* promoter (Figure 3). As a point of reference, the *GAL10* promoter has a dynamic range of over 1000 [28] and an expression level similar to the *TEF1* promoter [29]. The *TEF1* promoter has previously been noted for its stable, high level of expression [29] and it was interesting that we observed a ~10 fold decrease in *TEF1* promoter mediated GFP expression between the early exponential phase and the diauxic shift (Figure 3). This difference could possibly be due to the fact that previous observations relied on the beta-galactosidase reporter gene, which has a half-life of around 20 hours in *S. cerevisiae* [30]. In contrast, our system utilised a destabilized version of the γ EGFP gene that has a half-life of ~20 minutes [25], making it far more sensitive to any decreases in gene expression levels. The *TEF1* gene encodes a translational elongation factor that coordinates the positioning of aminoacylated tRNAs at ribosomes for the elongation of polypeptides [31]. Decreased expression from the *TEF1* promoter is consistent with the overall decrease in gene expression during and after the diauxic shift in *S. cerevisiae* [32,33].

Testing sucrose mediated regulation of RNA interference

Many of the most useful applications for dynamic regulatory systems require both the up-regulation and down-

regulation of gene expression. To address this issue we sought to integrate the sucrose mediated induction system with a recently developed *S. cerevisiae* RNA interference (RNAi) system [5,34,35]. *S. cerevisiae* does not have a native system for RNA interference, but when the Argonaute and Dicer genes from *Saccharomyces castelii* are expressed, mRNAs can be specifically targeted for degradation by expressing complementary full-length antisense RNA [35] (Figure 4a), or double-stranded hairpin RNA [5,34,35].

Previous efforts at implementing dynamic control of RNAi resulted in significant leaking of target-gene repression [5]. With the tight repression of expression from the *SUC2* promoter in the presence of glucose (Figure 2), we hypothesised that sucrose regulation would enable dynamic control of RNAi without non-induced repression occurring. The system was tested by expressing a destabilized GFP gene from the strong constitutive *TEF1* promoter, and targeting it for repression by expressing the entire GFP ORF in the antisense direction from the *SUC2* promoter in an RNAi capable strain (Argonaute and Dicer genes present) (Figure 4b). This strain was grown on medium containing a mixture of 1% glucose and 1% sucrose so that the *SUC2* promoter would activate GFP repression only after the glucose in the growth medium was consumed as previously demonstrated (Figure 2). GFP expression levels were normalised to a 'control' strain which was identical except for the absence of the GFP antisense construct that facilitates RNAi. During the glucose consumption phase (0–8 hr; Figure 2b) there was no significant difference





Figure 3 Comparison of TEF1 and SUC2 promoter strengths. GFP fluorescence (a) and population density (b) were measured for P_{TEF1} -GFP and P_{SUC2} -GFP expressing strains in 1% glucose, 1% sucrose containing medium over 48 hours. Mean and standard deviation for triplicate cultivations are shown.



between RNAi and control strains, suggesting that little or no non-specific repression was occurring. Between 11 and 13 hours, GFP levels dropped sharply, indicating rapid activation of the repression module, with fluorescence levels reduced to 30% of the control strain (Figure 4c). Repression was maintained with expression levels being further reduced to around 17% of the control strain by 56 hours. The two strains were cultivated in the same media and there was no difference in growth dynamics (Figure 4d).

Conclusions

The dynamic control of gene expression is a core component of synthetic biology, and has great potential for the improvement of microbial cell factories. Dynamic

control can be used to optimise a trade-off between growth based physiology and an organism's engineered function [36]. By using dynamic regulation, any engineered functions that compete with normal growth-based physiology can be delayed until a population has reached a high density. Such functions are common, as even the most basic applications such as heterologous protein production can draw resources such as ATP, translation machinery, and amino acids away from native processes. In metabolic engineering the diversion of carbon flux, redox power, and ATP towards the production of target metabolites can reduce or potentially eliminate growth [1,2]. In support of this concept the *SUC2* promoter has previously been used for high-level expression of amylase enzymes [37,38].

Although purified sucrose is not commonly used as an industrial carbon source, it is often hydrolysed into its constituent monosaccharides glucose and fructose. It is possible that the use of the *SUC2* induction system would reduce the need for pre-hydrolysis of sucrose in industrial processes. For future directions it would also be interesting to test the sucrose induction system on an industrial carbon source such as sugarcane syrup at a much higher population density.

We present a simple dynamic regulatory system which does not require the addition of any expensive inducer compounds to the media, has a high dynamic range, low noise levels, maintains expression for prolonged periods, and can be coupled to RNAi to enable the repression of gene expression. We predict that this system will be of great utility for a diverse array of future applications in yeast cell factories.

Plasmids P_{TEF1} -yEGFPCLN2PEST-pRS406, P_{SUC2} -yEGFPCLN2PEST-pRS406, and P_{SUC2} -GFPantisense-CYC1t-pRS413 will be made available from AddGene (www.addgene.org/).

Materials and methods

Media

Strain pre-cultures were grown in chemically defined liquid CBS medium with 5 g/L ammonium sulfate, 20 g/L glucose, vitamins and trace elements as described previously [39]. For GFP expression experiments CBS media with the indicated concentrations of glucose and/or sucrose was used. During strain construction purified amino acids (Sigma) were used to complement appropriate auxotrophies in agar plates (same composition as chemically defined media above) while YPD supplemented with appropriate antibiotics was used during gene deletion procedures. *E. coli* DH5 α was used for plasmid propagation/storage and was grown in LB medium with kanamycin.

Strains and plasmids

Primers, plasmids, and strains used in this study are shown in (Table 1, Table 2, and Table 3), respectively. Annotated vector maps and sequences can be found in 'Additional file 2'. DNA manipulation and propagation were carried out using standard techniques [40] unless stated otherwise. All *S. cerevisiae* transformations were carried out using the lithium acetate method [41]. BAR1 and FUS1 genes were deleted in the base RNAi strain 'S01' as part of another related project which utilises pheromone quorum sensing for dynamic regulation (reference [42]), and are not important design features for this study. Strains transformed with yeast integrating plasmids were screened for correct single copy integration using PCR as previously described [43] except

Table 1 Primers

Primer name	5' to 3' sequence
MAL12F	TATTATctcgagACCAACCCGAAAATTCTTC
MAL12R	TATTATgaattcTTATGTAATTTAGTTACGCTTGAC
GPH1F	TATTATGatcgatTAGTTATCCGACTAGCAAG
GPH1R	TATTATgaattcTGTTCAAATAAAATTAAGTTG
GLC3F	TATTATctcgagCGGTGATTTACAAGAAGAGG
GLC3R	TATTATatcgatTTTATTCTTGACGGTCTTTATAC
SUC2F	TATTATctcgagACATACTAAGACATTTACCG
SUC2R	TATTATgaattcCATATACGTTAGTGAAGAAAAG
XhoI-pTEF1F	TATTATctcgagGCACACCCATAG
EcoRI-pTEF1R	TATTATgaattcTTGTAATTAATACTTAGATTAGATTG
GFP1F	TATTATgaattcCTATATTACTTGGGTATTGCC
GFP1R	TATTATcccgggTCTAAAGTGGAAGATTATTCAC
LEU2A2	ATAGAATTGTGTAGAATTGCAG
LEU2D2	ATGAAATGAACATTGATTTACTATC
CYC1tF	TATAATtctagaACAGGCCCTTTCTCTTTGT
CYC1tR	TATTATgagctcACGATGAGAGTGTAACTGC
Apal-SUC2F	TATTATgggcccACATACTAAGACATTTACCG
ClaI-SUC2R	TATTATatcgatCATATACGTTAGTGAAGAAAAG

Restriction enzyme sites are shown in lower case.

using primers LEU2A2 and LEU2D2 to screen to LEU2 locus.

All putative sucrose-responsive promoters were PCR amplified from CENP.K2-1c genomic DNA, and contain ~700 bp upstream of the start codon of the native gene. *MAL12* (XhoI/EcoRI), *GPH1* (ClaI/EcoRI), *GLC3* (XhoI/ClaI), and *SUC2* (XhoI/EcoRI) promoter regions were cloned 5' of the start codon in the GFPCLN2PEST-pRS406 plasmid using the indicated restriction enzyme combinations. The P_{TEF1} -yEGFPCLN2PEST-pRS406 plasmid was made by inserting the *TEF1* promoter amplified from pSF019 using primers 9 and 10 into the yEGFPCLN2PEST-pRS406 plasmid with XhoI/EcoRI.

The GFP silencing construct was expressed using the *SUC2* promoter on the low copy number pRS413 plasmid. To construct this expression cassette a *CYC1* terminator region was PCR amplified from genomic DNA using CYC1tF and CYC1tR primers and cloned into pRS413 using XbaI and SacI. The *SUC2* promoter was then amplified using Apal-SUC2F and ClaI-SUC2R and inserted 5' of *CYC1t* in pRS413 to create P_{SUC2} -CYC1t-pRS413. The GFP silencing construct was made by cloning the full length GFPCLN2PEST ORF sequence in the antisense direction between the *SUC2* promoter and *CYC1* terminator in P_{SUC2} -CYC1t-pRS413 using GFP1F/GFP1R primers and EcoRI/XmaI restriction enzymes to create the P_{SUC2} -GFP1i-CYC1t-pRS413 plasmid.

Table 2 Plasmids

Name	Details	Origin
pRS406	URA3 integrating vector	[45], Euroscarf
pSF019	TEF1 promoter containing vector	[29]
yEGFPCLN2PEST-pRS406	Destabilized GFP base plasmid	[36]
P _{MAL12} -yEGFPCLN2PEST-pRS406	MAL12 promoter driven GFP expression	This study
P _{GLC3} -yEGFPCLN2PEST-pRS406	GLC3 promoter driven GFP expression	This study
P _{GPH1} -yEGFPCLN2PEST-pRS406	GPH1 promoter driven GFP expression	This study
P _{SUC2} -yEGFPCLN2PEST-pRS406	SUC2 promoter driven GFP expression	This study
P _{TEF1} -yEGFPCLN2PEST-406	Constitutive TEF1 promoter driven GFP expression	This study
pRS413	Yeast centromeric plasmid with HIS3 selection marker	[45], Euroscarf
CYC1t-pRS413	CYC1 terminator	This study
P _{SUC2} -CYC1t-pRS413	SUC2 promoter, CYC1 terminator	This study
P _{SUC2} -GFPantisense-CYC1t-pRS413	SUC2 regulated expression of an antisense RNAi construct for GFP	This study
pRS404-PTEF -Ago1-CYC1t	TRP1 integrating vector with constitutive Argonaute expression	[5]
pRS405-PTEF-Dcr1-CYC1t	LEU2 integrating vector with constitutive Dicer expression	[5]
pUG6	LoxP-KanMX-LoxP cassette	Euroscarf
pUG66	LoxP-Ble-LoxP cassette	Euroscarf

Fermentation conditions

All growth experiments were carried out in baffled screw top flasks shaking at 200 rpm, 30°C with medium comprising 10% of the total flask volume. Individual colonies were transferred to liquid media and pre-cultured for approximately 15 hours. Cultures were then passaged into a second pre-culture and grown to mid exponential phase (OD_{660nm} between 1 and 6) prior to inoculation of experimental cultures at an OD_{660nm} of 0.1 for the initial

promoter comparison (Figure 1) and 0.4 for the time course experiments (Figure 2).

Analytics

GFP fluorescence was measured using an Accuri C6 flow cytometer (BD Biosciences) as in [36]. HPLC was used to measure sucrose, fructose, and glucose as previously described [44].

Table 3 *S. cerevisiae* strains used in this study

Name	Genotype	Notes	Origin
CEN.PK113-5D	MATa; ura3-52; MAL2-8 ^C ; SUC2	Haploid MATa lab strain with uracil auxotrophy	Euroscarf
CEN.PK113-7D	MATa; MAL2-8 ^C ; SUC2	Prototrophic haploid MATa lab strain	Euroscarf
S01	CEN.PK2-1c, bar1Δ	<i>BAR1</i> gene deleted	This study
S02	CEN.PK2-1c, bar1Δ, fus1::KanMX, trp1::pRS404-PTEF-Ago1	<i>BAR1</i> and <i>FUS1</i> deleted, Argonaute gene integrated at <i>TRP1</i>	This study
S03	CEN.PK2-1c, bar1Δ, fus1::KanMX, trp1::pRS404-PTEF-Ago1, leu2::pRS405-PTEF-Dcr1	RNAi capable base strain with Argonaute and Dicer integration	This study
GFP01	CEN.PK113-5D, bar1::phleo, fus1::KanMX, ura3-52::P _{TEF1} -GFPCLN2PEST-ADH1t-pRS406	constitutive destabilized GFP expression	This study
GFP02	CEN.PK113-5D, ura3::P _{MAL12} -GFPCLN2PEST-ADH1t-pRS406	<i>MAL12</i> regulated GFP expression	This study
GFP03	CEN.PK113-5D, ura3::P _{GLC3} -GFPCLN2PEST-ADH1t-pRS406	<i>GLC3</i> regulated GFP expression	This study
GFP04	CEN.PK113-5D, ura3::P _{GPH1} -GFPCLN2PEST-ADH1t-pRS406	<i>GPH1</i> regulated GFP expression	This study
GFP05	CEN.PK113-5D, ura3::P _{SUC2} -GFPCLN2PEST-ADH1t-pRS406	<i>SUC2</i> regulated GFP expression	This study
GFP06	CEN.PK2-1c, bar1Δ, fus1::KanMX, ura3::P _{TEF1} -yEGFPCLN2PEST-406, trp1::pRS404-PTEF-Ago1, leu2::pRS405-PTEF-Dcr1, pRS413	RNAi capable, constitutive GFP expressing control strain	This study
GFP07	CEN.PK2-1c, bar1Δ, fus1::KanMX, ura3::P _{TEF1} -yEGFPCLN2PEST-406, trp1::pRS404-PTEF-Ago1, leu2::pRS405-PTEF-Dcr1, P _{SUC2} -GFPantisense-CYC1t-pRS413	<i>SUC2</i> promoter mediated repression of GFP using RNAi	This study

Statistical analysis

Mean GFP fluorescence values from triplicate experiments were compared using a two-sided, unpaired student's *t*-test with equal variance. Null hypotheses were rejected when *p*-values were ≤ 0.05 .

Additional files

Additional file 1: Figure S1. Supplementary material.
Additional file 2: Annotated plasmid maps and sequences.

Competing interests

The authors declare no competing interests.

Authors' contributions

TCW conceived of the study and designed experiments. TCW and MIE performed experiments. TCW, MIE, LKN, and CEV wrote and edited the manuscript. LKN and CEV participated in the design and coordination of the project. All authors read and approved the final manuscript.

Acknowledgements

TCW was supported by an Australian Postgraduate Award and the AIBN Top Up Scholarship. We thank Manuel Plan of the Metabolomics Australia Queensland node for technical assistance with HPLC.

Received: 20 January 2015 Accepted: 6 March 2015

Published online: 01 April 2015

References

- Lee SK et al. Metabolic engineering of microorganisms for biofuels production: from bugs to synthetic biology to fuels. *Curr Opin Biotechnol*. 2008;19(6):556–63.
- Keasling JD. Synthetic Biology for Synthetic Chemistry. *ACS Chem Biol*. 2008;3(1):64–76.
- Hong K-K, Nielsen J. Metabolic engineering of *Saccharomyces cerevisiae*: a key cell factory platform for future biorefineries. *Cell Mol Life Sci*. 2012;69(16):2671–90.
- Lohr D, Venkov P, Zlatanova J. Transcriptional regulation in the yeast GAL gene family: a complex genetic network. *FASEB J*. 1995;9(9):777–87.
- Drinnenberg IA et al. RNAi in Budding Yeast. *Science*. 2009;326(5952):544–50.
- Westfall PJ, Pitera DJ, Lenihan JR, Eng D, Woollard FX, Regentin R, et al. Production of amorpha-4-diene in yeast, and its conversion to dihydroartemisinin acid, precursor to the antimalarial agent artemisinin. *Proc Natl Acad Sci U S A*. 2012;109(3):E111–8.
- Gancedo JM. Yeast Carbon Catabolite Repression. *Microbiol Mol Biol Rev*. 1998;62(2):334–61.
- Hashimoto H et al. Regulation of expression of the galactose gene cluster in *Saccharomyces cerevisiae*. *Mol Gen Genet MGG*. 1983;191(1):31–8.
- Wishart JA et al. Doxycycline, the drug used to control the tet-regulatable promoter system, has no effect on global gene expression in *Saccharomyces cerevisiae*. *Yeast*. 2005;22(7):565–9.
- Weinhandl K et al. Carbon source dependent promoters in yeasts. *Microb Cell Fact*. 2014;13(1):5.
- Hubmann G, Thevelein J, Nevoigt E. Natural and Modified Promoters for Tailored Metabolic Engineering of the Yeast *Saccharomyces cerevisiae*. In: Mapelli V, editor. *Yeast Metabolic Engineering*. New York: Springer; 2014. p. 17–42.
- Bruschi M et al. A transferable sucrose utilization approach for non-sucrose-utilizing *Escherichia coli* strains. *Biotechnol Adv*. 2012;30(5):1001–10.
- Sabri S, Nielsen LK, Vickers CE. Molecular Control of Sucrose Utilization in *Escherichia coli* W, an Efficient Sucrose-Utilizing Strain. *Appl Environ Microbiol*. 2013;79(2):478–87.
- Carlson M. Glucose repression in yeast. *Curr Opin Microbiol*. 1999;2(2):202–7.
- Bu Y, Schmidt MC. Identification of cis-acting elements in the SUC2 promoter of *Saccharomyces cerevisiae* required for activation of transcription. *Nucleic Acids Res*. 1998;26(4):1002–9.
- Fleming AB, Pennings S. Tup1-Ssn6 and Swi-Snf remodelling activities influence long-range chromatin organization upstream of the yeast SUC2 gene. *Nucleic Acids Res*. 2007;35(16):5520–31.
- Lutfiyya LL, Johnston M. Two zinc-finger-containing repressors are responsible for glucose repression of SUC2 expression. *Mol Cell Biol*. 1996;16(9):4790–7.
- Ozcan S et al. Expression of the SUC2 gene of *Saccharomyces cerevisiae* is induced by low levels of glucose. *Yeast*. 1997;13(2):127–37.
- Cheng Q, Michels CA. MAL11 and MAL61 encode the inducible high-affinity maltose transporter of *Saccharomyces cerevisiae*. *J Bacteriol*. 1991;173(5):1817–20.
- Yao B, Marmur J, Sollitti P. Construction of glucose-repressible yeast expression vectors. *Gene*. 1993;137(2):223–6.
- Francois J, Parrou JL. Reserve carbohydrates metabolism in the yeast *Saccharomyces cerevisiae*. *FEMS Microbiol Rev*. 2001;25(1):125–45.
- Sunnarborg SW et al. Expression of the yeast glycogen phosphorylase gene is regulated by stress-response elements and by the HOG MAP kinase pathway. *Yeast*. 2001;18(16):1505–14.
- Thon VJ et al. Coordinate regulation of glycogen metabolism in the yeast *Saccharomyces cerevisiae*. Induction of glycogen branching enzyme. *J Biol Chem*. 1992;267(21):15224–8.
- Basso TO et al. Engineering topology and kinetics of sucrose metabolism in *Saccharomyces cerevisiae* for improved ethanol yield. *Metab Eng*. 2011;13(6):694–703.
- Mateus C, Avery SV. Destabilized green fluorescent protein for monitoring dynamic changes in yeast gene expression with flow cytometry. *Yeast*. 2000;16(14):1313–23.
- Hua Lin K et al. Control of gene expression from the SUC2 promoter of *Saccharomyces cerevisiae* with the aid of a glucose analyser. *Appl Microbiol Biotechnol*. 1989;32(3):313–6.
- Galdieri L et al. Transcriptional Regulation in Yeast during Diauxic Shift and Stationary Phase. *OMICS: A Journal of Integrative Biology*. 2010;14(6):629–38.
- St John TP, Davis RW. The organization and transcription of the galactose gene cluster of *Saccharomyces*. *J Mol Biol*. 1981;152(2):285–315.
- Partow S et al. Characterization of different promoters for designing a new expression vector in *Saccharomyces cerevisiae*. *Yeast*. 2010;27(11):955–64.
- Bachmair A, Finley D, Varshavsky A. In vivo half-life of a protein is a function of its amino-terminal residue. *Science*. 1986;234(4773):179–86.
- Schirmaier F, Philippson P. Identification of two genes coding for the translation elongation factor EF-1 alpha of *S. cerevisiae*. *EMBO J*. 1984;3(13):3311–5.
- Warner JR, Vilardell J, Sohn JH. Economics of Ribosome Biosynthesis. *Cold Spring Harb Symp Quant Biol*. 2001;66:567–74.
- Werner-Washburne M et al. Stationary phase in the yeast *Saccharomyces cerevisiae*. *Microbiol Mol Biol Rev*. 1993;57(2):383–401.
- Crook NC, Schmitz AC, Alper HS. Optimization of a yeast RNA interference system for controlling gene expression and enabling rapid metabolic engineering. *ACS Synth Biol*. 2014;3(5):307–13.
- Si T et al. RNAi-Assisted Genome Evolution in *Saccharomyces cerevisiae* for Complex Phenotype Engineering. *ACS Synthetic Biology*, 2014. (<http://pubs.acs.org/doi/abs/10.1021/sb500074a>). The DOI is 10.1021/sb500074a.
- Williams TC, Nielsen LK, Vickers CE. Engineered quorum sensing using pheromone-mediated cell-to-cell communication in *Saccharomyces cerevisiae*. *ACS Synth Biol*. 2013;2(3):136–49.
- Cha HJ et al. Enhancement, by succinate addition, of the production of cloned glucoamylase from recombinant yeast using a SUC2 promoter. *Process Biochem*. 1998;33(3):257–61.
- Zhang X et al. Enhancement of production of cloned α -amylase by lactic acid feeding from recombinant *Saccharomyces cerevisiae* using a SUC2 promoter. *Biotechnol Lett*. 2001;23(4):259–62.
- Verduyn C et al. Effect of benzoic acid on metabolic fluxes in yeasts: A continuous-culture study on the regulation of respiration and alcoholic fermentation. *Yeast*. 1992;8(7):501–17.
- Sambrook J, Russell DW. *Molecular cloning, a laboratory manual*. New York: Cold Spring Harbour Laboratory Press; 2001.
- Gietz RD, Schiestl RH. Quick and easy yeast transformation using the LiAc/PEG carrier DNA/PEG method. *Nat Protocols*. 2007;2:35–7.
- Stansfield I, Stark MJR. *2 Yeast Genetics and Strain Construction*. In: Ian S, Michael JRS, editors. *Methods in Microbiology*. London, UK: Academic Press; 2007. p. 23–43.

43. Arifin Y et al. Deletion of *cscR* in *Escherichia coli* W improves growth and poly-3-hydroxybutyrate (PHB) production from sucrose in fed batch culture. *J Biotechnol.* 2011;156(4):275–8.
44. Williams TC, Aversch NJH, Winter G, Plan MR, Vickers CE, Nielsen LK, et al. Quorum-Sensing Linked RNA interference for Dynamic Metabolic Pathway Control in *Saccharomyces cerevisiae*. *Metab Eng.* 2015. (<http://www.sciencedirect.com/science/article/pii/S1096717615000348>). The DOI is 10.1016/j.ymben.2015.03.008.
45. Sikorski RS, Hieter P. A System of Shuttle Vectors and Yeast Host Strains Designed for Efficient Manipulation of DNA in *Saccharomyces cerevisiae*. *Genetics.* 1989;122(1):19–27.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

