



REVIEW

Recent advances in the study of HPV-associated carcinogenesis

Liyan Jin, Zhi-Xiang Xu[✉]

Division of Hematology and Oncology, Comprehensive Cancer Center, University of Alabama at Birmingham, Birmingham 35205, USA

Human papillomaviruses (HPVs) cause virtually all cervical cancers, the second leading cause of death by cancer among women, as well as other anogenital cancers and a subset of head and neck cancers. Approximately half of women, who develop cervical cancer die from it. Despite the optimism that has accompanied the introduction of prophylactic vaccines to prevent some HPV infections, the relatively modest uptake of the vaccine, especially in the developing world, and the very high fraction of men and women who are already infected, means that HPV-associated disease will remain as a significant public health problem for decades. In this review, we summarize some recent findings on HPV-associated carcinogenesis, such as miRNAs in HPV-associated cancers, implication of stem cells in the biology and therapy of HPV-positive cancers, HPV vaccines, targeted therapy of cervical cancer, and drug treatment for HPV-induced intraepithelial neoplasias.

KEYWORDS human papillomavirus (HPV); carcinogenesis; vaccine; miRNA; cancer stem cell (CSC); cervical intraepithelial neoplasias (CIN); targeted therapy

INTRODUCTION

Human papillomaviruses (HPVs) are a group of small, non-enveloped, double-stranded DNA tumor viruses, categorized to the *papillomaviridae* family. Approximately 200 types of HPVs have been identified. HPVs are not only species specific but also display a tropism for squamous epithelia. A large number of HPVs infect cutaneous epithelia, whereas other groups infect mucosal epithelia (Bravo and Féllez-Sánchez, 2015). The mucosal HPVs are classified as “high-risk” and “low-risk,” depending on the capability of the viruses to cause the malignant progression of the lesions. Low-risk HPVs, such as HPV type 6 (HPV6) and HPV11, cause genital warts. High-risk HPVs, such as HPV16 and HPV18, trigger squamous intraepithelial lesions, which may progress to ma-

lignant foci (zur Hausen, 2009). High-risk HPVs are associated with greater than 99% of cervical carcinomas, a portion of additional anogenital tumors and approximately a quarter of oropharyngeal tumors (Walboomers et al., 1999; Mirghani et al., 2015). This is the combined action of two viral oncoproteins, E6 and E7, which subvert cellular regulatory pathways controlling cell cycle and cell survival (McLaughlin-Drubin and Munger, 2009; Moody and Laimins, 2010). The E7 oncoprotein binds to more than 20 cellular targets and interferes with numerous cellular processes, leading to deregulated cell cycle progression, malignant transformation, centrosome amplification, DNA damage, anoikis loss and anchorage-independent cell growth, immune surveillance evasion and persistent infection. The E6 oncoprotein abrogates cell growth arrest and apoptosis, induces genomic instability and somatic mutations, activates telomerase and the telomerase reverse transcriptase to promote immortalization, disrupts cell polarity, prevents anoikis, and allows cellular growth without attachment to extracellular matrix (ECM). Together, these observations demonstrate that E6 and E7 are multifunctional proteins driving oncogenic transformation and tumorigenicity of the HPVs.

Received: 21 March 2015, Accepted: 16 April 2015

Published online: 20 April 2015

✉ Correspondence:

Phone: +1-205-934-1868, Fax: +1-205-934-1870,

Email: zhixiangxu@uabmc.edu

ORCID: 0000-0003-0072-0334

MOLECULAR ACTIVITIES OF HIGH-RISK HPV E6/7

The most prominent targets of high-risk E6 and E7 proteins are tumor suppressor genes p53 and pRb family pocket proteins (McLaughlin-Drubin and Munger, 2009; Moody and Laimins, 2010). The E7 interacts with a large number of host proteins (White et al., 2012b). In particular, the high-risk HPV E7 protein targets the pRb family proteins for degradation, thereby inhibiting pRb-mediated repression of E2F-responsive genes. In addition, it inhibits the cyclin-dependent kinase (CDK) inhibitors p21 and p27, and activates both cyclin A/CDK2 and cyclin E/CDK2 (Nguyen and Munger, 2008). E7 also interacts with histone deacetylases (HDACs) to affect cellular gene expression (Brehm et al., 1999; Longworth and Laimins, 2004). Therefore, E7 subverts cell cycle control and induces hyperproliferation. Overexpression of E7 stimulates centrosome amplification through the enhancement of CDK2 activity and interaction with γ -tubulin, contributing to the accumulation of chromosomal alterations and increased risk of genomic instability (Duensing et al., 2000). Moreover, E7 interacts with p600, preventing anoikis and rendering the cell anchorage-independent growth (Huh et al., 2005). Lastly, E7 inactivates interferon regulatory factor 1 (IRF1), contributing to the evasion of immune surveillance and the establishment of a persistent infection (Park et al., 2000).

The E6 proteins of various HPV types interact with a variety of host proteins (Howie et al., 2009). p53 is among the most important targets. The high-risk HPV E6 protein forms a trimeric complex with E6-associated protein (E6AP) and p53 resulting in the degradation of p53 (Scheffner et al., 1993; McLaughlin-Drubin and Munger, 2009; Moody and Laimins, 2010). E6 interacts with the histone acetyl transferases p300, CREB binding protein (CBP), and alteration/deficiency in activation-3 (ADA3) to prevent p53 acetylation, hence suppressing the transcription of p53-responsive genes (Thomas and Chiang, 2005). Thus, p53-dependent cellular responses to aberrant proliferation, genomic instability, and mutations are suppressed by E6. Through the interaction with IRF3 (Ronco et al., 1998), E6 would interrupt the interferon response. E6/E7 inhibits growth-suppressive cytokines-induced apoptosis by interaction with and suppression of the TNF- α -FADD (FAS-associated protein with death domain)-caspase 8 signaling, and by the degradation of BAX and BAK (Boccardo et al., 2004; Garnett et al., 2006; Liu et al., 2008; Underbrink et al., 2008). In addition, E6 activates the telomerase reverse transcriptase (TERT) and telomerase and hence prevents the telomere shortening in response to persistent proliferation and in turn promoting immortalization (Klingelhutz et al., 1996; Xu et al., 2013). Moreover, E6 mediates the degradation of several PDZ domain containing host proteins, leading

to the loss of cell polarity and inducing hyperplasia (Pim et al., 2012).

In a recent report, White *et al.* (2012a) applied a mass spectrometry-based platform to systematically identify and characterize interactions between HPV oncoproteins and host cellular proteins. They found that HPV E6 interacts with host proteins in a genus and species common or specific manner. The E6 interaction data set not only contains previously reported interaction proteins of E6, such as p300/CBP, E6AP, and p53, but also newly identified proteins that bind to and interact with E6, such as Ccr4-Not complex. Ccr4-Not acts as a deadenylase conserved from yeast to humans, and affects mRNA metabolism (Collart and Panasenko, 2012). In addition, it possesses an ubiquitin ligase function linking to ubiquitylation and the proteasome. These findings, on the one hand, provide a comprehensive database for studies on the diverse biology of the HPVs. On the other hand, they suggest that current understanding for the functions of HPV oncoproteins is incomplete. Thus, a continued exploration for the multifaceted roles of HPV oncoproteins in driving proliferation and carcinogenesis is necessary.

MIRNAS IN HPV-ASSOCIATED CANCERS

microRNAs (miRNAs) are noncoding regulatory RNAs of 18–25 nucleotides in size. They are derived from RNA polymerase II transcripts of coding or noncoding genes (Bartel, 2004). miRNAs expression is tissue- or differentiation-specific. They modulate gene expression at the posttranscriptional level by base-pairing with complementary nucleotide sequences of target mRNAs, leading to the degradation of mRNA or translational suppression (Lewis et al., 2005; Bueno et al., 2010). As part of the transcriptome, the small non-coding RNA species have attracted much attention due to both their genesis and their ability to regulate gene expression.

miRNA expressions have been frequently found near fragile sites in chromosomes or integration sites of high-risk HPVs (Georgakilas et al., 2014). Integration of HPV oncogenes may alter miRNA expression via deletion, amplification, or genomic rearrangement (TCGA, 2015). A large number of miRNA genes are regulated by transcription factors c-Myc, p53, and E2F, which are targeted by oncogenic HPV E6 and E7 (Zheng and Wang, 2011). For example, E6-mediated degradation of p53 reduces the expression of miR-34a (Wang et al., 2009; Li et al., 2010) and miR-23b (Au Yeung et al., 2011) at the transcriptional level. In addition, E6 and E7 oncoproteins interact with multiple cellular factors, and these interactions could lead to increased or decreased expression of cellular miRNAs. In a recent report, 13 host miRNAs were found to be specifically regulated by HPV16 and HPV18 in organotypic raft cultures of foreskin and vag-

inal keratinocytes with a miRNA array assay in combination with small RNA sequencing (Wang et al., 2014b). The increase of miR-16, miR-25, miR-92a, and miR-378 and the decrease of miR-22, miR-27a, miR-29a, and miR-100 could be attributed to and mediate the functions of viral oncoprotein E6, E7, or both. The authors suggest that an expression ratio ≥ 1.5 of miR-25/92a group over miR-22/29a group might be used to distinguish cervical cancers from normal cervix (Wang et al., 2014b).

IMPLICATION OF STEM CELLS IN THE BIOLOGY AND THERAPY OF HPV-ASSOCIATED CANCERS

Stem cells are a group of undifferentiated cells that act as a reservoir for new cells in order to replace defective or necrotic cells. An essential characteristic of stem cells is the ability to self-renew and to differentiate into diverse types of cells. Current researches regarding cancer stem cells (CSCs) demonstrate that these cells maintain the capabilities of normal tissue stem cells, such as unlimited self-regeneration, and differentiation into various cell types. CSCs initiate malignant tumors on a single cell basis via symmetric proliferation, and express cancer stem cell specific markers. These characteristics of CSCs are responsible for tumor maintenance and metastasis and possibly also for the resistance toward chemotherapy and radiation therapy.

Researchers have characterized the surface markers for cervical cancer stem cells, e.g. p63, cytokeratin 17 (CK17), Nanog, Musashi-1 (Msi1), Nucleostemin (NS), CD49f, ALDH1, CD44, and a few more others. Nanog, NS, and Msi1 were found to be highly expressed in cervical carcinomas relative to normal cervix and were proposed to be involved in carcinogenesis of the cervix (Ye et al., 2008). Embryonic stem cells (ESCs) markers (Sox2, Oct4) and the Wnt signal pathway (β -catenin) are crucial for the progression of various human malignancies. Ji *et al.* (2014) showed that Sox2 and Oct4 are highly expressed in cervical squamous cell carcinomas and that Wnt signal (beta-catenin) is activated. Thus, Sox2 could be a novel predictor for poor prognosis of such lesions (Ji et al., 2014). Using a different approach, Lopez *et al.* (2012) investigated the tumor initiating cells in cervical carcinoma cell lines, HeLa and SiHa, and found a high level of CD49f in these cells, whereas Gu *et al.* (2011) showed that cancer initiating cells in HeLa cell line exhibit a CD44 (high) / CD24 (low) expression pattern.

HPVs infect stem cells, which are speculated to be the origination of the cervical epithelial cancers. Stem cells could also be sources for latently infected cells that can persist for a long period. The HPV oncogenic E6 and E7 modulate multiple cellular pathways with parallel

roles in the carcinogenic process. Michael *et al.* (2013) examined the effects of HPV16 E6/E7 on stem cells located in the bulge of hair follicles. The results showed that expression of HPV16 oncogenes reduces the number of quiescent cells, a typical feature of stem cells within hair follicles, whereas stem cell markers in the follicles are also decreased. The authors suggest that HPV infection may induce aberrant mobilization of the stem cells. These effects may play a role in viral life cycle and/or ensuing carcinogenesis.

Tang *et al.* (2013) applied *in vitro* and *in vivo* analysis to determine if CSCs of head and neck squamous cell carcinoma (HNSCC) are affected by HPVs. It shows that HPV status does not correlate with the proportion of CSCs present in HNSCC. The HPV positive cells and those transduced with HPV E6/E7 possess a greater clonogenicity than HPV-negative cells. CSCs of the HNSCC are more resistant to cisplatin than non-CSCs. Consistently, HPVs do not affect the response of CSCs to the treatment of cisplatin, further supporting the notion that the functions of HPV are not overlapped with CSCs.

Taken together, characterization of stem cells in cervix provides new insights into the mechanisms by which cervical cancer is developed. However, it remains largely unknown what the role of HPV oncogenic E6/E7 may play in the formation of cervix CSCs and the potential therapeutic significance of CSCs may exist in HPV positive cancers.

HPV VACCINES

Vaccines have been developed to prevent infections with certain types of HPVs. Currently, there are a quadrivalent HPV vaccine and a bivalent HPV vaccine licensed in the United States. The former produces immunity against HPV 6, 11, 16, and 18, whereas the latter is administered to prevent infections of HPV 16 and 18. Both vaccines are based on virus-like particles of the L1 capsid protein. The vaccines are highly efficient and immunogenic if given before exposure to these types of HPVs (Schiller and Lowy, 2012). Both vaccines are administered in a 3-dose series. The protective duration of the bivalent vaccine is over 10 years and counting. Recent studies suggest that 2-dose provide effective protection as well (Romanowski et al., 2014; Dobson et al., 2013). To increase the protection, a VLP based nine-valent prophylactic HPV vaccine (HPV 6/11/16/18/31/33/45/52/58) (Serrano et al., 2012) has been approved by FDA. A cross-reactive prophylactic L2-based vaccine is also under development (Wang et al., 2104 a). Several therapeutic vaccines are undergoing clinical trials in women with cervical cancers (Hibbitts et al., 2012; Tran et al., 2014). These vaccines are developed to produce an immune reaction against HPV16 E6 or E7 protein. Thus, the en-

hanced immunity might kill the cancer cells or stop them from growing.

TARGET THERAPY

Based on international guidelines, treatment of cervical cancer (CC) consists of surgery in early stages and of chemoradiation in locally advanced disease. Metastatic disease is usually treated with palliative chemotherapeutic regimens. Cytostatic drugs exhibit substantial side effects and limited efficacy. Thus, the discovery of new anticancer agents, interfering with molecular targets expressed in the microenvironment of the cancer or by the tumor cell itself, represents an opportunity challenging the cancer. Vascular endothelial growth factor (VEGF) promotes angiogenesis in cervical cancer, leading to disease progression. Bevacizumab, a humanized anti-VEGF monoclonal antibody, was tested in patients with recurrent, persistent, or metastatic cervical cancer in conjunction with chemotherapies. The increase in time to progression or survival was 3 to 4 months relative to chemotherapy alone (Wright et al., 2006; Tewari et al., 2014). As another example, pazopanib, a multi-tyrosine kinase inhibitor (multi TKI), has been used to treat kidney and ovarian cancer. Recent clinical trials confirmed the activity of the antiangiogenesis agents in advanced and recurrent cervical cancer. However, the benefits in progression free survival only amounted to 3 months (Monk et al., 2010).

DRUG TREATMENT FOR HPV-INDUCED INTRAEPITHELIAL NEOPLASIAS

Standard treatment of cervical pre-cancer (such as cervical intraepithelial neoplasia; CIN) includes cryotherapy, laser treatment, and conization. Researchers are asking whether CIN could be treated with pharmaceutical agents. Del Priore *et al.* (2010) treated CIN2 and CIN3 patients for 12 weeks with diindolylmethane (DIM), a component of indole-3-carbinol (I3C) found in Brassica vegetables. The subjects were evaluated every 3–4 months with Pap smear, HPV, colposcopy, biopsy and physical examination at follow-up. The results showed that oral administration of DIM leads to a high rate of clinically significant improvement in confirmed CIN 2 or 3 lesions in a one-year follow-up. However, there is no statistically significant difference between DIM- and placebo-treated groups. In another study, a randomized controlled trial was applied to evaluate a topical treatment for CIN 2+ using anti-viral drug cidofovir (Van Pachterbeke *et al.*, 2009). Although regression was more frequently achieved than the placebo group as judged by histology and by *in situ* hybridization, the more sensitive Hybrid Capture 2 assay did not reveal significant differ-

ence between the two groups of patients. The authors, therefore, concluded that it is a promising candidate for topical chemotherapy but it cannot replace conization at this juncture.

The most promising agent appears to be imiquimod, an immune response modifier. It is a patient-applied cream used to treat certain types of skin diseases, superficial malignant melanomas, as well as genital warts (*condylo-mata acuminata*). Several groups applied imiquimod to subjects with CIN 2 and 3 and showed a higher histologic regression as compared with placebo treatment subjects (Grimm *et al.*, 2012). More importantly, HPV clearance rates are increased in the imiquimod group (60%) compared with the placebo group (14%). In patients with HPV-16 infection, complete remission rates are 47% in the imiquimod group compared with 0% in the placebo group. Consistently, administrations of imiquimod alone or in combination with HPV therapeutic vaccination in patients with anal and vulval intraepithelial neoplasia produced similar results (Hibbitts *et al.*, 2010; Daayana *et al.*, 2010). There was a substantially increased local infiltration of CD8 and CD4 T cells in lesion responders. Taken together, current clinical trials suggest that topical imiquimod 5% cream may be beneficial in anogenital intraepithelial neoplasias.

ACKNOWLEDGMENTS

We greatly appreciate Dr. Louise T. Chow for critical reading of the article. The work in Xu lab was supported by a grant from National Cancer Institute R01CA133053, Career Development Awards from the Cervical Cancer SPORE NCI P50CA098252 and from the STI CRC NIH U19 AI 113212, and the UAB Comprehensive Cancer Center Pilot Program Project grant.

COMPLIANCE WITH ETHICS GUIDELINES

The authors declare that they have no conflict of interest. This article does not contain any studies with human or animal subjects performed by any of the authors.

REFERENCES

- Au Yeung CL, Tsang TY, Yau PL, Kwok TT. 2011. Human papillomavirus type 16 E6 induces cervical cancer cell migration through the p53/microRNA-23b/urokinase-type plasminogen activator pathway. *Oncogene*, 30: 2401–2410.
- Bartel DP. 2004. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell*, 116: 281–297.
- Boccardo E, Noya F, Broker TR, Chow LT, Villa LL. 2004. HPV-18 confers resistance to TNF-alpha in organotypic cultures of human keratinocytes. *Virology*, 328: 233–243.
- Bravo IG, Féllez-Sánchez M. 2015. Papillomaviruses: Viral evolution, cancer and evolutionary medicine. *Evol Med Public Health*, 2015: 32–51.

- Brehm A, Nielsen SJ, Miska EA, McCance DJ, Reid JL, Bannister AJ, Kouzarides T. 1999. The E7 oncoprotein associates with Mi2 and histone deacetylase activity to promote cell growth. *EMBO J*, 18: 2449–2458.
- Bueno MJ, Gómez de Cedrón M, Laresgoiti U, Fernández-Piqueras J, Zubiaga AM, Malumbres M. 2010. Multiple E2F-induced microRNAs prevent replicative stress in response to mitogenic signaling. *Mol Cell Biol*, 30: 2983–2995.
- Collart MA, Panasenko OO. 2012. The Ccr4--not complex. *Gene*, 492: 42–53.
- Daayana S, Elkord E, Winters U, Pawlita M, Roden R, Stern PL, Kitchener HC. 2010. Phase II trial of imiquimod and HPV therapeutic vaccination in patients with vulval intraepithelial neoplasia. *Br J Cancer*, 102:1129–1136.
- Del Priore G, Gudipudi DK, Montemarano N, Restivo AM, Malanowska-Stega J, Arslan AA. 2010. Oral diindolylmethane (DIM): pilot evaluation of a nonsurgical treatment for cervical dysplasia. *Gynecol Oncol*, 116:464–467.
- Dobson SR, McNeil S, Dionne M, Dawar M, Ogilvie G, Kraiden M, Sauvageau C, Scheifele DW, Kollmann TR, Halperin SA, Langley JM, Bettinger JA, Singer J, Money D, Miller D, Naus M, Marra F, Young E. 2013. Immunogenicity of 2 doses of HPV vaccine in younger adolescents vs 3 doses in young women: a randomized clinical trial. *JAMA*, 309: 1793–1802.
- Duensing S, Lee LY, Duensing A, Basile J, Piboonniyom S, Gonzalez S, Crum CP, Munger K. 2000. The human papillomavirus type 16 E6 and E7 oncoproteins cooperate to induce mitotic defects and genomic instability by uncoupling centrosome duplication from the cell division cycle. *Proc Natl Acad Sci U S A*, 97: 10002–10007.
- Garnett TO, Filippova M, Duerksen-Hughes PJ. 2006. Accelerated degradation of FADD and procaspase 8 in cells expressing human papilloma virus 16 E6 impairs TRAIL-mediated apoptosis. *Cell Death Differ*, 13: 1915–1926.
- Georgakilas AG, Tsantoulis P, Kotsinas A, Michalopoulos I, Townsend P, Gorgoulis VG. 2014. Are common fragile sites merely structural domains or highly organized “functional” units susceptible to oncogenic stress? *Cell Mol Life Sci*, 71: 4519–4544.
- Grimm C, Polterauer S, Natter C, Rahhal J, Heffler L, Tempfer CB, Heinze G, Stary G, Reinthaller A, Speiser P. 2012. Treatment of cervical intraepithelial neoplasia with topical imiquimod: a randomized controlled trial. *Obstet Gynecol*, 120:152–159.
- Gu W, Yeo E, McMillan N, Yu C. 2011. Silencing oncogene expression in cervical cancer stem-like cells inhibits their cell growth and self-renewal ability. *Cancer Gene Ther*, 18: 897–905.
- Hibbitts S. 2010. TA-CIN, a vaccine incorporating a recombinant HPV fusion protein (HPV16 L2E6E7) for the potential treatment of HPV16-associated genital diseases. *Curr Opin Mol Ther*, 12: 598–606.
- Howie HL, Katzenellenbogen RA, Galloway DA. 2009. Papillomavirus E6 proteins. *Virology*, 384: 324–334.
- Huh KW, DeMasi J, Ogawa H, Nakatani Y, Howley PM, Münger K. 2005. Association of the human papillomavirus type 16 E7 oncoprotein with the 600-kDa retinoblastoma protein-associated factor, p60. *Proc Natl Acad Sci U S A*, 102: 11492–11497.
- Ji J, Wei X, Wang Y. 2014. Embryonic stem cell markers Sox-2 and OCT4 expression and their correlation with WNT signal pathway in cervical squamous cell carcinoma. *Int J Clin Exp Pathol*, 7: 2470–2476.
- Klingelutz AJ, Foster SA, McDougall JK. 1996. Telomerase activation by the E6 gene product of human papillomavirus type 16. *Nature*, 380: 79–82.
- Lajer CB, Garnæs E, Friis-Hansen L, Norrild B, Therkildsen MH, Glud M, Rossing M, Lajer H, Svane D, Skotte L, Specht L, Buchwald C, Nielsen FC. 2012. The role of miRNAs in human papilloma virus (HPV)-associated cancers: bridging between HPV-related head and neck cancer and cervical cancer. *Br J Cancer*, 106: 1526–1534.
- Lewis BP, Burge CB, Bartel DP. 2005. Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell*, 120: 15–20.
- Li B, Hu Y, Ye F, Li Y, Lv W, Xie X. 2010. Reduced miR-34a expression in normal cervical tissues and cervical lesions with high-risk human papillomavirus infection. *Int J Gynecol Cancer*, 20: 597–604.
- Liu HC, Chen GG, Vlantis AC, Tse GM, Chan AT, van Hasselt CA. 2008. Inhibition of apoptosis in human laryngeal cancer cells by E6 and E7 oncoproteins of human papillomavirus 16. *J Cell Biochem*, 103:1125–1143.
- Longworth MS, Laimins LA. 2004. The binding of histone deacetylases and the integrity of zinc finger-like motifs of the E7 protein are essential for the life cycle of human papillomavirus type 31. *J Virol*, 78:3533–3541.
- Lopez J, Poitevin A, Mendoza-Martinez V, Perez-Plasencia C, Garcia-Carranca A. 2012. Cancer-initiating cells derived from established cervical cell lines exhibit stem-cell markers and increased radioresistance. *BMC Cancer*, 12:48.
- McLaughlin-Drubin ME, Munger K. 2009. Oncogenic activities of human papillomaviruses. *Virus Res*, 143: 195–208.
- Michael S, Lambert PF, Strati K. 2013. The HPV16 oncogenes cause aberrant stem cell mobilization. *Virology*, 443: 218–225.
- Mirghani H, Amen F, Moreau F, Lacau St, Guily J. 2015. Do high-risk human papillomaviruses cause oral cavity squamous cell carcinoma? *Oral Oncol*, 51: 229–236.
- Monk BJ, Mas Lopez L, Zarba JJ, Oaknin A, Tarpin C, Termrungruanglert W, Alber JA, Ding J, Stutts MW, Pandite LN. 2010. Phase II, open-label study of pazopanib or lapatinib monotherapy compared with pazopanib plus lapatinib combination therapy in patients with advanced and recurrent cervical cancer. *J Clin Oncol*, 28: 3562–3569.
- Moody CA, Laimins LA. 2010. Human papillomavirus oncoproteins: pathways to transformation. *Nat Rev Cancer*, 10:550–560.
- Nguyen CL, Münger K. 2008. Direct association of the HPV16 E7 oncoprotein with cyclin A/CDK2 and cyclin E/CDK2 complexes. *Virology*, 380: 21–25.
- Park JS, Kim EJ, Kwon HJ, Hwang ES, Namkoong SE, Um SJ. 2000. Inactivation of interferon regulatory factor-1 tumor suppressor protein by HPV E7 oncoprotein. Implication for the E7-mediated immune evasion mechanism in cervical carcinogenesis. *J Biol Chem*, 275: 6764–6769.
- Pim D, Bergant M, Boon SS, Ganti K, Kranjec C, Massimi P, Subbaiah VK, Thomas M, Tomaić V, Banks L. 2012. Human papillomaviruses and the specificity of PDZ domain targeting. *FEBS J*, 279: 3530–3537.
- Romanowski B, Schwarz TF, Ferguson LM, Ferguson M, Peters K, Dionne M, Schulze K, Ramjattan B, Hillemanns P, Behre U, Suryakiran P, Thomas F, Struyf F. 2014. Immune response to the HPV-16/18 AS04-adjuvanted vaccine administered as a 2-dose or 3-dose schedule up to 4 years after vaccination: results from a randomized study. *Hum Vaccin Immunother*, 10: 1155–1165.
- Ronco LV, Karpova AY, Vidal M, Howley PM. 1998. Human papillomavirus 16 E6 oncoprotein binds to interferon regulatory factor-3 and inhibits its transcriptional activity. *Genes Dev*, 12: 2061–2072.
- Scheffner M, Huibregtse JM, Vierstra RD, Howley PM. 1993. The HPV-16 E6 and E6-AP complex functions as a ubiquitin-protein ligase in the ubiquitination of p53. *Cell*, 75: 495–505.
- Schiller JT, Lowy DR. 2012. Understanding and learning from the

- success of prophylactic human papillomavirus vaccines. *Nat Rev Microbiol*, 10: 681–692.
- Serrano B, Alemany L, Tous S, Bruni L, Clifford GM, Weiss T, Bosch FX, de Sanjosé S. 2012. Potential impact of a nine-valent vaccine in human papillomavirus related cervical disease. *Infect Agent Cancer*, 7:38.
- Tang AL, Owen JH, Hauff SJ, Park JJ, Papagerakis S, Bradford CR, Carey TE, Prince ME. 2013. Head and neck cancer stem cells: the effect of HPV--an in vitro and mouse study. *Otolaryngol Head Neck Surg*, 149: 252–260.
- Tewari KS, Sill MW, Long HJ 3rd, Penson RT, Huang H, Ramondetta LM, Landrum LM, Oaknin A, Reid TJ, Leitao MM, Michael HE, Monk BJ. 2014. Improved survival with bevacizumab in advanced cervical cancer. *N Engl J Med*, 370: 734–743.
- The Cancer Genome Atlas Network (TCGA). 2015. Comprehensive genomic characterization of head and neck squamous cell carcinomas. *Nature*, 517: 576–582.
- Thomas MC, Chiang CM. 2005. E6 oncoprotein represses p53-dependent gene activation via inhibition of protein acetylation independently of inducing p53 degradation. *Mol Cell*, 17: 251–264.
- Tran NP, Hung CF, Roden R, Wu TC. 2014. Control of HPV infection and related cancer through vaccination. *Recent Results Cancer Res*, 193: 149–171.
- Underbrink MP, Howie HL, Bedard KM, Koop JI, Galloway DA. 2008. E6 proteins from multiple human betapapillomavirus types degrade Bak and protect keratinocytes from apoptosis after UVB irradiation. *J Virol*, 82: 10408–10417.
- Van Pachterbeke C, Bucella D, Rozenberg S, Manigart Y, Gilles C, Larsimont D, Vanden Houte K, Reynders M, Snoeck R, Bossens M. 2009. Topical treatment of CIN 2+ by cidofovir: results of a phase II, double-blind, prospective, placebo-controlled study. *Gynecol Oncol*, 115: 69–74.
- Walboomers JM, Jacobs MV, Manos MM, Bosch FX, Kummer JA, Shah KV, Snijders PJ, Peto J, Meijer CJ, Muñoz N. 1999. Human papillomavirus is a necessary cause of invasive cervical cancer worldwide. *J Pathol*, 189: 12–19.
- Wang JW, Jagu S, Wang C, Kitchener HC, Daayana S, Stern PL, Pang S, Day PM, Huh WK, Roden RB. 2014a. Measurement of neutralizing serum antibodies of patients vaccinated with human papillomavirus L1 or L2-based immunogens using furin-cleaved HPV Pseudovirions. *PLoS One*, 9: e101576.
- Wang X, Wang HK, Li Y, Hafner M, Banerjee NS, Tang S, Briskin D, Meyers C, Chow LT, Xie X, Tuschl T, Zheng ZM. 2014b. microRNAs are biomarkers of oncogenic human papillomavirus infections. *Proc Natl Acad Sci U S A*, 111: 4262–4267.
- Wang X, Wang HK, McCoy JP, Banerjee NS, Rader JS, Broker TR, Meyers C, Chow LT, Zheng ZM. 2009. Oncogenic HPV infection interrupts the expression of tumor-suppressive miR-34a through viral oncoprotein E6. *RNA*, 15: 637–647.
- White EA, Kramer RE, Tan MJ, Hayes SD, Harper JW, Howley PM. 2012a. Comprehensive analysis of host cellular interactions with human papillomavirus E6 proteins identifies new E6 binding partners and reflects viral diversity. *J Virol*, 86: 13174–13186.
- White EA, Sowa ME, Tan MJ, Jeudy S, Hayes SD, Santha S, Münger K, Harper JW, Howley PM. 2012b. Systematic identification of interactions between host cell proteins and E7 oncoproteins from diverse human papillomaviruses. *Proc Natl Acad Sci U S A*, 109: 260–267.
- Wright JD, Viviano D, Powell MA, Gibb RK, Mutch DG, Grigsby PW, Rader JS. 2006. Bevacizumab combination therapy in heavily pretreated, recurrent cervical cancer. *Gynecol Oncol*, 103: 489–93.
- Xu M, Katzenellenbogen RA, Grandori C, Galloway DA. 2013. An unbiased in vivo screen reveals multiple transcription factors that control HPV E6-regulated hTERT in keratinocytes. *Virology*, 446: 17–24.
- Ye F, Zhou C, Cheng Q, Shen J, Chen H. 2008. Stem-cell-abundant proteins Nanog, Nucleostemin and Musashi1 are highly expressed in malignant cervical epithelial cells. *BMC Cancer*, 8: 108–112.
- Zheng ZM, Wang X. 2011. Regulation of cellular miRNA expression by human papillomaviruses. *Biochim Biophys Acta*, 1809: 668–677.
- zur Hausen H. 2009. Papillomaviruses in the causation of human cancers - a brief historical account. *Virology*, 384: 260–265.