

# Striatal Morphology is Associated with Tobacco Cigarette Craving

Amy C Janes<sup>\*1</sup>, Min Tae M Park<sup>2</sup>, Stacey Farmer<sup>1</sup> and M Mallar Chakravarty<sup>2,3</sup>

<sup>1</sup>McLean Imaging Center, McLean Hospital, Harvard Medical School, Belmont, MA, USA; <sup>2</sup>Cerebral Imaging Centre, Douglas Mental Health University Institute, Montreal, QC, Canada; <sup>3</sup>Department of Psychiatry, McGill University, Montreal, QC, Canada

The striatum has a clear role in addictive disorders and is involved in drug-related craving. Recently, enhanced striatal volume was associated with greater lifetime nicotine exposure, suggesting a bridge between striatal function and structural phenotypes. To assess this link between striatal structure and function, we evaluated the relationship between striatal morphology and this brain region's well-established role in craving. In tobacco smokers, we assessed striatal volume, surface area, and shape using a new segmentation methodology coupled with local shape indices. Striatal morphology was then related with two measures of craving: state-based craving, assessed by the brief questionnaire of smoking urges (QSU), and craving induced by smoking-related images. A positive association was found between left striatal volume and surface area with both measures of craving. A more specific relationship was found between both craving measures and the dorsal, but not in ventral striatum. Evaluating dorsal striatal subregions showed a single relationship between the caudate and QSU. Although cue-induced craving and the QSU were both associated with enlarged striatal volume and surface area, these measures were differentially associated with global or more local striatal volumes. We also report a connection between greater right striatal shape deformations and cue-induced craving. Shape deformations associated with cue-induced craving were specific to striatal subregions involved in habitual responding to rewarding stimuli, which is relevant given the habitual nature of cue-induced craving. The current findings confirm a relationship between striatal function and morphology and suggest that variation in striatal morphology may be a biomarker for craving severity.

*Neuropsychopharmacology* (2015) **40**, 406–411; doi:10.1038/npp.2014.185; published online 20 August 2014

## INTRODUCTION

The connection between striatal function and addiction is well founded as this brain region has a major role in subjective drug craving (Volkow *et al*, 2006) and the maintenance of drug-cue associations (See *et al*, 2007). In relation to tobacco smoking specifically, the striatum mediates the rewarding effect of smoking (Barrett *et al*, 2004), the motivation to smoke (Le Foll *et al*, 2014), and damage to this region due to stroke disrupts nicotine dependence (Gaznick *et al*, 2014). The striatum also shows persistent reactivity to smoking-related cues even during abstinence, which may contribute to relapse risk during a quit attempt (Janes *et al*, 2009; McClernon *et al*, 2009). Recently, larger striatal volume has been linked with greater lifetime exposure to nicotine (Das *et al*, 2011), which calls into question whether there is a link between striatal function and structural phenotype. Although previous studies have evaluated striatal volume in the context of addiction (Churchwell *et al*, 2012; Das *et al*, 2011; Ersche

*et al*, 2012), anomalies in subcortical shape are beginning to emerge as a robust endophenotype in neuropsychiatric disorders (Shaw *et al*, 2014), suggesting that this measure may provide additional insight into the relationship between striatal morphology and substance abuse.

To assess the link between craving and striatal anatomy, we used a new segmentation methodology coupled with local shape indices along the striatal surface to estimate striatal volume (Chakravarty *et al*, 2013), surface area, and shape (Raznahan *et al*, 2014). For volumetric analyses, we assessed both total striatal volume as well as striatal subregions to get a sense of both global and localized structural effects. These measures provide a comprehensive evaluation of striatal morphology not easily captured by standard volumetric analyses alone. We then assessed the link between striatal morphology and two measures of cigarette craving: baseline subjective craving as measured by the brief questionnaire of smoking urges (QSU; Cox *et al*, 2001) and craving induced by exposure to smoking-related cues (Janes *et al*, 2013). Although both measures evaluate craving, the QSU assesses craving state prior to exposure to smoking cues, whereas cue-induced craving measures the amount of craving induced by visual stimuli associated with nicotine reward. We hypothesize that smokers who report more intense cigarette craving will have larger striatal volume, which may partly explain the connection between

\*Correspondence: Dr AC Janes, Brain Imaging Center, Mail Stop 310, McLean Hospital, 115 Mill Street, Belmont, MA 02478, USA, Tel: +1 617 855 3244, Fax: +1 617 855 2770,

E-mail: ajanes@mclean.harvard.edu

Received 21 May 2014; revised 27 June 2014; accepted 15 July 2014; accepted article preview online 24 July 2014

larger striatal volume and lifetime cigarette use (Das *et al*, 2011). In addition, we will build upon the existent literature by providing the first analysis that links cigarette craving and more nuanced measures of striatal morphology including surface area and shape assessments.

## MATERIALS AND METHODS

### Participants

Twenty-five nicotine-dependent participants (6 left-handed, 15 women/10 men) participated in all study measures at the McLean Imaging Center of McLean Hospital. Participants were  $26.3 \pm 5.3$  (mean  $\pm$  SD) years old and had an average of  $6.4 \pm 4.1$  pack-years of smoking experience. Pack-years of smoking use were calculated by multiplying the average number of packs of cigarettes smoked per day by the number of years the participant smoked. Inclusion criteria included smoking  $\geq 10$  cigarettes/day over the past 6 months and on average, participants smoked  $14.2 \pm 3.7$  cigarettes per day. All participants were moderately to heavily nicotine dependent as indicated by an average Fagerstrom test for nicotine dependence (Heatherton *et al*, 1991) score of  $6.12 \pm 1.1$ . Participants were assessed by the Structured Clinical Interview for DSM and met criteria for current nicotine dependence. Participants had an average expired carbon monoxide level of  $24.4 \pm 12.4$  ppm and  $15 \pm 2.1$  years of education. Participants were excluded if they had a lifetime diagnosis of the following conditions: organic mental disorder, bipolar depression, or schizophrenia spectrum disorder or depressive symptoms in the past 6 months. Participants were excluded for current substance use disorder other than nicotine dependence. Smokers also were excluded for pregnancy, recent drug use, or excessive alcohol use. Participants were recruited using online advertisements and fliers posted in the Boston metropolitan area. All participants provided both verbal and written informed consent prior to participating in the study and the institutional review board at McLean Hospital approved this study.

### Magnetic Resonance Imaging

All participants smoked one of their own cigarettes immediately following signing the informed consent to standardize the time as a cigarette was last smoked. MRI scanning began  $\sim 1.5$  h after smoking this cigarette. Scans were acquired on a Siemens Trio 3 Tesla scanner (Erlangen, Germany) with a 32-channel head coil. Multiecho multiplanar rapidly acquired gradient-echo (ME-MPRAGE) structural images were acquired with the following parameters (TR = 2.1 s, TE 3.3 ms, slices = 128, matrix =  $256 \times 256$ , flip angle =  $7^\circ$ , resolution =  $1.0 \text{ mm} \times 1.0 \text{ mm} \times 1.33 \text{ mm}$ ).

### Craving Measures

Craving was assessed just prior to scanning (1.5 h after smoking) using the 10-item QSU (Cox *et al*, 2001), which is a standard tobacco craving measure. Following scanning (3 h after smoking), participants were shown smoking and neutral images and were asked to rate the amount of craving

they experienced when viewing each image on a five-point scale. Smoking and neutral images were comprised of images from various sources including the International Smoking Image Series (Gilbert and Rabinovich, 1999) and those used in our prior work (Janes *et al*, 2013) and by Kober *et al* (2010). To control for visual characteristics, all images were converted to gray scale. The smoking images represented people smoking, hands holding cigarettes, and items such as cigarettes. The neutral images, which lacked the smoking element, were matched for content and included images of people, hands holding objects such as paintbrushes, and items such as pens. One hundred and forty-four images of each type were presented. The average difference in craving to smoking vs neutral images was then calculated and was associated with brain morphology. Average reported subjective craving between smoking and neutral images was compared using a paired *t*-test.

### Striatal Volume and Surface Area

Neuroanatomical segmentations are typically limited in accuracy by errors in the transformation estimation, irreconcilable neuroanatomical differences between the neuroanatomy of the template and the subject, and resampling errors after the application of the transformation to the labels (Collins and Pruessner, 2010; Heckemann *et al*, 2006). MAGEt Brain limits these errors by using a modified 'multi-atlas'-based strategy while still relying on a single well-defined template. As input, the algorithm uses a single atlas containing the striatum, which was previously defined using a three-dimensional reconstruction of serial histological data (Chakravarty *et al*, 2006) that was warped to an MRI template (Holmes *et al*, 1998). The atlas is first customized to a subset of the data set (21 subjects, based on an arbitrary subsample of the cohort being studied) using a nonlinear transformation estimated in a region of interest defined around the subcortical structures (Chakravarty *et al*, 2008, 2009, 2012). This set of subjects now acts as a set of templates to which all other subjects are now warped, providing 21 candidate segmentations for each subject and effectively uses the neuroanatomical variability of the cohort being analyzed to improve overall segmentation accuracy. The final segmentation is decided upon by using a voxel-wise majority vote (ie, the label occurring most frequently at a specific voxel location is retained; Collins *et al*, 1995). In our original paper, we demonstrated high overlap between our method and gold standards manually derived on MRI data (Dice Kappa of 0.861 when using MAGEt Brain with the ANIMAL algorithm (Collins *et al*, 1995) for nonlinear transformation estimation; Chakravarty *et al*, 2013). As in our recent work (Raznahan *et al*, 2014), we also quantified local surface area. A surface-based model was warped to each subject using all 21 transformation estimated through the MAGEt Brain segmentation procedure. To ensure homology between vertices, all surface vertices were redefined using a Voronoi diagram (see Lyttleton *et al*, 2009). The median position was taken for each coordinate at each vertex in order to create a 'median surface' representation. This can be considered to be the surface-based homolog to majority-vote label fusion measures and has the benefit of filtering noise and outliers from the surface-based representation. Total surface area was

then estimated as the sum of the surface area for each polygon in the surface. If a significant relationship between craving and total striatal morphology was identified, a striatal subregion analysis was then conducted to identify a more localized effect. Striatal subregion volumes were estimated using the MAGE<sub>T</sub> Brain algorithm as well. In this case, the striatal mask was subdivided in the original histological data set in order to identify the ventral striatum and the caudate and putamen.

### Vertex-Wise Analyses of Striatal Shape: Surface Displacement

Shape analysis was carried out using a surface-based methodology proposed by Lerch *et al* (2008). First, surface-based representations of the striatum defined on the input atlas were estimated using the marching cubes algorithm (Lorensen and Cline, 1987) and morphologically smoothed using the AMIRA software package (Visage Imaging, San Diego, CA, USA). The resulting surfaces are defined by ~6300 vertices per striatum. The nonlinear portions of the 21 transformations that map each subject to the input atlas were concatenated and then averaged across the template library in order to limit the effects of noise and error and to increase precision and accuracy (Frey *et al*, 2011; Dorr *et al*, 2008; Borghammer *et al*, 2010). The dot product between the nonlinear deformation vector (of the inverse of the averaged atlas-to-subject transformation) and the striatum surface normal (a unit vector describing the direction perpendicular to the surface) is estimated and provides a local measure of inward or outward displacement along the normal (as first described in Lerch *et al* (2008); and as in the method described in Magon *et al*, (2014)). Prior to estimating the dot product between the normal to the surface and the deformation field, we explicitly modeled and removed global linear effects not originally accounted for in the initial linear transformations. This ensures that the contributions of overall differences in brain volume are minimized in this analysis.

### Statistics

RMINC (<https://github.com/mcvaneede/RMINC>), a statistical image analysis software package built to work in the R environment, was used for all surface-based vertex-wise analyses. A linear model was used and all imaging results were corrected using a 10% false discovery rate (FDR) correction (Genovese *et al*, 2002). All analyses with respect to PSYRATS subscales were conducted using the values of the assessment. SPSS was used to conduct correlation and regression analyses. Correlations were conducted to evaluate the relationship between QSU and cue-induced craving measures, whereas regression analyses were used to evaluate associations between brain morphology and craving measures. For the volumetric analyses, total brain volume, age, and sex were included as control variables.


## RESULTS

### Subjective Craving and Smoking History

Participants reported experiencing significantly greater craving when viewing smoking *vs* neutral images ( $t=7.6$ ,  $p<0.0001$ ). Smoking *vs* neutral cue-induced craving was positively correlated with craving as measured by the QSU ( $r=0.42$ ,  $p=0.035$ ).

### Striatal Morphology

**Striatal volume.** A significant positive association was found between left striatal volume and craving as measured by the QSU ( $t=2.6$ ,  $p=0.016$ ; Figure 1) and craving induced by exposure to smoking *vs* neutral cues ( $t=2.3$ ,  $p=0.032$ ; Figure 1). Given this significant association, a follow-up subregion analysis was conducted. Dorsal, but not ventral, striatal volume was related with both the QSU ( $t=2.3$ ,  $p=0.026$ ) and smoking *vs* neutral cue-induced craving ( $t=2.3$ ,  $p=0.03$ ). Craving measured by the QSU was significantly associated with the caudate ( $t=3.3$ ,  $p=0.004$ ), but not the putamen. Smoking *vs* neutral cue-induced craving was not related to either dorsal striatal



**Figure 1** Left striatal volume and surface area relationships with craving. A positive relationship was found between craving (1. QSU and 2. cue-induced) and left striatal volume and surface area.



subregion. No significant associations were found between right total striatal volume and either craving measure.

**Surface area.** A positive association was found between the left striatal surface area QSU ( $t=2.5$ ,  $p=0.02$ ; Figure 1) and smoking *vs* neutral cue-induced craving ( $t=0.02$ ,  $p=0.007$ , Figure 1).

**Striatal shape.** A significant relationship was found between smoking *vs* neutral cue-induced craving and right striatal shape deformations for both local contractions ( $t$ -statistic range:  $-2.7 < t < -3.95$ ) and expansions ( $t$ -statistic range:  $2 < t < 3.56$ ) (FDR corrected 10%; Figure 2). Similar trends were found for the left striatum, but were not significant (Figure 2). No relationship was found between striatal shape and QSU.

## DISCUSSION


These current findings show a relationship between cigarette craving and striatal morphology. Both subjective craving, as measured by the QSU, and craving induced by exposure to smoking cues were associated with larger left striatal volume and surface area. Cue-induced craving was significantly related with more profound right striatal shape deformations. Given the striatum's functional role in craving (Volkow *et al*, 2006), and cue-reactivity (See *et al*, 2007) the current findings suggest a link between striatal morphology and its underlying function. In addition, our novel surface area and shape findings provide more nuanced insight into the relationship between striatal morphology and craving.

The relationship between craving and left lateralized striatal volume and surface area is consistent with previous findings linking stimulant abuse with greater left striatal volume (Das *et al*, 2011; Churchwell *et al*, 2012, Ersche *et al*, 2012). The connection between the left striatum and

substance abuse may be tied to the more dominant role of the left hemisphere in reward and approach behavior (Davidson and Irwin, 1999). In fact, the left striatum shows greater activation following unexpected reward delivery (McClure *et al*, 2003) and sustained activation following reward feedback (Delgado *et al*, 2000). Greater positive incentive motivation also is associated with higher dopamine D2 receptor availability in the left relative to the right striatum (Tomer *et al*, 2008). It is tempting to speculate that the relationship between left striatal volume and craving is due to greater left striatal D2 receptor availability. Such a hypothesis is plausible given that D2 binding potential is positively associated with striatal volume (Woodward *et al*, 2009), and craving (Volkow *et al*, 2006).

When assessing more localized left striatal subregions, both the QSU and smoking cue-induced craving measures were associated with the dorsal, but not ventral striatum. On further inspection, only the QSU was associated with more localized volume in the caudate, indicating that the specificity of the volumetric association depends on the type of craving evaluated. This specific relationship between caudate volume and QSU is consistent with the caudate's role in craving (Volkow *et al*, 2006) and reward processing (Delgado *et al*, 2000). In contrast, smoking cue-induced craving is associated with enhanced dorsal striatal volume, which includes both the caudate and putamen. These findings suggest that craving, induced by exposure to smoking-related cues may involve more distributed striatal structures, whereas subjective craving, independent of cue-exposure, is associated with more localized caudate morphology.

A relationship between smoking cue-induced craving and right dorsal striatal shape was found independent of overall striatal volume, specifically in the ventral anterior putamen and the right caudate tail. The ventral anterior putamen not only is larger in occasional stimulant users (Mackey *et al*, 2014), but this striatal subregion receives projections from the insula (Chikama *et al*, 1997), a brain region implicated in maintaining nicotine dependence (Naqvi *et al*, 2007) and smoking cue-reactivity (Janes *et al*, 2010). The insula and striatum may interact to mediate nicotine dependence, as damage to both of these brain regions disrupts smoking (Gaznick *et al*, 2014), and suggests a possible link between ventral anterior putamen deformations and cue-reactivity due to striatal-insula interactions. The fact that cue-reactivity was associated with shape differences in the caudate tail is particularly interesting given the role of this striatal subregion in visual habits (Fernandez-Ruiz *et al*, 2001). This is relevant to the current findings owing to the habitual nature of smoking cue-reactivity (Everitt *et al*, 2001). The caudate tail facilitates visual habits as this striatal subregion interacts with inferior temporal brain areas as part of the 'visual' corticostriatal loop, thus providing a 'link in a visual pathway underlying oculomotor control and visuo-motor associations' (Saint-Cyr *et al*, 1990, pg.1). Owing to this link, the caudate tail facilitates automatic responding toward environmental stimuli, which have been consistently rewarded (Yamamoto *et al*, 2013). The caudate tail most likely maintains the automatic and habitual responding to the smoking-related images used in the current study, as these images are 'consistently rewarded visual stimuli' owing to their repeated pairing with the reinforcing experience of nicotine. As subcortical shape



**Figure 2** Cue-induced craving and striatal shape deformations. A significant association was found between smoking cue-induced craving and both local (a) contractions and (b) expansions in the right striatum. Colored regions indicate the range of  $t$ -statistics and regions of significance after FDR correction ( $q=0.1$ ), only applicable for the right striatum. An arbitrary threshold ( $t=2$ ) is chosen for trend-level results for the left striatum, which are not significant after correction.

measures are beginning to emerge as a robust endophenotype in neuropsychiatric disorders (Shaw *et al*, 2014), it is possible that individuals with the variability in striatal shape, we report, might be predisposed to form stronger visual habits. However, this possibility is speculative and requires additional testing.

Although the current work shows a clear link between striatal morphology and craving, additional research will aid in our understanding of how striatal morphology contributes to addiction and other habitual behaviors. For instance, future studies should consider different populations of smokers, as we restricted our sample to a relatively young, healthy population. Whether variation in striatal morphology continues to be associated with craving following brain changes associated with aging remains to be determined. Whereas our sample size is comparable to prior anatomical studies of nicotine dependence (Brody *et al*, 2004), research involving larger sample sizes may address some of these additional questions. In addition, the link between craving and striatal morphology prompts the question of whether variation in striatal morphology could be linked with differences in other 'visual habits' relevant to normal learning or other forms of craving and compulsive disorders. For example, enhanced striatal surface area has been associated with obsessive-compulsive disorder (OCD; Shaw *et al*, 2014), suggesting that larger striatal surface area may be a feature of compulsive disorders including OCD and addiction. Finally, our research focused only on the relationship between striatal morphology and cigarette craving as previous research has shown a clear link between stimulant abuse and enhanced striatal volume when substance abusers and healthy controls were compared (Das *et al*, 2011; Churchwell *et al*, 2012, Ersche *et al*, 2012). It is unclear whether variability in striatal morphology, in non-abusing populations also may explain individual differences in craving severity for other, more natural, reinforcers. Collectively, our findings indicate that striatal morphology is related to this brain regions' role in craving and that morphological assessment of the striatum, beyond standard volumetric analysis, provides more nuanced information regarding the relationship between striatal morphology and function.

## FUNDING AND DISCLOSURE

This work was supported by the National Institute on Drug Abuse Grant K01DA029645 (AJ). The other authors declare no conflict of interest.

## REFERENCES

- Barrett SP, Boileau I, Okker J, Pihl RO, Dagher A (2004). The hedonic response to cigarette smoking is proportional to dopamine release in the human striatum as measured by positron emission tomography and [<sup>11</sup>C]raclopride. *Synapse* 54: 65–71.
- Borghammer P, Ostergaard K, Cumming P, Gjedde A, Rodell A, Hall N *et al* (2010). A deformation-based morphometry study of patients with early-stage Parkinson's disease. *Eur J Neurol* 17: 314–320.
- Chakravarty MM, Bertrand G, Hodge CP, Sadikot AF, Collins DL (2006). The creation of a brain atlas for image guided neurosurgery using serial histological data. *Neuroimage* 30: 359–376.
- Chakravarty MM, Broadbent S, Rosa-Neto P, Lambert CM, Collins DL (2009). Design, construction, and validation of an MRI-compatible vibrotactile stimulator intended for clinical use. *J Neurosci Methods* 184: 129–135.
- Chakravarty MM, Felsky D, Tampakeras M, Lerch JP, Mulsant BH, Kennedy J L *et al* (2012). DISC1 and Striatal Volume: A Potential Risk Phenotype For mental Illness. *Front Psychiatry* 3: 57.
- Chakravarty MM, Sadikot AF, Germann J, Bertrand G, Collins DL (2008). Towards a validation of atlas warping techniques. *Med Image Anal* 12: 713–726.
- Chakravarty MM, Sadikot AF, Germann J, Hellier P, Bertrand G, Collins DL (2009). Comparison of piece-wise linear, linear, and nonlinear atlas-to-patient warping techniques: analysis of the labeling of subcortical nuclei for functional neurosurgical applications. *Hum Brain Mapp* 30: 3574–3595.
- Chakravarty MM, Steadman P, van Eede MC, Clacott RD, Gu V, Shaw P *et al* (2013). Performing label-fusion-based segmentation using multiple automatically generated templates. *Hum Brain Mapp* 34: 2635–2654.
- Chikama M, McFarland NR, Amaral DG, Haber SN (1997). Insular cortical projections to functional regions of the striatum correlate with cortical cytoarchitectonic organization in the primate. *J Neurosci* 17: 9686–9705.
- Churchwell JC, Care PD, Ferrett HL, Stein DJ, Yurgelun-Todd DA (2012). Abnormal striatal circuitry and intensified novelty seeking among adolescents who abuse methamphetamine and cannabis. *Dev Neurosci* 34: 310–317.
- Collins DL, Holmes CJ, Peters TM, Evans AC (1995). Automatic 3-D model-based neuroanatomical segmentation. *Hum Brain Mapp* 3: 190–208.
- Collins DL, Pruessner JC (2010). Towards accurate, automatic segmentation of the hippocampus and amygdala from MRI by augmenting ANIMAL with a template library and label fusion. *Neuroimage* 52: 1355–1366.
- Cox LS, Tiffany ST, Christen AG (2001). Evaluation of the brief questionnaire of smoking urges (QSU-brief). in laboratory and clinical settings. *Nicotine Tob Res* 3: 7–16.
- Das D, Cherbuin N, Anstey KJ, Sachdev PS, Eastaer S (2011). Lifetime cigarette smoking is associated with striatal volume measures. *Addict Biol* 17: 817–825.
- Davidson RJ, Irwin W (1999). The functional neuroanatomy of emotion and affective style. *Trends Cog Sci* 3: 11–21.
- Delgado MR, Nystrom LE, Fissell C, Noll DC, Fiez JA (2000). Tracking the hemodynamic responses to reward and punishment in the striatum. *J Neurophysiol* 84: 3072–3077.
- Dorr AE, Lerch JP, Spring S, Kabani N, Henkelman RM (2008). High resolution three-dimensional brain atlas using an average magnetic resonance image of 40 adult C57Bl/6j mice. *Neuroimage* 42: 60–69.
- Ersche KD, Jones PS, Williams GB, Turton AJ, Robbins TW, Bullmore ET (2012). Abnormal brain structure implicated in stimulant drug addiction. *Science* 335: 601–604.
- Everitt BJ, Dickinson A, Robbins TW (2001). The neuropsychological basis of addictive behavior. *Brain Res Rev* 36: 129–138.
- Fernandez-Ruiz J, Wang J, Aigner TG, Mishkin M (2001). Visual habit formation in monkeys with neurotoxic lesions of the ventro-caudal neostriatum. *Proc Natl Acad Sci USA* 98: 4196–4201.
- Frey S, Pandya DN, Chakravarty MM, Bailey L, Petrides M, Collins DL (2011). An MRI based average macaque monkey stereotaxic atlas and space (MNI monkey space). *Neuroimage* 55: 1435–1442.
- Gaznick N, Tranel D, McNutt A, Bechara A (2014). Basal ganglia plus insula damage yields stronger disruption of smoking addiction than basal ganglia damage alone. *Nicotine Tob Res* 16: 445–453.

- Genovese CR, Lazar NA, Nichols T (2002). Thresholding of statistical maps in functional neuroimaging using the false discovery rate. *Neuroimage* 15: 870–878.
- Gilbert DG, Rabinovich NE (1999). *International smoking image series (with neutral counterparts), Version 1.2*. Integrative Neuroscience Laboratory, Department of Psychology, Southern Illinois University: Carbondale, IL, USA.
- Heatherton TF, Kozlowski LT, Frecker RC, Fagerstrom KO (1991). The Fagerstrom Test For Nicotine Dependence: A revision of the Fagerstrom Tolerance Questionnaire. *Br J Addict* 86: 1119–1127.
- Heckemann RA, Hajnal JV, Aljabar P, Rueckert D, Hammers A (2006). Automatic anatomical brain MRI segmentation combining label propagation and decision fusion. *Neuroimage* 33: 115–126.
- Holmes CJ, Hoge R, Collins L, Woods R, Toga AW, Evans AC (1998). Enhancement of MR images using registration for signal averaging. *J Comput Assist Tomogr* 22: 324–333.
- Janes AC, Frederick BB, Richardt S, Burbridge C, Merlo-Pich E, Reshaw PF et al (2009). Brain fMRI reactivity to smoking-related images before and during extended smoking abstinence. *Exp Clin Psychopharmacol* 17: 365–373.
- Janes AC, Pizzagalli DA, Richardt S, Frederick BB, Chuzi S, Pachas G et al (2010). Brain reactivity to smoking cues prior to smoking cessation predicts ability to maintain tobacco abstinence. *Biol Psychiatry* 67: 722–729.
- Janes AC, Ross RS, Farmer S, Frederick BB, Nickerson LD, Lukas SE et al (2013). Memory retrieval of smoking-related images induce greater insula activation as revealed by an fMRI-based delayed matching to sample task. *Addict Biol* (e-pub ahead of print 22 November 2013; doi:10.1111/abd.12112).
- Kober H, Mende-Siedlecki P, Kross EF, Weber J, Mischel W, Hart CL et al (2010). Prefrontal-striatal pathway underlies cognitive regulation of craving. *Proc Natl Acad Sci USA* 107: 14811–14816.
- Le Foll B, Guranda M, Wilson AA, Houle S, Rusjan PM, Wing VC et al (2014). Elevation of dopamine induced by cigarette smoking: novel insights from a [(11)C]-(+)-PHNO PET study in humans. *Neuropsychopharmacology* 39: 415–424.
- Lerch JP, Carroll JB, Spring S, Bertram LN, Schwab C, Hayden MR et al (2008). Automated deformation analysis in the YAC128 Huntington disease mouse model. *Neuroimage* 39: 32–39.
- Lorenson WE, Cline HE (1987). Marching cubes: a high resolution 3D surface construction algorithm. *AMC SIGGRAPH Computer Graphics* 21: 163–169.
- Lytelton OC, Karama S, Ad-Dab'bagh Y, Zatorre RJ, Carbonell F, Worsley K et al (2009). Positional and surface area asymmetry of the human cerebral cortex. *Neuroimage* 46: 895–903.
- Mackey S, Stewart JL, Connly CG, Tapert SF, Paulus MP (2014). A voxel-based morphometry study of young occasional users of amphetamine-type stimulants and cocaine. *Drug and Alcohol Depend* 135: 104–111.
- Magon S, Chakravarty MM, Amann M, Weier K, Maegelin Y, Andelova M et al (2014). Label-fusion-segmentation and deformation-based shape analysis of deep gray matter in multiple sclerosis: the impact of thalamic subnuclei on disability. *Hum Brain Mapp* 35: 4193–4203.
- McClernon FJ, Kozink RV, Lutz AM, Rose JE (2009). 24-h smoking abstinence potentiates fMRI-BOLD activation to smoking cues in cerebral cortex and dorsal striatum. *Psychopharmacology* 204: 25–35.
- McClure SM, Berns GS, Montague R (2003). Temporal prediction errors in a passive learning task activate human striatum. *Neuron* 38: 339–346.
- Naqvi NH, Rudrauf D, Damasio H, Bechara A (2007). Damage to the insula disrupts addiction to cigarette smoking. *Science* 315: 531–534.
- Raznahan A, Shaw PW, Lerch JP, Clasen LS, Greenstein D, Berman R et al (2014). Longitudinal four-dimensional mapping of subcortical anatomy in human development. *Proc Natl Acad Sci USA* 111: 1592–1597.
- Saint-Cyr JA, Ungerleider LG, Desimone R (1990). Organization of visual cortical inputs to the striatum and subsequent outputs to the pallidum-nigral complex in the monkey. *J Comp Neurol* 298: 129–156.
- See RE, Elliott JC, Feltenstein MW (2007). The role of dorsal vs ventral striatal pathways in cocaine-seeking behavior after prolonged abstinence in rats. *Psychopharmacology* 194: 321–331.
- Shaw P, Sharp W, Sudre G, Wharton A, Greenstein D, Raznahan A et al (2014). Subcortical and cortical morphological anomalies as an endophenotype in obsessive-compulsive disorder. *Mol Psychiatry* e-pub ahead of print 11 February 2014 doi:10.1038/mp.2014.3.
- Tomer R, Goldstein RZ, Wang G, Wong C, Volkow ND (2008). Incentive motivation is associated with striatal dopamine asymmetry. *Biol Psychology* 22: 98–101.
- Volkow ND, Wang G, Telang F, Fowler JS, Logan J, Childress AR et al (2006). Cocaine cues and dopamine in dorsal striatum: Mechanism of craving in cocaine addiction. *J Neurosci* 26: 6583–6588.
- Woodward ND, Zald DH, Ding Z, Riccardi P, Ansari MS, Baldwin RM et al (2009). Cerebral morphology and dopamine D2/D3 receptor distribution in humans: a combined [18F]fallypride and voxel-based morphometry study. *Neuroimage* 46: 31–38.
- Yamamoto S, Kim HF, Hikosaka O (2013). Reward value-contingent changes of visual responses in primate caudate tail associated with a visuomotor skill. *J Neurosci* 3: 11227–11238.