



Published in final edited form as:

J Cogn Neurosci. 2014 September ; 26(9): 1981–1991. doi:10.1162/jocn_a_00607.

Disposed to distraction: Genetic variation in the cholinergic system influences distractibility but not time-on-task effects

Anne S. Berry¹, Elise Demeter², Surya Sabhapathy³, Brett A. English⁴, Randy D. Blakely⁴, Martin Sarter^{1,5}, and Cindy Lustig^{1,5}

¹Neuroscience Program, University of Michigan

²Center for Cognitive Neuroscience, Duke University

³University of Michigan Medical School

⁴Department of Pharmacology, Vanderbilt University School of Medicine

⁵Psychology Department, University of Michigan

Abstract

Both the passage of time and external distraction make it difficult to keep attention on the task at hand. We tested the hypothesis that time-on-task and external distraction pose independent challenges to attention, and that the brain's cholinergic system selectively modulates our ability to resist distraction. Participants with a polymorphism limiting cholinergic capacity (Ile89Val variant (rs1013940) of the choline transporter gene *SLC5A7*) and matched controls completed self-report measures of attention and a laboratory task that measured decrements in sustained attention with and without distraction. We found evidence that distraction and time-on-task effects are independent and that the cholinergic system is strongly linked to greater vulnerability to distraction. Ile89Val participants reported more distraction during everyday life than controls, and their task performance was more severely impacted by the presence of an ecologically valid video distractor (similar to a television playing in the background). These results are the first to demonstrate a specific impairment in cognitive control associated with the Ile89Val polymorphism, and add to behavioral and cognitive neuroscience studies indicating the cholinergic system's critical role in overcoming distraction.

“Pay attention!” You may have given yourself this exhortation during a long, boring drive as you realized both your mind and your car were drifting off the road, or received it from your conversation partner when they noticed your eyes wandering towards an attractive stimulus walking by. Both extended task periods and external distractors challenge our ability to remain focused on the task at hand, and individuals with psychiatric disorders such as schizophrenia and attention deficit hyperactivity disorder (ADHD) may be especially vulnerable to such challenges. Effective treatment of these vulnerabilities will require a better understanding of whether they reflect general declines in cognitive control or more specific processing deficits, and of the underlying neural systems that may be targets for

pharmaceutical intervention. Here we present evidence that performance declines caused by time-on-task demands and external distraction are independent, and that the cholinergic system plays an especially important role in resisting external distraction.

Intuitively, one might expect that any demand on cognitive control, regardless of its source, would reduce performance, and that compounding such demands would have an especially detrimental effect. That is, external distractors might be especially tempting when one is already having difficulty maintaining attention on an attended task. In turn, competition from external distractors may make it especially difficult to maintain focus, causing steeper time-on-task declines. However, the reality is much more complex. Vulnerability to external distraction is typically increased by cognitive load, but decreased by perceptual load (Lavie, Hirst, de Fockert, & Viding, 2004; Lavie, 2010). Time-on-task performance declines have been attributed by some to overloads of attention and by others to underloads of attention (see discussion by Pattyn, Neyt, Herdericlx, & Soetens, 2008), suggesting opposite predictions for the effects of adding an additional cognitive load in the form of external distraction.

Previous attempts to test potential interactions between distraction and time-on-task effects have yielded ambiguous results. Rosenberg, Noonan, DeGutis, & Esterman (2013) used a continuous performance test (detect female target faces versus male nontarget faces) that showed significant declines in both accuracy and response-time stability over a 12-minute period, and manipulated whether or not the background consisted of distracting scenes. This manipulation did not affect performance overall or the slope of time-on-task declines, but did alter individual-differences factors: Only in the distraction condition did mindfulness scores predict time-on-task declines. In another study, Breckel, Giessing, & Thiel (2011) added motion distractors (moving vs stationary dots) to a simple signal detection task. However, these “distractors” did not affect time-on-task declines on the minutes scale, and paradoxically alleviated performance declines associated with long inter-target intervals on no-distractor (stationary) trials. The distraction manipulation also had no effect on brain activity (measured using fMRI) related to time-on-task effects and inter-target interval. Looking in the other direction, activity related to processing the motion distractors was not affected by time-on-task, and there were mixed results in regions involved in motion processing (some areas increasing, others decreasing) as a function of inter-target interval.

In short, the bulk of the evidence from these studies suggests that time-on-task and external distractor effects are independent. However, the lack of an overall performance decrement as a result of the nominal distractor manipulations in these studies reduces the support for this conclusion. That is, without a reliable main effect of distraction, it is hard to know whether distractor effects might interact with time-on-task effects. The neuroimaging data are also somewhat ambiguous, as it is hard to distinguish which changes represent the engagement of cognitive control to inhibit distractor processing from those that may represent increased distractor processing because such inhibition has failed (see also Demeter, Hernandez-Garcia, Sarter, & Lustig, 2011).

We therefore took a different approach, applying a manipulation that creates robust distractor effects to a task with rapid time-on-task performance declines, and asked whether

time-on-task and distractor effects might be related to different neuromodulatory systems. This approach also has a translational advantage: Developing effective drug treatments for specific aspects of attentional function will require understanding their links to the neuromodulator systems targeted by such treatments. Time-on-task effects have been linked to genetic variation in the dopaminergic system (Lim et al., 2012); here we asked whether distraction effects might be linked to genetic variation in the cholinergic system.

The basal forebrain cholinergic system projects throughout neocortex and modulates several neurocognitive functions, including both perceptual and goal-driven aspects of attention (e.g., Rokem, Landau, Garg, Prinzmetal, & Silver, 2010; Silver, Shenhav, & D'Esposito, 2008; see Demeter & Sarter, 2013 for a recent review). Although the cholinergic system has been traditionally described as a diffuse neuromodulatory system contributing to arousal, plasticity, and improved signal-to-noise ratio (see Picciotto, Higley, & Mineur, 2012 for a recent review of this evidence), current models suggest that acetylcholine is also capable of mediating specific cognitive operations through spatially restricted signaling (Hasselmo & Sarter, 2011).

In particular, rodent studies indicate that right-lateralized frontoparietal cholinergic circuitry mediates the ability to resist distraction (Broussard, Karelina, Sarter, & Givens, 2009; Gill, Sarter, & Givens, 2000; St. Peters, Demeter, Lustig, Bruno, & Sarter, 2011). When humans are tested in the same paradigm, they show increased right prefrontal activation during the distraction condition, paralleling rodents' increase in right prefrontal acetylcholine levels. Translational studies using human clinical populations (e.g., Demeter, Guthrie, Taylor, Sarter, & Lustig, 2013; Pa et al., 2013) and animal models (see reviews by Sarter, Martinez, & Kozak, 2009; Lustig, Kozak, Sarter, Young, & Robbins, 2012) also suggest that cholinergic pathways play a critical role in the increased vulnerability to distraction seen in neurological disorders including schizophrenia and mild cognitive impairment.

We tested the hypothesis that increased vulnerability to distraction would be associated with genetic variation affecting the high-affinity choline transporter (CHT), which is responsible for transporting choline into the nerve terminal for the synthesis of acetylcholine (ACh). Mice with reduced CHT expression have normal ACh release and choline clearance at baseline, but are impaired on both these measures in response to task demands (Parikh, St. Peters, Blakely, & Sarter, 2013). In humans, the Ile89Val variant (rs1013940) of the CHT gene *SLC5A7* reduces the rate of choline transport by approximately 40–60% compared to the major allele (Okuda, Okamura, Kaitsuka, Haga, & Gurwitz, 2002). The frequency of the Ile89Val variant in normal Caucasian subjects, ~6% (English et al., 2009), is equivalent to a prevalence in this population in the United States of over 10 million individuals according to recent census data. This variant occurs more often in ADHD patients than controls, and has been linked to greater symptom severity in depressed patients (English et al., 2009; Hahn et al., 2008).

The link to depression suggests that individuals with the Ile89Val allele might be more likely to engage in rumination and mind-wandering, but based on previous human and animal data linking the cholinergic system to distraction, we expected an increased susceptibility to distraction to be their major difference from control participants without the

allele. To preview our results, Ile89Val participants showed an increased vulnerability to distraction on both self-report measures and laboratory task performance, but were spared on other measures of attentional control, including those involved in maintaining performance over time. These results indicate that time-on-task demands and external distraction tax at least partially dissociable components of cognitive control, and that the cholinergic system plays a particularly important role in resisting distraction.

Experiment 1

METHODS

Participants—617 individuals recruited from the greater Ann Arbor community completed the Poor Attentional Control questionnaire (Huba, Singer, Aneshensel, & Antrobus, 1982; see description below) and contributed saliva samples for genotyping. From this larger pool, 67 Ile89Val heterozygotes were identified and compared to age, gender, and education-matched controls. See Table 1 for demographics.

Genotyping methods—CHT SNP genotyping was carried out using the procedures described in English et al. (2009). Briefly, DNA was extracted from saliva samples using a commercial DNA isolation kit (Genra Systems, Minneapolis, MN) as previously described (Mazei-Robinson, Couch, Shelton, Stein, & Blakely, 2005). An allelic discrimination assay was performed in the Vanderbilt Center for Human Genetics Research DNA Resources Core using TaqMan® SNP Genotyping Assay reagents (Applied Biosystems, Inc). Four nanograms (ng) of DNA were used as template in reactions containing 1X TaqMan® Universal PCR Master Mix, and 900 nM forward (5'-TGTACCAGGTTATGGCCTAGCTT-3') and reverse (5'-ACTGAGATTTGCACTTTCACCTACCT-3') amplification primers, 200 nM VIC® (5'-CAGGCACCAATTGGATA-3') and FAM® (5'-AGGCACCAGTTGGATA-3') dye-labeled probes. Thermal cycling (95°C for 10 min, followed by 50 cycles of 92°C for 15 sec and 60°C for 1 min) and product detection were accomplished using the ABI 7900HT Real-Time PCR System (ABI). Quality control analyses showed that the results were consistent with recommended guidelines (e.g., Edenberg & Liu, 2009). The call rate (rate of at which samples could be successfully assigned a genotype) in our sample was >95% with TaqMan®, with 100% consistency when TaqMan® was compared with gel-based genotyping. No-call TaqMan® samples were reanalyzed using gel-based genotyping.

Attention questionnaire measure

Trait attention: Participants completed 36 items from the Imaginal Processes Inventory (Singer & Antrobus, 1970). Each item consisted of a statement about cognitive function in everyday life (ex. “I find it difficult to concentrate when the TV or radio is on”), and participants rated the degree to which they identified with each statement on a scale from 1 to 5. Our analyses focus on the 15 items that make up the Poor Attentional Control (PAC) subscale identified in a later factor analysis (Huba et al., 1982).

The PAC has good internal consistency (coefficient alpha = .83) and test-retest reliability ($r = .73$; see also Tanaka & Huba, 1985–1986). It can be subdivided into subscales (5

questions each) of distractibility, mind-wandering, and boredom. Although Huba et al. (1982) do not provide psychometric data on these subscales, analyses of a large dataset from our lab ($N = 510$; see Berry, Li, Lin, & Lustig, in press) indicate good internal consistency within subscales (mind-wandering coefficient $\alpha = .84$, distraction coefficient $\alpha = .79$, boredom coefficient $\alpha = .77$). The subscales also have reasonable discriminant validity (average correlation between subscale total and items not in that subscale all $r < .49$ compared to items in that subscale all $r > .72$).

RESULTS AND DISCUSSION

As illustrated in Table 1, Ile89Val participants reported experiencing more distractibility and mind-wandering in everyday life than did controls, but not more boredom. The groups' equivalence on boredom is important, as it indicates that Ile89Val participants were not simply interpreting the scale differently and marking all items higher than controls.

The Ile89Val participants' higher ratings for distraction were in line with our predictions, based on the cholinergic system's role in resisting distraction as described in the Introduction. The higher ratings for mind-wandering were somewhat less expected. However, in everyday life it may be difficult to disentangle the subjective experiences of mind-wandering and distraction. For example, in factor analyses "...I notice all the other things around me I could be doing" groups with the distraction factor and involves external distractors whereas "...thoughts unrelated to my work always creep in" groups with the mind-wandering factor and involves only internal experiences. Despite these distinctions on a principled basis (external vs internal) and in large-scale factor analyses, it is easy to see how they might overlap or become confused in individual subjects' self-reports.

We therefore conducted a second experiment with a clearer operationalization of external distraction versus other, presumably internal challenges to attentional control and performance, including those that result from extended time on task. In particular, when asked about distractibility, Experiment 1 participants regardless of genotype gave high ratings to items such as "Faced with a tedious job, I notice all the other things around me I could be doing", and "I find it difficult to concentrate when the TV or radio is on." These responses guided the design of the experimental task we used in Experiment 2.

Specifically, we added an external distractor to the Continuous Temporal Expectancy Test (CTET; O'Connell et al., 2009) and examined its effect on overall performance and time-on-task declines. The CTET is a duration judgment task: On most trials, a grid pattern rotates after a standard duration (800 ms), but on target trials it takes slightly longer (1020 ms). Because the grid pattern is the same for target and nontarget trials, there is no bottom-up perceptual change signaling the presence of a target. Instead, duration judgments require continuous, focused attention to time and are exquisitely sensitive to fluctuations in attention; in some cases performance declines may occur in as few as four trials (Zakay & Block, 1997; Penney, Gibbon, & Meck, 2000; Lustig & Meck, 2005). O'Connell et al. found that the ability to detect the target declined linearly over 3 minutes, and that failures to detect the target were predicted up to 20 seconds beforehand by trends in alpha-band activity (measured using electroencephalography (EEG)) thought to index mind-wandering.

The CTET thus constituted the “tedious task” described in the self-report data. To mimic the distraction caused by the TV or radio, next to the main task computer we placed a laptop playing a series of distracting video clips. This compelling distractor, similar to the situations that participants described as most distracting in everyday life, allowed us to test for potential interactions between distraction and time-on-task demands with greater sensitivity than previous studies that did not find overall distractor effects. To preview our results, we found that both distraction and time-on-task had robust but independent effects on performance, and that Ile89Val participants were especially susceptible to distraction.

Experiment 2

METHODS

Participants—32 Ile89Val heterozygotes agreed to return for additional testing including the CTET with distractor and more questionnaire measures. The control group consisted of 32 individuals homozygous for the major allele and matched with the heterozygotes on age, gender, education, and PAC scores. (Table 2). We matched participants on PAC score to reduce potential concerns that a finding of increased distractibility by Ile89Val participants on the laboratory task might be an artifact of selection bias. That is, if we had *not* matched the samples for this experiment on PAC score, there might have been concerns that we happened to pick low-distractibility participants from the control population and high-distractibility participants from the Ile89Val population, and thereby inflated our chances of finding a group difference on the laboratory task. Instead, by matching the groups on PAC score, we conducted a conservative test, as any selection bias created using this method would be against our preferred hypothesis. (i.e., we have likely picked control participants relatively high in the distractibility distribution for their population, and Ile89Val participants relatively low in the distractibility distribution for their population) Our results may thus under-estimate the size of the group difference in the distraction effect. Ile89Val has been associated with ADHD and depression, so to maintain sample size, we matched subjects as closely as possible on history of diagnosis rather than excluding on this basis. (Analyses excluding these subjects gave similar results with slightly reduced power.)

Additional screening measures

Depression: Hahn et al. (2008) found that Ile89Val was associated with depression severity (though not incidence) in a clinical population. We therefore matched participants on depression ratings using the Patient Health Questionnaire-9 (PHQ-9; Kroenke, Spitzer, & Williams, 2001). Participants respond to each of 9 symptom questions using a scale of 0–3. Possible scores range from 0–27, with 0 indicating no self-reported experience of depression and 27 reflecting severe depression symptoms.

Sleep and other health measures: Because the cholinergic system is also involved in the sleep/wake cycle, and sleep quality can affect attention, we wanted to ensure that our groups were also matched on this front. The Pittsburgh Sleep Quality Index (PSQI; Buysse, Reynolds, Monk, Berman, & Kupfer, 1989) consists of 10 multi-component questions. Scores range from 0–21 with 0 indicating the best sleep quality. The Berlin Sleep Questionnaire (Netzer, Stoobs, Netzer, Clark, & Strohl, 1999) consists of 10 questions

relating to snoring, sleep problems, and blood pressure, as well as age and body mass index. Participants are considered at high risk for sleep apnea if they have a positive score (more than 2 points) in two categories. Participants also completed the standard health, demographics, and screening form used in our laboratory, which asks questions about medications and medical or psychiatric conditions that could affect performance.

CTET with video distractor—CTET procedures generally followed those used by O’Connell et al. (2009), with some modifications (e.g., target and standard durations, response window) based on pilot testing to make it feasible for subjects with a wide range of ability. The task was presented on a Dell PC using Presentation software (Neurobehavioral Systems, Inc.). On each trial, participants were presented with a black and white 10x10 grid of square tiles (15 mm² each) divided diagonally into black and white halves. On standard trials, the grid randomly changed rotation (90, 180, or 270°) after 800 ms; on target trials it rotated after 1070 ms. (Figure 1). Participants were instructed to press the spacebar as soon as they detected the target.

The distraction manipulation was implemented using a laptop oriented 32° to the left of the CTET task and 65 cm from the participant. In the No Distractor condition, the laptop was silent and displayed a gray screen. In the Distractor condition, the laptop played a series of 30 second video clips from various sources (e.g., cartoons, movies, sports) with sound presented via headphones. None of the videos contained music or other obviously rhythmic content, or overtly violent or sexual content.

Responses were recorded as correct (hits) up to 1.5 seconds following target offset. Responses outside this window were coded as false alarms (FAs). Participants received feedback at the end of each run. Before beginning the experiment, participants were given 6 practice runs that contained 3 targets each. For the first practice run, the duration difference between target and standard trials was exaggerated to ensure participants understood the task rules (standard: 800 ms; target: 1600 ms). Participants were informed that the timing parameters of the subsequent 5 practice runs would be more challenging and identical to the test runs (standard: 800 ms; target: 1070 ms). To ensure participants were able to discriminate between standard and target durations, they were required to continue practice until they achieved 100% (3/3 targets detected) if they had not already reached this criterion in the mandated 5 practice runs.

Participants then performed 9 task runs (4 minutes each) with 24 targets per run.¹ Stimuli were pseudo-randomly intermixed, with 4–8 target trials presented per minute, and 7–14 standard duration stimuli presented between each target. Participants took a 1 minute break between each experimental run. For each participant, 4 runs were presented in the No Distractor condition, and 4 runs were presented in the Distractor condition. Distractor and No Distractor runs alternated and their order was counterbalanced across participants. Participants completed 1 additional “Boring Distractor” run for piloting purposes that is not

¹In this and other studies, time-on-task effects occur *within* runs; there are no systematic differences *between* runs (i.e., performance is not worse in run 5 than run 1; see also Berry et al., in press; O’Connell et al., 2009).

included in the present analysis. The single Boring Distractor run was inserted as either the 1st, 3rd, 6th, or 9th run.

Post-experiment questionnaire

Distractor video content: After the CTET, participants completed a surprise 15-item multiple-choice quiz assessing memory for the video content. Similar procedures have revealed relationships between memory for the distracting stimuli and the impact of distraction on performance, as well as experimental group differences (Berry et al., in press; Biss, Ngo, Hasher, Campbell, & Rowe, 2013; Clapp & Gazzaley, 2012; Gazzaley, Cooney, Rissman, & D’Esposito, 2005).

State attention: After completing the CTET and the quiz for memory of the distractors, participants rated the level of distractibility, mind-wandering, and boredom they experienced during the task. These questions were intended to be similar in form and content to items from the PAC trait attention measure. Participants were given five statements and asked to rate the degree to which they identified with each statement on a scale from 1 to 5 (see Table 3). Questions 1, 2, and 4 measured mind-wandering, question 3 measured boredom, and question 5 measured distractibility. We emphasized mind-wandering (3 questions) when constructing the questionnaire as that had been the focus of the original O’Connell et al. (2009) paper. However, subsequent analyses pooling across CTET experiments in our lab show that question 4 consistently shows the highest correlations with the PAC mind-wandering score and with CTET performance. For that reason, and so that an equal number of items are used for the mind-wandering, distractibility, and boredom scores, as in prior work (Berry et al., in press), we focus on question 4 as the “state” measure of mind-wandering.

RESULTS AND DISCUSSION

Consistent with previous papers (Berry et al., in press; O’Connell et al., 2009), analyses of CTET performance focused on hits (correct target detections); analyses of d' measures yields similar results. Greenhouse-Geisser sphericity correction was applied as needed for reporting p values but degrees of freedom are reported as integers in the text for easier reading. Effect sizes are reported using η^2_G (Bakeman, 2005), which gives smaller values than the frequently-used η^2_P but is preferable as it reduces error when comparing across studies (Fritz, Morris, & Richler, 2012).

As seen in Figure 2, external distraction and time-on-task effects were independent, and heterozygotes showed a specific sensitivity to distraction. ANOVA results with within-subjects factors distraction (No Distractor, Distractor), time (Minute 1, 2, 3, 4), and between-subjects factor genotype (Control, Ile89Val) revealed main effects of distraction, $F(1, 62) = 59.27, p < .0005, \eta^2_G = .07$, and time-on-task, $F(3, 186) = 21.35, p < .0005, \eta^2_G = .03$, that did not interact, $F < 1$. (Figure 2). Of primary interest, heterozygotes were more sensitive than controls to distraction, $F(1, 62) = 11.65, p = .001, \eta^2_G = .01$, but not time-on-task, $F < 1$.² The three-way interaction between distraction, time-on-task, and genotype did not approach significance, $F < 1$. Though groups were matched on PAC distractibility, the CTET with distraction was sensitive to group differences in distractor vulnerability

suggesting that the PAC and laboratory performance measures captured at least partially distinct aspects of distractibility.

False alarm (FA) rates were low and did not differ across groups. An ANOVA with within-subjects factors distraction (No Distractor, Distractor) and time (Minute 1, 2, 3, 4), and between-subjects factor genotype (Control, Ile89Val) revealed only a main effect of distraction, $F(1, 62) = 5.10$, $p = .03$, $\eta^2_G < 0.01$, such that FA rates were higher during distraction (No Distractor $M = .94\%$, $SD = 1.7$; Distractor $M = 1.15\%$, $SD = 2.04$). There was no effect of time or genotype, and no interactions (all $F < 1$).

Post-task ratings of subjective experience generally followed the patterns seen for ratings of everyday attentional function (PAC scores) in Experiment 1: Ile89Val participants tended to give higher ratings for items concerning mind-wandering and distractibility, but not boredom. (Table 3). Although these effects are small, they are notable given that in selecting our participants, we had matched Ile89Val and control participants as closely as possible in their PAC subjective ratings of everyday attention.

Thus, even though the Ile89Val participants tested in this experiment were selected to give similar ratings as controls for attentional function in everyday life, they still tended to show an increased subjective as well as objective susceptibility to distraction during the experimental task. In addition Ile89Val participants tended to have better memory for the video contents. Together with the CTET performance data, these patterns suggest that Ile89Val individuals are more susceptible to distraction, but may benefit from better memory for nominally irrelevant information.

To determine how task performance related to subjective measures of attention, we examined how CTET performance correlated with the self-report measures of attention in everyday life and state of attention during the task. To reduce the number of comparisons and ensure replication across datasets, unless otherwise noted we restrict our analyses to those of theoretical interest and also examined in a previous experiment using an undergraduate sample (Berry et al., in press). In the text we note the consistencies and discrepancies compared to Berry et al. so that the reader can weigh results accordingly. Full correlation tables for the combined group and each subgroup are available upon request.

We first asked how performance correlated with the PAC “trait” measures of attention, with particular interest in those measures related to distraction. (Figure 3). Across both groups, there was a small-to-moderate correlation such that participants who reported more distractibility in everyday life also showed a greater distractor-related drop in performance on the CTET, $r = .25$, $p = .05$. In a previous experiment using an undergraduate sample (Berry et al., in press), this correlation was in the same direction, but did not reach significance. The greater variance in the distraction effect in the present study (due to Ile89Val participants’ distractor vulnerability) may have contributed to this difference. In our undergraduate sample, self-reported mind-wandering negatively correlated with overall performance in the No Distractor condition, but this relationship did not reach significance

²As a check of reliability, we split each group in half (even/odd pairs) and re-ran the analysis on the subsamples; effects were in the same direction and of similar effect size as those reported here.

in the current sample ($r = -.19, p = .13$). Thus, across studies, correlations between performance and self-reported everyday attentional function are of modest size but in the expected direction.

Likewise consistent with the Berry et al. (in press) undergraduate sample, in this community sample stronger correlations were found between task performance and post-task ratings. (Table 3, Figure 3). Target detection in the No Distractor condition was negatively related to difficulty keeping attention focused ($r = -.40, p = .001$). Greater performance declines as a result of the video distractor correlated with greater self-rated distraction by the videos, $r = .27, p = .03$, and to better memory for the videos' content, $r = .33, p = .008$.

A reviewer noted that the mean group differences in the size of the distractor effect on performance might have inflated the size of the cross-group correlations with this effect. To examine this possibility, we calculated the correlations with this factor for each group. For the correlation with trait distractibility as measured by the PAC subscale, the difference between the groups was relatively small ($r = .16$ for controls, $r = .27$ for Ile89Val participants), perhaps because we had matched the groups on PAC scores. However, for both self-rated distraction ($r = -.01$ for controls, $r = .34$ for Ile89Val participants) and memory for the videos' content ($r = -.004$ for controls, $r = .53$ for Ile89Val participants), correlations with the distractor effect on performance were largely driven by Ile89Val participants. In other words, in addition to being more vulnerable to distraction's detrimental effects on performance, Ile89Val participants were also more sensitive to its subjective effects and its potential benefits to later memory.

GENERAL DISCUSSION

The present results reveal three major findings. First, although both distraction and time-on-task reduce performance, their effects are independent. This seems inconsistent with simple "underload" or "overload" resource explanations of vigilance decrements. Instead, these different types of demand appear to tap dissociable attention control processes. Although neuroimaging studies suggest a core network involved in stable task control (e.g., Dosenbach et al., 2006), recent studies focused on time-on-task or fatigue effects show substantial variance in the neural correlates of such effects indicating fatigue of specific task-dependent processes (e.g., Lim et al., 2010; Persson, Larsson, & Reuter-Lorenz, 2013). The task used here may provide a useful tool for examining these processes in clinical and individual-differences research.

Second, in a community-based sample, genetic variation in cholinergic function was related to self-reported everyday (trait) distractibility and mind-wandering, but not boredom. There are legitimate concerns about the degree to which a single nucleotide polymorphism (SNP) can be linked to cognitive processes, as genes are not solitary mediators of behavior. Instead, gene-behavior relationships likely take the form of complex interactions that include other genes and the environment (Thomas, 2010). However, the pattern found here, especially increased distractibility, was predicted *a priori* from prior research and can be interpreted in the context of evidence about its neurobiological impact. When first describing the Ile89Val SNP, Okuda et al. (2002) noted that the effects of reduced choline transport might be

especially evident under demanding conditions, consistent with rodent studies that show that responding to the demands imposed by distraction depends critically on the right basal forebrain cholinergic system (Gill et al., 2000; St. Peters et al., 2011). These findings, as well as parallel human neuroimaging studies showing increased right prefrontal activation in the same conditions that produce increased right prefrontal ACh in rodents, converge to predict the present association between Ile89Val and distractibility.

Third, even after matching participants on self-report measures of distraction in everyday life, Ile89Val participants showed a specific vulnerability to distraction on a laboratory task (the CTET). A distraction score of 12% discriminated the groups with 63% sensitivity and 72% specificity. Conceptually replicating the group differences on the trait measures, when asked about their attentional state during the task, Ile89Val participants gave higher ratings on items concerning distractibility and mind-wandering, but not boredom. The equivalent state and trait boredom scores, No Distractor task performance, and time-on-task declines, as well as superior performance on the memory test, indicate that Ile89Val participants are not generally impaired but have a particular vulnerability to distraction.

To summarize, the association between Ile89Val and distractibility is predicted from molecular, systems, and cognitive neuroscience, and replicates across self-report measures of everyday attention, laboratory task performance, and self-report ratings of attention during the task. In contrast, the link to mind-wandering is more tenuous. It is not clear how mind-wandering relates to the mechanism (attenuated cholinergic response to high task demands) indicated by molecular and cellular studies, and although Ile89Val participants consistently indicated higher rates of mind-wandering on the self-report measures, they did not show worse task performance overall or steeper time-on-task declines. A potential reason for the discrepancy between the self-report and performance measures is that while the former ask about off-track thoughts, those thoughts may not always disrupt performance. Esterman, Noonan, Rosenberg, & DeGutis (2013) recently reported different neural correlates and timescales for vigilance declines, off-track thought, and performance lapses. Of particular interest, default-network activity putatively related to off-track thought had a nonlinear relationship to performance lapses depending on the stability of goal-directed attention. Future investigations using experience-sampling techniques and neural measures with Ile89Val participants may resolve this issue.

The more robust link between Ile89Val and distractibility found here is to our knowledge the first evidence linking this polymorphism to a specific cognitive process, and the first indicating its role in normal cognitive variation beyond clinical conditions such as ADHD. Besides their vulnerability to the distractor's detrimental effects, Ile89Val subjects' better memory for the distractor is reminiscent of findings from older adults showing that increased processing of nominally irrelevant information provides an advantage if that information becomes useful later on (Biss et al., 2013; Clapp & Gazzaley, 2012; Gazzaley et al., 2005). Increased processing of irrelevant information has been linked to creativity in ADHD (White & Shah, 2011); an interesting question for future studies is whether this advantage extends to Ile89Val.

As noted above, understanding the role that any SNP, including Ile89Val, plays in normal cognition and cognitive disorders requires consideration of its interaction with other genetic and environmental factors. For example, variation at a nearby SNP (G to T substitution at CHT 3'UTR) has been linked to corticolimbic reactivity (Neumann et al., 2006), and this variation paired with Ile89Val was especially common in the combined subtype of ADHD (English et al., 2009). Recent evidence from a mouse model suggests that reduced CHT function in combination with dopamine depletion may explain why some Parkinson's patients show pronounced cognitive declines (Zurkovsky et al., 2013) and links between the amyloid precursor protein and CHT indicate a potential role in Alzheimer's disease (Wang, Yang, Wang, & Zheng, 2007).

The present results suggest that the cholinergic system plays an important and specific role in susceptibility to distraction in everyday life (Experiment 1) and on a laboratory task (Experiment 2). They add to a growing body of research that integrates findings from several areas of psychology and neuroscience to clarify links between cognitive functions and the neural systems that underlie them (see also Howe et al., 2013). This integrative approach holds promise for improving our understanding of how genes influence cognition and behavior, and for developing more effective, precisely-targeted treatments when these processes are disordered (Insel et al., 2010).

Acknowledgments

We thank Mina Raj for assistance in creating the stimuli for Experiment 2, Patricia J. Deldin for advising on depression and sleep screening measures and serving as a clinical contact for the depression screen, and John Jonides for comments on the manuscript. We are also grateful to the participants, especially those who returned for Experiment 2, and the many research assistants who facilitated data collection.

Funding

The study was funded by NIH grant 1R01MH086530. A.B. and E.D. were both funded by National Science Foundation Graduate Fellowships.

References

- Bakeman R. Recommended effect size statistics for repeated measures designs. *Behavior Research Methods*. 2005; 37:379–384. [PubMed: 16405133]
- Berry AS, Li X, Lin Z, Lustig C. Shared and distinct factors driving attention and temporal processing across modalities. *Acta Psychologica*. in press.
- Biss RK, Ngo JKW, Hasher L, Campbell KL, Rowe G. Distraction can reduce age-related forgetting. *Psychological Science*. 2013; 24:448–455. [PubMed: 23426890]
- Breckel TPK, Giessing C, Thiel CM. Impact of brain networks involved in vigilance on processing irrelevant visual motion. *Neuro Image*. 2011; 55:1754–1762. [PubMed: 21255659]
- Broussard JI, Karelina K, Sarter M, Givens B. Cholinergic optimization of cue-evoked parietal activity during challenged attentional performance. *European Journal of Neuroscience*. 2009; 29:1711–1722. [PubMed: 19419433]
- Buysse DJ, Reynolds CF 3rd, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Research*. 1989; 28:193–213. [PubMed: 2748771]
- Clapp WC, Gazzaley A. Distinct mechanisms for the impact of distraction and interruption on working memory in aging. *Neurobiology of Aging*. 2012; 33:134–148. [PubMed: 20144492]

- Demeter E, Guthrie SK, Taylor SF, Sarter M, Lustig C. Increased distractor vulnerability but preserved vigilance in patients with schizophrenia: Evidence from a translational Sustained Attention Task. *Schizophrenia Research*. 2013; 144:136–141. [PubMed: 23374860]
- Demeter E, Hernandez-Garcia L, Sarter M, Lustig C. Challenges to attention: A continuous arterial spin labeling (ASL) study of the effects of distraction on sustained attention. *Neuroimage*. 2011; 54:1518–1529. [PubMed: 20851189]
- Demeter E, Sarter M. Leveraging the cortical cholinergic system to enhance attention. *Neuropharmacology*. 2013; 64:294–304. [PubMed: 22796110]
- Dosenbach NUF, Visscher KM, Palmer ED, Miezin FM, Wenger KK, Kang HSC, et al. A core system for the implementation of task sets. *Neuron*. 2006; 50:799–812. [PubMed: 16731517]
- Edenberg HJ, Liu Y. Laboratory methods for high-throughput genotyping. *Cold Spring Harbor protocols*, 2009. 2009 pdb top62.
- English BA, Hahn MK, Gizer IR, Mazei-Robison M, Steele A, Kurnik DM, et al. Choline transporter gene variation is associated with attention-deficit hyperactivity disorder. *Journal of Neurodevelopmental Disorders*. 2009; 1:252–263. [PubMed: 21547719]
- Esterman M, Noonan SK, Rosenberg M, DeGutis J. In the zone or zoning out? Tracking behavioral and neural fluctuations during sustained attention. *Cerebral Cortex*. 2013
- Fritz CO, Morris PE, Richler JJ. Effect Size Estimates: Current Use, Calculations, and Interpretation. *Journal of Experimental Psychology-General*. 2012; 141:2–18. [PubMed: 21823805]
- Gazzaley A, Cooney JW, Rissman J, D'Esposito M. Top-down suppression deficit underlies working memory impairment in normal aging. *Nature Neuroscience*. 2005; 8:1298–1300.
- Gill TM, Sarter M, Givens B. Sustained visual attention performance-associated prefrontal neuronal activity: Evidence for cholinergic modulation. *Journal of Neuroscience*. 2000; 20:4745–4757. [PubMed: 10844044]
- Hasselmo ME, Sarter M. Modes and models of forebrain cholinergic neuromodulation of cognition. *Neuropsychopharmacology: official publication of the American College of Neuropsychopharmacology*. 2011; 36:52–73. [PubMed: 20668433]
- Hahn MK, Blackford JU, Haman K, Mazei-Robison M, English BA, Prasad HC, et al. Multivariate permutation analysis associates multiple polymorphisms with subphenotypes of major depression. *Genes Brain and Behavior*. 2008; 7:487–495.
- Howe WM, Berry AS, Francois J, Gilmour G, Carp JM, Tricklebank M, et al. Prefrontal cholinergic mechanisms instigating shifts from monitoring for cues to cue-guided performance: Converging electrochemical and fMRI evidence from rats and humans. *Journal of Neuroscience*. 2013; 33:8742–8752. [PubMed: 23678117]
- Huba, GJ.; Singer, JL.; Aneshensel, CS.; Antrobus, JS. *Short Imaginal Processes Inventory*. Port Huron, MI: Research Psychologists Press; 1982.
- Insel T, Cuthbert B, Garvey M, Heinssen R, Pine DS, Quinn K, et al. Research Domain Criteria (RDoC): Toward a new classification framework for research on mental disorders. *American Journal of Psychiatry*. 2010; 167:748–751. [PubMed: 20595427]
- Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *Journal of General Internal Medicine*. 2001; 16:606–613. [PubMed: 11556941]
- Lavie N. Attention, distraction, and cognitive control under load. *Current Directions in Psychological Science*. 2010; 19:143–148.
- Lavie N, Hirst A, de Fockert JW, Viding E. Load theory of selective attention and cognitive control. *Journal of Experimental Psychology-General*. 2004; 133:339–354. [PubMed: 15355143]
- Lim J, Ebstein R, Tse CY, Monakhov M, Lai PS, Dinges DF, et al. Dopaminergic polymorphisms associated with time-on-task declines and fatigue in the psychomotor vigilance test. *PloS one*. 2012; 7:e33767. [PubMed: 22438994]
- Lim J, Wu WC, Wang JJ, Detre JA, Dinges DF, Rao HY. Imaging brain fatigue from sustained mental workload: An ASL perfusion study of the time-on-task effect. *Neuroimage*. 2010; 49:3426–3435. [PubMed: 19925871]
- Lustig C, Kozak R, Sarter M, Young JW, Robbins TW. CNTRICS final animal model task selection: Control of attention. *Neuroscience and biobehavioral reviews*. 2012

- Lustig C, Meck WH. Chronic treatment with haloperidol induces deficits in working memory and feedback effects of interval timing. *Brain and Cognition*. 2005; 58:9–16. [PubMed: 15878723]
- Mazei-Robinson MS, Couch RS, Shelton RC, Stein MA, Blakely RD. Sequence variation in the human dopamine transporter gene in children with attention deficit hyperactivity disorder. *Neuropharmacology*. 2005; 49:724–736. [PubMed: 16171832]
- Netzer NC, Stoohs RA, Netzer CM, Clark K, Strohl KP. Using the Berlin Questionnaire to identify patients at risk for the sleep apnea syndrome. *Annals of Internal Medicine*. 1999; 131:485–491. [PubMed: 10507956]
- Neumann SA, Brown SM, Ferrell RE, Flory JD, Manuck SB, Hariri AR. Human choline transporter gene variation is associated with corticolimbic reactivity and autonomic-cholinergic function. *Biological Psychiatry*. 2006; 60:1155–1162. [PubMed: 16876130]
- O’Connell RG, Dockree PM, Robertson IH, Bellgrove MA, Foxe JJ, Kelly SP. Uncovering the neural signature of lapsing attention: Electrophysiological signals predict errors up to 20 s before they occur. *Journal of Neuroscience*. 2009; 29:8604–8611. [PubMed: 19571151]
- Okuda T, Okamura M, Kaitsuka C, Haga T, Gurwitz D. Single nucleotide polymorphism of the human high affinity choline transporter alters transport rate. *Journal of Biological Chemistry*. 2002; 277:45315–45322. [PubMed: 12237312]
- Pa J, Berry AS, Compagnone M, Boccanfuso J, Greenhouse I, Rubens MT, et al. Cholinergic enhancement of functional networks in older adults with mild cognitive impairment. *Annals of Neurology*. 2013; 73:762–773. [PubMed: 23447373]
- Parikh V, St Peters M, Blakely RD, Sarter M. The presynaptic choline transporter imposes limits on sustained cortical acetylcholine release and attention. *Journal of Neuroscience*. 2013; 33:2326–2337. [PubMed: 23392663]
- Pattyn N, Neyt X, Herdericlx D, Soetens E. Psychophysiological investigation of vigilance decrement: Boredom or cognitive fatigue? *Physiology & Behavior*. 2008; 93:369–378. [PubMed: 17999934]
- Penney TB, Gibbon J, Meck WH. Differential effects of auditory and visual signals on clock speed and temporal memory. *Journal of Experimental Psychology-Human Perception and Performance*. 2000; 26:1770–1787. [PubMed: 11129373]
- Persson J, Larsson A, Reuter-Lorenz PA. Imaging fatigue of interference control reveals the neural basis of executive resource depletion. *Journal of Cognitive Neuroscience*. 2013; 25:338–351.
- Picciotto MR, Higley MJ, Mineur YS. Acetylcholine as a neuromodulator: cholinergic signaling shapes nervous system function and behavior. *Neuron*. 2012; 76:116–129. [PubMed: 23040810]
- Rokem A, Landau AN, Garg D, Prinzmetal W, Silver MA. Cholinergic enhancement increases the effects of voluntary attention but does not affect involuntary attention. *Neuropsychopharmacology: official publication of the American College of Neuropsychopharmacology*. 2010; 35:2538–2544. [PubMed: 20811340]
- Rosenberg M, Noonan S, DeGutis J, Esterman M. Sustaining visual attention in the face of distraction: a novel gradual-onset continuous performance task. *Attention, perception & psychophysics*. 2013; 75:426–439.
- Sarter M, Martinez V, Kozak R. A neurocognitive animal model dissociating between acute illness and remission periods of schizophrenia. *Psychopharmacology*. 2009; 202:237–258. [PubMed: 18618100]
- Silver MA, Shenhav A, D’Esposito M. Cholinergic enhancement reduces spatial spread of visual responses in human early visual cortex. *Neuron*. 2008; 60:904–914. [PubMed: 19081383]
- Singer, JL.; Antrobus, JS. *Manual for the Imaginal Processes Inventory*. Princeton, NJ: Educational Testing Service; 1970.
- St Peters M, Demeter E, Lustig C, Bruno JP, Sarter M. Enhanced control of attention by stimulating mesolimbic-cortical cholinergic circuitry. *Journal of Neuroscience*. 2011; 31:9760–9771. [PubMed: 21715641]
- Tanaka JS, Huba GJ. Longitudinal stability of three second-order daydreaming factors. *Imagination, Cognition and Personality*. 1985–86; 5:181–206.
- Thomas D. Gene-environment-wide association studies: emerging approaches. *Nature Reviews Genetics*. 2010; 11:259–272.

- Wang B, Yang L, Wang Z, Zheng H. Amyloid precursor protein mediates presynaptic localization and activity of the high-affinity choline transporter. *Proceedings of the National Academy of Sciences of the United States of America*. 2007; 104:14140–14145. [PubMed: 17709753]
- White HA, Shah P. Creative style and achievement in adults with attention-deficit/hyperactivity disorder. *Personality and Individual Differences*. 2011; 50:673–677.
- Zakay D, Block RA. Temporal cognition. *Current Directions in Psychological Science*. 1997; 6:12–16.
- Zurkovsky L, Bychkov E, Tsakem EL, Siedlecki C, Blakely RD, Gurevich EV. Cognitive effects of dopamine depletion in the context of diminished acetylcholine signaling capacity in mice. *Disease Models & Mechanisms*. 2013; 6:171–183. [PubMed: 22864020]




Figure 1. CTET with video distractor

As shown in (a) each trial consisted of a black and white grid made up of squares divided into triangles. At the end of the trial, the triangles rotated (90,180, or 270 degrees, chosen randomly) to start the next trial. The participant’s task was to press the spacebar when they realized that the grid had taken longer than usual (1070 ms rather than the standard 800 ms) to rotate. (b) The distractor manipulation was implemented using a laptop oriented 32° to the left of the main task computer. In the No Distractor condition, the laptop was silent and displayed a gray screen. In the Distractor condition, it played video clips with sound.




Figure 2. CTET performance with and without video distractor

Markers represent mean proportion of hits for each minute; error bars represent standard error of the mean. Both groups exhibit time-on-task and distraction effects that do not interact, and the distractor effect is larger for IleVal89 participants.




Figure 3. Correlations between distractor-related performance declines in the CTET, self-rated “trait” distractibility measured using the PAC, self-rated “state” distractibility measured using the post-task questionnaire, and performance on the post-task quiz for memory of the content of the distracting videos

Larger numbers indicate greater distractibility. If the data from the control subject in the lower left corner are excluded, the correlations remain in the same direction as indicated in the main text at reduced effect size ($r = .20$ for trait distraction, $r = .22$ for state distraction, and $r = .29$ for distractor memory).

Table 1

Demographics and self-reported everyday attention function (PAC measures; Huba et al., 1982) for Ile89Val participants and controls. Each group included 67 participants (41 females, 26 males).

	Control	Ile89Val	t-test	Effect size (Cohen's d)
Age (yrs)				
M	42.49	42.33	t < 1	d = .01
SD	17.20	17.33	p = .96	
Edu (yrs)				
M	16.08	16.50	t < 1	d = .14
SD	2.50	3.49	p = .43	
Distractibility				
M	13.43	15.16	t = 2.36	d = .41
SD	4.04	4.43	p = .02	
Mind-wandering				
M	12.93	14.45	t = 2.16	d = .36
SD	3.46	4.77	p = .05	
Boredom				
M	12.64	12.97	t < 1	d = .09
SD	3.26	4.12	p = .61	

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2

Demographic and PAC questionnaire measures for participants completing the CTET with distraction task. t-test and Cohen's d refer to the comparison between control and Ile89Val participants. Each group included 19 females, 13 males for a total n = 32 per group.

	Control	Ile89Val	t-test	Effect size (Cohen's d)
<i>Demographic and health measures</i>				
Age (yrs)				
M	45.13	44.66	t < 1	d = .03
SD	16.85	16.99	p = .91	
Edu (yrs)				
M	16.73	16.53	t < 1	d = .06
SD	2.57	3.59	p = .80	
History of ADHD/depression/anxiety (n)	6	7	-	-
Psychotropic medications (n)	6	5	-	-
PHQ-9 depression score				
M	3.50	3.47	t < 1	d = .01
SD	3.97	4.12	p = .98	
Pittsburgh Sleep Quality Index				
M	5.91	4.75	t = 1.17	d = .29
SD	4.32	3.54	p = .25	
<i>Self-report everyday attention measures (PAC)</i>				
Distractibility				
M	13.50	14.59	t = 1.01	d = .25
SD	4.21	4.45	p = .32	
Mind-wandering				
M	12.91	13.38	t < 1	d = .11
SD	3.77	4.47	p = .65	
Boredom				
M	12.94	12.25	t < 1	d = .18
SD	3.30	4.21	p = .47	

Table 3

Post-experiment questionnaire measures for participants completing the CTET with distraction task. t-test and Cohen's d refer to the comparison between control and lle89Val participants.

<i>Post-task questionnaire</i>				
1. At times of this task, it was hard for me to keep my mind from wandering				
M	2.63	3.53	t = 3.02	d = .75
SD	1.21	1.19	p = .004	
2. (reverse scored) During the task, my thoughts seldom shifted from the subject in front of me.				
M	2.81	3.31	1.7	d = .43
SD	1.12	1.23	p = .09	
3. I was easily bored during this task.				
M	2.47	2.63	t < 1	d = .12
SD	1.19	1.43	p = .64	
4. I had difficulty in keeping my attention focused on this long, tedious task				
M	2.59	3.09	t = 1.76	d = .44
SD	0.98	1.28	p = .08	
5. No matter how hard I tried to concentrate, I felt easily distracted by the videos playing				
M	2.47	3.09	t = 2.11	d = .53
SD	1.16	1.20	p = .04	
Recognition memory for distractor content (% correct)				
M	0.60	0.71	t = .193	d = .46
SD	0.26	0.22	p = .06	