

# Chronic kidney disease in disadvantaged populations

G. Garcia-Garcia<sup>1</sup>, V. Jha<sup>2,3,4</sup> on behalf of the World Kidney Day Steering Committee\*

<sup>1</sup>Nephrology Service, Hospital Civil de Guadalajara, University of Guadalajara Health Sciences Center, Guadalajara, Mexico

<sup>2</sup>Postgraduate Institute of Medical Education and Research, Chandigarh, India

<sup>3</sup>George Institute for Global Health, New Delhi, India

<sup>4</sup>University of Oxford, Oxford, UK

## Abstract

The increased burden of chronic kidney disease (CKD) in disadvantaged populations is due to both global factors and population-specific issues. Low socioeconomic status and poor access to care contribute to health care disparities and exacerbate the negative effects of genetic or biological predisposition. Provision of appropriate renal care to these populations requires a two-pronged approach: expanding the reach of dialysis through development of low-cost alternatives that can be practiced in remote locations, and implementation and evaluation of cost-effective prevention strategies. Kidney transplantation should be promoted by expansion of deceased donor transplant programs and use of inexpensive, generic immunosuppressive drugs. The message of World Kidney Day 2015 is that a concerted attack against the diseases that lead to end-stage renal disease, by increasing community outreach, better education, improved economic opportunity, and access to preventive medicine for those at highest risk, could end the unacceptable relationship between CKD and disadvantage in these communities.

## Introduction

*"Of all of the forms of inequality, injustice in health is the most shocking and inhumane."*

Dr. Martin Luther King, Jr.

March 12, 2015 will mark the 10th anniversary of World Kidney Day (WKD), an initiative of the International Society of Nephrology and the International Federation of Kidney Foundations. Since its inception in 2006, WKD has become the most successful effort ever mounted to raise awareness among decision makers and the general public about the importance of kidney disease. Each year, WKD reminds us that kidney disease is common, harmful, and treatable. The focus of WKD 2015 is on chronic kidney disease (CKD) in disadvantaged populations. This article reviews the key links between poverty and CKD and the consequent implications for the prevention of kidney disease and the care of kidney patients in these populations.

CKD is increasingly recognized as a global public health problem and a key determinant of the poor health outcomes. There is compelling evidence that disadvantaged communities, i.e., those from low-resource racial and minority ethnic

communities and/or indigenous and socially disadvantaged backgrounds, suffer from marked increases in the burden of unrecognized and untreated CKD. Although the entire populations of some low- and middle-income countries could be considered disadvantaged, further discrimination on the basis of local factors creates a position of extreme disadvantage for certain population groups (peasants, those living in some rural areas, women, the elderly, religious minorities, etc.). The fact that, even in developed countries, racial and ethnic minorities bear a disproportionate burden of CKD and have worse outcomes suggests that there is much to learn beyond the traditional risk factors contributing to CKD-associated complications (1).

About 1.2 billion people live in extreme poverty worldwide. Poverty negatively influences healthy behaviors, health care access, and environmental exposure, all of which contribute to health care disparities (Table 1) (2). The poor are more susceptible to disease because of lack of access to goods and services, in particular clean water and sanitation, information about preventive behaviors, adequate nutrition, and reduced access to health care (3).

Correspondence: World Kidney Day: <[info@worldkidneyday.org](mailto:info@worldkidneyday.org)>.

\*Members of the World Kidney Day Steering Committee: P.K.T. Li, G. Garcia-Garcia, W.G. Couser, T. Erk, E. Zakharova, L. Segantini, P. Shay, M.C. Riella, C. Osafo, S. Dupuis, Ch. Kernahan.

Received November 11, 2014. Accepted November 11, 2014. First published online March 6, 2015.

**Table 1.** Possible mechanisms by which poverty increases the burden of disease.

Health behavior	Access to health care	Biological factors	Environmental factors
Lack of information on preventive behaviors	Lack of access to health care	Low birth weight	Increased exposure to pollutants
Lack of knowledge on how best to respond to an episode of illness	Greater distance from health care providers	Genetic predisposition	Increased exposure to communicable diseases
Health beliefs and unhealthy behaviors	Lack of out-of-pocket resources	Cumulative biological risk profiles Inadequate nutrition	Lack of clean water and sanitation

### CKD in developed countries

In the United States, ethnic minorities have a higher incidence of end-stage renal disease (ESRD). Despite similar prevalence rates for early stages of CKD (4), poor outcomes such as ESRD are 1.5-4 times higher (2,5-7) among minorities (i.e., African American, Hispanic, and Native American). Poverty further increases the disparity in ESRD rates, with African Americans being at greater risk (8). In the United Kingdom, rates of treated ESRD are higher in ethnic minority groups and with increasing social deprivation (9). Similarly in Singapore, the CKD prevalence is higher among Malays and Indians compared to the Chinese, with socioeconomic and behavioral factors accounting for 70-80% of the excess risk (10).

ESRD incidence is also higher among the less advantaged indigenous populations in developed countries. Canadian First Nations people experience ESRD at rates 2.5-4 times higher than the general population (11). In Australia, the increase in the number of indigenous people starting renal replacement therapy (RRT) over the past 25 years exceeded that of the non-indigenous population by 3.5-fold, largely because of a disproportionate (>10-fold) difference in ESRD due to type II diabetic nephropathy, a disease largely attributable to lifestyle issues such as poor nutrition and lack of exercise (12). Indigenous populations also have a higher incidence of ESRD due to glomerulonephritis and hypertension (13). Compared to the US general population, the ESRD incidence rate is higher in Guam and Hawaii, where the proportion of indigenous people is high, again driven primarily by diabetic ESRD (14). Native Americans have a greater prevalence of albuminuria and higher ESRD incidence rates (15-18). Nearly three-quarters of all incident ESRD cases among this population have been attributable to type II diabetes.

### CKD in developing countries

Poverty-related factors, such as infectious diseases secondary to poor sanitation, inadequate supply of safe water, environmental pollutants, and high concentrations of disease-transmitting vectors, continue to play an important role in the development of CKD in low-income

countries. Although rates of diabetic nephropathy are rising, chronic glomerulonephritis and interstitial nephritis are among the principal causes of CKD in many countries. Of note is the emergence of human immunodeficiency virus (HIV) associated nephropathy as the major cause of CKD in Sub-Saharan Africa (19).

A high prevalence of CKD of unknown etiology has been reported in rural agricultural communities from Central America, Egypt, India, and Sri Lanka. Male farm workers are affected disproportionately, and the clinical presentation is suggestive of interstitial nephritis, confirmed on renal biopsies. The strong association with farm work has led to suggestions that exposure to agrochemicals, dehydration, and consumption of contaminated water might be responsible (20). Additionally, the use of traditional herbal medications is common and frequently associated with CKD among the poor (21,22). In Mexico, CKD prevalence among the poor is two- to three-fold higher than the general population, and the etiology is unknown in 30% of ESRD patients (23-26).

### Low birth weight and risk of CKD in disadvantaged populations

An association between low birth weight (LBW) due primarily to nutritional factors and kidney disease has been described in disadvantaged populations. In Australia, the frequency of LBW in the Aboriginal population is more than double that in the non-Aboriginal population. The high prevalence of albuminuria in this population has been linked to low nephron number related to LBW (27,28). Morphometric studies of kidney biopsies in the Aboriginals show glomerulomegaly, perhaps secondary to nephron deficiency, which might predispose to glomerulosclerosis (29,30). A correlation between LBW and CKD has also been described in poor African Americans and Caucasians living in the southeastern United States (31). Similarly, in an Indian cohort, LBW and early malnutrition were associated with later development of metabolic syndrome, diabetes, and diabetic nephropathy (32). The finding of a high prevalence of proteinuria, elevated blood pressure, and CKD of unknown etiology in South Asian children may also be explained by this mechanism (33,34).

## Disparities in access to RRT

A recent analysis shows that, globally, there were 2.6 million people on dialysis in 2010, with 93% in high- or upper-middle-income countries. By contrast, the number of people requiring RRT was estimated at 4.9-9 million, suggesting that at least 2.3 million died prematurely because of lack of access to RRT. Even though diabetes and hypertension increase the burden of CKD, the current provision of RRT is linked largely to two factors: per capita gross national product, and age, suggesting that poverty is a major disadvantage for receiving RRT. By 2030, the number of people receiving RRT around the world is projected to increase to 5.4 million. Most of this increase will be in the developing countries of Asia and Africa (35).

Access to RRT in the emerging world is dependent mostly on the health care expenditures and economic strength of individual countries, with the relationship between income and access to RRT being almost linear in low- and middle-income countries (19,36). In Latin America, RRT prevalence and kidney transplantation rates correlate significantly with gross national income and health expenditure (37), while in India and Pakistan, less than 10% of all ESRD patients have access to RRT (38). Additionally, developing countries have low transplant rates because of a combination of low levels of infrastructure; geographical remoteness; lack of legislation governing brain death; religious, cultural, and social constraints; and commercial incentives that favor dialysis (39).

There are also differences in utilization of renal replacement modalities between indigenous and non-indigenous groups in the developed countries. In Australia and New Zealand, the proportion of people receiving home dialysis is considerably lower among indigenous people. At the end of 2007 in Australia, 33% of non-indigenous people requiring RRT were receiving home-based dialysis therapies, in contrast to 18% of Aboriginal people. In New Zealand, home-based dialysis was utilized by 62% of the non-indigenous RRT population but only by 42% of Maori/Pacific Islanders (12). The rate of kidney transplantation is also lower among disadvantaged communities. Maori and Pacific people are only 25% as likely to get a transplant as European New Zealanders, and the proportion of indigenous people who underwent transplantation and had a functioning kidney transplant is lower among Aboriginal Australians (12%) compared to non-indigenous Australians (45%). In the United Kingdom, white individuals from socially deprived areas, South Asians, and blacks were all less likely to receive a preemptive renal transplant or living donor transplant than their more affluent white counterparts (9). A multinational study found that, when compared with white patients, the likelihood of receiving a transplant for Aboriginal patients

was 77% lower in Australia and New Zealand, and 66% lower in Canadian First Nations individuals (40).

Disparities in renal care are more evident in developing nations. Data from India shows that there are fewer nephrologists and nephrology services in the poorer states. As a result, people living in these states are likely to receive less care (41). In Mexico, the fragmentation of the health care system has resulted in unequal access to RRT. In the state of Jalisco, the acceptance and prevalence rates in the more economically advantaged insured population were higher (327 per million population [pmp] and 939 pmp, respectively) than for patients without medical insurance (99 pmp and 166 pmp, respectively). The transplant rate also was dramatically different, at 72 pmp for those with health insurance and 7.5 pmp for those without it (42).

## Bidirectional relationship between poverty and CKD

In addition to having a higher disease burden, the poor have limited access to resources for meeting treatment costs. A large proportion of patients who are forced to meet the expensive ESRD treatment costs by incurring out-of-pocket expenditures get pushed into extreme poverty. In one Indian study, over 70% of patients undergoing kidney transplantation experienced catastrophic health care expenditures (43). Entire families felt the impact of this, including job losses and interruptions in the education of children.

## Outcomes

Overall mortality rates among those who do receive RRT are higher in the indigenous, minorities, and uninsured populations, even after adjustment for comorbidities. The hazard ratios for death on dialysis relative to the non-indigenous group are 1.4 for Aboriginal Australians and New Zealand Maori (44). The Canadian First Nations patients achieve target levels for blood pressure and mineral metabolism less frequently (45). In the United States, living in predominantly black neighborhoods was associated with higher than expected mortality rates on dialysis and increased time to transplantation (46). Similarly, black patients on peritoneal dialysis had a higher risk of death or technique failure compared to whites (47).

In Mexico, mortality of patients on peritoneal dialysis is three-fold higher among the uninsured population compared to Mexican patients receiving treatment in the United States, and the survival rate is significantly lower than the insured Mexican population (48), while in India almost two-thirds of the patients are unable to continue dialysis beyond the first 3 months due to financial reasons (49).

## References

- Pugsley D, Norris KC, Garcia-Garcia G, Agodoa L. Global approaches for understanding the disproportionate burden of chronic kidney disease. *Ethn Dis* 2009; 19: S1-S2.
- Crews DC, Charles RF, Evans MK, Zonderman AB, Powe NR. Poverty, race, and CKD in a racially and socio-economically diverse urban population. *Am J Kidney Dis* 2010; 55: 992-1000, doi: 10.1053/j.ajkd.2009.12.032.
- Sachs JD. *Macroeconomics and health: Investing in health for economic development. Report of the Commission on Macroeconomics and Health*. Geneva: WHO; 2001.
- Kalantar-Zadeh K, Block G, Humphreys MH, Kopple JD. Reverse epidemiology of cardiovascular risk factors in maintenance dialysis patients. *Kidney Int* 2003; 63: 793-808, doi: 10.1046/j.1523-1755.2003.00803.x.
- Hsu CY, Lin F, Vittinghoff E, Shlipak MG. Racial differences in the progression from chronic renal insufficiency to end-stage renal disease in the United States. *J Am Soc Nephrol* 2003; 14: 2902-2907, doi: 10.1097/01.ASN.000000915.86.46532.B4.
- Norris K, Nissenson AR. Race, gender, and socioeconomic disparities in CKD in the United States. *J Am Soc Nephrol* 2008; 19: 1261-1270, doi: 10.1681/ASN.2008030276.
- Bruce MA, Beech BM, Crook ED, Sims M, Wyatt SB, Flessner MF, et al. Association of socioeconomic status and CKD among African Americans: the Jackson Heart Study. *Am J Kidney Dis* 2010; 55: 1001-1008, doi: 10.1053/j.ajkd.2010.01.016.
- Volkova N, McClellan W, Klein M, Flanders D, Kleinbaum D, Soucie JM, et al. Neighborhood poverty and racial differences in ESRD incidence. *J Am Soc Nephrol* 2008; 19: 356-364, doi: 10.1681/ASN.2006080934.
- Caskey FJ. Renal replacement therapy: can we separate the effects of social deprivation and ethnicity? *Kidney Int Suppl* 2013; 3: 246-249, doi: 10.1038/kisup.2013.24.
- Sabanayagam C, Lim SC, Wong TY, Lee J, Shankar A, Tai ES. Ethnic disparities in prevalence and impact of risk factors of chronic kidney disease. *Nephrol Dial Transplant* 2010; 25: 2564-2570, doi: 10.1093/ndt/gfq084.
- Gao S, Manns BJ, Culleton BF, Tonelli M, Quan H, Crowshoe L, et al. Prevalence of chronic kidney disease and survival among aboriginal people. *J Am Soc Nephrol* 2007; 18: 2953-2959, doi: 10.1681/ASN.2007030360.
- McDonald S. Incidence and treatment of ESRD among indigenous peoples of Australasia. *Clin Nephrol* 2010; 74 (Suppl 1): S28-S31.
- Collins JF. Kidney disease in Maori and Pacific people in New Zealand. *Clin Nephrol* 2010; 74 (Suppl 1): S61-S65.
- Weil EJ, Nelson RG. Kidney disease among the indigenous peoples of Oceania. *Ethn Dis* 2006; 16: S2-S30.
- United States Renal Data System. *USRDS 2006 Annual Data Report. National Institute of Diabetes and Digestive and Kidney Diseases*. National Institutes of Health; 2013.
- Kasiske BL, Rith-Najarian S, Casper ML, Croft JB. American Indian heritage and risk factors for renal injury. *Kidney Int* 1998; 54: 1305-1310, doi: 10.1046/j.1523-1755.1998.00106.x.
- Nelson RG, Morgenstern H, Bennett PH. An epidemic of proteinuria in Pima Indians with type 2 diabetes mellitus. *Kidney Int* 1998; 54: 2081-2088, doi: 10.1046/j.1523-1755.1998.00191.x.
- Scavini M, Shah VO, Stidley CA, Tentori F, Paine SS, Harford AM, et al. Kidney disease among the Zuni Indians: the Zuni Kidney Project. *Kidney Int Suppl* 2005; S126-S131, doi: 10.1111/j.1523-1755.2005.09721.x.
- Jha V, Garcia-Garcia G, Iseki K, Li Z, Naicker S, Plattner B, et al. Chronic kidney disease: global dimension and perspectives. *Lancet* 2013; 382: 260-272, doi: 10.1016/S0140-6736(13)60687-X.
- Almaguer M, Herrera R, Orantes CM. Chronic kidney disease of unknown etiology in agricultural communities. *MEDICC Rev* 2014; 16: 9-15.
- Ulasi II, Ijoma CK, Onodugo OD, Arodiwe EB, Ifebunandu NA, Okoye JU. Towards prevention of chronic kidney disease in Nigeria; a community-based study in Southeast Nigeria. *Kidney Int Suppl* 2013; 3: 195-201, doi: 10.1038/kisup.2013.13.
- Otieno LS, McLigeyo SO, Luta M. Acute renal failure following the use of herbal remedies. *East Afr Med J* 1991; 68: 993-998.
- Obrador GT, Garcia-Garcia G, Villa AR, Rubilar X, Olvera N, Ferreira E, et al. Prevalence of chronic kidney disease in the Kidney Early Evaluation Program (KEEP) Mexico and comparison with KEEP US. *Kidney Int Suppl* 2010; S2-S8, doi: 10.1038/ki.2009.540.
- Gutierrez-Padilla JA, Mendoza-Garcia M, Plascencia-Perez S, Renoirte-Lopez K, Garcia-Garcia G, Lloyd A, et al. Screening for CKD and cardiovascular disease risk factors using mobile clinics in Jalisco, Mexico. *Am J Kidney Dis* 2010; 55: 474-484, doi: 10.1053/j.ajkd.2009.07.023.
- Garcia-Garcia G, Gutierrez-Padilla AJ, Chavez-Iniguez J, Perez-Gomez HR, Mendoza-Garcia M, Gonzalez-De la Pena Mdel, et al. Identifying undetected cases of chronic kidney disease in Mexico. Targeting high-risk populations. *Arch Med Res* 2013; 44: 623-627, doi: 10.1016/j.arcmed.2013.10.007.
- Amato D, Alvarez-Aguilar C, Castaneda-Limonos R, Rodriguez E, Avila-Diaz M, Arreola F, et al. Prevalence of chronic kidney disease in an urban Mexican population. *Kidney Int Suppl* 2005; S11-S17, doi: 10.1111/j.1523-1755.2005.09702.x.
- Hoy W, McDonald SP. Albuminuria: marker or target in indigenous populations. *Kidney Int Suppl* 2004; S25-S31, doi: 10.1111/j.1523-1755.2004.09207.x.
- McDonald SP, Maguire GP, Hoy WE. Renal function and cardiovascular risk markers in a remote Australian Aboriginal community. *Nephrol Dial Transplant* 2003; 18: 1555-1561, doi: 10.1093/ndt/gfg199.
- Hoy WE, Samuel T, Mott SA, Kincaid-Smith PS, Fogo AB, Dowling JP, et al. Renal biopsy findings among Indigenous Australians: a nationwide review. *Kidney Int* 2012; 82: 1321-1331, doi: 10.1038/ki.2012.307.
- Hoy WE, Hughson MD, Zimanyi M, Samuel T, Douglas-Denton R, Holden L, et al. Distribution of volumes of individual glomeruli in kidneys at autopsy: association with age, nephron number, birth weight and body mass index. *Clin Nephrol* 2010; 74 (Suppl 1): S105-S112.
- Lackland DT, Bendall HE, Osmond C, Egan BM, Barker DJ. Low birth weights contribute to high rates of early-onset chronic renal failure in the Southeastern United States. *Arch*

- Intern Med* 2000; 160: 1472-1476, doi: 10.1001/archinte.160.10.1472.
32. Bhargava SK, Sachdev HS, Fall CH, Osmond C, Lakshmy R, Barker DJ, et al. Relation of serial changes in childhood body-mass index to impaired glucose tolerance in young adulthood. *N Engl J Med* 2004; 350: 865-875, doi: 10.1056/NEJMoa035698.
  33. Jafar TH, Chaturvedi N, Hatcher J, Khan I, Rabbani A, Khan AQ, et al. Proteinuria in South Asian children: prevalence and determinants. *Pediatr Nephrol* 2005; 20: 1458-1465, doi: 10.1007/s00467-005-1923-8.
  34. Jafar TH, Islam M, Poulter N, Hatcher J, Schmid CH, Levey AS, et al. Children in South Asia have higher body mass-adjusted blood pressure levels than white children in the United States: a comparative study. *Circulation* 2005; 111: 1291-1297, doi: 10.1161/01.CIR.0000157699.87728.F1.
  35. Liyanage T, Ninomiya T, Jha V, Patrice HM, Okpechi I, Zhao M, et al. Worldwide access to treatment for end stage kidney disease: a systematic review. *Lancet* 2015 (in press).
  36. Barsoum RS. Chronic kidney disease in the developing world. *N Engl J Med* 2006; 354: 997-999, doi: 10.1056/NEJMp058318.
  37. Cusumano AM, Garcia-Garcia G, Gonzalez-Bedat MC, Marinovich S, Lugon J, Poblete-Badal H, et al. Latin American Dialysis and Transplant Registry: 2008 prevalence and incidence of end-stage renal disease and correlation with socioeconomic indexes. *Kidney Int Suppl* 2013; 3: 153-156, doi: 10.1038/kisup.2013.2.
  38. Jha V. Current status of end-stage renal disease care in India and Pakistan. *Kidney Int Supplements* 2013; 3: 157-160, doi: 10.1038/kisup.2013.3.
  39. Garcia GG, Harden PN, Chapman JR. World Kidney Day 2012: the global role of kidney transplantation. *Am J Kidney Dis* 2012; 59: 319-324, doi: 10.1053/j.ajkd.2012.01.004.
  40. Yeates KE, Cass A, Sequist TD, McDonald SP, Jardine MJ, Trpeski L, et al. Indigenous people in Australia, Canada, New Zealand and the United States are less likely to receive renal transplantation. *Kidney Int* 2009; 76: 659-664, doi: 10.1038/ki.2009.236.
  41. Jha V. Current status of chronic kidney disease care in southeast Asia. *Semin Nephrol* 2009; 29: 487-496, doi: 10.1016/j.semnephrol.2009.06.005.
  42. Garcia-Garcia G, Monteon-Ramos JF, Garcia-Bejarano H, Gomez-Navarro B, Reyes IH, Lomeli AM, et al. Renal replacement therapy among disadvantaged populations in Mexico: a report from the Jalisco Dialysis and Transplant Registry (REDTJAL). *Kidney Int Suppl* 2005; S58-S61, doi: 10.1111/j.1523-1755.2005.09710.x.
  43. Ramachandran R, Jha V. Kidney transplantation is associated with catastrophic out of pocket expenditure in India. *PLoS One* 2013; 8: e67812, doi: 10.1371/journal.pone.0067812.
  44. McDonald SP, Russ GR. Burden of end-stage renal disease among indigenous peoples in Australia and New Zealand. *Kidney Int Suppl* 2003; S123-S127, doi: 10.1046/j.1523-1755.63.s83.26.x.
  45. Chou SH, Tonelli M, Bradley JS, Gourishankar S, Hemmelgarn BR. Quality of care among Aboriginal hemodialysis patients. *Clin J Am Soc Nephrol* 2006; 1: 58-63, doi: 10.2215/CJN.00560705.
  46. Rodriguez RA, Sen S, Mehta K, Moody-Ayers S, Bacchetti P, O'Hare AM. Geography matters: relationships among urban residential segregation, dialysis facilities, and patient outcomes. *Ann Intern Med* 2007; 146: 493-501, doi: 10.7326/0003-4819-146-7-200704030-00005.
  47. Mehrotra R, Story K, Guest S, Fedunyszyn M. Neighborhood location, rurality, geography, and outcomes of peritoneal dialysis patients in the United States. *Perit Dial Int* 2012; 32: 322-331, doi: 10.3747/pdi.2011.00084.
  48. Garcia-Garcia G, Briseno-Renteria G, Luquin-Arellan VH, Gao Z, Gill J, Tonelli M. Survival among patients with kidney failure in Jalisco, Mexico. *J Am Soc Nephrol* 2007; 18: 1922-1927, doi: 10.1681/ASN.2006121388.
  49. Parameswaran S, Geda SB, Rathi M, Kohli HS, Gupta KL, Sakhuja V, et al. Referral pattern of patients with end-stage renal disease at a public sector hospital and its impact on outcome. *Natl Med J India* 2011; 24: 208-213.