

# Metabolic Engineering of Mannitol Production in *Lactococcus lactis*: Influence of Overexpression of Mannitol 1-Phosphate Dehydrogenase in Different Genetic Backgrounds

H. Wouter Wisselink,<sup>1,2</sup> Astrid E. Mars,<sup>1,2</sup> Pieter van der Meer,<sup>2</sup> Gerrit Eggink,<sup>1,2</sup> and Jeroen Hugenholtz<sup>1,3\*</sup>

Wageningen Centre for Food Sciences<sup>1</sup> and Wageningen University and Research Centre—Agrotechnology and Food Innovations,<sup>2</sup> 6700 AA Wageningen, and NIZO Food Research, 6710 BA Ede,<sup>3</sup> The Netherlands

Received 22 November 2003/Accepted 12 March 2004

**To obtain a mannitol-producing *Lactococcus lactis* strain, the mannitol 1-phosphate dehydrogenase gene (*mltD*) from *Lactobacillus plantarum* was overexpressed in a wild-type strain, a lactate dehydrogenase (LDH)-deficient strain, and a strain with reduced phosphofructokinase activity. High-performance liquid chromatography and <sup>13</sup>C nuclear magnetic resonance analysis revealed that small amounts (<1%) of mannitol were formed by growing cells of *mltD*-overexpressing LDH-deficient and phosphofructokinase-reduced strains, whereas resting cells of the LDH-deficient transformant converted 25% of glucose into mannitol. Moreover, the formed mannitol was not reutilized upon glucose depletion. Of the metabolic-engineering strategies investigated in this work, *mltD*-overexpressing LDH-deficient *L. lactis* seemed to be the most promising strain for mannitol production.**

Mannitol is a sugar alcohol that is produced by a wide variety of organisms. It is assumed to have several beneficial effects as a food additive. It can serve as an antioxidant (5, 6, 24, 25) and as a low-calorie sweetener, which can replace sucrose (7, 10). Efiuvwevwe and coworkers (11) showed that mannitol has an osmoprotectant and antioxidant effect on the dairy lactic acid bacterium *Lactococcus lactis* subjected to decreased water activity and that mannitol enhances survival during drying of starter cells. The viability of starter cultures of *L. lactis*, which are extensively used in the dairy industry, may thus be enhanced by mannitol production in these strains. In addition, the use of mannitol-producing *L. lactis* strains may result in fermented products with extra nutritional value.

Mannitol biosynthesis in homofermentative lactic acid bacteria, such as *L. lactis*, starts with the glycolysis intermediate fructose 6-phosphate (Fig. 1). Mannitol 1-phosphate dehydrogenase (MPDH) (EC 1.1.1.17) catalyzes the reduction of fructose 6-phosphate, and also the reverse reaction, the oxidation of mannitol 1-phosphate (4, 12). Mannitol 1-phosphate is dephosphorylated to mannitol by mannitol phosphatase activity. Although the gene encoding MPDH (*mltD*) has been reported for *L. lactis* IL-1403 (2), mannitol production by *L. lactis* and other homofermentative lactic acid bacteria is not very likely. Presumably, the *mltD* gene, which is located in a mannitol operon, is not transcribed due to catabolite repression, as described for *Bacillus stearothermophilus* (14, 15), and no mannitol production can take place during growth on certain sugar substrates. Therefore, overexpression of MPDH might be an important step for mannitol synthesis in *L. lactis*.

In contrast to some heterofermentative lactic acid bacteria, mannitol production by homofermentative lactic acid bacteria

is not very common. However, mannitol production by *L. lactis* was observed by Neves and coworkers (22). In resting high-density cell suspensions of a lactate dehydrogenase (LDH)-deficient *L. lactis* strain, high levels of intracellular mannitol were produced. Upon glucose depletion, mannitol was remethylated. Also an LDH-negative mutant of *Lactobacillus plantarum* produced small amounts of mannitol (12). In these cases, mannitol production was believed to be an alternative pathway to regenerate NAD<sup>+</sup> instead of lactate formation.

Metabolic engineering can be a helpful tool to achieve mannitol production in *L. lactis*. In this report, *mltD* from *L. plantarum* was cloned and overexpressed in *L. lactis*. The involvement of fructose 6-phosphate as a substrate of MPDH could imply that the accumulation of fructose 6-phosphate, such as was reported in *L. lactis* with reduced phosphofructokinase (PFK) activity (1), could coincide with mannitol production. Also, alternative NAD<sup>+</sup> regeneration via MPDH, as reported in an LDH-deficient *L. lactis* strain (22), could contribute to mannitol production. Therefore, *mltD* was overexpressed in different genetic backgrounds: the parental *L. lactis* strain, NZ9000; an LDH-deficient strain, NZ9010; and a strain with reduced PFK activity, HWA217. Also, a comparison was made of mannitol production by growing cells and by high-density resting cells.

## MATERIALS AND METHODS

***L. lactis* strains, plasmids, and media.** The *L. lactis* strains and plasmids used in this study are listed in Table 1. The *L. lactis* strains were grown at 30°C in M17 broth (Oxoid) supplemented with 0.5% glucose. For (semi)anaerobic cultivations, cells were grown in batch cultures in 50-ml tubes without aeration. When cells were grown aerobically, shaking flasks with baffles were used. When applicable, chloramphenicol and erythromycin were supplemented at 10 and 5 µg/ml, respectively. Growth was followed by measuring the optical density at 600 nm (OD<sub>600</sub>) with an Ultrospec 2000 spectrophotometer (Pharmacia Biotech). For inducing MPDH activity, 1 ng of nisin/ml was added to a growing culture at an OD<sub>600</sub> of 0.5.

\* Corresponding author. Mailing address: NIZO Food Research, Kernhemseweg 2, 6718 ZB Ede, The Netherlands. Phone: (31) 318 659511. Fax: (31) 317 650400. E-mail: Jeroen.Hugenholtz@nizo.nl.




FIG. 1. Proposed pathway for hexose metabolism of homofermentative lactic acid bacteria. (1) Phosphoenolpyruvate-dependent sugar phosphotransferase system (PTS). (2) Mannitol-specific PTS. (3) Phosphoglucose isomerase. (4) Mannitol 1-phosphate dehydrogenase. (5) Mannitol 1-phosphatase. (6) 6-Phosphofructokinase. (7) Fructose-diphosphatase. (8) Fructose 1,6-diphosphate aldolase. (9) Triosephosphate isomerase. (10) Glyceraldehyde 3-phosphate dehydrogenase and phosphoglycerate kinase. (11) Phosphoglyceromutase and enolase. (12) Pyruvate kinase. (13) Lactate dehydrogenase. (14) Pyruvate-formate lyase. (15) Acetaldehyde dehydrogenase and alcohol dehydrogenase. (16) Pyruvate dehydrogenase. (17) Acetate kinase. (18)  $\alpha$ -Acetolactate synthase. (19)  $\alpha$ -Acetolactate decarboxylase. (20) 2,3-Butanediol dehydrogenase.

**Construction of plasmid pNZmt1D.** The gene encoding mannitol 1-phosphate dehydrogenase (*mt1D*) from *L. plantarum* was cloned into the nisin-inducible expression vector pNZ8148 (Table 1). For this, *mt1D* was amplified by PCR from *L. plantarum* genomic DNA (accession no. NP 784055) (19) using the primers MPDHLP-1FW (5'-TCGTACCATGGTAGACGTACATTTTG-3') and MPDHLP-3RV (5'-GTCACTAGACTACTTTGCTGCAGCTAAG-3'), with introduced *Nco*I and *Xba*I digestion sites, respectively (underlined). *Nco*I-*Xba*I-digested *mt1D* was cloned into pNZ8148, resulting in pNZmt1D containing *mt1D* fused to the *nisA* promoter. The sequence of *mt1D* was verified by sequencing the cloned PCR product (Eurogentec, Seraing, Belgium). pNZmt1D was cloned into the *L. lactis* strains NZ9000, NZ9010, and HWA217 (Table 1). Plasmid pNZ9530, containing *nisR* and *nisK* genes, was cotransformed in *L. lactis* HWA217 to make nisin induction of *mt1D* possible. The *nisRK* genes, coding for the histidine protein kinase NisK and the response regulator NisR, are the only *nis* genes required for *nisA* promoter activation on the pNZmt1D plasmid (18).

**Analysis of fermentation products and glucose consumption.** During the growth experiments, samples were taken from the *L. lactis* cultures and centrifuged for 1 min at 10,000  $\times$  g, and the supernatants were stored at  $-20^{\circ}\text{C}$  until they were analyzed. In the supernatant, lactate, acetate, formate, glucose, mannitol, ethanol, 2,3-butanediol, and acetoin were analyzed by high-performance liquid chromatography (HPLC). Separation was performed with a 30-cm-long ION-300 ion exclusion column (Alltech, Breda, The Netherlands) at a flow rate of 0.4 ml/min and a temperature of  $90^{\circ}\text{C}$ . The eluent consisted of 3 mM sulfuric acid. The products were detected on a refractive-index detector (Waters 410).

The distribution of glucose to different fermentation products was calculated as the slope of the product concentration versus the consumed glucose.

**Preparation of cell extracts.** Cell extracts were prepared by disruption of cells by glass beads. A cell culture (50 ml) was centrifuged ( $4^{\circ}\text{C}$ ; 20 min at 2,000  $\times$  g) and washed with 50 mM morpholineethanesulfonic acid (MES) buffer (pH 7.0). The cells were resuspended in 2 ml of 50 mM MES buffer (pH 7.0). For cell disruption, 1 ml of cell suspension was added to 1.0 g of 0.1-mm-diameter zirconium-silica beads (BioSpec Products, Inc.) in a 2-ml Eppendorf cup, and the cells were disrupted by vigorous shaking at  $4^{\circ}\text{C}$  for 5 min. Cell debris was removed by centrifugation ( $4^{\circ}\text{C}$ ; 2 min at 10,000  $\times$  g), and the supernatant was used for all enzyme assays. The protein contents of the extracts were determined by the bicinchoninic acid protein assay (Pierce) with bovine serum albumin as the standard.

**Enzyme assays.** Cell cultures (50 ml) were harvested for enzyme assays at an  $\text{OD}_{600}$  of  $\pm 1.3$  or 2 h after induction with nisin, and cell extracts were prepared as described above. LDH activity was determined by the method of Hillier and Jago (16). PFK activity was assayed according to the method of Grobbs et al. (13), with the modification that 5 mM fructose 6-phosphate was used to initiate the reaction. The PFK activities of extracts with high MPDH activity could not be determined, as fructose 6-phosphate is a substrate for both PFK and MPDH. Mannitol 1-phosphate oxidation by MPDH was determined in a reaction mixture containing 25 mM Tris-HCl, pH 8.0, and 1.5 mM NAD. The reaction was started with the addition of 1 mM mannitol 1-phosphate. The reduction of fructose 6-phosphate was assayed in 25 mM sodium phosphate buffer, pH 6, with 0.15 mM NADH. Fructose 6-phosphate (1 mM) was used to initiate the reaction. LDH, PFK, and MPDH activities were determined from the rate of NADH oxidation or formation at  $30^{\circ}\text{C}$  by measuring the absorbance at 340 nm (Ultrospec 2000).

Mannitol 1-phosphatase activity was determined in a 1-ml reaction mixture containing 0.1 to 0.2 mg of protein enzyme extract/ml, 50 mM MES buffer (pH 7.0), 10 mM  $\text{MgCl}_2$ , and 3 mM mannitol 1-phosphate. The amounts of inorganic phosphate formed were determined after 0, 60, 120, and 180 min of incubation at  $30^{\circ}\text{C}$  by a modified protocol of the Sigma inorganic phosphate kit. At the above time points, 200- $\mu\text{l}$  samples were taken, and the reaction was stopped with the addition of 40  $\mu\text{l}$  of acid molybdate solution. Ten microliters of Fiske & SubbaRow reducer solution was added to 200  $\mu\text{l}$  of the clear centrifuged solution in a 96-well microplate, and the absorbance at 655 nm was measured with a microplate reader (3550-UV; Bio-Rad).

**NMR experiments.** Cells grown on M17 broth were harvested at an  $\text{OD}_{600}$  of  $\sim 1.5$ , centrifuged, washed, and resuspended in 50 mM potassium phosphate buffer (pH 6.5) to an  $\text{OD}_{600}$  of  $\pm 35$ .  $^{13}\text{C}$  nuclear magnetic resonance (NMR) spectra were taken using a Bruker AMX-400Wb spectrometer at 100.62 MHz. All experiments were carried out at  $30^{\circ}\text{C}$  in a 10-mm NMR tube. Cell suspension (4 ml) was placed in the NMR tube, and an initial spectrum was acquired. At time zero, 20 mM [ $1\text{-}^{13}\text{C}$ ]glucose (Campro Scientific, Veenendaal, The Netherlands) was supplied in the NMR tube.  $^{13}\text{C}$ -NMR spectra were acquired for 104 s (64 scans). Chemical shifts were referred to the  $\beta\text{-C1}$  of D-glucose (96.6 ppm). Resonances in the spectra (Fig. 2) were identified by spiking with the pure (unlabeled) materials.

**Quantification of products by  $^{13}\text{C}$ -NMR.** Glucose, lactate, acetoin, 2,3-butanediol, ethanol, and mannitol were quantified during the consumption of [ $1\text{-}^{13}\text{C}$ ]glucose. Due to the fast pulsing conditions and short repetition times, the in vivo NMR spectra were not fully relaxed, and therefore there was not a direct correlation between peak intensities and concentrations. To correct for satura-

TABLE 1. Bacterial strains and plasmids used in this study

Strains and plasmids	Characteristics	Reference(s) or source
<b>Strains</b>		
<i>L. lactis</i> NZ9000	MG1363 <i>pepN::nisRK</i>	9, 20a
<i>L. lactis</i> NZ9010	NZ9000 <i>ldh::ery</i> Ery <sup>r</sup>	3, 17
<i>L. lactis</i> HWA217	Reduced PFK activity	1
<b>Plasmids</b>		
pNZ8148	pNZ8048 derivative; Cm <sup>r</sup> ; lactococcal cloning and expression vector with <i>nisA</i> promoter upstream of a multiple cloning site	9, 20a
pNZmt1D	pNZ8148 carrying <i>L. plantarum mt1D</i> gene	This work
pNZ9530	Ery <sup>r</sup> <i>nisRK</i>	18




FIG. 2.  $^{13}\text{C}$ -NMR spectra during metabolism of  $[1\text{-}^{13}\text{C}]$ glucose by high-density suspensions of nongrowing cells of the *L. lactis* strains NZ9000/pNZmtlD (A), HWA217/pNZmtlD (B), NZ9010 (C), and NZ9010/pNZmtlD (D). The spectra were recorded until no changes could be detected. The resonances of  $\alpha$  and  $\beta$  C1 of D-glucose were detected at 92.8 and 96.6 ppm.  $\blacklozenge$ ,  $^{13}\text{C}$ -labeled lactate;  $\bullet$ , acetoin;  $\times$ , 2,3-butanediol;  $\blacktriangle$ , ethanol;  $\blacksquare$ , mannitol.

tion, the saturation recovery method was used by performing relaxation measurements.

## RESULTS

In this work, we used different metabolic-engineering approaches to induce mannitol production in *L. lactis*. Basically, we used the glycolytic model of *L. lactis* (17; <http://jjj.biochem.sun.ac.za>) to predict the most efficient metabolic-engineering strategy. The model predicted that knocking out the LDH gene and decreasing the PFK activity could play a significant role in mannitol production in *L. lactis* by increasing the intracellular NADH and fructose 6-phosphate levels. Mannitol production in three *L. lactis* strains was determined: the wild-type strain, NZ9000, the LDH-deficient strain NZ9010 (17), and the PFK-reduced strain HWA217 (1). Enzyme activities in mid-log-phase cultures were measured, fermentation products in supernatants of batch grown cultures were analyzed, and  $^{13}\text{C}$ -NMR spectra were recorded during  $[1\text{-}^{13}\text{C}]$ glucose consumption by cell suspensions.

**MPDH enzyme activity in *L. lactis*.** Both *L. lactis* strains NZ9000 and HWA217 that were grown on glucose did not

show any MPDH activity, while cell extracts of *L. lactis* NZ9010 contained very low MPDH activity (Table 2). Since MPDH activity is essential for mannitol production, the corresponding gene of *L. plantarum* (*mtlD*) was overexpressed in all three *L. lactis* strains using pNZmtlD (Table 1). The transformants were grown on glucose and induced with nisin, and the cells were harvested after 2 h of induction. In all three hosts, the introduction of pNZmtlD led to a large increase in MPDH activity (Table 2). The low MPDH activity exhibited by the noninduced culture can be explained by the very low residual activity of the nisin promoter in the pNZ8148 expression vector under noninduced conditions (8, 9).

**Mannitol production by *mtlD* constructs.** The effect of overexpression of *mtlD* on the formation of fermentation products was determined by cultivating the strains in batch cultures on M17 medium under oxic or anoxic conditions. To induce MPDH activity, 1 ng of nisin/ml was added to growing cultures of the strains containing the pNZmtlD plasmid, and the fermentation products were determined over time in the culture supernatant (Table 3). Furthermore, in vivo  $^{13}\text{C}$ -NMR measurements were performed with suspensions of glucose-grown

TABLE 2. Enzyme activities determined in crude cell extracts of the various *L. lactis* strains grown on glucose

<i>L. lactis</i> strain	Nisin (ng/ml)	Activity <sup>a</sup>				
		LDH	PFK	MPDH		MP
				(F6P → M1P)	(M1P → F6P)	
NZ9000	0	13.5 ± 3.5	1.6 ± 0.6	ND	ND	4.2 ± 1.5
NZ9000/pNZmt1D	0	15.8 ± 2.7	1.8 ± 0.6	ND	0.03 ± 0.01	5.2 ± 1.0
NZ9000/pNZmt1D	1	16.1 ± 2.8	– <sup>b</sup>	6.2 ± 0.4	20.9 ± 0.3	3.5 ± 0.3
NZ9010	0	ND	1.5 ± 0.5	0.02 ± 0.02	≤0.01	4.9 ± 0.4
NZ9010/pNZmt1D	0	ND	1.2 ± 0.6	0.03 ± 0.02	0.03 ± 0.02	5.6 ± 0.3
NZ9010/pNZmt1D	1	ND	–	9.8 ± 1.6	30.3 ± 8.3	3.6 ± 0.2
HWA217/pNZ9530	0	17.0 ± 5.2	0.6 ± 0.4	ND	ND	8.6 ± 1.9
HWA217/pNZ9530/pNZmt1D	0	19.2 ± 1.2	0.3 ± 0.2	0.04 ± 0.01	0.12 ± 0.09	7.5 ± 0.8
HWA217/pNZ9530/pNZmt1D	1	17.7 ± 6.0	–	2.5 ± 0.9	8.4 ± 2.3	6.7 ± 2.4

<sup>a</sup> In micromoles per minute per milligram of protein (nanomoles per minute per milligram of protein for mannitol 1-phosphatase [MP]). ND, not detected. Cell extracts were made from cultures during exponential growth at an OD<sub>600</sub> of 1.3 to 1.6 or after 2 h of induction with nisin (*mtlD* overexpression strains).

<sup>b</sup> –, not determined.

cells (Fig. 2 and Table 4). The consumption of [1-<sup>13</sup>C]glucose by the suspensions was monitored until no changes in the spectra were observed. Resonances in the spectra of these measurements (Fig. 2) were identified as lactate (20.7 ppm), acetoin (18.8 ppm), ethanol (17.5 ppm), 2,3-butanediol (17.4 ppm), and mannitol (63.9 ppm) by spiking with the pure (unlabeled) materials.

**Overexpression of *mtlD* in parental *L. lactis*.** Wild-type *L. lactis* NZ9000 showed a typical homolactic fermentation pattern, with lactate as the main fermentation product (Table 3). Overexpression of *mtlD* in NZ9000 resulted in the same homolactic pattern (Table 3), and no mannitol was detected during growth. However, <sup>13</sup>C-NMR analysis of [1-<sup>13</sup>C]glucose-consuming cell suspensions of nisin-induced *L. lactis* NZ9000/pNZmt1D revealed the accumulation of a small amount of mannitol (Fig. 2A). The conversion rate of glucose into mannitol was 0.02 mol of mannitol per mol of glucose (Table 4).

**MtlD overexpression in an *L. lactis* strain with reduced PFK activity.** Growth of *L. lactis* with reduced PFK activity (strain HWA217) resulted in a mainly homolactic fermentation pattern with small amounts of acetate and formate as side products, as described by Andersen et al. (1). Introduction of pNZmt1D in HWA217 resulted in mannitol production during growth of 0.005 and 0.008 mol of mannitol per mol of consumed glucose (Table 3). The mannitol production was not dramatically improved when the MPDH activity was increased

by nisin induction (Table 2). Analysis of glucose consumption by nongrowing cell suspensions confirmed the homolactic pattern of HWA217 (spectra not shown). <sup>13</sup>C-NMR analysis of suspensions of *L. lactis* HWA217/pNZ9530/pNZmt1D showed that mannitol and ethanol accumulated (spectra not shown). In this strain, 1% of the [1-<sup>13</sup>C]glucose is converted into [1-<sup>13</sup>C]mannitol by cell suspensions (Table 4).

**Overexpression of *mtlD* in LDH-deficient *L. lactis*.** HPLC analysis of the LDH-deficient strain *L. lactis* NZ9010 showed a mixed-type fermentation pattern when the bacteria were grown on glucose under aerobic and (semi)anaerobic conditions (Table 3), similar to the fermentation patterns of the LDH-deficient strains described by Bongers et al. (3). No mannitol could be detected during growth. The introduction of pNZmt1D in strain NZ9010 resulted in conversion rates of 0.6 and 1.1%, for the uninduced and the induced growing cultures, respectively (Table 3). Production of mannitol was not observed during growth under aerobic conditions (Table 3).

<sup>13</sup>C-NMR analysis of the glucose metabolism of *L. lactis* NZ9010 suspensions did not show large amounts of mannitol (Fig. 2C); only 2% of the <sup>13</sup>C-labeled glucose was converted into mannitol (Table 4). Furthermore, glucose consumption by the cell suspension of *L. lactis* NZ9010 was much slower than that of the other strains, since only 44% of the glucose was metabolized during 45 min, whereas the others strains converted 65 to 88% of the glucose in the same time or less (Table

TABLE 3. Amounts of fermentation products formed per mole of consumed glucose during growth on 0.5% glucose

<i>L. lactis</i> strain	ox	Nisin (ng/ml)	Amt of product (mol mol of glucose <sup>-1</sup> )								
			Lactate	Formate	Acetate	Ethanol	Acetoin	2,3-Butanediol	Pyruvate	Mannitol	C recovery
NZ9000			1.71								0.86
NZ9000/pNZmt1D			1.74								0.87
NZ9000/pNZmt1D		1	1.74								0.87
NZ9010			0.13	1.1	0.15	1.1	0.28	0.07			1.04
NZ9010	+ <sup>a</sup>		0.02		0.09	0.12	0.57	0.03	0.26		0.85
NZ9010/pNZmt1D			0.06	1.0	0.37	0.77	0.17	0.16		0.006	0.94
NZ9010/pNZmt1D		1	0.08	0.93	0.15	0.87	0.24	0.15		0.011	0.95
NZ9010/pNZmt1D	+		0.02		0.22		0.58		0.16		0.78
NZ9010/pNZmt1D	+	1	0.02		0.21		0.60		0.15		0.79
HWA217/pNZ9530			1.86	0.12	0.01						0.94
HWA217/pNZ9530/pNZmt1D			1.83		0.01					0.005	0.93
HWA217/pNZ9530/pNZmt1D		1	1.87		0.01					0.008	0.95

<sup>a</sup> ox +, oxic growth.



TABLE 4. Formation of  $^{13}\text{C}$ -labeled fermentation products during  $[1-^{13}\text{C}]$ glucose consumption at  $30^\circ\text{C}$  by high-density suspensions of nongrowing *L. lactis* cells

<i>L. lactis</i> strain	$^{13}\text{C}$ glucose consumed (%)	Amt of product (mol mol of $[1-^{13}\text{C}]$ glucose $^{-1}$ ) <sup>a</sup>						
		Lactate	Acetate	Ethanol	Acetoin	2,3-Butanediol	Mannitol	$^{13}\text{C}$ recovery
NZ9000	70	0.68						0.68
NZ9000/pNZmt1D	85	0.62					0.02	0.64
NZ9010	44	0.17		0.22		0.24	0.02	0.65
NZ9010/pNZmt1D	100	0.07 (0.3)	(0.14)	0.03 (0.11)	0.14 (0.22)	0.07 (0.15)	0.15 (0.25)	0.46 (0.90)
HWA217/pNZ9530	65	0.49						0.49
HWA217/pNZ9530/pNZmt1D	88	0.53		0.02			0.01	0.56

<sup>a</sup> Spectra were recorded for 46.2 min, except for the spectrum of NZ9010/pNZmt1D, which was recorded for 420 min. The values in parentheses are concentrations measured in the supernatants by HPLC.

4). Much larger amounts of mannitol were produced by the nisin-induced cell suspensions of *L. lactis* NZ9010/pNZmt1D (Fig. 2D). A conversion rate as high as 15% (0.15 mol of  $^{13}\text{C}$ -labeled-mannitol per mol of  $[1-^{13}\text{C}]$ glucose) was obtained (Table 4). During glucose consumption, a rapid increase in mannitol was observed, and upon glucose exhaustion, the mannitol produced was not reutilized (Fig. 3).

The large difference in mannitol production between growing and nongrowing cells might be explained by intracellular accumulation of mannitol in the resting cells. To determine this, supernatant samples from cell suspensions converting  $[1-^{13}\text{C}]$ glucose were analyzed by HPLC. The measured concentrations of all fermentation products in the supernatants were higher than the concentrations that were measured with  $^{13}\text{C}$ -NMR. The extracellular mannitol production was corrected to 25% (Fig. 4 and Table 4). This suggested the loss of  $^{13}\text{C}$  label during the experiment, for example, via  $\text{CO}_2$  formation. HPLC analysis of cell extracts of glucose-consuming cell suspensions confirmed that mannitol was excreted and not accumulated in the cells (data not shown).

To investigate the mannitol-utilizing abilities of *L. lactis* NZ9010/pNZmt1D, glucose-precultured NZ9000 and NZ9010/

pNZmt1D cells were grown anaerobically on M17 broth supplemented with mannitol. Both NZ9000 and NZ9010/pNZmt1D strains showed a lag phase of  $\sim 24$  h before growth on mannitol was observed (Fig. 5).

## DISCUSSION

We investigated the effect of overexpression of the *L. plantarum* *mtlD* gene in three *L. lactis* hosts: the parental strain NZ9000, the LDH-deficient strain NZ9010, and the PFK-reduced strain HWA217. The mannitol-producing capacities of the constructed transformants were determined by HPLC analysis of supernatants of growing cultures, in vivo  $^{13}\text{C}$ -NMR analysis of glucose-consuming cell suspensions, and enzyme activity measurements. By combining these data, the effect of overexpression of *mtlD* in different genetic backgrounds on the mannitol-producing capacity of *L. lactis* could be determined.

When *mtlD* was overexpressed in wild-type *L. lactis* NZ9000, no mannitol was detected in supernatants of growing cells (Table 3). Although both MPDH and mannitol 1-phosphatase activities were present in cell extracts (Table 2), no mannitol production could be observed. Apparently, the glycolytic flux




FIG. 3. Formation of  $^{13}\text{C}$ -labeled fermentation products during  $[1-^{13}\text{C}]$ glucose consumption by high-density suspensions of nongrowing cells of *L. lactis* NZ9010/pNZmt1D. The starting concentration of  $[1-^{13}\text{C}]$ glucose is 20 mM.  $\circ$ , glucose;  $\blacklozenge$ , lactate;  $\bullet$ , acetoin;  $\times$ , 2,3-butanediol;  $\blacktriangle$ , ethanol;  $\blacksquare$ , mannitol.




FIG. 4. Mannitol production and glucose consumption by high-density suspensions of nongrowing cells of *L. lactis* NZ9010/pNZmt1D.  $\blacksquare$ , mannitol, and  $\bullet$ , glucose measured by HPLC in the supernatant;  $\square$ ,  $^{13}\text{C}$ -labeled mannitol, and  $\circ$ , glucose measured by  $^{13}\text{C}$ -NMR analysis.




FIG. 5. Growth, mannitol consumption, and product formation by *L. lactis* NZ9000 (A) and NZ9010/pNZ-*mtlD* (B) cultured under anaerobic conditions. M17 broth supplemented with 75 mM mannitol was inoculated with 2% (vol/vol) glucose-grown overnight culture. ■, mannitol; ◆, lactate; ▲, ethanol; ○, formate; □, acetate; ●, OD<sub>600</sub>.

in the growing cells is too high, and subsequently, the fructose 6-phosphate concentration is too low to enable a flux to mannitol. Also, NAD is regenerated by LDH, so the organism has no need to regenerate NAD by an alternative pathway, such as mannitol synthesis via MPDH. Under nongrowing conditions, *mtlD* overexpression resulted in a small amount of mannitol (Fig. 2A). About 2% of the glucose was converted to mannitol by this transformant (Table 4). Due to the low glycolytic flux in the nongrowing cells (20), the fructose 6-phosphate concentration might have been just high enough for MPDH to convert fructose 6-phosphate into mannitol 1-phosphate.

Increasing the intracellular fructose 6-phosphate concentration by lowering the PFK activity might be a strategy to enable mannitol production in *L. lactis*. Andersen and coworkers (1) showed that glycolytic fluxes were reduced in growing cells of an *L. lactis* mutant with 40% of the PFK activity of the MG1363 wild type. In this strain, sugar phosphates, such as glucose 6-phosphate and fructose 6-phosphate, accumulated. The increase in the fructose 6-phosphate concentration might contribute to a flux toward mannitol 1-phosphate via MPDH. Indeed, when *mtlD* is overexpressed in the PFK-reduced *L. lactis* strain HWA217, mannitol production is observed in the supernatants of growing cultures (Table 3). Since no mannitol was produced by the *mtlD*-overexpressing parental strain NZ9000, the results may imply that the reduction in PFK has resulted in the increased flux toward mannitol. However, nongrowing cells of MPDH-overexpressing HWA217 produced amounts of mannitol similar to those produced by the parental strain with high MPDH overexpression. Since PFK reduction increases the fructose 6-phosphate pool (1), it can be assumed that further increase of the fructose 6-phosphate pool does not increase the conversion of glucose into mannitol by nongrowing cells.

Another strategy to increase the flux to mannitol is to increase the NADH pool by knocking out the LDH activity, which is mainly responsible for the regeneration of NAD<sup>+</sup> in *L. lactis*. Neves et al. (22) showed that nongrowing cells of an LDH-deficient *L. lactis* strain transiently produced mannitol

intracellularly under anaerobic conditions to relieve the pressure to regenerate NAD<sup>+</sup>. In contrast to the LDH-deficient strain of Neves and coworkers (22), no high (intracellular) mannitol levels were produced by nongrowing cells of NZ9010 (Fig. 2C and Table 4). Moreover, both anaerobic and aerobic growing cultures of NZ9010 produced no mannitol at all. The formation of ethanol and 2,3-butanediol under anaerobic conditions suggests that the LDH-deficient strain has used these pathways to regenerate NAD<sup>+</sup>. Analysis of the fermentation products of the LDH-deficient strains showed that under anaerobic conditions lactate is formed. This can be attributed to transcriptional activation of the alternative LDH gene *ldhB* under anaerobic conditions (3).

When *mtlD* was overexpressed in NZ9010, mannitol production by growing cultures could be observed (Table 3). Both nisin-induced and uninduced cultures produced mannitol in small amounts: ~1% of the glucose was converted into mannitol. Apparently, the low MPDH activity in the noninduced cultures, due to residual activity of the nisin promoter in pNZ8148, is sufficient to accomplish a flux to mannitol. Nisin-induced overexpression of *mtlD* did not result in much higher mannitol production. Apparently, MPDH activity is not rate limiting in mannitol production. Noting that mannitol 1-phosphate activity is rather low in the different *L. lactis* strains (Table 2), this enzyme might have the highest control of the mannitol synthesis pathway.

Larger amounts of mannitol were detected by <sup>13</sup>C-NMR in the NZ9010/pNZ*mtlD* cell suspensions (Fig. 2D and Table 4). About 25% of the glucose was converted into mannitol, and conversion rates of glucose into acetoin, ethanol, and 2,3-butanediol were lower than in the NZ9010 strain. This implies that NAD<sup>+</sup> regeneration resorted to mannitol production via the introduced MPDH activity. In theory, up to 66.7% of glucose can be converted into mannitol when no NAD<sup>+</sup> is regenerated through lactate, ethanol, or 2,3-butanediol formation. These high conversion rates cannot be expected, considering the regained lactate production and the high ethanol pro-

duction by the LDH-deficient *L. lactis* strain. In addition, the fairly low mannitol 1-phosphatase activities in all three strains might be limiting for high glucose-mannitol conversions.

We showed that the mannitol concentration remains constant (Fig. 3), even after glucose exhaustion. This is in contrast with the observations of Neves et al. (22), which clearly demonstrated reutilization of the mannitol produced. Also, in the supernatants of growing cultures used in this work, mannitol was still present after 24 h, when glucose had already been depleted for at least 10 h (data not shown). For *L. lactis* IL-1403, a putative catabolite-responsive element (CRE) site in the promoter region of the mannitol operon was identified (accession no. AE006241 and AE006242), which suggests a possible involvement of CcpA in the regulation of the mannitol operon (21), and thus transcription of the genes involved in mannitol transport and metabolism will be derepressed when glucose is absent and mannitol is present. Moreover, *L. lactis* MG1363 and the LDH-deficient variant of Neves et al. are capable of growing on mannitol (23), so one would expect that mannitol is consumed when glucose is depleted. Apparently, the disruption of the LDH gene in the LDH-deficient strain used in this work did not induce the expression of genes coding for mannitol transport and utilization, unlike the mutant of Neves et al. (22, 23). In contrast to the findings of Neves and coworkers (23), the LDH-deficient strain did not grow better than the parental strain (Fig. 5). The 24-h lag phase of *L. lactis* NZ9010/pNZmtID growing on mannitol (Fig. 5) emphasizes that mannitol utilization genes have to be induced prior to growth on mannitol. Hence, immediate reutilization of mannitol by high-density resting cell suspensions of NZ9010/pNZmtID would not be expected.

The great difference in mannitol production between the growing and the nongrowing cells may be caused by accumulation of NADH in the resting cells. In the resting cells, where no ATP is needed for biomass production, the ATP demand is low (20), so ATP-generating steps, such as the conversion of phosphoenolpyruvate to pyruvate, are less important. Still, NADH is generated in glycolysis, and hence the nongrowing cells might give priority to regeneration of NAD<sup>+</sup> over ATP generation. Because of the LDH deficiency, MPDH takes a major part in NADH oxidation in the NZ9010/pNZmtID strain.

Our work presented here showed that *mtlD* overexpression, PFK reduction, and LDH deficiency have contributed to mannitol production in *L. lactis*. The most promising combination for mannitol production in *L. lactis* was *mtlD* overexpression in an LDH-deficient background. Despite the capability of growing on mannitol, no concomitant mannitol reutilization was observed in the LDH-deficient mutant, which is very desirable for the possibilities of mannitol in food products.

#### ACKNOWLEDGMENT

We thank Peter Ruhdal Jensen for providing the *L. lactis* HWA217 strain.

#### REFERENCES

- Andersen, H. W., C. Solem, K. Hammer, and P. R. Jensen. 2001. Twofold reduction of phosphofructokinase activity in *Lactococcus lactis* results in strong decreases in growth rate and in glycolytic flux. *J. Bacteriol.* **183**:3458–3467.
- Bolotin, A., P. Wincker, S. Mauger, O. Jaillon, K. Malarme, J. Weissenbach, S. D. Ehrlich, and A. Sorokin. 2001. The complete genome sequence of the lactic acid bacterium *Lactococcus lactis* ssp *lactis* IL1403. *Genome Res.* **11**:731–753.
- Bongers, R. S., M. H. N. Hoefnagel, M. J. C. Starrenburg, M. A. J. Siemerink, J. G. A. Arends, J. Hugenholtz, and M. Kleerebezem. 2003. IS981-mediated adaptive evolution recovers lactate production by *ldhB* transcription activation in a lactate dehydrogenase-deficient strain of *Lactococcus lactis*. *J. Bacteriol.* **185**:4499–4507.
- Chakravorty, M. 1964. Metabolism of mannitol and introduction of mannitol 1-phosphate dehydrogenase in *Lactobacillus plantarum*. *J. Bacteriol.* **87**:1246–1248.
- Chaturvedi, V., A. Bartiss, and B. Wong. 1997. Expression of bacterial *mtlD* in *Saccharomyces cerevisiae* results in mannitol synthesis and protects a glycerol-defective mutant from high-salt and oxidative stress. *J. Bacteriol.* **179**:157–162.
- Chaturvedi, V., B. Wong, and S. L. Newman. 1996. Oxidative killing of *Cryptococcus neoformans* by human neutrophils. Evidence that fungal mannitol protects by scavenging reactive oxygen intermediates. *J. Immunol.* **156**:3836–3840.
- Debord, B., C. Lefebvre, A. M. Guyot-Hermann, J. Hubert, R. Bouche, and J. C. Guyot. 1987. Study of different forms of mannitol: comparative behaviour under compression. *Drug Dev. Ind. Pharm.* **13**:1533–1546.
- de Ruyter, P. G. A., O. P. Kuipers, M. M. Beerthuyzen, I. van Alen-Boerrigter, and W. M. de Vos. 1996. Functional analysis of promoters in the nisin gene cluster of *Lactococcus lactis*. *J. Bacteriol.* **178**:3434–3439.
- de Ruyter, P. G. A., O. P. Kuipers, and W. M. de Vos. 1996. Controlled gene expression systems for *Lactococcus lactis* with the food-grade inducer nisin. *Appl. Environ. Microbiol.* **62**:3662–3667.
- Dwivedi, B. K. 1978. Low calorie and special dietary foods. CRC Press, Inc., West Palm Beach, Fla.
- Efiuvwevwe, B. J. O., L. G. M. Gorris, E. J. Smid, and E. P. W. Kets. 1999. Mannitol-enhanced survival of *Lactococcus lactis* subjected to drying. *Appl. Microbiol. Biotechnol.* **51**:100–104.
- Ferain, T., A. N. Schanck, and J. Delcour. 1996. <sup>13</sup>C nuclear magnetic resonance analysis of glucose and citrate end products in an *ldhL-ldhD* double-knockout strain of *Lactobacillus plantarum*. *J. Bacteriol.* **178**:7311–7315.
- Grobben, G. J., M. R. Smith, J. Sikkema, and J. A. M. De Bont. 1996. Influence of fructose and glucose on the production of exopolysaccharides and the activities of enzymes involved in the sugar metabolism and the synthesis of sugar nucleotides in *Lactobacillus delbrueckii* subsp. *bulgaricus* NCFB 2772. *Appl. Microbiol. Biotechnol.* **46**:279–284.
- Henstra, S. A., R. H. Duurkens, and G. T. Robillard. 2000. Multiple phosphorylation events regulate the activity of the mannitol transcriptional regulator MtlR of the *Bacillus stearothermophilus* phosphoenolpyruvate-dependent mannitol phosphotransferase system. *J. Biol. Chem.* **275**:7037–7044.
- Henstra, S. A., M. Tuinhof, R. H. Duurkens, and G. T. Robillard. 1999. The *Bacillus stearothermophilus* mannitol regulator, MtlR, of the phosphotransferase system. *J. Biol. Chem.* **274**:4754–4763.
- Hillier, A. J., and G. R. Jago. 1982. L-Lactate dehydrogenase, FDP-activated, from *Streptococcus cremoris*. *Methods Enzymol.* **89**:362–367.
- Hoefnagel, M. H. N., M. J. C. Starrenburg, D. E. Martens, J. Hugenholtz, M. Kleerebezem, S. Van Il, R. Bongers, H. V. Westerhoff, and J. L. Snoep. 2002. Metabolic engineering of lactic acid bacteria, the combined approach: kinetic modelling, metabolic control and experimental analysis. *Microbiology* **148**:1003–1013.
- Kleerebezem, M., M. M. Beerthuyzen, E. E. Vaughan, W. M. de Vos, and O. P. Kuipers. 1997. Controlled gene expression systems for lactic acid bacteria: transferable nisin-inducible expression cassettes for *Lactococcus*, *Leuconostoc*, and *Lactobacillus* spp. *Appl. Environ. Microbiol.* **63**:4581–4584.
- Kleerebezem, M., J. Boekhorst, R. van Kranenburg, D. Molenaar, O. P. Kuipers, R. Leer, R. Turchini, S. A. Peters, H. M. Sandbrink, M. W. E. J. Fiers, W. Stiekema, R. M. K. Lankhorst, P. A. Bron, S. M. Hoffer, M. N. N. Groot, R. Kerkhoven, M. de Vries, B. Ursing, W. M. de Vos, and R. J. Siezen. 2003. Complete genome sequence of *Lactobacillus plantarum* WCFS1. *Proc. Natl. Acad. Sci. USA* **100**:1990–1995.
- Koebmann, B. J., H. W. Andersen, C. Solem, and P. R. Jensen. 2002. Experimental determination of control of glycolysis in *Lactococcus lactis*. *Antonie van Leeuwenhoek* **82**:237–248.
- 20a. Kuipers, O. P., P. G. A. de Ruyter, M. Kleerebezem, and W. M. de Vos. 1998. Quorum sensing-controlled gene expression in lactic acid bacteria. *J. Biotechnol.* **64**:15–21.
- Luesink, E. J., R. E. M. A. Van Herpen, B. P. Grossiord, O. P. Kuipers, and W. M. de Vos. 1998. Transcriptional activation of the glycolytic *las* operon and catabolite repression of the *gal* operon in *Lactococcus lactis* are mediated by the catabolite control protein CcpA. *Mol. Microbiol.* **30**:789–798.
- Neves, A. R., A. Ramos, C. Shearman, M. J. Gasson, J. S. Almeida, and H. Santos. 2000. Metabolic characterization of *Lactococcus lactis* deficient in lactate dehydrogenase using in vivo C-13-NMR. *Eur. J. Biochem.* **267**:3859–3868.
- Neves, A. R., A. Ramos, C. Shearman, M. J. Gasson, and H. Santos. 2002. Catabolism of mannitol in *Lactococcus lactis* MG1363 and a mutant defective in lactate dehydrogenase. *Microbiology* **148**:3467–3476.
- Shen, B., R. G. Jensen, and H. J. Bohner. 1997. Mannitol protects against oxidation by hydroxyl radicals. *Plant Physiol.* **115**:527–532.
- Smirnov, N., and Q. J. Cumbes. 1989. Hydroxyl radical scavenging activity of compatible solutes. *Phytochemistry* **28**:1057–1060.