



Published in final edited form as:

AIDS. 2015 June 19; 29(10): 1263–1265. doi:10.1097/QAD.0000000000000735.

## Soluble CD14 is a nonspecific marker of monocyte activation

Carey L. Shive<sup>a</sup>, Wei Jiang<sup>b</sup>, Donald D. Anthony<sup>a,c</sup>, and Michael M. Lederman<sup>a</sup>

<sup>a</sup>Center for AIDS Research, Case Western Reserve University School of Medicine, Cleveland, Ohio

<sup>b</sup>Department of Microbiology and Immunology, Division of Infectious Diseases, Medical University of South Carolina, Charleston, South Carolina

<sup>c</sup>Veterans Administration Medical Center, Cleveland, Ohio, USA

### Abstract

Soluble CD14 is associated with morbidity and mortality in HIV disease. It is a co-receptor for lipopolysaccharide (LPS) that is released from monocytes upon activation. We demonstrate here, that inflammatory cytokines can induce the release of sCD14 in peripheral blood mononuclear cell cultures from healthy donors, and that TLR ligands other than LPS can cause a decrease in the monocyte cell surface expression of CD14. Thus, sCD14 is a marker of monocyte activation, not restricted to activation by LPS.

Elevated plasma levels of soluble CD14 (sCD14) are associated with poor prognosis in HIV-infected patients and are a strong predictor of morbidity and mortality [1–4]. Increased plasma sCD14 levels can persist even in treated HIV-infected patients, with durable control of viremia [5, 6], and this is associated with diminished CD4<sup>+</sup> T-cell restoration [5]. Additionally, plasma levels of sCD14 have been associated with cardiovascular disease in HIV infection [7], as well as in HIV-uninfected patients [8]. Elevated plasma levels of sCD14 are also observed in pediatric inflammatory lung diseases [9], in chronic hepatitis C virus (HCV) and hepatitis B virus (HBV) infections [10,11], and in rheumatoid arthritis [12].

CD14 is a myeloid differentiation marker found primarily on monocytes and macrophages, although low levels are also found on neutrophils. CD14 can exist as a GPI-anchored membrane protein or as one of two soluble isoforms that can be generated either by cleavage from the surface of the cell or released from intracellular pools [12–16]. CD14 is a co-receptor for lipopolysaccharide (LPS) – a cell wall component of gram-negative bacteria [17,18] – and it has been suggested that high levels of sCD14 in the plasma are reflective of LPS exposure [10,19]. Yet, CD14 does not bind only LPS; it can also bind Gram-positive cell wall components, LAM and HSP60 [20,21], as well as endogenous lipids [22,23].

Copyright © 2015 Wolters Kluwer Health, Inc. All rights reserved.

Correspondence to Michael M. Lederman, MD, Center for AIDS Research, Case Western Reserve University, 2061 Cornell Road, Cleveland, OH 44106, USA. lederman.michael@clevelandactu.org.

**Conflicts of interest:** The authors have no conflicting interests to declare.

Upon cell activation, CD14 surface expression decreases on monocytes and sCD14 is released. Although release of sCD14 is often accompanied by a decrease in surface expression of CD14, this is not always the case. This suggests that sCD14 can be generated by a mechanism other than cleavage from the cell surface [14]. Although LPS is a very potent activator of monocytes and induces protease-dependent generation of sCD14, we demonstrate here that a variety of TLR ligands can cause a down-modulation of CD14 on monocytes. We stimulated peripheral blood mononuclear cells (PBMCs) from healthy donors overnight with medium alone or medium supplemented with LPS (20 ng/ml; Sigma, St Louis, Missouri, USA), flagellin (1 µg/ml; InvivoGen, San Diego, California, USA), CpG oligodeoxynucleotides (CpG 2395 3 µg/ml; InvivoGen), or a combination of all three TLR ligands. The following day, the cells were washed and stained with anti-CD14-PE antibodies (BD, San Jose, California, USA). Events were acquired on an LSRII flow cytometer and analyzed for expression of CD14. The fluorescence of CD14 expression after stimulation is shown in (Fig. 1a). Stimulation with each individual TLR ligand caused a decrease in CD14 surface expression, and the greatest decrease was seen when all three ligands were combined.

Recognizing that the activation of monocytes by LPS induces inflammatory cytokines, we stimulated PBMCs from healthy donors with LPS (100 ng/ml; Sigma), recombinant human interleukin (IL)-6 (100 ng/ml; R&D Systems, Minneapolis, Minnesota, USA), or recombinant human IL-1β (10 ng/ml; R&D Systems). After 48 h of stimulation, we collected the culture supernatant and measured the concentration of sCD14 by ELISA (R&D Systems). We show that PBMCs stimulated with either IL-6 or IL-1β can induce sCD14 at levels comparable to, or only slightly less than concentrations induced by LPS (Fig. 1b). Similar findings were reported by Bas *et al.* [12] after stimulation of the human hepatoma cell line, HepG2. Though the levels they measured were lower than what we demonstrated, they saw the release of sCD14 from HepG2 cells after stimulation with IL-6, but saw slight decreases in sCD14 levels in HepG2 hepatoma cultures stimulated with IL-1β [12]. Marcos *et al.* [9] also saw increased culture supernatant levels of sCD14 after 40h stimulation of PBMCs with various TLR ligands, including LPS and CpG oligodeoxynucleotides. Bas *et al.* examined plasma levels of sCD14, IL-6, and C-reactive protein from arthritis patients, and found elevated plasma levels of sCD14 not only in patients with infection-mediated arthritis but also in patients with both crystal-mediated and rheumatoid arthritis [12]. This suggests that, *in vivo*, a number of inflammatory stimuli may cause elevated plasma levels of sCD14 and the presences of LPS is not required.

In conclusion, although LPS is a potent monocyte activator that binds to CD14 and induces cellular release of sCD14, other TLR ligands such as flagellin and CpG oligodeoxynucleotides can also induce release of sCD14 as can inflammatory cytokines such as IL-6 and IL-1β. Therefore, sCD14 should be considered a marker of monocyte activation that does not necessarily reflect monocyte activation by LPS.

## Acknowledgments


Funding sources: This work was supported by the National Institutes of Health grant AI091526 awarded to Wei Jiang; the National Institutes of Health (grants AI105937, AI 076174 and AI 68636), the CWRU Center for AIDS

Research (grant AI 36219), and the Fasenmyer Foundation awarded to Michael M. Lederman; the Veteran Affairs Merit 1 grant 01CX001104-01 awarded to Donald D. Anthony.

## References

1. Sandler NG, Wand H, Roque A, et al. Plasma levels of soluble CD14 independently predict mortality in HIV infection. *J Infect Dis.* 2011; 203:780–790. [PubMed: 21252259]
2. Tenorio AR, Zheng Y, Bosch RJ, et al. Soluble markers of inflammation and coagulation but not T-cell activation predict non-AIDS-defining morbid events during suppressive antiretroviral treatment. *J Infect Dis.* 2014
3. Hunt PW, Sinclair E, Rodriguez B, et al. Gut Epithelial Barrier Dysfunction and Innate Immune Activation Predict Mortality in Treated HIV Infection. *J Infect Dis.* 2014
4. Lien E, Aukrust P, Sundan A, Muller F, Froland SS, Espevik T. Elevated levels of serum-soluble CD14 in human immunodeficiency virus type 1 (HIV-1) infection: correlation to disease progression and clinical events. *Blood.* 1998; 92:2084–2092. [PubMed: 9731066]
5. Lederman MM, Calabrese L, Funderburg NT, et al. Immunologic failure despite suppressive antiretroviral therapy is related to activation and turnover of memory CD4 cells. *J Infect Dis.* 2011; 204:1217–1226. [PubMed: 21917895]
6. Mendez-Lagares G, Romero-Sanchez MC, Ruiz-Mateos E, et al. Long-term suppressive combined antiretroviral treatment does not normalize the serum level of soluble CD14. *J Infect Dis.* 2013; 207:1221–1225. [PubMed: 23322858]
7. Kelesidis T, Kendall MA, Yang OO, Hodis HN, Currier JS. Biomarkers of microbial translocation and macrophage activation: association with progression of subclinical atherosclerosis in HIV-1 infection. *J Infect Dis.* 2012; 206:1558–1567. [PubMed: 23066162]
8. Anker SD, Egerer KR, Volk HD, Kox WJ, Poole-Wilson PA, Coats AJ. Elevated soluble CD14 receptors and altered cytokines in chronic heart failure. *Am J Cardiol.* 1997; 79:1426–1430. [PubMed: 9165177]
9. Marcos V, Latzin P, Hector A, et al. Expression, regulation and clinical significance of soluble and membrane CD14 receptors in pediatric inflammatory lung diseases. *Respir Res.* 2010; 11:32. [PubMed: 20302606]
10. Sandler NG, Koh C, Roque A, et al. Host response to translocated microbial products predicts outcomes of patients with HBV or HCV infection. *Gastroenterology.* 2011; 141:1220–1230 e1-3. [PubMed: 21726511]
11. Schlatter DM, Sugalski JM, Chen Y, et al. Plasma proteome analysis reveals overlapping, yet distinct mechanisms of immune activation in chronic HCV and HIV infections. *J Acquir Immune Defic Syndr.* 2013; 63:563–571. [PubMed: 23507661]
12. Bas S, Gauthier BR, Spentato U, Stingelin S, Gabay C. CD14 is an acute-phase protein. *J Immunol.* 2004; 172:4470–4479. [PubMed: 15034063]
13. Bazil V, Strominger JL. Shedding as a mechanism of down-modulation of CD14 on stimulated human monocytes. *J Immunol.* 1991; 147:1567–1574. [PubMed: 1880416]
14. Bufler P, Stiegler G, Schuchmann M, et al. Soluble lipopolysaccharide receptor (CD14) is released via two different mechanisms from human monocytes and CD14 transfectants. *Eur J Immunol.* 1995; 25:604–610. [PubMed: 7533093]
15. Durieux JJ, Vita N, Popescu O, et al. The two soluble forms of the lipopolysaccharide receptor, CD14: characterization and release by normal human monocytes. *Eur J Immunol.* 1994; 24:2006–2012. [PubMed: 7522157]
16. Stelter F, Pfister M, Bernheiden M, et al. The myeloid differentiation antigen CD14 is N- and O-glycosylated. Contribution of N-linked glycosylation to different soluble CD14 isoforms. *Eur J Biochem.* 1996; 236:457–464. [PubMed: 8612616]
17. Kitchens RL, Thompson PA. Modulatory effects of sCD14 and LBP on LPS-host cell interactions. *J Endotoxin Res.* 2005; 11:225–229. [PubMed: 16176659]
18. Wright SD, Ramos RA, Tobias PS, Ulevitch RJ, Mathison JC. CD14, a receptor for complexes of lipopolysaccharide (LPS) and LPS binding protein. *Science.* 1990; 249:1431–1433. [PubMed: 1698311]

19. Brechley JM, Price DA, Schacker TW, et al. Microbial translocation is a cause of systemic immune activation in chronic HIV infection. *Nat Med.* 2006; 12:1365–1371. [PubMed: 17115046]
20. Kol A, Lichtman AH, Finberg RW, Libby P, Kurt-Jones EA. Cutting edge: heat shock protein (HSP) 60 activates the innate immune response: CD14 is an essential receptor for HSP60 activation of mononuclear cells. *J Immunol.* 2000; 164:13–17. [PubMed: 10604986]
21. Pugin J, Heumann ID, Tomasz A, et al. CD14 is a pattern recognition receptor. *Immunity.* 1994; 1:509–516. [PubMed: 7534618]
22. Wang PY, Kitchens RL, Munford RS. Phosphatidylinositides bind to plasma membrane CD14 and can prevent monocyte activation by bacterial lipopolysaccharide. *J Biol Chem.* 1998; 273:24309–24313. [PubMed: 9733716]
23. Yu B, Hailman E, Wright SD. Lipopolysaccharide binding protein and soluble CD14 catalyze exchange of phospholipids. *J Clin Invest.* 1997; 99:315–324. [PubMed: 9006000]



**Fig. 1. Soluble CD14 is generated by monocyte activation, not only by lipopolysaccharide stimulation**

(a) PBMCs from healthy donors were stimulated overnight with medium alone or medium supplemented with LPS (20 ng/ml), flagellin (1  $\mu$ g/ml), CpG-2395 (3  $\mu$ g/ml), or a combination of the three TLR ligands. Samples were stained and then acquired by flow cytometry, and expression of surface CD14 is shown as the geometric mean fluorescence intensity. (b) PBMCs from healthy donors ( $n = 3$ ) were stimulated for 48h with medium alone or medium supplemented with LPS (100 ng/ml), human IL-6 (100 ng/ml), or IL-1 $\beta$  (10ng/ml). After 48h, supernatant was measured for sCD14 levels by ELISA. IL, interleukin; LPS, lipopolysaccharide; PBMC, peripheral blood mononuclear cell.