



HHS Public Access

Author manuscript

Bull Menninger Clin. Author manuscript; available in PMC 2015 June 07.

Published in final edited form as:

Bull Menninger Clin. 2014 ; 78(3): 243–252. doi:10.1521/bumc.2014.78.3.243.

Perceptions about e-cigarette safety may lead to e-smoking during pregnancy

Selina Baeza-Loya,

Menninger Department of Psychiatry and Behavioral Sciences, Baylor College of Medicine, Houston, Texas

Rice University, Houston, Texas

Humsini Viswanath, MPH,

Menninger Department of Psychiatry and Behavioral Sciences, Baylor College of Medicine, Houston, Texas

Asasia Carter, BA,

Menninger Department of Psychiatry and Behavioral Sciences, Baylor College of Medicine, Houston, Texas

David L. Molfese, PhD,

Menninger Department of Psychiatry and Behavioral Sciences, Baylor College of Medicine, Houston, Texas

Kenia M. Velasquez, MS,

Menninger Department of Psychiatry and Behavioral Sciences, Baylor College of Medicine, Houston, Texas

Diana Helis Henry Medical Research Foundation, New Orleans, Louisiana

Philip R. Baldwin, PhD,

Menninger Department of Psychiatry and Behavioral Sciences, Baylor College of Medicine, Houston, Texas

Daisy G. Y. Thompson-Lake, MS,

Menninger Department of Psychiatry and Behavioral Sciences, Baylor College of Medicine, Houston, Texas

Carla Sharp, PhD,

Department of Psychology, University of Houston, Houston, Texas

J. Christopher Fowler, PhD,

Menninger Department of Psychiatry and Behavioral Sciences, Baylor College of Medicine, Houston, Texas

Richard De La Garza II, PhD, and

Copyright © 2014 The Menninger Foundation

Correspondence may be sent to Ramiro Salas at the Baylor College of Medicine, Menninger Department of Psychiatry and Behavioral Sciences, Houston TX 77030; rsalas@bcm.edu.

Selina Baeza-Loya and Humsini Viswanath contributed equally to this article.

Menninger Department of Psychiatry and Behavioral Sciences, Baylor College of Medicine, Houston, Texas

Ramiro Salas, PhD

Menninger Department of Psychiatry and Behavioral Sciences, Baylor College of Medicine, Houston, Texas

Abstract

Electronic cigarettes (e-cigarettes) are nicotine-delivery devices that are increasingly used, especially by young people. Because e-cigarettes lack many of the substances found in regular tobacco, they are often perceived as a safer smoking alternative, especially in high-risk situations such as pregnancy. However, studies suggest that it is exposure to nicotine that is most detrimental to prenatal development. The authors studied perceptions of tobacco and e-cigarette health risks using a multiple-choice survey. To study the perceived safety of e-cigarettes versus tobacco cigarettes, 184 modified Global Health Youth Surveys (WHO, <http://www.who.int/tobacco/surveillance/gyts/en/>) were completed electronically or on paper. Age range, smoking status, and perceptions about tobacco cigarettes and e-cigarettes were studied. The results verified that younger people use e-cigarettes more than older people. Tobacco cigarettes were perceived as more harmful than e-cigarettes to health in general, including lung cancer and pregnancy. Although more research is necessary, the authors postulate that the perception that e-cigarettes are safer during pregnancy may induce pregnant women to use these devices more freely. Given that nicotine is known to cause fetal harm, pregnant mothers who smoke e-cigarettes could cause even greater harm to the fetus because e-cigarettes are perceived as being safer than tobacco cigarettes. Until more data about the effects of nicotine during pregnancy are available, the authors advocate for labeling of e-cigarettes as potentially harmful, at least during pregnancy.

Electronic cigarettes (e-cigarettes) are devices that provide nicotine to users through vaporization of a nicotinic solution (Cahn & Siegel, 2011). In the past few years, e-cigarettes have become increasingly popular among adolescents, young adults, and tobacco cigarette smokers who want to quit or look for nicotine alternatives without the harmful side effects of tobacco (King, Alam, Promoff, Arrazola, & Dube, 2013). E-cigarettes have surpassed alternative tobacco products in online popularity (Ayers, Ribisl, & Brownstein, 2011), indicating an increase in both awareness and interest in these products.

Detailed research on the patterns of e-cigarette use is necessary and ongoing. The possible health effects of e-cigarette use are poorly understood, but educated guesses can be made using our knowledge of nicotine's effects on the body. Most studies regarding the safety of e-cigarettes compare the risks of using e-cigarettes relative to tobacco cigarettes. Young people in focus groups were unsure if alternative nicotine products, including e-cigarettes, were less harmful than cigarettes due to the lack of information about these products (Choi, Fabian, Mottey, Corbett, & Forster, 2012). Because e-cigarettes are thought to lack many of the harmful constituents associated with regular tobacco, including carcinogens known to cause diseases such as lung cancer (Hecht, 2012), people may reach the conclusion that e-cigarettes are a safer smoking alternative.

Given that individuals' perceptions affect their behavior (Ajzen, 2011), the perception that e-cigarettes are generally less harmful than tobacco cigarettes may lead to an increase in e-cigarette use during pregnancy. However, the safety of nicotine replacement devices, such as e-cigarettes or nicotine patches, is not supported by evidence from animal studies, which indicate that the dose of nicotine presents a great hazard to prenatal development (Wickstrom, 2007). Tobacco cigarette smoking is injurious during prenatal development because of direct delivery of nicotine to the fetus. Spontaneous abortions, neonatal intensive care unit admissions, perinatal and natal deaths from Sudden Infant Death Syndrome (SIDS), increased risk of learning disabilities, behavioral problems, attention-deficit/hyperactivity disorder (ADHD), and increased risk of addiction later in life may be attributed to the effects of nicotine during prenatal development (Ernst, Moolchan, & Robinson, 2001; Slotkin, 1998; Slotkin, Tate, Cousins, & Seidler, 2006). Given that these are nicotine's effects, it follows that e-cigarettes will probably cause similar effects to fetuses.

The primary aim of this study was to determine possible differences among the perceptions of the health risks of tobacco cigarettes and e-cigarettes. We were particularly interested in the perceived safety of the use of e-cigarettes during pregnancy.

Methods

Survey

A cross-sectional study was carried out using a version of the Global Youth Tobacco Survey (WHO, <http://www.who.int/tobacco/surveillance/gyts/en/>) modified to include questions about both tobacco cigarettes and e-cigarettes. The first section contained questions about participants' use of tobacco cigarettes and/or e-cigarettes. The next section included questions about the participants' perceptions of tobacco cigarettes and e-cigarettes. The following questions were of particular interest because of the potential health implications: "Do you think that tobacco cigarettes are harmful for pregnant women to use?" compared to "Do you think that e-cigarettes are harmful for pregnant women to use?" And "Do you think that tobacco cigarettes can give you lung cancer?" compared to "Do you think that e-cigarettes can give you lung cancer?" Each question could be answered "definitely yes," "probably yes," "probably not," or "definitely not."

Participants

Participants (184) were recruited during the year 2013 via electronic means (e.g., e-mail, Facebook) and word-of-mouth. Consent was included as part of the questionnaire. The survey was administered anonymously either electronically or on paper. No identifiable information was collected. Demographic information related to participant age range was acquired for the following age groups: 18–20, 21–25, 26–30, and 31+ years old (the 21–25 and 26–30 groups were pooled during analysis to equalize the number of respondents per group). This study was reviewed and approved by the Baylor College of Medicine Institutional Review Board.

Data analyses

No participant had missing data in any of the scores reported here. Scores of 0, 1, 2, and 3 were assigned to the answers “definitely yes,” “probably yes,” “probably not,” and “definitely not,” respectively. The Wilcoxon Sign Rank test was used to compare answer distributions.

Results

In total, 184 people responded to the questionnaire. Of those, 68 were young adults (18–20 years old), 55 were adults (21–30 years old), and 61 were older adults (31 or older). Across age groups, 109 respondents had smoked at least one tobacco cigarette and 40 had tried an e-cigarette.

As shown in Figure 1, we found that younger respondents tended to have tried e-cigarettes more and tobacco cigarettes less than older adults ($p < 10^{-12}$). This finding has been reported elsewhere (Choi & Forster, 2013; Pepper et al., 2013). When we examined health perceptions, we found that people of all ages perceive e-cigarettes as significantly less dangerous than tobacco cigarettes. This was true for questions about lung cancer (Figure 2A, $p < 10^{-33}$), harm during pregnancy (Figure 2B, $p < 10^{-32}$), and difficulty quitting (Figure 2C, $p < 10^{-10}$). On other health-related questions, similar results were obtained. Interestingly, there were no clear perception differences between age groups.

To verify that perceptions correlate with behavior in this setting, we divided the sample into “tried e-cigarettes” ($N = 40$) and “have not tried e-cigarettes” ($N = 144$) and compared perceptions of the link between smoking e-cigarettes and lung cancer. Responders who had tried e-cigarettes were more likely to answer “definitely not” and “probably not” to the question “Do you think that e-cigarettes can give you lung cancer?” than responders who had not tried e-cigarettes ($p < 10^{-4}$).

Discussion

Our findings suggest that e-cigarettes are perceived as less harmful than tobacco cigarettes. Given the vast literature on the deleterious effects of tobacco (Centers for Disease Control and Prevention, 2004; Talhout et al., 2011), it is not surprising that most people think tobacco-free e-cigarettes are safer than tobacco cigarettes. The link between tobacco and lung cancer/respiratory diseases was acknowledged by survey respondents. This relationship has been well documented (Hecht, 2012). Less well known is the potential relationship between nicotine and lung cancer (Catassi, Servent, Paleari, Cesario, & Russo, 2008). Thus, e-cigarettes may be perceived to have much lower effects on lung cancer than tobacco cigarettes, if any. Another interesting question concerns differences in nicotine addiction in e-cigarettes and tobacco cigarettes. Nicotine is known to be the main addictive component of tobacco (Picciotto & Mineur, 2014), but other components present in tobacco cigarettes may also promote addiction, including monoamine oxidase inhibitors (Guillem, Vouillac, Koob, Cador, & Stinus, 2008; Malin et al., 2013). Whether the nicotinic solution used in e-cigarettes could be used to deliver other addictive compounds beyond nicotine is an open

question. Regardless, we show here that e-cigarettes are perceived as less addictive than tobacco cigarettes.

We sought to determine if men and women perceive e-cigarettes as less harmful than tobacco cigarettes in the context of pregnancy. Perceptions that e-cigarettes are safer may lead women to trade tobacco cigarettes for e-cigarettes during pregnancy, eliminating intake of tobacco-related carcinogens while maintaining nicotine habits. Unfortunately, many of the negative fetal effects of tobacco have been specifically linked to nicotine, rather than to tobacco more generally (Ernst et al., 2001; Slotkin, 1998; Slotkin et al., 2006). Thus, e-cigarettes may be at least as deleterious as tobacco cigarettes, if not more harmful due to the potential for increased use during pregnancy in women who perceive e-cigarettes as a safer smoking alternative.

Although we did not collect demographic information related to level of education, our 184-person sample likely includes a high percentage of undergraduates, graduate students, and postdoctoral fellows living near and working in a major medical center. In this highly educated population, e-cigarettes were perceived to be safer than tobacco cigarettes. It is thus likely that similar perceptions on the safety of e-cigarettes exist in the general population, who do not have ready access to information about the negative effects of nicotine on fetal development, lung cancer, and other health risks. However, it is possible that our sample is skewed in the other direction, which highlights the need for further study on the possible effects of e-cigarettes on public health and perceptions of associated risks of e-cigarette smoking.

More research is necessary to understand the possible effects of e-cigarettes on public health. In the meantime, we advocate for a label on e-cigarettes to warn against e-smoking at least during pregnancy.

Acknowledgments

This work was supported by NIH grants DA026539 and DA09167. Additional funding and support were provided by the McNair Medical Institute and the Brain and Behavior Foundation. We acknowledge the support of the Diana Helis Henry Medical Research Foundation through its direct engagement in the continuous active conduct of medical research in conjunction with the Baylor College of Medicine Menninger Clinic, the Department of Psychiatry, and the Duncan Cancer Center. The authors would like to thank Eduardo Aramayo for technical help.

References

- Ajzen I. The theory of planned behaviour: Reactions and reflections. *Psychology & Health*. 2011; 26(9):1113–1127. [PubMed: 21929476]
- Ayers JW, Ribisl KM, Brownstein JS. Tracking the rise in popularity of electronic nicotine delivery systems (electronic cigarettes) using search query surveillance. *American Journal of Preventive Medicine*. 2011; 40(4):448–453. [PubMed: 21406279]
- Cahn Z, Siegel M. Electronic cigarettes as a harm reduction strategy for tobacco control: A step forward or a repeat of past mistakes? *Journal of Public Health Policy*. 2011; 32(1):16–31. [PubMed: 21150942]
- Catassi A, Servent D, Paleari L, Cesario A, Russo P. Multiple roles of nicotine on cell proliferation and inhibition of apoptosis: Implications on lung carcinogenesis. *Mutation Research*. 2008; 659(3): 221–231. [PubMed: 18495523]

- Centers for Disease Control and Prevention. Surgeon General's report—The health consequences of smoking. 2004. Retrieved from http://www.cdc.gov/tobacco/data_statistics/sgr/2004/index.htm
- Choi K, Fabian L, Mottey N, Corbett A, Forster J. Young adults' favorable perceptions of snus, dissolvable tobacco products, and electronic cigarettes: Findings from a focus group study. *American Journal of Public Health*. 2012; 102(11):2088–2093. [PubMed: 22813086]
- Choi K, Forster J. Characteristics associated with awareness, perceptions, and use of electronic nicotine delivery systems among young US Midwestern adults. *American Journal of Public Health*. 2013; 103(3):556–561. [PubMed: 23327246]
- Ernst M, Moolchan ET, Robinson ML. Behavioral and neural consequences of prenatal exposure to nicotine. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2001; 40(6):630–641. [PubMed: 11392340]
- Guillem K, Vouillac C, Koob GF, Cador M, Stinus L. Monoamine oxidase inhibition dramatically prolongs the duration of nicotine withdrawal-induced place aversion. *Biological Psychiatry*. 2008; 63(2):158–163. [PubMed: 17643399]
- Hecht SS. Lung carcinogenesis by tobacco smoke. *International Journal of Cancer. Journal international du cancer*. 2012; 131(12):2724–2732. [PubMed: 22945513]
- King BA, Alam S, Promoff G, Arrazola R, Dube SR. Awareness and ever-use of electronic cigarettes among U.S. adults, 2010–2011. *Nicotine & Tobacco Research*. 2013; 15(9):1623–1627. [PubMed: 23449421]
- Malin DH, Moon WD, Goyarzu P, Barclay E, Magallenes N, Vela AJ, Mills WR. Inhibition of monoamine oxidase isoforms modulates nicotine withdrawal syndrome in the rat. *Life Sciences*. 2013; 93(12–14):448–453. [PubMed: 23988853]
- Pepper JK, Reiter PL, McRee AL, Cameron LD, Gilkey MB, Brewer NT. Adolescent males' awareness of and willingness to try electronic cigarettes. *Journal of Adolescent Health*. 2013; 52(2):144–150. [PubMed: 23332477]
- Picciotto MR, Mineur YS. Molecules and circuits involved in nicotine addiction: The many faces of smoking. *Neuropharmacology*. 2014; 76(Part B):545–553. [PubMed: 23632083]
- Slotkin TA. Fetal nicotine or cocaine exposure: Which one is worse? *Journal of Pharmacology and Experimental Therapeutics*. 1998; 285(3):931–945. [PubMed: 9618392]
- Slotkin TA, Tate CA, Cousins MM, Seidler FJ. Prenatal nicotine exposure alters the responses to subsequent nicotine administration and withdrawal in adolescence: Serotonin receptors and cell signaling. *Neuropsychopharmacology*. 2006; 31(11):2462–2475. [PubMed: 16341021]
- Talhout R, Schulz T, Florek E, van Benthem J, Wester P, Opperhuizen A. Hazardous compounds in tobacco smoke. *International Journal of Environmental Research and Public Health*. 2011; 8(2): 613–628. [PubMed: 21556207]
- Wickstrom R. Effects of nicotine during pregnancy: Human and experimental evidence. *Current Neuropharmacology*. 2007; 5(3):213–222. [PubMed: 19305804]




Figure 1.

Percent of “Yes” answers to the questions “Have you ever tried a tobacco cigarette, even one or two puffs?” (dark bars) and “Have you ever tried an e-cigarette, even one or two puffs?” (light bars) according to age range: 18–20 years old ($n = 68$), 21–30 years old ($n = 55$), 31+ years old ($n = 61$).




Figure 2. Percent of answers to three representative health-related questions, separated by age groups: A, Do you think that tobacco cigarettes (or e-cigarettes) can give you lung cancer?; B, Do you think that tobacco cigarettes (or e-cigarettes) are harmful for pregnant women to use?; C, Once someone has started smoking tobacco cigarettes (or e-cigarettes), how easy or hard do you think it is to quit? The p values were calculated by comparing the distributions using the Wilcoxon sign rank test.