

Draft Genome Sequence of *Stenotrophomonas maltophilia* Strain UV74 Reveals Extensive Variability within Its Genomic Group

Oscar Conchillo-Solé,^a Daniel Yero,^{a,b} Xavier Coves,^{a,b} Pol Huedo,^{a,b} Sònia Martínez-Servat,^{a,b} Xavier Daura,^{a,c} Isidre Gibert^{a,b}

Institut de Biotecnologia i de Biomedicina (IBB), Universitat Autònoma de Barcelona (UAB), Bellaterra (Cerdanyola del Vallès), Barcelona, Spain^a; Departament de Genètica i de Microbiologia, Universitat Autònoma de Barcelona (UAB), Bellaterra (Cerdanyola del Vallès), Barcelona, Spain^b; Catalan Institution for Research and Advanced Studies (ICREA), Barcelona, Spain^c

O.C.-S. and D.Y. contributed equally to this work.

We report the draft genome sequence of *Stenotrophomonas maltophilia* UV74, isolated from a vascular ulcer. This draft genome sequence shall contribute to the understanding of the evolution and pathogenicity of this species, particularly regarding isolates of clinical origin.

Received 6 May 2015 Accepted 7 May 2015 Published 11 June 2015

Citation Conchillo-Solé O, Yero D, Coves X, Huedo P, Martínez-Servat S, Daura X, Gibert I. 2015. Draft genome sequence of *Stenotrophomonas maltophilia* strain UV74 reveals extensive variability within its genomic group. *Genome Announc* 3(3):e00611-15. doi:10.1128/genomeA.00611-15.

Copyright © 2015 Conchillo-Solé et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Xavier Daura, xavier.daura@uab.cat, or Isidre Gibert, isidre.gibert@uab.cat.

Although *Stenotrophomonas maltophilia* is an uncommon agent in human bacterial diseases, it can lead to serious nosocomial infections in immunocompromised patients (1, 2). The high genetic diversity exhibited by *S. maltophilia* clinical isolates represents a major limitation for the study of its epidemiology (3, 4). A more alarming characteristic is that most isolates are found to have intrinsic or acquired resistance mechanisms to a large number of antibiotic classes (5). Moreover, the formation of biofilms through quorum-sensing (QS) signals reduces antimicrobial effectiveness (5). *S. maltophilia* strain UV74 was isolated from a vascular ulcer in the Hospital Municipal de Badalona (Barcelona, Spain) during 2009. UV74 is a multidrug-resistant (MDR) organism, showing resistance to amikacin, imipenem, tetracycline, kanamycin, ciprofloxacin, and norfloxacin (6). This isolate also showed significantly high virulence in a zebrafish infection model and formed biofilm and adhered to HeLa cells (6). In this isolate, the diffusible signal factor (DSF)-mediated QS system is regulated by a new *rpf* cluster variant (7).

Genomic DNA was extracted with a GenElute bacterial genomics DNA kit (Sigma-Aldrich). Two different DNA fragment libraries were created and whole-genome sequencing was performed using Illumina MySeq technology at the genomics core facility of Universitat Autònoma de Barcelona, obtaining two different samples of reads. Paired-end reads were scanned for adapters using the NCBI UniVec library and the UCSC blat program (8). Detected adapters were selected and removed with Cutadapt (9) and Skewer (10) and the remaining reads were trimmed to different sizes according to the quality results obtained with FastQC. All possible combinations of trimmed reads from both samples were assembled *de novo* with VelvetOptimiser 2.2.5 (11), relying on Velvet 1.2.0 (12). Contig reorder and improvement was performed with the programs ABACAS and IMAGE, respectively, from the PAGIT package version 1 (13). As a last step, contigs were reordered once again with Mauve (14), using the *S. maltophilia* D457 complete genome (15) as the template. The final assembly

resulted in 179 contigs (GC content of 66.65%) with an N_{50} contig size of 44,960 nucleotides and covering a total of 4,889,583 bp. The average length of the contigs is 27.3 kb, and the largest contig is 283,791 bp long.

Genome annotation was performed by the NCBI Prokaryotic Genome Annotation Pipeline version 2.10 (rev. 463717), and 4,546 genes were predicted, of which there are 4,382 coding gene sequences (CDS), 89 pseudogenes, 6 rRNAs (5S, 16S, and 23S), 68 tRNAs, and 1 noncoding RNA (ncRNA). Previous multilocus sequence typing (MLST) analysis revealed that UV74 belongs to a new sequence type (<http://pubmlst.org/smaltophilia/>), being genetically very similar to the model MDR strain D457 (15) and the clinical isolate M30 (16), all three clustering in the same genomic group. Among the predicted CDS of UV74, 3,653 (83.4%) are shared by all three strains, with 299 CDS (6.8%) being exclusive to UV74. Despite the genetic similarity between the three strains in this genomic group, most of their core proteins (77%) showed some degree of variability. Consequently, this new genome may contribute to our understanding of how these bacteria evolve to adapt to different environments.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number [LBFT00000000](https://www.ncbi.nlm.nih.gov/nuclink/LBFT00000000). The version described in this paper is version [LBFT01000000](https://www.ncbi.nlm.nih.gov/nuclink/LBFT01000000).

ACKNOWLEDGMENTS

This work has been supported in part by the Seventh Research Framework Programme of the European Union (Health F3. 2009-223101), the Spanish MICINN (BFU2010-17199), and the Catalan AGAUR (2014SGR-1280).

We thank A. Calderón and T. Falgueras, from Badalona Serveis Assistencials (Hospital Municipal de Badalona), for providing the strain.

REFERENCES

1. Berg G, Martínez JL. 2015. Friends or foes: can we make a distinction between beneficial and harmful strains of the *Stenotrophomonas malto-*

- philia* complex? Front Microbiol 6:241. <http://dx.doi.org/10.3389/fmicb.2015.00241>.
2. Brooke JS. 2012. *Stenotrophomonas maltophilia*: an emerging global opportunistic pathogen. Clin Microbiol Rev 25:2–41. <http://dx.doi.org/10.1128/CMR.00019-11>.
 3. Gherardi G, Creti R, Pompilio A, Di Bonaventura G. 2015. An overview of various typing methods for clinical epidemiology of the emerging pathogen *Stenotrophomonas maltophilia*. Diagn Microbiol Infect Dis 81: 219–226. <http://dx.doi.org/10.1016/j.diagmicrobio.2014.11.005>.
 4. Valdezate S, Vindel A, Martín-Dávila P, Del Saz BS, Baquero F, Cantón R. 2004. High genetic diversity among *Stenotrophomonas maltophilia* strains despite their originating at a single hospital. J Clin Microbiol 42: 693–699. <http://dx.doi.org/10.1128/JCM.42.2.693-699.2003>.
 5. Brooke JS. 2014. New strategies against *Stenotrophomonas maltophilia*: a serious worldwide intrinsically drug-resistant opportunistic pathogen. Expert Rev Anti Infect Ther 12:1–4. <http://dx.doi.org/10.1586/14787210.2014.864553>.
 6. Ferrer-Navarro M, Planell R, Yero D, Mongiardini E, Torrent G, Huedo P, Martínez P, Roher N, Mackenzie S, Gibert I, Daura X. 2013. Abundance of the quorum-sensing factor Ax21 in four strains of *Stenotrophomonas maltophilia* correlates with mortality rate in a new zebrafish model of infection. PLoS One 8:e67207. <http://dx.doi.org/10.1371/journal.pone.0067207>.
 7. Huedo P, Yero D, Martínez-Servat S, Estibariz I, Planell R, Martínez P, Ruyra A, Roher N, Roca I, Vila J, Daura X, Gibert I. 2014. Two different *rpf* clusters distributed among a population of *Stenotrophomonas maltophilia* clinical strains display differential diffusible signal factor production and virulence regulation. J Bacteriol 196:2431–2442. <http://dx.doi.org/10.1128/JB.01540-14>.
 8. Kent WJ. 2002. BLAT—the BLAST-like alignment tool. Genome Res 12: 656–664. <http://dx.doi.org/10.1101/gr.229202>.
 9. Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. 17:10–12.
 10. Jiang H, Lei R, Ding S-W, Zhu S. 2014. Skewer: a fast and accurate adapter trimmer for next-generation sequencing paired-end reads. BMC Bioinformatics 15:182. <http://dx.doi.org/10.1186/1471-2105-15-182>.
 11. Zerbino DR. 2010. Using the Velvet *de novo* assembler for short-read sequencing technologies. Curr Protoc Bioinformatics Chapter 11:Unit 11.5. <http://dx.doi.org/10.1002/0471250953.bi1105s31>.
 12. Zerbino DR, Birney E. 2008. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. Genome Res 18:821–829. <http://dx.doi.org/10.1101/gr.074492.107>.
 13. Swain MT, Tsai IJ, Assefa SA, Newbold C, Berriman M, Otto TD. 2012. A post-assembly genome-improvement toolkit (PAGIT) to obtain annotated genomes from contigs. Nat Protoc 7:1260–1284. <http://dx.doi.org/10.1038/nprot.2012.068>.
 14. Darling AC, Mau B, Blattner FR, Perna NT. 2004. Mauve: multiple alignment of conserved genomic sequence with rearrangements. Genome Res 14:1394–1403. <http://dx.doi.org/10.1101/gr.2289704>.
 15. Lira F, Hernández A, Belda E, Sánchez MB, Moya A, Silva FJ, Martínez JL. 2012. Whole-genome sequence of *Stenotrophomonas maltophilia* D457, a clinical isolate and a model strain. J Bacteriol 194:3563–3564. <http://dx.doi.org/10.1128/JB.00602-12>.
 16. Huedo P, Conchillo-Solé O, Yero D, Martínez-Servat S, Daura X, Gibert I. 2014. Draft genome sequence of *Stenotrophomonas maltophilia* strain M30, isolated from a chronic pressure ulcer in an elderly patient. Genome Announc 2(3):e00576-14. <http://dx.doi.org/10.1128/genomeA.00576-14>.