

Two different evolutionary lines of filamentous phages in *Ralstonia solanacearum*: their effects on bacterial virulence

Ahmed Askora¹ and Takashi Yamada^{2*}

¹ Department of Microbiology and Botany, Faculty of Science, Zagazig University, Zagazig, Egypt, ² Department of Molecular Biotechnology, Graduate School of Advanced Sciences of Matter, Hiroshima University, Higashi-Hiroshima, Japan

OPEN ACCESS

Edited by:

Frank T. Robb,
University of Maryland, USA

Reviewed by:

Imke Schroeder,
University of California, Los Angeles,
USA

Steven P. T. Hooton,
Novolytics, UK

*Correspondence:

Takashi Yamada,
Department of Molecular
Biotechnology, Graduate School
of Advanced Sciences of Matter,
Hiroshima University,
1-3-1 Kagamiyama,
Higashi-Hiroshima 739-8530, Japan
tayamad@hiroshima-u.ac.jp

Specialty section:

This article was submitted to
Evolutionary and Genomic
Microbiology,
a section of the journal
Frontiers in Genetics

Received: 01 April 2015

Accepted: 03 June 2015

Published: 18 June 2015

Citation:

Askora A and Yamada T (2015) Two
different evolutionary lines
of filamentous phages in *Ralstonia
solanacearum*: their effects on
bacterial virulence.
Front. Genet. 6:217.
doi: 10.3389/fgene.2015.00217

The integration and excision of various filamentous phage genomes into and out of their host chromosomes occurs by site-specific recombination. The mechanisms proposed for these events include reactions mediated by phage-encoded recombinases and host recombination systems. Site-specific integration of filamentous phages plays a vital role in a variety of biological functions of the host, such as phase variation of certain pathogenic bacterial virulence factors. The importance of these filamentous phages in bacterial evolution is rapidly increasing with the discovery of new phages that are involved in pathogenicity. Studies of the diversity of two different filamentous phages infecting the phytopathogen *Ralstonia solanacearum* provide us with novel insights into the dynamics of phage genomes, biological roles of prophages, and the regulation and importance of phage–host interactions.

Keywords: filamentous phage, integration, pathogenic bacteria, virulence change

Filamentous Phages and Pathogenic Bacteria

Bacteriophages of the genus *Inovirus* are filamentous particles containing a circular single-stranded (ss) DNA genome. This kind of phage does not lyse host cells, but it establishes a persistent association with the host, producing and releasing phage particles from the growing and dividing host cells. The genome of inoviruses, represented by the *Escherichia coli* F-pilus-specific phage Ff (f1, fd, or M13), is generally organized in a modular structure in which functionally related genes are grouped together (Horiuchi, 1997; Rakonjac et al., 2011; Mai-Prochnow et al., 2015). Three functional modules are always present: the replication module (R), the structural module (S), and the assembly and secretion module (A-S; **Figure 1A**). The R module contains the genes encoding rolling-circle DNA replication and ssDNA-binding proteins pII, pV, and pX (Horiuchi, 1997). The S module contains genes for the major (pVIII) and minor coat proteins (pIII, pVI, pVII, and pIX). The gene *gIII* encodes the host recognition or adsorption protein pIII (Wang et al., 2006). The A-S module contains the genes for morphogenesis and extrusion of the phage particles (*gI* and *gIV*; Marvin, 1998). The gene *gIV* encodes protein pIV, an aqueous channel (secretin) in the outer membrane, through which phage particles exit from the host cells (Marciano et al., 1999). Although some phages encode their own secretins, others use host products (Davis et al., 2000). For the general infection cycle of inoviruses, see recent reviews (Rakonjac et al., 2011; Mai-Prochnow et al., 2015).

In pathogenic bacteria of either animals or plants, filamentous phage infection has been demonstrated to affect virulence. Examples include (i) enhancing production of virulence factors




FIGURE 1 | Continued

FIGURE 1 | Continued

Diversity of genomic arrangement in filamentous phages of *Ralstonia solanacearum*. (A) For ϕ RSM-type and ϕ RSS-type phages, gene insertion sites are shown along the linear genomic map of *Escherichia coli* phage M13 (Model and Russel, 1988; Marvin, 1998). Arrows indicate the direction of transcription and represent open reading frames (ORFs) or genes. The functional modules for replication (R), structure (S), and assembly and secretion (A-S) are indicated according to the M13 model. ORF sizes (in amino acids) are in parentheses. IG, intergenic region. (B) Genomic organization of ϕ RSS-type phages. According to the *E. coli* M13-model, ORFs identified in the phage genome are grouped into the R, S, and AS functional modules. IG, large intergenic region. ϕ RSS0, ϕ RSS2, ϕ RSS3, and ϕ RSS4 were derived from prophages of strains C319, M4S, MAFF106611, and MAFF211271, respectively. PE226 is a phage of Korean strains of *R. solanacearum* (Murugaiyan et al., 2010). ORFs shown in pink are homologous to ϕ RSS1 ORFs, and those in green are homologous to ϕ RSM-type ORFs. (C) Genomic organization of ϕ RSM-type phages. ϕ RSM1 and ϕ RS603 were isolated from soil (Kawasaki et al., 2007; Bich Van et al., 2014). ϕ RSM3 and ϕ RSM6 are prophages of strains MAFF730139 and CMR15 (phylo type III, Remenant et al., 2010), respectively. NP204 is similar to a phage found in the human virome (*Ralstonia* phage 1 NP2014, accession no. AHI87735.1). ORFs shown in green, red, and black are genes encoding an integrase (Int), transcriptional repressor, and ϕ RSS1-ORF11-like ORF, respectively.

such as extracellular polysaccharides (EPSs) in Xf- or Lf-infected *Xanthomonas campestris* (Kamionten and Wakimoto, 1982; Tseng et al., 1990), (ii) induction of biofilm formation in Pf4-producing *Pseudomonas aeruginosa* (Webb et al., 2004; Rice et al., 2009), and (iii) reduced twitching motility in ϕ RSM-infected *Ralstonia solanacearum* (Addy et al., 2012a) and in XacF1-infected *X. citri* (Ahmad et al., 2014). These are likely caused by changes in the host cell surface where phage proteins are secreted and filamentous particles are assembled. More direct involvement of filamentous phages in host virulence is well characterized in *Vibrio cholerae*. The pathogenicity of this severe diarrheal disease-causing bacterium depends on two key virulence factors, the toxin co-regulated pilus and cholera toxin. Cholera toxin genes are encoded on the filamentous phage CTX ϕ and are introduced into bacterial cells by phage integration mediated by the host *dif*/XerCD recombinase system (Huber and Waldor, 2002; Davis and Waldor, 2003). Also, the filamentous prophage MDA was found at multiple sites in the host chromosome associated with invasive isolates of *Neisseria meningitidis* (Bille et al., 2005). The prophage Ypf ϕ was reported to contribute to the pathogenicity of the plague bacillus, *Yersinia pestis* (Derbise et al., 2007). The acquisition of the filamentous phage CUS-1 encoding *puvA* was thought to contribute to the expression of a high-virulence phenotype in *Escherichia coli* O18:K1:H7 (Gonzalez and Allen, 2003). In these cases, filamentous phages with genes encoding toxins, virulence-enhancing factors, or host fitness factors were integrated into the host genome by various mechanisms. For other examples of filamentous phages infecting pathogenic bacteria, see the recent review by Ilyina (2015).

Different Strategies for Filamentous Phage DNA Integration into the Host Genome

To date, four different integration mechanisms used by filamentous phages have been described (Table 1). Well-characterized filamentous coliphages, such as M13 and fd, typically do not take a lysogenic replication cycle and replicate exclusively as an episome in their host bacteria (Model and Russel, 1988; Rakonjac et al., 2011). Some filamentous phages, including CTX ϕ of *V. cholerae*, accomplish site-specific integration into the *dif* site of the bacterial chromosome by

using the host XerC/D recombination system (Huber and Waldor, 2002). Filamentous phages such as VEJ ϕ of *Vibrio parahaemolyticus* (Campos et al., 2010); Cf1c, Cf1t, Cf16v1, and ϕ Lf of *X. campestris* (Campos et al., 2003); Xf1c and XacF1 of *X. citri* (Ahmad et al., 2014); Xf ϕ f1 of *Xylella fastidiosa* (de Mello Varani et al., 2008); and Ypf ϕ of *Yersinia pestis* (Lesterlin et al., 2004) also seem to use the host XerC/D recombination for their integration. In contrast, ϕ RSM1 and ϕ RSM3 of *R. solanacearum* encode a site-specific integrase (Int) of the resolvase/invertase subfamily of serine recombinases (Askora et al., 2011). This kind of serine recombinase mediates recombination involving the process of double-strand breakage followed by rotation and religation. Both integrative and excisive recombination reactions were catalyzed by ϕ RSM-Int (Askora et al., 2011). The phage *attP* corresponded to the 13 b sequence at the 5' of serine tRNA (UCG) of the host. The same unit of integration (Int-*attP*) was also found in a *R. pickettii* 12J filamentous prophage and in *Burkholderia pseudomallei* 668 prophage. A different strategy to integrate DNA into the host genome by filamentous phages may be via transposases. Kawai et al. (2005) observed filamentous prophages integrated into the chromosome of *Neisseria* species. Each prophage copy of the neisserial filamentous phage (*Nf*) was flanked by a duplication of the 5'-CT and carried an open reading frame (ORF) encoding a transposase homolog (*pivNM/irg*), suggesting the transposase-mediated integration of *Nf* DNA into host bacterial chromosomes. Bille et al. (2005) actually showed that the integration of *Nf* DNA is mediated by its own transposases (*pivNM/irg*). Meanwhile, Webb et al. (2004) and Mooij et al. (2007) characterized two filamentous prophages, Pf4 and Pf5, in the genome of *P. aeruginosa* PAO1 and PA14, respectively. Both prophages were integrated into tRNA genes of their host, probably in a reaction mediated by their own Int from the tyrosine-recombinase family. Thus, at least four different strategies for the integration of filamentous bacteriophage DNA into the host chromosome are known (Askora et al., 2012; Table 1).

Structural and Biological Diversity of Two Different Filamentous Phages Infecting *Ralstonia solanacearum*

Ralstonia solanacearum is a Gram-negative β -proteobacterium that causes bacterial wilt disease in many important crops

TABLE 1 | Comparison of site-specific recombination systems in filamentous phages.

Phage	Recombinase	Target sequence <i>attP</i>	Host	Reference
CTX ϕ	XerC/XerD	<i>dif</i>	<i>Vibrio cholerae</i>	Huber and Waldor (2002)
ϕ RSM	Resolvase/Invertase	Ser tRNA (3' -13 bp)	<i>Ralstonia solanacearum</i>	Askora et al. (2009)
Nf	Transposase	20-bp inverted repeat (dRSS3)	<i>Neisseria</i> sp.	Kawai et al. (2005)
Pf4	Tyrosine recombinase	Gly tRNA (3' -27 bp)	<i>Pseudomonas aeruginosa</i>	Webb et al. (2004)
ϕ RSS1	XerC/XerD	<i>dif</i>	<i>R. solanacearum</i>	Yamada (2013)
XacF1	XerC/XerD	<i>dif</i>	<i>Xanthomonas campestris</i>	Ahmad et al. (2014)

including tomato, potato, tobacco, eggplant, banana, ginger, and mulberry. Because of its wide geographic distribution and unusually broad host range (more than 50 plant families), it is responsible for significant crop losses worldwide (Hayward, 2000; Denny, 2006). Filamentous phages that were found to infect strains of *R. solanacearum* were classified into two groups, ϕ RSS-type and ϕ RSM-type phages. ϕ RSS1 is a representative of ϕ RSS-type phages and is a relatively small particle (1.1 μ m in length) containing an ssDNA genome of 6,662 nt (with a GC content of 62.6%) encoding 11 ORFs (Kawasaki et al., 2007). Genomic DNA of these types of phage was frequently found integrated in the host genome; 23 of 24 strains tested (all isolated in Japan) showed positive hybridization signals in Southern blot analysis (Yamada et al., 2007). Some prophage sequences were determined (Figure 1B). ϕ RSS0, ϕ RSS2, and ϕ RSS3 were derived from prophages of strain C319, M4S, and MAFF106611, respectively (Yamada, 2012). Compared with the M13 gene organization, additional genes are inserted within or next to the A-S module in these ϕ RSS genomes (Figure 1A). In the case of ϕ RSS0, a putative regulatory gene (with similarity to transcriptional repressors; *ORF13*) is inserted in the reverse orientation with two unknown ORFs (*ORF10* and *ORF11*). There is an *R. solanacearum dif* sequence within *ORF13* that serves as an *attP* site for integration into the host genome by host XerC/D recombinases (Yamada, 2013). In the case of ϕ RSS3, an additional gene encoding a transposase (IS4 family) was located in the reverse orientation. This may function for integration of the phage DNA in some occasions like the *Neisseria* cases described above (Kawai et al., 2005). Therefore, these ϕ RSS variations represent the possibility of functional equipment at this genomic region with genes for host fitness, integration, and regulatory functions (Figure 1A).

Another type of filamentous phage of *R. solanacearum* revealed a different story of evolution. ϕ RSM1, the first phage to be classified as a member of the ϕ RSM-type phages is a longer filamentous particle (1.5 μ m in length) containing ssDNA of 9,004 nt (with a GC content of 59.9%) as the genome (Kawasaki et al., 2007; Yamada et al., 2007). A total of 15 ORFs were found on the ϕ RSM1 genome including five extra genes in addition to M13-core genes. The extra genes are inserted within the R module or between the R and S modules (Figure 1C). Two of these extra genes (*orf14* and *orf15*) encode a DNA resolvase/invertase-like serine recombinase functioning as an Int (Askora et al., 2011) and a transcriptional repressor (Addy et al., 2012a), respectively. There was an *attP* site between *orf14* and *orf15* (Figure 1C). The function of the other extra genes is not known. In contrast to ϕ RSS phages, the integration of

ϕ RSM-type phage DNA into the genome of strains isolated in Japan was not frequent; 6 of 24 strains tested showed positive signals in genomic Southern blot analysis. However, genomic sequences of *R. solanacearum* strains and related β -proteobacteria in the databases frequently showed ϕ RSM-like prophage sequences. A comparison of those sequences revealed the genomic diversity of ϕ RSM-type phages as shown in Figure 1C and Supplementary Figure S1. Only one gene encoding a putative repressor (corresponding to ϕ RSM1 *ORF15*) is located within the extra region in the R module of *R. solanacearum* phage ϕ RS603 (Bich Van et al., 2014), whereas ϕ RSM6 in strain CMR15 (phylotype III) contained an Int gene (*ORF14*) in addition to the repressor gene (*ORF15*; Askora et al., 2014). Like ϕ RSM1, ϕ RSM3, ϕ RSM4 in strain UW551 (phylotype II), ϕ RSM5 in strain IPO1609 (phylotype II), and ϕ RSM7 in strain Y45 (phylotype IB) contained three genes within this region (*ORF13*, *ORF14*, and *ORF15*) with the same organization (Figure 1C). However, it is noteworthy that there are two different regulatory systems, where the amino acid sequence of *ORF15* and its upstream regulatory nucleotide sequence are different in phages infecting different phylotypes (Askora et al., 2014). ϕ RSM1, ϕ RSM3, and Y45, which infect strains of phylotype I, share similar regulatory systems, whereas ϕ RSM5, ϕ RSM6, and ϕ RSM7, which infect strains of phylotypes II or III, contained another system. Very similar ϕ RSM sequences were also found in the genomes of *R. syzygii* and *R. pickettii* (Askora et al., 2014). This kind of phage may have an extensive host range in β -proteobacteria. Interestingly, a ϕ RSM homolog was found in the human virome (*Ralstonia* phage 1 NP2014, accession no. AHI87735.1) as shown in Figure 1C. *Ralstonia* phage 1 NP2014 possesses a circular ssDNA genome that is highly homologous to those of ϕ RSM1 and ϕ RSM3. *Ralstonia* phage 1 NP2014 contains a unique *ORF11* with high similarity to ϕ RSS0 *ORF11* (Figure 1B).

As described above, two groups of filamentous phages of *R. solanacearum* have used different mechanisms for the evolution of genomic arrangements (Figure 1A). However, there may have been some opportunities for them to infect the same host cells by chance, which would have made it possible for the two types of phage to hybridize. Actually, such forms were detected (Figure 1B). A prophage (ϕ RSM4) found in strain MAFF211271 showed a gene arrangement with the ϕ RSM-type R module containing genes for an Int and regulator and with ϕ RSS-type S and A-S modules (Yamada, 2012). A smaller filamentous phage, PE226, was isolated with Korean strains and showed a gene arrangement with a ϕ RSS-type R module and ϕ RSM-type S and A-S modules (Murugaiyan et al., 2010). Therefore,

further genomic diversity by mixing these two types of phage gene arrangement is not surprising.

Filamentous Phage Diversity and Effects on the Host Virulence and Evolution in *R. solanacearum*

Both ϕ RSS-type and ϕ RSM-type filamentous phages affect the host physiology including virulence. ϕ RSS1-infected cells showed enhanced virulence on tobacco (Yamada et al., 2007) and tomato plants (Addy et al., 2012b). The virulence-enhancing effects by ϕ RSS1 infection can be explained as follows: surface-associated ϕ RSS1 particles (or phage proteins) may change the surface nature (hydrophobicity) of host cells to generate a high local cell density, resulting in early activation of *phcA*, the global virulence regulator, or lack of *orf13*, which is absent from the ϕ RSS1 genome (Addy et al., 2012b). The reduced virulence observed for ϕ RSS0-infected cells may be caused by the function(s) of ORF13 encoded by ϕ RSS0 (Yamada, 2013). Contrasting to the ϕ RSS1 effects, upon infection by ϕ RSM phages, the host cells showed loss of virulence phenotypes (Addy et al., 2012a). This loss of virulence effect of ϕ RSM infection can be explained in three ways: (i) reduced twitching motility and reduced amounts of type IV pili (Tfp), (ii) lower levels of β -1,4-endoglucanase (Egl) activity and EPS production, and (iii) reduced expression of certain virulence/pathogenicity genes (*egl*, *pehC*, *phcA*, *phcB*, *pilT*, and *hrpB*) in the infected cells (Addy et al., 2012a).

Thus, phages sometimes help host bacteria infect plants by enhancing bacterial virulence, and they sometimes interrupt bacterial infection of plants by repressing host genes involved in virulence. Such contradictory effects of these phages largely depend on the phage state, for example, replicating freely in the host, existing as a stable prophage (with *Int*), or expressing a special transcriptional regulator (Yamada, 2013). In general, the phage-encoded regulator somehow affects the expression of host genes involved in virulence, mostly through repression, in both ϕ RSS-type and ϕ RSM-type phages. However, integration into the host genome may cause a change in the regulatory function (namely direct effects on the host gene expression may be relaxed). As described above, cell surface changes

caused by filamentous phage secretion affect quorum sensing, twitching motility, and biofilm formation. Depending on the lifestyle of host bacterial cells in the environment, phage effects are different, and an advantageous state of cells with phage will be selected under the conditions. Cells whose virulence is enhanced by phage will predominate in the pathogenic stage. Similar types of phage involvement in host virulence regulation may be universal because ϕ RSS- or ϕ RSM-related sequences are frequently found in various bacterial genomic sequences, including *R. picketti* (accession no. CP001645), *R. syzygii* (FR854090), *Burkholderia rhizoxinica* (FR687359), *Pectobacterium wasabiae* (CP001790), and *Erwinia carotovora* (BX950851). The diversity observed in the genome arrangement and biological effects of filamentous phages infecting the phytopathogen *R. solanacearum* will serve as a good reference to consider interactions between various pathogenic bacteria and their phages.

Hypothesis

Filamentous phages are widely disseminated and exist as prophage states in different strains of pathogenic bacteria. They might evolve rapidly and play roles in the introduction of new genes into their hosts. Therefore, it is highly likely that filamentous phages are mediating the ecological adaptation and virulence of their hosts and thus play significant roles in the evolution of bacterial species.

Acknowledgments

This study was supported in part by the JSPS Postdoctoral Fellowship for Foreign Researchers (P13086 to AA); the Research and Development Projects for Application in Promoting New Policy of Agriculture, Forestry, and Fisheries (No. 250037B to TY); and JST/BIOTEC Strategic Research Cooperative Program on Biotechnology (A1200357 to TY).

Supplementary Material

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fgene.2015.00217>

References

- Addy, H. S., Askora, A., Kawasaki, T., Fujie, M., and Yamada, T. (2012a). Loss of virulence of the phytopathogen *Ralstonia solanacearum* through infection by ϕ RSM filamentous phages. *Phytopathology* 102, 469–477. doi: 10.1094/PHYTO-11-11-0319-R
- Addy, H. S., Askora, A., Kawasaki, T., Fujie, M., and Yamada, T. (2012b). The filamentous phage ϕ RSS1 enhances virulence of phytopathogenic *Ralstonia solanacearum* on tomato. *Phytopathology* 102, 244–251. doi: 10.1094/PHYTO-10-11-0277
- Ahmad, A. A., Askora, A., Kawasaki, T., Fujie, M., and Yamada, T. (2014). A novel filamentous phage causes loss of virulence to *Xanthomonas axonopodis* pv *citri* the causative agent of citrus canker disease. *Front. Microbiol.* 5:321. doi: 10.3389/fmicb.2014.00321
- Askora, A., Abdel-Haliem, M. E. F., and Yamada, T. (2012). Site-specific recombination systems in filamentous phages. *Mol. Genet. Genom.* 287, 525–530. doi: 10.1007/s00438-012-0700-1
- Askora, A., Kawasaki, T., Fujie, M., and Yamada, T. (2011). Resolvase-like serine recombinase mediates integration/excision in the bacteriophage ϕ RSM. *J. Biosci. Bioeng.* 111, 109–116. doi: 10.1016/j.jbiosc.2010.10.001
- Askora, A., Kawasaki, T., Fujie, M., and Yamada, T. (2014). Insight into the diversity of ϕ RSM phages infecting strains of the phytopathogen *Ralstonia solanacearum* complex: regulation and evolution. *Mol. Genet. Genomics*, 289, 589–598. doi: 10.1007/s00438-014-0835-3
- Askora, A., Kawasaki, T., Usami, S., Fujie, M., and Yamada, T. (2009). Host recognition and integration of filamentous phage ϕ RSM in the phytopathogen, *Ralstonia solanacearum*. *Virology* 384, 69–76. doi: 10.1016/j.virol.2008.11.007

- Bich Van, T. T., Yoshida, S., Miki, K., Kondo, A., and Kamei, K. (2014). Genomic characterization of ϕ RS603, a filamentous bacteriophage that is infectious to the phytopathogen *Ralstonia solanacearum*. *Microbiol. Immunol.* 58, 697–700. doi: 10.1111/1348-0421.12203
- Bille, E., Zahar, J. R., Perrin, A., Morelle, S., Kriz, P., Jolley, K. A., et al. (2005). A chromosomally integrated bacteriophage in invasive meningococci. *J. Exp. Med.* 201, 1905–1913. doi: 10.1084/jem.20050112
- Campos, J., Martinez, E., Izquierdo, Y., and Fando, R. (2010). VEJ ϕ , a novel filamentous phage of *Vibrio cholerae* able to transduce the cholera toxin genes. *Microbiology* 156, 108–115. doi: 10.1099/mic.0.032235-0
- Campos, J., Martinez, E., Suzarte, E., Rodriguez, B. E., Marrero, K., Silva, Y., et al. (2003). A novel filamentous phage of *Vibrio cholerae*, integrates into the same chromosomal site as CTX ϕ . *J. Bacteriol.* 185, 5685–5696. doi: 10.1128/JB.185.19.5685-5696.2003
- Davis, B. M., Lawson, E. H., Sandkvist, M., Sozhamannan, S., Ali, A., and Waldor, M. K. (2000). Convergence of the secretory pathways for cholera toxin and the filamentous phage, CTX ϕ . *Science* 288, 333–335. doi: 10.1126/science.288.5464.333
- Davis, B. M., and Waldor, M. K. (2003). Filamentous phages linked to virulence of *Vibrio cholerae*. *Curr. Opin. Microbiol.* 6, 35–42. doi: 10.1016/S1369-5274(02)00005-X
- de Mello Varani, A., Souza, R. C., Nakaya, H. I., de Lima, W. C., de Almeida, P., Kitajima, E. W., et al. (2008). Origins of the *Xylella fastidiosa* prophage-like regions and their impact in genome differentiation. *PLoS ONE* 3:e4059. doi: 10.1371/journal.pone.0004059
- Denny, T. P. (2006). “Plant pathogenic *Ralstonia* species” in *Plant-Associated Bacteria*, ed. S. S. Gnanamanickam (Amsterdam: Springer) 573–644. doi: 10.1007/978-1-4020-4538-7_16
- Derbise, A., Chenal-Francois, V., Pouillot, F., Fayolle, C., Prevost, M. C., Medigue, C., et al. (2007). A horizontally acquired filamentous phage contributes to the pathogenicity of the plague *Bacillus*. *Mol. Microbiol.* 63, 1145–1157. doi: 10.1111/j.1365-2958.2006.05570.x
- Gonzalez, E. T., and Allen, C. (2003). Characterization of a *Ralstonia solanacearum* operon required for polygalacturonate degradation and uptake of galacturonic acid. *Mol. Plant Microbe Interact.* 16, 536–544.
- Hayward, A. C. (2000). “*Ralstonia solanacearum*,” in *Encyclopedia of Microbiology*, Vol. 4, ed. J. Lederberg (San Diego, CA: Academic Press), 32–42.
- Horiuchi, K. (1997). Initiation mechanisms in replication of filamentous phage DNA. *Genes Cells* 2, 425–432. doi: 10.1046/j.1365-2443.1997.1360334.x
- Huber, K. E., and Waldor, M. K. (2002). Filamentous phage integration requires the host recombinases XerC and XerD. *Nature* 417, 656–659. doi: 10.1038/nature00782
- Ilyina, T. S. (2015). Filamentous bacteriophages and their role in the virulence and evolution of pathogenic bacteria. *Mol. Genet. Microbiol. Virol.* 30, 1–9. doi: 10.3103/S0891416815010036
- Kamiyama, H., and Wakimoto, S. (1982). Effect of the infection with filamentous phage Xf-2 on the properties of *Xanthomonas campestris* var *oryzae*. *Ann. Phytopathol. Soc. Japan* 47, 627–636. doi: 10.3186/jjphytopath.47.627
- Kawai, M., Uchiyama, I., and Kobayashi, I. (2005). Genome comparison in silico in *Neisseria* suggests integration of filamentous bacteriophages by their own transposase. *DNA Res.* 12, 389–401. doi: 10.1093/dnares/dsi021
- Kawasaki, T., Nagata, S., Fujiwara, A., Satsuma, H., Fujie, M., Usami, S., et al. (2007). Genomic characterization of the filamentous integrative bacteriophage ϕ RSS1 and ϕ RSM1, which infect *Ralstonia solanacearum*. *J. Bacteriol.* 189, 5792–5802. doi: 10.1128/JB.00540-07
- Lesterlin, C., Barre, F. X., and Cornet, F. (2004). Genetic recombination and the cell cycle: what we have learned from chromosome dimers. *Mol. Microbiol.* 54, 1151–1160. doi: 10.1111/j.1365-2958.2004.04356.x
- Mai-Prochnow, A., Hui, J. G. K., Kjelleberg, S., Rakonjac, J., McDougald, D., and Rice, S. A. (2015). Big things in small packages: the genetics of filamentous phage and effects on fitness of their host. *FEMS Microbiol. Rev.* doi: org/10.1093/femsre/fuu007 [Epub ahead of print].
- Marciano, D. K., Russel, M., and Simon, S. M. (1999). An aqueous channel for filamentous phage export. *Science* 284, 1516–1519. doi: 10.1126/science.284.5419.1516
- Marvin, D. A. (1998). Filamentous phage structure, infection and assembly. *Curr. Opin. Struct. Biol.* 8, 150–158. doi: 10.1016/S0959-440X(98)80032-8
- Model, P., and Russel, M. (1988). “Filamentous bacteriophages,” in *The Bacteriophages*, Vol. 2, ed. R. Calendar (New York, NY: Plenum Press) 375–456.
- Mooij, M. J., Drenkard, E., Llamas, M. A., Vandenbroucke-Grauls, C. M. J. E., Savelkoul, P. H. M., Ausubel, F. M., et al. (2007). Characterization of the integrated filamentous phage Pf5 and its involvement in small-colony formation. *Microbiology* 153, 1790–1798. doi: 10.1099/mic.0.2006/003533-0
- Murugaiyan, S., Bae, J. Y., Wu, J., Lee, S. D., Um, H. Y., Choi, H. K., et al. (2010). Characterization of filamentous bacteriophage PE226 infecting *Ralstonia solanacearum* strains. *J. Appl. Microbiol.* 110, 296–303. doi: 10.1111/j.1365-2672.2010.04882.x
- Rakonjac, J., Bennet, N. J., Spagnuolo, J., Gagic, D., and Russel, M. (2011). Filamentous bacteriophage: biology, phage display and nanotechnology applications. *Curr. Issues Mol. Biol.* 13, 51–76.
- Remenant, B., Coupat-Goutaland, B., Guidot, A., Cellier, G., Wicker, E., Allen, C., et al. (2010). Genomes of three tomato pathogens within the *Ralstonia solanacearum* species complex reveal significant evolutionary divergence. *BMC Genomics* 11:319. doi: 10.1186/1471-2164-11-379
- Rice, S. A., Tan, C. H., Mikkelsen, P. J., Kung, V., Woo, J., Tay, M., et al. (2009). The biofilm life cycle and virulence of *Pseudomonas aeruginosa* are dependent on a filamentous prophage. *ISME J.* 3, 271–282. doi: 10.1038/ismej.2008.109
- Tseng, Y. H., Lo, M. C., Lin, K. C., Pan, C. C., and Chang, R. Y. (1990). Characterization of filamentous bacteriophage ϕ Lf from *Xanthomonas campestris* pv. *campestris*. *J. Gen. Virol.* 71, 1881–1884. doi: 10.1099/0022-1317-71-8-1881
- Wang, Y. A., Yu, X., Overman, S., Tsuboi, M., Tomas, G. J. Jr. and Egelman, E. H. (2006). The structure of a filamentous bacteriophage. *J. Mol. Biol.* 361, 209–215. doi: 10.1016/j.jmb.2006.06.027
- Webb, J. S., Lau, M., and Kjelleberg, S. (2004). Bacteriophage and phenotypic variation in *Pseudomonas aeruginosa* biofilm development. *J. Bacteriol.* 186, 8066–8073. doi: 10.1128/JB.186.23.8066-8073.2004
- Yamada, T. (2012). “Bacteriophages of *Ralstonia solanacearum*: their diversity and utilization as biocontrol agents in agriculture,” in *Bacteriophages*, ed. I. Kurtboke (Rijeka: InTech-Open Access Publisher), 113–139.
- Yamada, T. (2013). Filamentous phages of *Ralstonia solanacearum*: double-edged swords for pathogenic bacteria. *Front. Microbiol.* 4:325. doi: 10.3389/fmicb.2013.00325
- Yamada, T., Kawasaki, T., Nagata, S., Fujiwara, A., Usami, S., and Fujie, M. (2007). New bacteriophages that infect the phytopathogen *Ralstonia solanacearum*. *Microbiology* 153, 2630–2639. doi: 10.1099/mic.0.2006/001453-0

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Askora and Yamada. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.