



HHS Public Access

Author manuscript

Int J Dev Neurosci. Author manuscript; available in PMC 2015 August 01.

Published in final edited form as:

Int J Dev Neurosci. 2014 August ; 36: 38–44. doi:10.1016/j.ijdevneu.2014.05.006.

Perinatal α -linolenic acid availability alters the expression of genes related to memory and to epigenetic machinery, and the *Mecp2* DNA methylation in the whole brain of mouse offspring

Fuli He¹, Daniel S. Lupu, and Mihai D. Niculescu*

Department of Nutrition and Nutrition Research Institute, University of North Carolina at Chapel Hill, Kannapolis, NC 28081, USA

Abstract

Many animal and human studies indicated that dietary ω -3 fatty acids could have beneficial roles on brain development, memory, and learning. However, the exact mechanisms involved are far from being clearly understood, especially for α -linolenic acid (ALA), which is the precursor for the ω -3 elongation and desaturation pathways. This study investigated the alterations induced by different intakes of flaxseed oil (containing 50% ALA), during gestation and lactation, upon the expression of genes involved in neurogenesis, memory-related molecular processes, and DNA methylation, in the brains of mouse offspring at the end of lactation (postnatal day 19, P19). In addition, DNA methylation status for the same genes was investigated. Maternal flaxseed oil supplementation during lactation increased the expression of *Mecp2*, *Ppp1cc*, and *Reelin*, while decreasing the expression of *Ppp1cb* and *Dnmt3a*. *Dnmt1* expression was decreased by postnatal flaxseed oil supplementation but this effect was offset by ALA deficiency during gestation. *Mecp2* DNA methylation was decreased by maternal ALA deficiency during gestation, with a more robust effect in the lactation-deficient group. In addition, linear regression analysis revealed positive correlations between *Mecp2*, *Reelin*, and *Ppp1cc*, between *Gadd45b*, *Bdnf*, and *Creb1*, and between *Egr1* and *Dnmt1*, respectively. However, there were no correlations, in any gene, between DNA methylation and gene expression. In summary, the interplay between ALA availability during gestation and lactation differentially altered the expression of genes involved in neurogenesis and memory, in the whole brain of the offspring at the end of lactation. The *Mecp2* epigenetic status was correlated with ALA availability during gestation. However, the epigenetic status of the genes investigated was not associated with transcript levels, suggesting that either the regulation of these genes is not necessarily under epigenetic control, or that the whole brain model is not adequate for the exploration of epigenetic regulation in the context of this study.

Keywords

Flaxseed oil; α -Linolenic acid; Perinatal nutrition; Brain development; DNA methylation; Lactation

© 2014 ISDN. Published by Elsevier Ltd. All rights reserved.

*Corresponding author at: UNC Nutrition Research Institute, Rm. 2104, 500 Laureate Way, Kannapolis, NC 28081, USA. Tel.: +1 704 250 5029. address: Mihai_Niculescu@unc.edu (M.D. Niculescu).

¹Present address: Center for Excellence in Post-harvest Technologies, North Carolina A&T State University, Kannapolis, NC 28081, USA.

1. Introduction

Memory is the process by which new information is being encoded, stored, and then retrieved (Abel and Lattal, 2001). Specificity in gene expression, protein synthesis and structural properties of neurons and synapses, is required for all three stages (Alberini, 2009). It has been acknowledged that dietary ω -3 polyunsaturated fatty acids (ω -3 PUFAs) could improve learning and memory, and also alter gene expression and proteome profiles in the whole brain or certain brain regions, such as cerebrum and hippocampus, therefore altering neurogenesis and synaptic plasticity (Kitajka et al., 2004; Lee et al., 2012).

α -Linolenic acid (ALA; 18:3 n-3) is the precursor for the ω -3 PUFAs elongation and desaturation pathways. This 18-carbon fatty acid possesses three double bonds and is commonly found in plant-derived dietary oils such as flaxseed, canola, and soybean oils. All the other common dietary ω -3 PUFAs have longer chains, and are found mainly in fish and fish oils, including the 20-carbon eicosapentaenoic acid (EPA; 20:5 n-3) and the 22-carbon docosahexaenoic acid DHA (Surette, 2013). Although numerous studies have defined the roles of DHA in brain development, less is known about the roles of ALA.

ALA availability has been shown to be important for brain development and its function during gestation and early postnatal life. During the perinatal period, the neonate's brain is experiencing a substantial acceleration in growth, cellular proliferation, and neuronal and glial differentiation (Hoffman et al., 2009). Also, perinatal ALA deficiency induced gene expression changes in the brain of adult rats (Kitajka et al., 2004). We have previously indicated that, in mice, ALA availability during gestation and lactation altered cell proliferation, early neuronal differentiation, and apoptosis, in the hippocampus of the offspring (Niculescu et al., 2011). These outcomes were associated with postnatal ALA supplementation, but were offset by gestational ALA deficiency. We also reported that the perinatal ALA availability was associated with epigenetic alterations in *Fads2* DNA methylation in maternal and offspring livers (Niculescu et al., 2013).

Recent studies indicated the importance of epigenetic mechanisms in the alteration of neuronal gene expression patterns, which are required for synaptic plasticity or memory formation (reviewed in Sultan and Day, 2011). Epigenetic mechanisms include interrelated processes such as DNA methylation, histone modifications and RNA interference (Wood et al., 2005; Feng et al., 2010; Lubin et al., 2011).

In the current study, we explored the impact of maternal perinatal ALA availability on the expression of memory-associated genes in offspring brains, and the associated modifications in DNA methylation. Because memory and learning are complex processes that are not confined to only a brain area (Vousden et al., 2014; van Groen et al., 2014), we sought to first determine, in a whole brain model, whether such alterations could be detected, and whether this approach could warrant additional and more specific explorations.

2. Materials and methods

All reagents were purchased from Sigma–Aldrich (St. Louis, MO, USA) if not otherwise specified.

2.1. Animals, diets, and tissue collection

This study was approved by the University of North Carolina Institutional Animal Care and Use Committee. This work is a continuation of our previously published studies (Niculescu et al., 2011, 2013). Briefly, mouse C57BL/6J females (10 wk old; Jackson Laboratory, Bar Harbor, ME, USA) were randomly assigned into two initial feeding groups (intervention 1) for 30 days prior, and during gestation. One group was fed a defined control diet (AIN-93G, DYETS, Bethlehem, PA) containing soybean oil as fat source (ALA control, C, $n = 12$), while the other group was given an AIG-93G modified diet (DYETS) with corn oil as the only source of fatty acids (ALA deficient, D, $n = 12$). After 30 days, the females were bred overnight with males that were maintained at all times on the C diet. One day prior to delivery date, the two groups of females were randomly split into two subgroups (intervention 2). Half from each group remained on the same diet ($n = 6$), and the other half were switched to a modified AIN-93G diet containing flaxseed oil (ALA supplemented, S, $n = 6$), until postnatal day 20 (P19, considering the date of delivery as P0). On P19, all mothers and their pups were sacrificed. The pups were decapitated, the brains were extracted, snap frozen in liquid nitrogen, and stored at -80°C . In this study, the brain samples from the male pups were used for further analysis. However, due to the lack of male pups, only $n = 5$ male brains were included here in the CC, CS and DS respectively, while the DD group had $n = 6$ samples. For further reference, each group is designated by two letters: first letter reflects the diet received prior and during gestation, and the second letter reflects the maternal diet received during lactation.

The fatty acid composition of each diet has been previously published (Niculescu et al., 2011). Particularly, the diets contained 18:2 n-6 (linoleic acid, LA) and 18:3 n-3 (ALA) with the following concentrations: D diet, 87.2 and 1.5 nmol/mg; C diet, 86.9 and 10.8 nmol/mg; S diet, 23.1 and 75.3 nmol/mg.

2.2. Real-time RT-PCR

RNA was extracted from the male offspring brains (all at P19) with a QIAcube instrument (Qiagen, Valencia, CA, USA) using AllPrep[®] DNA/RNA mini kit (Qiagen), according to manufacturer's protocol. After the concentration and quality were determined with a NanoPhotometer (Implen GmbH, Munich, Germany), real time reverse transcriptase PCR (RT-PCR) method was used for the assessment of gene expression. First, cDNA synthesis was performed using a QuantiTect reverse transcription kit (Qiagen), on an Eppendorf Mastercycler ProS (Eppendorf, Hamburg, Germany). The amplification was performed on an Eppendorf Mastercycler Realplex² ep Gradient S real-time PCR cycler, using the QuantiTect SYBR Green PCR kit (Qiagen). QuantiTect primer assays (Qiagen) were purchased for the following genes: *Dnmt1*, *Dnmt3a*, *Mecp2*, *Gadd45b*, *Ppp1cb*, *Ppp1cc*, *Ppp3ca*, *Arc*, *Bdnf*, *Creb1*, *Reelin*, *Egr1* and *18s* (used as internal reference for each sample). The real-time PCR reactions were run in triplicate, and data were retrieved as C_T

values normalized to 18S, then \log_2 transformed for subsequent statistical analysis. For each gene final data were expressed as ratios between each sample and the average of the CC group.

2.3. Bisulfite pyrosequencing

DNA from male offspring brains (all at P19) was extracted with a QIAcube instrument (Qiagen, Valencia, CA, USA) using AllPrep[®] DNA/RNA mini kit (Qiagen), following manufacturer's protocol. Up to 1 μg DNA from each sample was treated with sodium bisulfite using the same instrument and the corresponding protocol (EpiTect kit, Qiagen).

A pyrosequencing DNA methylation assay was designed for *Mecp2* promoter using PyroMark Assay Design 2.0 software (Qiagen). The other 8 PyroMark CpG assays were directly purchased from Qiagen. Primers and sequences to be analyzed are indicated in Table 1. The amplification of templates was performed on an Eppendorf Mastercycler ProS using a PyroMark PCR kit and following the recommended protocol (Qiagen). The PCR conditions were: 95 °C for 15 min; 45 cycles of 94 °C for 30s, 56°C for 30 s, 72°C for 30s; final extension at 72°C for 10 min, and 4°C hold. The biotinylated strand of the amplified DNA was subjected to pyrosequencing on a PyroMark MD machine (Qiagen), as described previously (Mehedint et al., 2010). Each sample and sequence were run in triplicate, and only pyrosequencing reactions that passed the quality test were included in the analysis. For each sample and sequence, percentage methylated cytosine for each CpG site was expressed as the average of triplicates, and the values across all sites were averaged to express the average methylation across each sequence.

2.4. Statistical analysis

Statistical analysis was carried out using both JMP 10 software (SAS, Cary, NC, USA) and MeV 4.9. (Dana-Farber Cancer Institute, Boston, MA, USA). MeV was used for the assessment of false discovery rates (FDR, Benjamini-Hochberg method) in the case of both gene expression and DNA methylation assessment. For each variable the p value ($p < 0.05$) was adapted accordingly to reflect a FDR of 5%. MeV was also used for Cluster Affinity Search Technique (CAST) on gene expression. For DNA methylation, because of its beta distribution, multiple comparisons were performed using JMP with the nonparametric Wilcoxon Each Pair test against adjusted p values calculated by the Kruskal–Wallis test (MeV). Gene expression was analyzed in JMP using two-way ANOVA (for the gestation and lactation factors), followed by Tukey HSD test for multiple comparisons, against adjusted p values. In addition, bivariate linear fit analysis was performed (JMP) for the correlation between two variables, with a threshold for significance of $p < 0.05$.

3. Results

3.1. Perinatal ALA availability alters the expression of genes associated with memory and DNA methylation

The expression of eight genes involved in memory and brain development, and of four genes involved in the regulation of DNA methylation, was determined in the whole brains of pups at the end of lactation (P19). The expression of three genes associated with memory was

altered by ALA supplementation during lactation (*Reelin*, *Ppp1cb*, and *Ppp1cc*), regardless the ALA status during pregnancy (Fig. 1A). The expression of *Reelin* was increased in both CS and DS groups as compared to CC (ratio 5.44 ± 1.65 SE, and ratio 7.96 ± 1.60 , respectively). While DS alterations were statistically significant, the CS alterations did not reach statistical significance ($p = 0.017$ vs. the FDR-corrected threshold $p = 0.008$). However, the expression values in both ALA supplemented groups were significantly higher than in the DD group. The expression of *Ppp1cb* was significantly lower in the CS group as compared to CC (ratio 0.23 ± 0.05 SE). *Ppp1cc* expression was higher in the CS and DS groups as compared to both CC and DD groups (ratio 5.70 ± 1.25 SE vs. CC, and ratio 6.72 ± 1.76 SE vs. CC, respectively).

Among the genes involved in the epigenetic regulation of gene expression, three genes (*Dnmt1*, *Dnmt3a*, and *Mecp2*) were altered by perinatal ALA availability (Fig. 1B). *Dnmt1* expression was decreased only in the CS group when compared to all other three treatment groups (ratio 0.29 ± 0.08 SE vs. CC). The expression of *Dnmt3a* was markedly lower in both ALA supplemented groups (CS and DS) as compared to either the CC or the DD groups (ratio 0.012 ± 0.003 SE vs. CC, and ratio 0.053 ± 0.03 SE vs. CC). The expression of *Mecp2* was higher in ALA-supplemented groups. However, only the DS group reached statistical significance against both CC and DD groups (ratio 6.27 ± 1.40 SE vs. CC), while the increased expression in the CS group (ratio 4.33 ± 1.44 SE vs. CC) was not statistically significant ($p = 0.055$ vs. the FDR-corrected threshold $p = 0.008$) when compared to the CC group, but only against the DD group.

3.2. Correlations between the expression of memory-associated genes and epigenetic regulators

We sought to determine whether the expression of genes involved in memory and neuronal development were correlated with the expression of epigenetic regulators. Using CAST analysis, we first identified clusters of genes that were candidates for such correlations. The identified clusters were then subject of linear regression for the assessment of statistical significance and coefficient of determination (R^2), against a significance threshold $p = 0.05$. Several clusters were identified.

Mecp2, *Reelin*, and *Ppp1cc* formed one cluster (Fig. 2), as both the expression of *Reelin* (Fig. 2A, $R^2 = 0.98$ and $p < 0.0001$) and *Ppp1cc* (Fig. 2B, $R^2 = 0.92$ and $p < 0.0001$) were highly correlated with *Mecp2*.

A second cluster consisted of *Creb1*, *Gadd45b*, and *Bdnf* (Fig. 3). Linear regression analysis indicated that both *Bdnf* (Fig. 3A, $R^2 = 0.65$ and $p < 0.0001$) and *Creb1* (Fig. 3B, $R^2 = 0.87$ and $p < 0.0001$) were associated with *Gadd45b*.

In addition, *Dnmt1* expression was significantly correlated with *Egr1* across all samples (Fig. 4, $R^2 = 0.86$ and $p < 0.0001$).

3.3. Effects of the maternal ALA treatment on DNA methylation in the offspring brains

The assessment of DNA methylation status was performed by bisulfite pyrosequencing for the genes and sequences indicated in Table 1. *Mecp2* promoter DNA methylation (Fig. 5)

was altered by perinatal ALA availability in the CS and DD groups. *Mecp2* promoter methylation in the DD group was significantly lower (% methylation 2.82 ± 0.35 SE) than in both CS and CC groups (% methylation 4.43 ± 0.19 SE, and 5.56 ± 0.44 SE, respectively), when tested against a FDR-adjusted value of $p = 0.022$. However, no differences between groups were noted for the *Mecp2* intron 1 methylation. For all other DNA methylation assays, no significant changes were noted when testing against FDR-adjusted p values (data not shown).

3.4. Lack of correlation between gene expression and DNA methylation

Using a linear regression model, we sought to determine whether gene expression correlates with the DNA methylation status for the genes presented in Table 1. Using a threshold for statistical significance of $p < 0.05$, no correlations were found for any of the genes, including *Mecp2* (data not shown).

4. Discussion

The findings of this study support the hypothesis that maternal ALA availability during pregnancy and lactation alters the expression of memory-associated genes and of those involved in the DNA methylation machinery, in the offspring brain at P19 (Fig. 1). In addition, ALA deficiency decreased the DNA methylation within the *Mecp2* promoter, but not within intron 1 (Fig. 5). Interestingly, among the six genes that had substantial alterations in their transcript levels, five (*Reelin*, *Ppp1cb*, *Ppp1cc*, *Dnmt3a*, and *Mecp2*) were dependent exclusively on postnatal maternal ALA availability (regardless gestational ALA intakes), while *Dnmt1* was decreased only in the CS group, indicating that postnatal ALA supplementation triggered its decreased expression, which was nevertheless offset by maternal ALA deficiency during gestation. The increased expression of *Ppp1cc*, *Reelin* and *Mecp2* in the CS and DS groups support, in part, our previous findings about the enhanced neuronal differentiation due to postnatal ALA supplementation (Niculescu et al., 2011).

Reelin has been associated with the maintenance of polarity in cortical neurons and in the structural development of juvenile prefrontal circuits and memory formation (Forster, 2014; Iafrati et al., 2014). Moreover, the cognitive decline registered in aged rats was also associated with a reduction of *Reelin* expression (Stranahan et al., 2011).

Ppp1cc encodes one of the catalytic subunit (PP1 γ) of nuclear protein-phosphatase 1 (Pp1), which was reported to decrease phosphorylation and ubiquitination of cAMP response element-binding protein (CREB), and thereby promote neurogenesis (Mu et al, 2013, 2011). While the role of the Ppp1cc subunit has not been sufficiently studied in brain, controversial roles have been assigned to the PP1 complex in regard to memory. In CD1 mice, PP1 levels were positively associated with olfactory memory (Winding et al., 2011). In contrast, increased PP1 activity was implicated in deficits in learning and memory induced by lead exposure in postnatal rats (Rahman et al., 2012), and in the epigenetic suppression of fear memory and synaptic plasticity in the amygdala (Koshibu et al., 2011). Moreover, the regulation of PP1 transcription could be epigenetically controlled by its DNA methylation (Miller and Sweatt, 2007). However, in our study, we did not find any DNA methylation

differences between groups, which indicated that *Ppp1cc* expression control might be independent of its DNA methylation status in the context of maternal ALA availability.

In contrast with the increased *Ppp1cc* expression by postnatal ALA supplementation, *Ppp1cb* was decreased in same groups (CS and DS). This gene encodes the PP1 catalytic subunit PP1 β (Saadat et al., 1994), and its nuclear inhibition was associated with increased memory performance (Graff et al., 2010). Again, our study did not identify any epigenetic modifications between groups.

The alterations induced by ALA availability to DNA methyltransferases (*Dnmt1* and *Dnmt3a*), and to *Mecp2* expression were more specific (Fig. 1B). *Dnmt1* expression was decreased only in the CS group but not in the DS group, indicating that the influence of postnatal ALA supplementation upon *Dnmt1* is offset by gestational ALA deficiency. *Dnmt1* is a maintenance DNA methyltransferase required for the propagation of DNA methylation patterns to replicated DNA (Jin and Robertson, 2013). Because *Dnmt1* is over expressed in post-mitotic neurons, then down-regulated in adult neurons (Inano et al., 2000), one could speculate that the *Dnmt1* reduced expression in the CS group could reflect the more advanced differentiated state of neurons, which we previously reported to be the case for hippocampus (Niculescu et al., 2011). However, in the same study, we also reported that the CS group had increased cellular proliferation. Therefore, in this study, it is difficult to interpret the biological meaning of *Dnmt1* decreased expression with ALA supplementation.

Dnmt3a expression was decreased in both postnatal ALA supplemented groups (CS and DS), irrespective of the ALA intakes during gestation. This suggests that, at P19, *Dnmt3a* expression in postnatal brains is not dependent upon gestational ALA levels. Because *Dnmt3a* is transiently expressed during neuronal differentiation (Watanabe et al., 2006), ALA-induced under-expression could suggest an increased differentiation state of the offspring brains exposed to postnatal ALA supplementation.

Mecp2 is an important epigenetic factor in the maintenance and development of the neurotransmission in cortical and hippocampal regions of the brain, through the regulation of synaptic gene expression (Na et al., 2013). Strongly involved in the development of central nervous system, *Mecp2* deficiency was associated in mouse models with reduced neuronal dendritic formation (Schule et al., 2008). In the present study *Mecp2* was over-expressed by postnatal ALA supplementation, and the increase was more robust statistically in the DS group (Fig. 1B). Because the increase in the CS group was not statistically significant due to one outlier, it is difficult to ascertain whether gestational ALA exposure was an independent factor. While *Mecp2* promoter DNA methylation was decreased in the DD group as compared with either CC or CS groups, but not when compared with the DS group, one might speculate that its methylation status would be dependent primarily on the gestational ALA availability, rather than its postnatal intake levels. Linear regression indicated no association between *Mecp2* expression levels and its DNA methylation status (data not shown).

Although the dietary treatments altered the expression of only six genes out of twelve, we sought to also investigate the relationship between the expression of genes, independent of

availability alters the content of EPA and DPA in the brains of the offspring (Niculescu et al., 2011), it is not clear whether ALA's role in altering the expression of genes is direct (via signaling), or due to alterations in other ω -3 species.

Our study suggested that ALA availability during gestation and lactation differentially altered the expression of genes involved in memory formation and neurogenesis, and in the epigenetic regulation of gene expression. Some of these alterations could be the result of the interplay between ALA availability in the two developmental periods, while others depended exclusively on postnatal ALA supplementation. With the notable exception of *Mecp2* promoter methylation, ALA availability did not alter the methylation of the aforementioned genes, as measured in P19 whole brain extracts. Also, *Mecp2* methylation was not correlated with its expression. Whether potential correlations between gene expression and DNA methylation could be altered by ALA availability in specific brain areas, it remains to be investigated further.

Acknowledgments

This work was funded, in part, by a grant to MDN from the UNC Center of Excellence in Children's Nutrition sponsored by Mead Johnson Nutrition.

References

- Abel T, Lattal KM. Molecular mechanisms of memory acquisition, consolidation and retrieval. *Curr Opin Neurobiol.* 2001; 11:180–187. [PubMed: 11301237]
- Alberini CM. Transcription factors in long-term memory and synaptic plasticity. *Physiol Rev.* 2009; 89:121–145. [PubMed: 19126756]
- Blondeau N, Widmann C, Lazdunski M, Heurteaux C. Activation of the nuclear factor- κ B is a key event in brain tolerance. *J Neurosci.* 2001; 21:4668–4677. [PubMed: 11425894]
- Cartron PF, Blanquart C, Hervouet E, Gregoire M, Vallette FM. HDAC1-mSin3a-NCOR1, Dnmt3b-HDAC1-Egr1 and Dnmt1-PCNA-UHRF1-G9a regulate the NY-ESO1 gene expression. *Mol Oncol.* 2013; 7:452–463. [PubMed: 23312906]
- Darios F, Davletov B. Omega-3 and omega-6 fatty acids stimulate cell membrane expansion by acting on syntaxin 3. *Nature.* 2006; 440:813–817. [PubMed: 16598260]
- Deng C, Lu Q, Zhang Z, Rao T, Attwood J, Yung R, Richardson B. Hydralazine may induce autoimmunity by inhibiting extracellular signal-regulated kinase pathway signaling. *Arthritis Rheum.* 2003; 48:746–756. [PubMed: 12632429]
- Feng J, Zhou Y, Campbell SL, Le T, Li E, Sweatt JD, Silva AJ, Fan G. Dnmt1 and Dnmt3a maintain DNA methylation and regulate synaptic function in adult forebrain neurons. *Nat Neurosci.* 2010; 13:423–430. [PubMed: 20228804]
- Forster E. Reelin, neuronal polarity and process orientation of cortical neurons. *Neuroscience.* 2014; 269C:102–111. [PubMed: 24657457]
- Gavin DP, Sharma RP, Chase KA, Matrisciano F, Dong E, Guidotti A. Growth arrest and DNA-damage-inducible, beta (GADD45b)-mediated DNA demethylation in major psychosis. *Neuropsychopharmacology.* 2012; 37:531–542. [PubMed: 22048458]
- Graff J, Koshibu K, Jouvenceau A, Dutar P, Mansuy IM. Protein phosphatase 1-dependent transcriptional programs for long-term memory and plasticity. *Learn Mem.* 2010; 17:355–363. [PubMed: 20592054]
- Hoffman DR, Boettcher JA, Diersen-Schade DA. Toward optimizing vision and cognition in term infants by dietary docosahexaenoic and arachidonic acid supplementation: a review of randomized controlled trials. *Prostaglandins Leukot Essent Fatty Acids.* 2009; 81:151–158. [PubMed: 19505812]

- Iafrati J, Orejarena MJ, Lassalle O, Bouamrane L, Chavis P. Reelin, an extracellular matrix protein linked to early onset psychiatric diseases, drives postnatal development of the prefrontal cortex via GluN2B-NMDARs and the mTOR pathway. *Mol Psychiatry*. 2014; 19:527.
- Inano K, Suetake I, Ueda T, Miyake Y, Nakamura M, Okada M, Tajima S. Maintenance-type DNA methyltransferase is highly expressed in postmitotic neurons and localized in the cytoplasmic compartment. *J Biochem*. 2000; 128:315–321. [PubMed: 10920268]
- Innis SM. Perinatal biochemistry and physiology of long-chain polyunsaturated fatty acids. *J Pediatrics*. 2003; 143:1–8.
- Jin B, Robertson KD. DNA methyltransferases, DNA damage repair, and cancer. *Adv Exp Med Biol*. 2013; 754:3–29. [PubMed: 22956494]
- Kitajka K, Sinclair AJ, Weisinger RS, Weisinger HS, Mathai M, Jayasooriya AP, Halver JE, Puskas LG. Effects of dietary omega-3 polyunsaturated fatty acids on brain gene expression. *Proc Natl Acad Sci U S A*. 2004; 101:10931–10936. [PubMed: 15263092]
- Koshibu K, Graff J, Mansuy IM. Nuclear protein phosphatase-1: an epigenetic regulator of fear memory and amygdala long-term potentiation. *Neuroscience*. 2011; 173:30–36. [PubMed: 21093547]
- Lee J, Park S, Lee JY, Yeo YK, Kim JS, Lim J. Improved spatial learning and memory by perilla diet is correlated with immunoreactivities to neurofilament and alpha-synuclein in hilus of dentate gyrus. *Proteome Sci*. 2012; 10:72. [PubMed: 23216756]
- Lemberger T, Parkitna JR, Chai M, Schutz G, Engblom D. CREB has a context-dependent role in activity-regulated transcription and maintains neuronal cholesterol homeostasis. *FASEB J*. 2008; 22:2872–2879. [PubMed: 18424767]
- Lu R, Wang X, Chen ZF, Sun DF, Tian XQ, Fang JY. Inhibition of the extracellular signal-regulated kinase/mitogen-activated protein kinase pathway decreases DNA methylation in colon cancer cells. *J Biol Chem*. 2007; 282:12249–12259. [PubMed: 17307743]
- Lubin FD, Gupta S, Parrish RR, Grissom NM, Davis RL. Epigenetic mechanisms: critical contributors to long-term memory formation. *Neuroscientist*. 2011; 17:616–632. [PubMed: 22130638]
- Ma DK, Jang MH, Guo JU, Kitabatake Y, Chang ML, Pow-Anpongkul N, Flavell RA, Lu B, Ming GL, Song H. Neuronal activity-induced Gadd45b promotes epigenetic DNA demethylation and adult neurogenesis. *Science*. 2009; 323:1074–1077. [PubMed: 19119186]
- Mehedint MG, Niculescu MD, Craciunescu CN, Zeisel SH. Choline deficiency alters global histone methylation and epigenetic marking at the Re1 site of the calbindin 1 gene. *FASEB J*. 2010; 24:184–195. [PubMed: 19752176]
- Miller CA, Sweatt JD. Covalent modification of DNA regulates memory formation. *Neuron*. 2007; 53:857–869. [PubMed: 17359920]
- Mu J, Krafft PR, Zhang JH. Hyperbaric oxygen therapy promotes neurogenesis: where do we stand? *Med Gas Res*. 2011; 1:14. [PubMed: 22146131]
- Mu J, Ostrowski RP, Soejima Y, Rolland WB, Krafft PR, Tang J, Zhang JH. Delayed hyperbaric oxygen therapy induces cell proliferation through stabilization of cAMP responsive element binding protein in the rat model of MCAo-induced ischemic brain injury. *Neurobiol Dis*. 2013; 51:133–143. [PubMed: 23146993]
- Na ES, Nelson ED, Kavalali ET, Monteggia LM. The impact of MeCP2 loss- orgain-of-function on synaptic plasticity. *Neuropsychopharmacology*. 2013; 38:212–219. [PubMed: 22781840]
- Niculescu MD, Lupu DS, Craciunescu CN. Maternal alpha-linolenic acid availability during gestation and lactation alters the postnatal hippocampal development in the mouse offspring. *Int J Dev Neurosci*. 2011; 29:795–802. [PubMed: 21964326]
- Niculescu MD, Lupu DS, Craciunescu CN. Perinatal manipulation of alphaslinolenic acid intake induces epigenetic changes in maternal and offspring livers. *FASEB J*. 2013; 27:350–358. [PubMed: 22997227]
- Nishioka C, Ikezoe T, Takeshita A, Yang J, Tasaka T, Yang Y, Kuwayama Y, Komatsu N, Togitani K, Koeffler HP, Taguchi H. ZD6474 induces growth arrest and apoptosis of human leukemia cells, which is enhanced by concomitant use of a novel MEK inhibitor, AZD6244. *Leukemia*. 2007; 21:1308–1310. [PubMed: 17361222]

- Rahman A, Khan KM, Al-Khaledi G, Khan I, Al-Shemary T. Over activation of hippocampal serine/threonine protein phosphatases PP1 and PP2A is involved in lead-induced deficits in learning and memory in young rats. *Neurotoxicology*. 2012; 33:370–383. [PubMed: 22387731]
- Ren J, Chung SH. Anti-inflammatory effect of alpha-linolenic acid and its mode of action through the inhibition of nitric oxide production and inducible nitric oxide synthase gene expression via NF-kappaB and mitogen-activated protein kinase pathways. *J Agric Food Chem*. 2007; 55:5073–5080. [PubMed: 17542608]
- Saadat M, Kakinoki Y, Mizuno Y, Kikuchi K, Yoshida MC. Chromosomal localization of human, rat, and mouse protein phosphatase type 1 beta catalytic subunit genes (PPP1CB) by fluorescence in situ hybridization. *Jpn J Genet*. 1994; 69:697–700. [PubMed: 7857673]
- Schule B, Armstrong DD, Vogel H, Oviedo A, Francke U. Severe congenital encephalopathy caused by MECP2 null mutations in males: central hypoxia and reduced neuronal dendritic structure. *Clin Genet*. 2008; 74:116–126. [PubMed: 18477000]
- Slack A, Pinard M, Araujo FD, Szyf M. A novel regulatory element in the dnmt1 gene that responds to co-activation by Rb and c-Jun. *Gene*. 2001; 268:87–96. [PubMed: 11368904]
- Stranahan AM, Haberman RP, Gallagher M. Cognitive decline is associated with reduced reelin expression in the entorhinal cortex of aged rats. *Cereb Cortex*. 2011; 21:392–400. [PubMed: 20538740]
- Sultan FA, Day JJ. Epigenetic mechanisms in memory and synaptic function. *Epigenomics*. 2011; 3:157–181. [PubMed: 22122279]
- Surette ME. Dietary omega-3 PUFA and health: stearidonic acid-containing seed oils as effective and sustainable alternatives to traditional marine oils. *Mol Nutr Food Res*. 2013
- van Groen T, Miettinen P, Kadish I. Axonal tract tracing for delineating interacting brain regions: implications for Alzheimer's disease-associated memory. *Future Neurol*. 2014; 9:89–98. [PubMed: 24678267]
- Veyrac A, Besnard A, Caboche J, Davis S, Laroche S. The transcription factor Zif268/Egr1, brain plasticity, and memory. *Prog Mol Biol Transl Sci*. 2014; 122:89–129. [PubMed: 24484699]
- Vousden DA, Epp J, Okuno H, Nieman BJ, Van Eede M, Dazai J, Ragan T, Bito H, Frankland PW, Lerch JP, Henkelman RM. Whole-brain mapping of behaviourally induced neural activation in mice. *Brain Struct Funct*. 2014
- Watanabe D, Uchiyama K, Hanaoka K. Transition of mouse de novo methyltransferases expression from Dnmt3b to Dnmt3a during neural progenitor cell development. *Neuroscience*. 2006; 142:727–737. [PubMed: 16973295]
- Winding C, Sun Y, Hoger H, Bubna-Littitz H, Pollak A, Schmidt P, Lubec G. Serine/threonine-protein phosphatase 1 alpha levels are paralleling olfactory memory formation in the CD1 mouse. *Electrophoresis*. 2011; 32:1675–1683. [PubMed: 21647921]
- Wood MA, Kaplan MP, Park A, Blanchard EJ, Oliveira AM, Lombardi TL, Abel T. Transgenic mice expressing a truncated form of CREB-binding protein (CBP) exhibit deficits in hippocampal synaptic plasticity and memory storage. *Learn Mem*. 2005; 12:111–119. [PubMed: 15805310]
- Zhubi A, Chen Y, Dong E, Cook EH, Guidotti A, Grayson DR. Increased binding of MeCP2 to the GAD1 and RELN promoters may be mediated by an enrichment of 5-hmC in autism spectrum disorder (ASD) cerebellum. *Transl Psychiatry*. 2014; 4:e349. [PubMed: 24448211]

Abbreviations

ALA	α -linolenic acid
P19	postnatal day 19
DHA	docosahexaenoic acid
DPA	docosapentaenoic acid
EPA	eicosapentaenoic acid


LA linoleic acid

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Fig. 1.**

Gene expression assessment. At the end of lactation (P19), gene expression for DNA methylation regulatory genes and memory-associate genes was measured in male offspring brains, as described in Section 2. (A) The expression of eight genes involved in memory and brain development is indicated for each treatment group (whole P19 male brains). Columns marked with different letters denote statistical significance as assessed by Tukey HSD test. The following p values, adjusted against 5% FDR were used: *Reelin* $p < 0.008$, *Ppp1cb* $p < 0.03$, and *Ppp1cc* $p < 0.008$. Error bars indicate standard error (SE). (B) The expression of four genes involved in epigenetic regulation was measured for each treatment group. Columns marked with different letters denote statistical significance as assessed by Tukey HSD test. The following p values, adjusted against 5% FDR were used: *Dnmt1* $p < 0.03$, *Dnmt3a* $p < 0.008$, and *Mecp2* $p < 0.008$. Error bars indicate standard error (SE).




Fig. 2. Correlations between *Mecp2*, *Reelin*, and *Ppp1cc* gene expression. CAST analysis followed by linear regression analysis (all samples) indicated that *Mecp2* expression was highly correlated with the expression of *Reelin* (panel A) and *Ppp1cc* (panel B). The inset at the top shows the cluster identified by CAST analysis. R^2 indicates the coefficient of determination for linear regression.




Fig. 3. Correlations between *Gadd45b*, *Bdnf*, and *Creb1* gene expression. CAST analysis followed by linear regression analysis (all samples) indicated that *Gadd45b* expression was highly correlated with the expression of *Bdnf* (panel A) and *Creb1* (panel B). The inset at the top shows the cluster identified by CAST analysis. R^2 indicates the coefficient of determination for linear regression.




Fig. 4. *Dnmt1* expression is correlated with *Egr1* expression. CAST analysis followed by linear regression analysis (all samples) indicated that *Dnmt1* expression was highly correlated with the expression of *Egr1*. The inset at the top shows the cluster identified by CAST analysis. R^2 indicates the coefficient of determination for linear regression.




Fig. 5. ALA availability alters *Mecp2* promoter DNA methylation but not its intron I methylation. DNA was extracted from P19 whole brains, and subject of bisulfite pyrosequencing. *Mecp2* DNA methylation was lower in the DD group, when compared to either the CC or the CS groups, indicating the overall ALA deficiency during gestation and lactation could trigger epigenetic alterations. Horizontal bars with asterisk above indicate statistical significance ($p < 0.05$) between the respective two groups, as assessed by the Wilcoxon Each Pair test against adjusted p values calculated by the Kruskal–Wallis test.

Table 1

Assay information for bisulfite pyrosequencing.

Gene & position	Assay	Sequence to analyze	Number of CpGs
Mecp2 promoter	<i>In-house designed</i>	F: AGTTTGGGTTTTATAATTAATGAAGGG R[5-Biotin]: ACCTTAACCATCCCACCTCACAATCTC S: AGGTGTAGTAGTATATAGG <i>Sequence to analyze:</i> TTGGTCGGGAGGGCGGGCGCGACGTTTGTCTGTCGGGG	7
Mecp2intron1	Mm_Mecp2_01_PM	CGCGCGCAGCCCCAACTGGCGAAGCCCAGACGA	5
Reelin promoter	Mm_Rehn_01_PM	GGACCCGACAGGCGAGCTTCGCCGGACTCTGTATTTACGCGT	6
Egr1 promoter	Mm_Egr1_08_PM	CTCCACCCTGCGACCCGCTCCGGCATCGCGAGCGC	6
Arc exon1	Mm_Arc_02_PM	AACTTGACGGCTACGTGCCACCCGGCGACTCACAGCGC	5
Ppp1cb promoter	Mm_Ppp1cb_01_PM	CGAAACGCCCGGTGACTCGTAGGTGAGAACGCCG	7
Ppp1cc intron1	Mm_Ppp1cc_03_PM	CGGCGCGCGGGTGGGTGGCGGGGCAGGCCCGCCGGT	7
Dnmt3a promoter	Mm_Dnmt3a_04_PM	CGCGTCGCCACCCACAGCCAGGTGCCCGGT	5
Dnmt1 promoter	Mm_Dnmt1_01_PM	GCGCATGCGCGCAACGGA	4

F: forward primer; R, reverse primer; S, sequencing primer. CpG denotes a cytosine followed by a Guanine (CpG site).