

RESEARCH ARTICLE

Open Access



α -Melanocyte-stimulating-hormone (α -MSH) modulates human chondrocyte activation induced by proinflammatory cytokines

Franco Capsoni^{1*}, Anna Maria Ongari¹, Caterina Lonati², Riccardo Accetta³, Stefano Gatti² and Anna Catania²

Abstract

Background: Alpha-melanocyte-stimulating-hormone (α -MSH) has marked anti-inflammatory potential. Proinflammatory cytokines are critical mediators of the disturbed cartilage homeostasis in osteoarthritis, inhibiting anabolic activities and increasing catabolic activities in chondrocytes. Since human chondrocytes express α -MSH receptors, we evaluated the role of the peptide in modulating chondrocyte production of pro-inflammatory cytokines, matrix metalloproteinases (MMPs), tissue inhibitors of MMPs (TIMPs), inducible nitric oxide synthase (iNOS) and nitric oxide (NO) in response to interleukin-1 β (IL-1 β) and tumor necrosis factor- α (TNF- α).

Methods: Human articular chondrocytes were obtained from osteoarthritic joint cartilage from subjects undergoing hip routine arthroplasty procedures. The cells were cultured with or without α -MSH in the presence of IL-1 β or TNF- α . Cell-free supernatants were collected and cells immediately lysed for RNA purification. Expression of cytokines, MMPs, TIMPs, iNOS was determined by Reverse Transcription Real-time Polymerase Chain Reaction and enzyme-linked immunosorbent assay. Griess reaction was used for NO quantification.

Results: Gene expression and secretion of IL-6, IL-8, MMP-3, MMP-13 were significantly increased in IL-1 β or TNF- α -stimulated chondrocytes; α -MSH did not modify the release of IL-6 or IL-8 while the peptide significantly reduced their gene expression on TNF- α -stimulated cells. A significant inhibition of MMP3 gene expression and secretion from IL-1 β or TNF α -stimulated chondrocytes was induced by α -MSH. On the other hand, α -MSH did not modify the release of MMP-13 by cytokine-stimulated chondrocyte but significantly decreased gene expression of the molecule on TNF- α -stimulated cells. Detectable amount of TIMP-3 and TIMP-4 were present in the supernatants of resting chondrocytes and a significant increase of TIMP-3 gene expression and release was induced by α -MSH on unstimulated cells. TIMP-3 secretion and gene expression were significantly increased in IL-1 β -stimulated chondrocytes and α -MSH down-regulated gene expression but not secretion of the molecule. TIMP-4 gene expression (but not secretion) was moderately induced in IL-1 β -stimulated chondrocytes with a down-regulation exerted by α -MSH. IL-1 β and TNF- α were potent stimuli for NO production and iNOS gene expression by chondrocytes; no inhibition was induced by α -MSH on cytokine-stimulated NO production, while the peptide significantly reduced gene expression of iNOS.

Conclusions: Our results underscore a potential anti-inflammatory and chondroprotective activity exerted by α -MSH, increasing TIMP-3 gene expression and release on resting cells and down- modulating TNF- α -induced activation of human chondrocytes. However, the discrepancy between the influences exerted by α -MSH on gene expression and protein release as well as the difference in the inhibitory pattern exerted by α -MSH in TNF- α - or IL-1 β -stimulated cells leave some uncertainty on the role of the peptide on chondrocyte modulation.

* Correspondence: franco.capsoni@unimi.it

¹Allergy, Clinical Immunology & Rheumatology Unit, Istituto Auxologico Italiano, IRCCS, University of Milan, Piazzale Brescia, 20 - 20149 Milano, Italy
Full list of author information is available at the end of the article

Background

Osteoarthritis (OA) is a chronic rheumatic disease characterized by cartilage degradation and loss, subchondral bone remodelling, and possible synovial inflammation. The precise cause of OA is unknown. A failure of chondrocytes to maintain the balance between synthesis and degradation of the cartilage extracellular matrix is considered a significant factor in the loss of cartilage [1].

The assumed mechanisms involved in chondrocyte dysregulation and/or apoptosis include mechanical stress, age-related functional changes and altered production of pro-inflammatory cytokines, predominantly interleukin-1 β (IL-1 β) and tumor necrosis factor- α (TNF- α), that induce production of oxygen radicals and proteinases such as matrix metalloproteinases (MMPs) and aggrecanases [2]. These observations suggested that anti-cytokine and anti-oxidant compounds could have chondroprotective effects providing novel therapeutic opportunities for OA treatment [3, 4].

α -Melanocyte-stimulating hormone (α -MSH) is an endogenous tridecapeptide that exerts multiple effects on host cells [5]. The natural peptide and its synthetic analogs inhibit inflammatory response in experimental models of acute and chronic disorders, including inflammatory bowel diseases, allergy, adjuvant arthritis, and sepsis [6–9]. α -MSH interacts with host cells through activation of four of the five recognized melanocortin receptors (MCR), specifically MCR 1, 3, 4 and 5 [9]. The anti-inflammatory action of the peptide depends primarily on inhibition of cytokine production by target cells, although other leukocyte functions, including reactive oxygen intermediate (ROI) production, nitric oxide (NO) generation and release of proteolytic enzymes, are likewise influenced [7–9].

In spite of substantial evidence suggesting a beneficial effect of melanocortin peptides in control of numerous inflammatory disorders and the observation that human chondrocytes express MCR [10], only recently the therapeutic potential of melanocortins in OA has been explored. Yoon et al. [11], using a human chondrosarcoma cell line (HTB-94) showed that α -MSH inhibited TNF- α -induced expression of MMP-13, through a decrease in mitogen-activated protein kinase (MAPK) p38 phosphorylation and subsequent activation of nuclear factor- κ B (NF- κ B). In human articular chondrocytes, α -MSH decreased IL-1 β and TNF- α mRNA but increased the secretion of TNF- α , IL-6 and Transforming Growth Factor- β 1 (TGF- β 1) [10]. In addition, in rodent chondrocytes, adrenocorticotropin (ACTH) treatment promoted development of the chondrocyte phenotype through activation of the MCR 3 [12]. Recently Kaneva et al. [13], using a human chondrocyte cell line (C-20/A4) demonstrated that melanocortin peptides strongly modulated chondrocyte function: the molecules inhibited

TNF- α -induced production of pro-inflammatory cytokines whilst increasing production of the anti-inflammatory and chondroprotective cytokine interleukin-10 (IL-10); decreased gene expression of MMPs; prevented chondrocyte apoptosis by inhibiting TNF- α induced caspase-3 and -7 activation.

The main obstacle in studies on human chondrocyte biology depends on the difficulty to obtain primary articular chondrocytes as these cells lose chondrocytic phenotype when expanded and cultured in monolayer. In the present study we evaluated the capacity of primary articular human chondrocytes to produce pro-inflammatory cytokines, MMPs, tissue inhibitors of MMPs (TIMPs) and NO in response to pro-inflammatory cytokines (IL-1 β and TNF- α). In parallel experiments the effects of α -MSH in modulation of mediator production were investigated.

Methods

Chondrocyte isolation and culture

Samples of adult human articular cartilage were harvested from subjects undergoing hip routine arthroplasty procedures at the Istituto Ortopedico Galeazzi IRCCS, Milan. This study did not undergo ethical approval since the cartilage was collected as waste material after receiving patients signed informed consent and the approval of the Institutional Review Board (IRCCS Istituto Ortopedico Galeazzi).

Minced cartilage fragments (2-3 mm²) were submitted to enzymatic digestion with 0.15 % collagenase type II (Worthington Biochemical Corporation, Lakewood, NJ, USA) overnight at 37 °C., as previously described [14]. Freshly isolated chondrocytes were plated for expansion at a density of 1x10⁴ cells/cm² and cultured with complete medium (CM) containing Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10 % fetal bovine serum (FBS, Lonza Group Ltd, Basel, CH), 1 mM sodium pyruvate, 100 mM HEPES buffer, 100 U/mL penicillin, 100 μ g/mL streptomycin, 0.29 mg/mL L-glutamine [15]. When the cells reached 80–90 % of confluence, they were detached with trypsin/EDTA (0.5 % trypsin/0.2 % EDTA and washed before performing the assays. Human chondrocytes were placed in 15 ml tissue culture tubes for 4 h with either phosphate buffered solution (PBS) (control) or α -MSH 10⁻⁶M. The tests employed the analogue Nle4,DPhe7- α -MSH (NDP-MSH), a non specific MC agonist that exerts similar effects relative to the natural α -MSH and is generally preferred for its greater chemical stability [16]. The cells were then cultured for 40 h in the presence or absence of 10 ng/mL human recombinant IL-1 β or 10 ng/mL human recombinant TNF- α [17]. In our preliminary experiments, these culture conditions had been found to be optimal to activate human chondrocytes. Where not otherwise specified, Sigma-Aldrich (Poole, Dorset, UK) products were used.

Cell-free supernatants were collected and stored at -20 °C and cells immediately lysed for RNA purification.

Cytokine quantification by ELISA

A qualitative commercial enzyme-linked immunosorbent assay test (Multi-analyte ELISArray kit (SABiosciences - QIAGEN, Maryland, USA) has been used to simultaneously profile the level of multiple cytokines (IL-1α, IL-1β, IL-2, IL-4, IL-6, IL-8, IL-10, IL-12, IL-17A, IFNγ, TNF-α, GM-CSF) in cell culture supernatants.

A cutoff of twice the absorbance value (read at 450 nm) of the negative control for every cytokine (A₄₅₀) was used and the results were reported as positive (A₄₅₀ ≥ specific cutoff) or negative (A₄₅₀ < specific cutoff).

Protein concentration of detectable cytokines IL-6, IL-8, MMP-3, MMP-13 (Boster Biological Technology, Fremont, CA, USA), TIMP-3 and TIMP-4 (R&D Systems, Minneapolis, MN, USA) was then determined in cell-free supernatants using quantitative commercially available ELISA kits.

Quantification of nitrites

The total NO production was measured using a commercial kit that involves the enzymatic conversion of nitrate to nitrite by the enzyme Nitrate Reductase (Enzo Life Sciences - Farmingdale, NY, USA). Briefly, 50 μl of the culture supernatant and 50 μl of Nitrate Reductase were mixed and incubated for 30 min at 37 °C in 96-well plates. One hundred microliters of the Griess reagent were added to the wells and incubated for further 10 min before reading the optical density at 540–570 nm. The nitrite concentration was calculated from a standard curve of sodium nitrate and expressed as μmole/l.

Table 1 Gene Symbols and Assay ID of the genes evaluated using Real-time RT-PCR

Gene name	Gene symbol	Assay ID
IL6	interleukin 6	Hs00174131_m1
IL8	interleukin 8	Hs00174103_m1
MMP3	matrix metalloproteinase 3	Hs00968305_m1
MMP13	matrix metalloproteinase 13	Hs00233992_m1
TIMP1	TIMP metalloproteinase inhibitor 1	Hs00171558_m1
TIMP3	TIMP metalloproteinase inhibitor 3	Hs00165949_m1
TIMP4	TIMP metalloproteinase inhibitor 4	Hs00162784_m1
ACTB	actin, beta	Hs99999903_m1
GAPDH	glyceraldehyde-3-phosphate dehydrogenase	Hs99999905_m1
RPLP0	ribosomal protein, large, P0	Hs99999902_m1
HPRT1	hypoxanthine phosphoribosyltransferase 1	Hs99999909_m1

Table 2 Effect of IL-1β and TNF-α on secretion of different molecules by human chondrocytes

Molecules in the Supernatants (ng/mL)	Stimulus		
	Medium	IL-1β	TNF-α
IL-6	3.0 ± 0.2	367 ± 4.9*	70 ± 3.7*
IL-8	0.0	781 ± 32*	120 ± 5.6*
MMP-3	2.0 ± 0.00	14500 ± 400*	8100 ± 50*
MMP-13	8.1 ± 0.5	71.0 ± 8.0*	35.3 ± 2.7*
TIMP-3	1.63 ± 0.12	2.71 ± 0.05°	1.23 ± 0.01 ns
TIMP-4	0.11 ± 0.01	0.12 ± 0.01 ns	0.12 ± 0.01 ns
NO (μM)	19.5 ± 3.5	102.8 ± 32.6*	95.8 ± 36.9*

Primary human chondrocytes were cultured for 40 h in medium or in the presence of 10 ng/mL IL-1β or TNF-α. The cell-free supernatants were collected and the concentration of interleukins, MMPs, TIMPs and NO were measured.

Data from 4 separate experiments (± S.E.M.) are shown

* p < 0.0001; ° p < 0.002; ns: not significant vs medium

RNA purification

Total RNA was isolated from chondrocyte lysates by anion exchange chromatography using RNeasy MinElute columns (RNeasy Mini Kit, Qiagen Inc., Hilden, Germany). Briefly, cell lysates were added (v/v) to 70 % ethanol and immediately transferred to an RNeasy column. DNase I treatment (Qiagen) was performed to remove genomic DNA contamination. After two washes in ethanol-based buffer (Buffer RPE, Qiagen), RNA were eluted in RNase-free water and then quantified by optical density measurement using Nanodrop ND-100 spectrophotometer (Nanodrop Technologies, Wilmington, DE). Each sample showed a 260/280 ratio between 1.8 and 2. RNA integrity was assessed by electrophoresis on denaturing agarose–formaldehyde gels.

Table 3 Effect of IL-1β and TNF-α on gene expression of different molecules in human chondrocytes

Relative gene expression for	Stimulus	
	IL-1β	TNF-α
IL-6	2767 ± 176*	680 ± 49*
IL-8	5090 ± 271*	1088 ± 78*
MMP-3	112 ± 10.0*	71 ± 6.5*
MMP-13	7.5 ± 2.0*	6.5 ± 0.5*
TIMP-3	3.4 ± 0.7°	1.0 ± 0.1 ns
TIMP-4	1.9 ± 0.2°	1.0 ± 0.05 ns
iNOS	1656 ± 100*	1663 ± 110*


Primary human chondrocytes were cultured for 40 h in medium or in the presence of 10 ng/mL IL-1β or TNF-α. RNA was purified and analyzed by Real-time RT-PCR. Gene expression data were calculated with the ΔCT method and using HPRT1 and GAPDH as reference genes. Relative expression was obtained comparing samples to untreated control whose gene expression was considered as baseline. Data from 4 separate experiments (± S.E.M.) are shown

* p < 0.0001; ° p < 0.02; ns: not significant vs untreated controls

Reverse Transcription Real-time Polymerase Chain Reaction (Real-time RT-PCR) analysis

Gene expression was evaluated by Real-Time Reverse Transcription-Polymerase Chain Reaction (RT-PCR) analysis. Two micrograms of total RNA were reverse transcribed to single stranded cDNA using SuperScript VILO cDNA Synthesis Kit (Life Technologies, Foster City, CA) according to the manufacturer's instructions. Real-time PCR based on TaqMan chemistry was performed with 20 ng of retrotranscribed RNA, 1X FastStart Universal Probe Master (Roche Diagnostics GmbH, Mannheim, Germany), and predesigned 900 nM primers and 250 nM probe mix (Taqman Gene Expression Assays, Life Technologies, (assay IDs are listed in Table 1) in a final volume of 10 μ l. Amplification reactions were carried out on an

ABI PRISM 7900HT sequence detection system (Applied Biosystems, Life Technologies) with the following thermal profile settings: an initial step of 10 min at 95°C to activate the FastStart Taq DNA polymerase, then 50 cycles of 95 °C for 10 s and 60 °C for 30 s. Three independent PCR amplification experiments were performed for each transcript. Fluorescence intensities were converted in threshold cycles (Ct) using ABI Prism SDS 2.3 software (Life Technologies); baseline and threshold were set by automatic analysis. Relative quantification of target gene expression was calculated with the Δ Ct method, using the average Ct across basal samples as calibrator for each gene. Analysis of reference genes stability, performed with the geNorm software, indicated HPRT1 and GAPDH as the most stable [18].



Statistical analysis

The data are expressed as the mean ± standard error of the mean (S.E.M.) from 4 determinations as indicated. Statistical analysis was done using Student’s *t* test for unpaired data, as appropriate. Differences were considered to be statistically significant at *P* < 0.05.


Results

Effect of pro-inflammatory cytokines on human chondrocytes

Of the twelve cytokines evaluated in the supernatants of IL-1β or TNF-α-stimulated chondrocytes with a qualitative ELISA test (IL-1α, IL-1β, IL-2, IL-4, IL-6, IL-8, IL-10, IL-12, IL-17α, IFNγ, TNF-α, GM-CSF), significant levels were observed only for IL-6 and IL-8 (not

shown). The protein concentration of these two cytokines is shown in Table 2. Gene expression of IL-6 and IL-8 was also significantly increased in IL-1β or TNF-α-stimulated chondrocytes (Table 3). Increased levels of MMP-3 and MMP-13 in the supernatants (Table 2) as well as enhanced gene expression (Table 3) were observed in cytokine-stimulated chondrocytes, with IL-1β being the most potent stimulus.

Low but detectable amounts of TIMP-4 (Table 2) were present in the supernatants of non-stimulated chondrocytes with no increased secretion after stimulation by IL-1β or TNF-α. TIMP-4 gene expression was moderately induced by IL-1β while there were no changes when the stimulus was TNF-α (Table 3). TIMP-3 was also present in the supernatants of resting



chondrocytes and its secretion (Table 2) as well as gene expression (Table 3) were significantly increased by IL-1 β but not by TNF- α stimulation. Both IL-1 β and TNF- α were potent stimuli for NO production (Table 2) and iNOS gene expression (Table 3).

Effect of α -MSH on human chondrocytes

Pre-treatment of chondrocytes with 10⁻⁶M α -MSH did not modify the release of IL-6 or IL-8 by IL-1 β (Fig. 1) or TNF- α (Fig. 2) stimulated chondrocytes. On the other hand, while only slightly decreased gene expression of IL-6 and IL-8 was observed when the cells were stimulated with IL-1 β (Fig. 1), the reduction became significant when the stimulus was TNF- α (Fig. 2)

A significant inhibition of MMP-3 gene expression and secretion from IL-1 β - or TNF- α -stimulated chondrocytes was induced by α -MSH (Fig. 3). On the other hand, α -MSH did not modify the release of MMP-13 by cytokine-stimulated chondrocyte but the peptide caused a decreased gene expression of the molecule that was significant for TNF- α -stimulated cells (Fig. 4)

α -MSH did not modify TIMPs secretion by cytokine-stimulated chondrocyte (Figs. 5 and 6) but the peptide down-regulated the IL-1 β -induced gene expression of TIMP-3 (Fig. 5) and TIMP-4 (Fig. 6). Of interest, a significant increase of TIMP-3 gene expression and release was induced by α -MSH (three fold increase) on unstimulated cells (Fig. 5).




Fig. 3 Effect of α -MSH on MMP-3 secretion (a) and gene expression (b) in IL-1 β or TNF- α -stimulated chondrocytes. Human primary chondrocytes were incubated for 4 h at 37 °C with 10⁻⁶M α -MSH and further cultured for 40 h in the presence or absence of 10 ng/mL IL-1 β or TNF- α . MMP-3 concentration in the supernatans (a) and gene expression (b) were then measured. Data from 4 separate experiments (\pm S.E.M.) are shown. * p < 0.0001 vs Medium; § p < 0.001 vs IL-1 β ; ° p < 0.01 vs TNF- α ; oo p < 0.001 vs TNF- α




Fig. 4 Effect of α -MSH on MMP-13 secretion (a) and gene expression (b) in IL-1 β or TNF- α -stimulated chondrocytes. Human primary chondrocytes were incubated for 4 h at 37 °C with 10⁻⁶M α -MSH and further cultured for 40 h in the presence or absence of 10 ng/mL IL-1 β or TNF- α . MMP-13 concentration in the supernatans (a) and gene expression (b) were then measured. Data from 4 separate experiments (\pm S.E.M.) are shown. * p < 0.0001 vs Medium; ^o p < 0.05 vs TNF- α

No inhibition was induced by α -MSH on cytokine-stimulated NO production but the peptide was able to reduce gene expression of iNOS2, the enzyme involved in inducible synthesis of nitric oxide (Fig. 7).

Discussion

The data show that stimulation of human primary chondrocytes with IL-1 β or TNF- α induces an inflammatory phenotype. Both cytokines induced gene expression and secretion of IL-6, IL-8, MMP-3 and MMP-13 associated with a significant production of NO. These results confirm a significant chondrocyte responsiveness to exogenous pro-inflammatory stimuli with amplification of

the inflammatory response and possible local cartilage degradation. Under our experimental conditions the anti-inflammatory peptide α -MSH did not modify release or gene expression of IL-6, IL-8, MMP-13, MMP-3 in resting chondrocytes. These results are not consistent with previous observations by Grässel et al. [10] who, using different chondrocyte culture conditions (micromass pellets) reported an α -MSH-induced down-regulation of IL-1 β and TNF- α gene expression and an increased secretion of IL-6 and TNF- α in human chondrocytes. Of interest, we observed that α -MSH was able to induce gene expression and secretion of TIMP-3 in resting cells. These results suggest a




Fig. 5 Effect of α -MSH on TIMP-3 secretion (a) and gene expression (b) in IL-1 β or TNF- α -stimulated chondrocytes. Human primary chondrocytes were incubated for 4 h at 37 °C with 10⁻⁶M α -MSH and further cultured for 40 h in the presence or absence of 10 ng/mL IL-1 β or TNF- α . TIMP-3 concentration in the supernatans (a) and gene expression (b) were then measured. Data from 4 separate experiments (\pm S.E.M.) are shown. * P < 0.002 vs Medium; ** p < 0.001 vs Medium; $^{\circ}p$ < 0.02 vs Medium; $^{\circ\circ}p$ < 0.002 vs Medium; $^{\S}p$ < 0.05 vs IL-1 β

potential role for α -MSH in the maintenance of cartilage homeostasis. Indeed the anticatabolic influence of TIMP-3 as an inhibitor of molecules implicated in cartilage degradation, including MMP and aggrecanase-2, is well recognized [19].

When the capacity of α -MSH to modulate IL-1 β or TNF- α -induced chondrocyte activation was evaluated, we observed that the peptide did not exert anti-oxidative activity on human chondrocytes as it did not modify cytokine-induced NO production. On the other hand, α -MSH significantly inhibited MMP-3 gene expression and secretion induced by both cytokines further suggesting an anticatabolic role of the peptide. In TNF- α -stimulated chondrocytes the peptide significantly reduced

gene expression of IL-6, IL-8, and MMP-13. Conversely, there was no substantial effect in IL-1 β -activated cells. These results are consistent with those of Yoon et al. [11] who showed that α -MSH inhibited TNF- α -induced MMP-13 gene expression in a human chondrosarcoma cell line.

The present data are partly consistent with those of Kaneva et al. [13] who reported an inhibitory effect of α -MSH on pro-inflammatory cytokine release (IL-1, IL-6, IL-8) and on MMP-1, MMP-3, MMP-13 gene expression in a TNF- α -stimulated human chondrocyte cell line.

Collectively, these results underscore a potential anti-inflammatory and chondroprotective activity exerted by α -MSH in TNF- α -stimulated human chondrocytes.




Fig. 6 Effect of α -MSH on TIMP-4 secretion (a) and gene expression (b) in IL-1 β or TNF- α -stimulated chondrocytes. Human primary chondrocytes were incubated for 4 h at 37 °C with 10^{-6} M α -MSH and further cultured for 40 h in the presence or absence of 10 ng/mL IL-1 β or TNF- α . TIMP-4 concentration in the supernatans (a) and gene expression (b) were then measured. Data from 4 separate experiments (\pm S.E.M.) are shown. * $p < 0.02$ vs Medium; ° $p < 0.05$ vs IL-1 β

In our experimental conditions, the α -MSH-induced inhibition of cytokine-stimulated gene expression of IL-6, IL-8 and MMP-13, and the down-regulation of IL-1 β -induced increase in TIMP-3 and TIMP-4 gene expression were not accompanied by a parallel reduction in protein release. A similar discordance between gene expression and protein secretion was previously reported by Grässel et al. [10] using α -MSH-treated micromass pellet cultures.

Although the discrepancy between influences on gene expression and protein release does not have a definite explanation, it may be secondary to an incomplete or

late inhibition of gene expression by α -MSH compared to the activation induced by pro-inflammatory cytokines *in vitro*.

The biological significance of the difference in the inhibitory pattern exerted by α -MSH in TNF- α -stimulated chondrocytes (inhibition of IL-6, IL-8 and MMPs gene expression) relative to IL-1 β -stimulated cells (inhibition of TIMP-3 and TIMP-4 gene expression) is likewise unclear. Both IL-1 β and TNF- α exert their biological responses mainly through the activation of nuclear factor- κ B (NF- κ B) signalling pathway and NF- κ B suppression is considered the key molecular mechanism of the anti-




Fig. 7 Effect of α -MSH on NO secretion and iNOS gene expression in IL-1 β or TNF- α -stimulated chondrocytes. Human primary chondrocytes were incubated for 4 h at 37 °C with 10⁻⁶M α -MSH and further cultured for 40 h in the presence or absence of 10 ng/mL IL-1 β or TNF- α . NO concentration in the supernatants (a) and iNOS gene expression (b) were then measured. Data from 4 separate experiments (\pm S.E.M.) are shown. * p < 0.0001 vs Medium; § p < 0.005 vs IL-1 β ; ° p < 0.005 vs TNF- α

inflammatory effect of α -MSH [9]. It is possible that different concentrations of the peptide are required to inhibit the activation induced by different cytokines in vitro. An interference exerted by α -MSH on membrane cytokine receptors could be also considered but, while it has been demonstrated that α -MSH potently and selectively reduce membrane binding of IL-1 β to T-cell clones, we have no knowledge of similar data for TNF- α .

Conclusions

The present results indicate that α -MSH exerts chondroprotective activity suggesting a potential role of α -MSH in

the maintenance of cartilage homeostasis although the role of the peptide in the control of chondrocyte activation remains to be defined.

Competing interests

The authors declare that they have no competing interests.

Authors' contribution

FC conceived and designed the study; performed statistical analysis and interpretation of data; drafted the manuscript; AMO provided study material; conducted the experimental assays; acquired the data and helped in statistical analysis; CL carried out the molecular genetic studies and helped to draft the manuscript; RA selected and recruited patients; SG and AC participated in study design and critically revised the manuscript for important intellectual content. All author read and approved the final manuscript.

Acknowledgments

This work was supported by the following research funds: 'Ricerca Corrente 2009', Istituto Ortopedico Galeazzi, Istituto Di Ricovero e Cura a Carattere Scientifico, Milan, Italy, and PUR 2008, University of Milan, Italy.

Author details

¹Allergy, Clinical Immunology & Rheumatology Unit, Istituto Auxologico Italiano, IRCCS, University of Milan, Piazzale Brescia, 20 - 20149 Milano, Italy. ²Center for Preclinical Investigation, Fondazione IRCCS Ospedale Maggiore Policlinico, Mangiagalli e Regina Elena, Milan, Italy. ³Traumatology and First Aid Unit, Istituto Ortopedico Galeazzi, IRCCS, Milan, Italy.

Received: 24 February 2015 Accepted: 8 June 2015

Published online: 21 June 2015

References

1. Goldring MB. Update on the biology of the chondrocyte and new approaches to treating cartilage diseases. *Best Pract Res Clin Rheumatol*. 2006;20:1003–25.
2. Aigner T, Sachse A, Gebhard PM, Roach HI. Osteoarthritis: Pathobiology-targets and ways for therapeutic intervention. *Adv Drug Deliv Rev*. 2006;58:128–49.
3. Kapoor M, Martel-Pelletier J, Lajeunesse D, Pelletier JP, Fahmi H. Role of proinflammatory cytokines in the pathophysiology of osteoarthritis. *Nat Rev Rheumatol*. 2011;7:33–42.
4. Martel-Pelletier J, Wildi LM, Pelletier JP. Future therapeutics for osteoarthritis. *Bone*. 2012;51:297–311.
5. Catania A. Neuroprotective actions of melanocortins: a therapeutic opportunity. *Trends Neurosci*. 2008;31:353–60.
6. Luger TA, Brzoska T. α -MSH related peptides: a new class of anti-inflammatory and immunomodulating drugs. *Ann Rheum Dis*. 2007;66(Suppl III):iii52–5.
7. Catania A. The melanocortin system in leukocyte biology. *J Leukoc Biol*. 2007;81:383–92.
8. Getting SJ. Targeting melanocortin receptors as potential novel therapeutics. *Pharmacol Ther*. 2006;111:1–15.
9. Brzoska T, Luger TA, Maaser C, Abels C, Böhm M. α -Melanocyte-Stimulating Hormone and related tripeptides: biochemistry, anti-inflammatory and protective effects in vitro and in vivo, and future perspectives for the treatment of immune-mediated inflammatory diseases. *Endocr Rev*. 2008;29:581–602.
10. Grässel S, Opolka A, Anders S, Straub RH, Grifka J, Luger TA, et al. The melanocortin system in articular chondrocytes melanocortin receptors, pro-opiomelanocortin, precursor proteases, and a regulatory effect of α -melanocyte-stimulating hormone on proinflammatory cytokines and extracellular matrix components. *Arthritis Rheum*. 2009;60:3017–27.
11. Yoon SW, Chun JS, Sung MH, Kim JY, Poo H. α -MSH inhibits TNF- α -induced matrix metalloproteinase-13 expression by modulating p38 kinase and nuclear factor κ B signalling in human chondrosarcoma HTB-94 cells. *Osteoarthritis Cartilage*. 2008;16:115–24.
12. Evans JF, Shen CL, Pollack S, Aloia JF, Yeh JK. Adrenocorticotropin Evokes Transient Elevations in Intracellular Free Calcium ($[Ca^{2+}]_i$) and Increases Basal $[Ca^{2+}]_i$ in Resting Chondrocytes through a Phospholipase C-Dependent Mechanism. *Endocrinology*. 2005;146:3123–32.
13. Kaneva MK, Kerrigan MJP, Greco P, Curley GP, Locke IC, Gettino SJ. Chondroprotective and anti-inflammatory role of melanocortin peptides in TNF- α activated human C-20/A4 chondrocytes. *Br J Pharmacol*. 2012;167:67–79.
14. Jakob M, Demarteau O, Schafer D, Stumm M, Heberer M, Martin I. Enzymatic digestion of adult human articular cartilage yields a small fraction of the total available cells. *Connect Tissue Res*. 2003;44:173–80.
15. Barbero A, Ploegert S, Heberer M, Martin I. Plasticity of clonal populations of dedifferentiated adult human articular chondrocytes. *Arthritis Rheum*. 2003;48:1315–25.
16. Sawyer TK, Sanfilippo PJ, Hruby VJ, Engel MH, Heward CB, Burnett JB, et al. 4-Norleucine, 7-D-phenylalanine- α -melanocyte-stimulating hormone: a highly potent α -melanotropin with ultralong biological activity. *Proc Natl Acad Sci U S A*. 1980;77:5754–8.
17. Terkeltaub R, Yang B, Lotz M, Liu-Bryan R. Chondrocyte AMP-activated protein kinase activity suppresses matrix degradation responses to proinflammatory cytokines interleukin-1 β and tumor necrosis factor α . *Arthritis Rheum*. 2011;63:1928–37.
18. Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, et al. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol*. 2002;3:research0034.
19. Sahebjam S, Khokha R, Mort JS. Increased collagen and aggrecan degradation with age in the joints of TIMP-3(–/–) mice. *Arthritis Rheum*. 2007;56:905–9.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

