

## Basic Study

## Altered distribution of regulatory lymphocytes by oral administration of soy-extracts exerts a hepatoprotective effect alleviating immune mediated liver injury, non-alcoholic steatohepatitis and insulin resistance

Tawfik Khoury, Ami Ben Ya'acov, Yehudit Shabat, Lidya Zolotarovya, Ram Snir, Yaron Ilan

Tawfik Khoury, Ami Ben Ya'acov, Yehudit Shabat, Lidya Zolotarovya, Ram Snir, Yaron Ilan, Gastroenterology and Liver Units, Department of Medicine, Hadassah-Hebrew University Medical Center, Jerusalem IL-91120, Israel

**Author contributions:** Khoury T and Ben Ya'acov A contributed equally to this work; Khoury T, Ben Ya'acov A, Shabat Y and Zolotarovya L conducted the experiments and summarized the data; Khoury T, Ben Ya'acov A, Snir R and Ilan Y prepared the manuscript.

**Supported by Grants from The Roaman-Epstein Liver Research Foundation** (partly, to Ilan Y).

**Institutional animal care and use committee:** All procedures involving animals were reviewed and approved by the Institutional Animal Care and Use committee of the Hebrew University Hadassah Medical Center (IACUC protocol number: MD-13-13733-3).

**Conflict-of-interest:** The authors declare no conflicting interest.

**Data sharing:** No additional data are available.

**Open-Access:** This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Correspondence to:** Tawfik Khoury, MD, Gastroenterology and Liver Units, Department of Medicine, Hadassah-Hebrew University Medical Center, PO Box 12000, Jerusalem IL-91120, Israel. [tawfikkhoury1@hotmail.com](mailto:tawfikkhoury1@hotmail.com)  
Telephone: +972-2-6777816  
Fax: +972-2-6431021

Received: November 11, 2014  
Peer-review started: November 12, 2014  
First decision: January 8, 2015  
Revised: February 8, 2015  
Accepted: March 30, 2015  
Article in press: March 31, 2015  
Published online: June 28, 2015

### Abstract

**AIM:** To determine the immune-modulatory and the hepatoprotective effects of oral administration of two soy extracts in immune mediated liver injury and non-alcoholic steatohepatitis (NASH).

**METHODS:** Two soy extracts, M1 and OS, were orally administered to mice with concanavalin A (ConA) immune-mediated hepatitis, to high-fat diet (HFD) mice and to methionine and choline reduced diet combined with HFD mice. Animals were followed for disease and immune biomarkers.

**RESULTS:** Oral administration of OS and M1 had an additive effect in alleviating ConA hepatitis manifested by a decrease in alanine aminotransferase and aspartate aminotransferase serum levels. Oral administration of the OS and M1 soy derived fractions, ameliorated liver injury in the high fat diet model of NASH, manifested by a decrease in hepatic triglyceride levels, improvement in liver histology, decreased serum cholesterol and triglycerides and improved insulin resistance. In the methionine and choline reduced diet combined with the high fat diet model, we noted a decrease in hepatic triglycerides and improvement in blood glucose levels and liver histology. The effects were associated

with reduced serum tumor necrosis factor alpha and alteration of regulatory T cell distribution.

**CONCLUSION:** Oral administration of the combination of OS and M1 soy derived extracts exerted an adjuvant effect in the gut-immune system, altering the distribution of regulatory T cells, and alleviating immune mediated liver injury, hyperlipidemia and insulin resistance.

**Key words:** Non-alcoholic steatohepatitis; Fatty liver; Regulatory T cells; Soy; Type 2 diabetes; ConA hepatitis

© **The Author(s) 2015.** Published by Baishideng Publishing Group Inc. All rights reserved.

**Core tip:** Oral administration of the combination of OS and M1 soy derived extracts exerted an adjuvant effect in the gut-immune system, altering the distribution of regulatory T cells, and alleviating immune mediated liver injury, hyperlipidemia and insulin resistance. Oral administration of these extracts had an additive effect in alleviating concanavalin A hepatitis and ameliorated liver injury in the high fat diet model of non-alcoholic steatohepatitis, and in the methionine and choline reduced diet combined with the high fat diet model. The effects were associated with reduced serum tumor necrosis factor alpha and alteration of regulatory T cell distribution.

Khoury T, Ben Ya'acov A, Shabat Y, Zolotarovya L, Snir R, Ilan Y. Altered distribution of regulatory lymphocytes by oral administration of soy-extracts exerts a hepatoprotective effect alleviating immune mediated liver injury, non-alcoholic steatohepatitis and insulin resistance. *World J Gastroenterol* 2015; 21(24): 7443-7456 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v21/i24/7443.htm> DOI: <http://dx.doi.org/10.3748/wjg.v21.i24.7443>

## INTRODUCTION

Non-alcoholic fatty liver disease (NAFLD) is increasingly becoming a major health burden<sup>[1,2]</sup>. Non-alcoholic steatohepatitis (NASH) is characterized by fatty infiltration of the liver with hepatocytes ballooning and lobular inflammation, and is associated with increased risk of cirrhosis and hepatocellular carcinoma<sup>[3]</sup>. Both NAFLD and NASH are associated with obesity and diabetes<sup>[4]</sup>.

Liver steatosis is associated with an increase in free circulating fatty acids<sup>[4-7]</sup>. Lipids normally undergo metabolism in the liver by  $\beta$ -oxidation. The metabolism of excess lipids via their incorporation into the very low density lipoprotein mechanism is down regulated in patients with NASH<sup>[4-7]</sup>. Accumulation of lipids in the liver is also associated with increased susceptibility to multiple oxidative factors. In addition, a reactive inflammatory process secondary to faulty lipid metabolism may lead to the secretion of cytokines

and adipokines including tumor necrosis factor alpha (TNF- $\alpha$ ). TNF- $\alpha$  further decreases the anti-oxidant capacity of the liver, promoting mitochondrial damage. Activation of the immune system and of hepatic stellate cells lead to liver inflammation and liver fibrosis in hepatic steatosis<sup>[4-7]</sup>. The increase in visceral adipose tissue is also associated with an increased release of free fatty acids, inducing a low grade inflammation state by cytokine release. This state is associated induced insulin resistance by down regulating the insulin receptor substrate 1 (IRS-1)<sup>[8]</sup>.

In patients with NASH an increased secretion of pro-inflammatory cytokines from adipose tissue (e.g., TNF- $\alpha$ ) was described<sup>[9,10]</sup>. An increased level of macrophages which further exacerbate the pro-inflammatory state was shown. Regulatory T cells (Tregs) can suppress several immune mediated disorders via alteration of effector T cells and the cytokine paradigm. In animal models of NASH, promotion of Tregs was associated with alteration of macrophage function and alteration of the cytokine paradigm from a pro- to an anti-inflammatory one<sup>[11,12]</sup>. A similar phenomenon was suggested in humans<sup>[13]</sup>. These effects were associated with alleviation of liver steatosis and insulin resistance.

Soy is part of the regular diet in many countries. Soybean is a rich source of vegetable protein, complex carbohydrates, polyunsaturated fat, soluble sugars and phytoestrogens<sup>[14-17]</sup>. Some soy proteins are also associated with fatty acids, saponins, isoflavones and phospholipids. Soy derived products are safe and were suggested beneficial in immune mediated disorders<sup>[14-17]</sup>. Soy products were reported to decrease the oxidative stress and the subsequent pro-inflammatory cytokine secretion and lipid peroxidation in the MCD diet model of NASH<sup>[18]</sup>. Specific components of soy, including isoflavones, can increase the anti-oxidant capacity of hepatocytes and decrease oxidative stress<sup>[17]</sup>. They also have beneficial effect in the treatment of gastric cancer<sup>[19]</sup>. One of the components of soy,  $\beta$ -glucosylceramide (GC), was shown to exert an immune modulatory effect in various models<sup>[20-22]</sup>, further supporting the potential to use soy derived extracts as immune modulators. Epidemiological studies have shown that soy products might decrease the morbidity of NASH<sup>[23]</sup> as well as lower serum lipids in the blood and liver, reduce liver uptake of lipids, improve anti-oxidant ability and insulin resistance<sup>[24]</sup>.

The aim of the present study was to determine the immune-mediated hepatoprotective effect of soy extracts. We have tested several soy extracts in three mice models: the concanavalin A (ConA) immune-mediated hepatitis model, and in two models of NASH. Oral administration of the combination of OS and M1 soy derived extracts exerted an adjuvant effect in the gut-immune system, altered the distribution of Tregs, alleviated immune mediated liver injury, hyperlipidemia, insulin resistance and liver damage

in the high fat diet (HFD), and in the methionine and choline reduced (HFD/MCR) model.

## MATERIALS AND METHODS

### Animals

All mice were maintained in the animal core of the Hebrew University Hadassah Medical School (Jerusalem, Israel). The mice were given the diets described below (HFD, HFD/MCR) and had free access to water and were maintained in a 12-h light-dark cycle. All experiments were performed in accordance with the guidelines of the Hebrew University-Hadassah Institutional Committee for Care and Use of Laboratory Animals with the approval of the Ethics Committee.

### Oral administration of soy extracts

Two soy extracts were received from Solbar Israel (CHS), and studied in the below described animal models. OS-fraction, derived from the solvent extraction of soybeans into oil, and contains mainly tri- and di-glycerides, free fatty acids and phosphatides; MI-fraction which is derived from aqueous-ethanol extraction left after the solvent extraction, and contains mainly isoflavones, sugars (oligo-, di-, mono-), and lipids (including phosphatides, phytosterols, saponins).

### Induction of ConA hepatitis

ConA (MP Biomedicals, United States) was dissolved in 50 mmol/L Tris pH 7, 150 mmol/L NaCl, 4 mmol/L CaCl<sub>2</sub> and was injected into the tail vein at a dose of 500 µg/mouse (15 mg/kg).

### Animal models of NASH

Two models of NASH were studied. Model A: Harlan, TD88137, in this diet 42% of the caloric intake is derived from fat and it resembles a severe form of the human fatty diet. The mice given this diet exhibited features of fatty liver with characteristics of NASH, weight gain, hyperlipidemia and insulin resistance. Model B: HFD with methionine and choline reduced diet (Harlan, TD10810), this diet outpointed the inflammatory elements of NASH and oxidative stress. It's advantageous over the methionine and choline deficient diet due to less weight reduction<sup>[25]</sup>.

### Assessment of the effect of oral soy extracts on liver damage in ConA hepatitis model

**Experimental groups:** Eight groups were examined, 11-12 wk old male mice were purchased from Harlan Laboratories (Jerusalem, Israel). Four to six mice were treated in each group. To induce autoimmune hepatitis, all mice were injected iv (tail vein) with ConA. All mice were treated orally for three days with OS and M-1 soy compounds prior to ConA injections. Mice were sacrificed 14 h after the ConA injection. Mice in control Group A were treated with PBS. Group B was treated with GC (25 microgram/mouse), a

natural β-glycosphingolipid, (Avanti Polar lipids, AL, United States). Group C was treated with 0.35 µg of dexamethasone. Group D was administered with 3 µg of M1 soy fraction. Group E with 3 µg OS of soy fraction. Groups F, G and H were treated with different combinations of OS and M1 soy-derived extracts in 0.3, 3, and 30 µg doses, for each fraction respectively.

**Histological examination of the liver:** Paraffin-embedded liver sections were prepared from each mouse. The livers were cut into 4-5 µm thin slices and stained with hematoxylin-eosin (HE). Slides were scored to assess the extent of liver damage as described<sup>[26]</sup>.

**Liver enzymes:** Serum was obtained from individual mice. The serum aspartate aminotransferase (AST) and alanine aminotransferase (ALT) levels were determined with an automatic analyzer.

### Assessment of the effect of oral soy extracts on liver damage in high fat diet model of NASH

**Experimental groups:** Six groups with five mice each, of 6-7 wk old male mice were obtained from Harlan Laboratories (Jerusalem, Israel). Mice received HFD diet in which 42% of calories are derived from fat, administered orally 3 times a week for 3 mo. Group A was treated with 30 µL of PBS. Group B with 3 µg of OS soy fraction. Group C with 25 µg GC. Group D with 3 µg of M1 soy fraction. Groups E and F were administered with combinations of soy products in doses of 3 µg and 0.3 µg for each component respectively.

### Assessment of the effect of oral soy extracts on liver damage in the HFD/MCR model of NASH

For the HFD/MCR model, four groups were included; each group contained 6 mice purchased from Harlan Laboratories (Jerusalem, Israel). Mice under this diet regimen were administered orally 3 times a week for 3 mo. Group A was treated with 30 µL of PBS. Group B given 0.3 µg of OS and 0.6 µg of M1. Group C was treated with 3 µg OS and 6 µg M1. Group D was treated with 9 µg OS and 18 µg M1.

### Measurement of lipid accumulation in the liver and plasma triglycerides

Plasma triglycerides (TGs) and total cholesterol were measured by the Reflovet Plus clinical chemistry analyzer (Roche Diagnostics, GmbH, Mannheim, Germany). TGs were extracted from aliquots of snap-frozen livers using a modification of the Folch method. Hepatic TG content was assayed spectrophotometrically using a GPO-Trinder kit (Sigma, Rehovot, Israel) and was normalized to the protein content in the homogenate. The HFD mice were weighed every 2 wk. Serum ALT, TGs and cholesterol were measured every 2 wk by the Reflovet device

while HFD/MCR mice were monitored for body weight, serum ALT, and triglycerides, every 4 wk.

#### **Evaluation of insulin action and sensitivity**

Insulin sensitivity was determined by oral glucose tolerance test. Mice underwent a glucose tolerance test after overnight fasting. Glucose was administered orally (1.25 g/kg body weight). Serum glucose measurements were performed every 15 min for 3 h (180 min). Glucose levels were measured by a standard glucometer. Fasting blood glucose (FBG) was monitored every two weeks. Serum insulin levels were determined using a commercially available ELISA kit (Mercodia AB; Uppsala, Sweden), according to the manufacturer's instructions.

#### **Histological analysis of hepatic tissues**

For histopathology, livers from individual mice were fixed in 10% formaldehyde solution and kept at room temperature until use. The tissue blocks were then embedded in paraffin, sectioned and stained with HE for morphological examination. Specimens were examined under a light microscope.

#### **Assessment of the effect of soy extracts on the systemic immune system**

The immune modulatory effect of soy extracts was determined by FACS analysis and serum cytokines.

#### **Isolation of splenocytes and hepatic lymphocytes**

Spleens and livers were kept in RPMI-1640 supplemented with FCS. Spleens were crushed through a 70  $\mu$ m nylon cell strainer<sup>[27]</sup> and centrifuged (1250 rpm for 7 min) to remove debris. Red blood cells were lysed with 1 mL of cold 155 mM ammonium chloride lysis buffer and immediately centrifuged (1250 rpm for 3 min). Splenocytes were then washed and resuspended in 1 mL RPMI + FCS. Any remains of connective tissue were removed. The viability, as assessed using trypan blue staining, was above 90%. For the isolation of hepatic lymphocytes, livers were first crushed through a stainless mesh (size 60, Sigma), and the cell suspension was placed in a 50 mL tube for 5 min so that cell debris could pellet. The cell suspension was slowly underlaid with 10 mL of lymphoprep (Ficoll, Axis-Shield PoC AS, Oslo, Norway) in a 50 mL tube. The tubes were then centrifuged at 1800 rpm for 18 min. Cells at the interface were collected and transferred to a new tube that was centrifuged again at 1800 rpm for 10 min. The resulting pellet of hepatocyte-depleted lymphocytes was resuspended in a final volume of 250  $\mu$ L. Approximately  $1 \times 10^6$  intrahepatic lymphocytes were recovered per mouse liver.

#### **Flow cytometry**

Flow cytometry was performed on splenocytes and hepatic lymphocytes, which were resuspended in 1 mL

of FACS buffer (PBS + 1% BSA + 0.1% sodium azide). Cells were stained with the diluted anti-LAP antibody (50  $\mu$ L /sample), FITC-conjugated anti-CD4/CD8 (0.5  $\mu$ L per sample), PE-conjugated anti-CD25/NK1.1 Pacific Blue-conjugated anti-CD3 (3  $\mu$ L per sample), and PerCP-conjugated anti-CD45 (2  $\mu$ L per sample). All stains were performed after blocking the Fc receptor with anti-mouse CD16/CD32 (BD Fc Block). Flow cytometry was performed using a FACScan flow cytometer and Cellquest software.

#### **Cytokine measurement**

Serum IFN- $\gamma$  levels were measured in each animal using a commercially available "sandwich" ELISA kit (Quantikine, R&D Systems, MN, United States).

#### **Statistical analysis**

All analysis was performed using Excel 2003 (Microsoft, Redmond, WA, United States). The variables were expressed as mean  $\pm$  SD. The comparison of two independent groups was performed using Student's *t*-test. All tests applied were two-tailed. *P* value of 0.05 or less was considered to be statistically significant.

## **RESULTS**


#### **Oral administration of soy-extracts (OS, M1) alleviated ConA-mediated immune hepatitis**

Figure 1A shows the effect of oral administration of the soy extracts (M1, OS) in immune-mediated hepatitis. All treated groups showed statistically significant improvement in the levels of liver enzymes, ALT and AST ( $P < 0.05$ ). No dose dependency was noted. Administration of GC or dexamethasone also significantly alleviated immune mediated liver damage ( $P < 0.01$ ,  $P < 0.004$ , respectively). Figure 1B shows that the hepatoprotective effect of the soy extracts was associated with a reduction in serum levels of pro-inflammatory cytokine IFN- $\gamma$ . This effect was most notable in mice treated with the combination of OS and M1 ( $P < 0.005$ ), for mice treated with 30  $\mu$ g of OS and M1 in lowering ( $P = 0.001$ ). Figure 1C shows the effect of oral administration of soy extracts on TNF- $\alpha$  serum levels. The 30 micrograms dose of OS and M, lowered the pro-inflammatory cytokines TNF- $\alpha$  and INF- $\gamma$  ( $P = 0.0001$  vs  $P = 0.00001$  and  $P < 0.005$  vs  $P = 0.001$  respectively).


#### **Oral administration of soy extracts alleviated metabolic syndrome in HFD model**


Oral M1 and OS soy extracts alleviated liver damage and insulin resistance. No significant changes were noted in body weight between all groups.

Figure 2A shows the effect of oral administration of soy extracts on serum cholesterol levels that were measured every two weeks. Administration of 0.3  $\mu$ g OS + 0.3  $\mu$ g M1 was associated with a significant decrease in serum levels noted as early as week 2,



**Figure 1** Effect of soy extracts on concanavalin A immune mediated liver injury. A: Combination of soy extracts in doses of 3 µg and 30 µg of both OS and M1, administered to mice injected with concanavalin A (ConA), significantly decrease aspartate aminotransferase and alanine aminotransferase serum levels (<sup>a</sup> $P < 0.04$ , <sup>b</sup> $P < 0.01$ , <sup>c</sup> $P < 0.004$  vs control); B: The combination of OS and M1 soy extracts in all doses and 3 µg M1 were associated with significant reduction of serum interferon gamma levels measured by ELISA at the end of the treatment period (<sup>a</sup> $P < 0.02$ , <sup>b</sup> $P < 0.005$ , <sup>c</sup> $P = 0.001$  vs control); C: All combinations of soy extracts were associated with a significant decrease in serum TNF- $\alpha$  serum levels measured by ELISA at the end of the study (<sup>a</sup> $P < 0.005$ , <sup>b</sup> $P = 0.0001$ , <sup>c</sup> $P = 0.00001$  vs control). AST: Aspartate aminotransferase; ALT: Alanine aminotransferase.





**Figure 2** Effect of soy extracts on a high fat diet mouse model of non-alcoholic steatohepatitis. A: Serum cholesterol levels were measured every two weeks in all groups. The combination of 0.3 μg of OS and M1 was associated with significant a decrease of serum cholesterol level compared to the control group (<sup>e</sup>*P* < 0.05, <sup>f</sup>*P* < 0.01 vs control); B: Serum triglyceride levels were measured every two weeks in all groups. The combination of 3 μg of each OS and M1 serum levels on week 12 (<sup>e</sup>*P* < 0.03 vs control); C: Fasting serum glucose levels were measured every two weeks in all groups. The combination of OS and M1 improved fasting glucose levels as evident from week number 6. A dose of 3 μg of M1 significantly improved fasting glucose level at week 12 (<sup>e</sup>*P* < 0.04, <sup>b</sup>*P* < 0.008, <sup>c</sup>*P* < 0.0005 vs control); D: Hepatic triglyceride content was measured in all mice at the end of the study. The combination of OS and M1, and M1 alone, were associated with significant reduction of hepatic fat content (<sup>e</sup>*P* < 0.05, <sup>b</sup>*P* < 0.003 vs control); E: Representative H&E (magnification × 10) sections from liver biopsies performed in all groups at the end of the treatment period; F: Serum TNF-α levels were measured by ELISA at the end of the study. A significant reduction was noted in mice treated with the combination of 3 μg M1 and 3 μg OS (<sup>e</sup>*P* < 0.05 vs control); G: FACS analysis was performed on splenocytes for NKT and CD8+CD25+Foxp3+ regulatory T lymphocytes. Both the M1 + OS combination and the M1 alone treated group showed a significant reduction in these subsets (<sup>a</sup>*P* < 0.002, <sup>b</sup>*P* < 0.0009, <sup>c</sup>*P* = 0.0003 vs control).

in comparison with the control group ( $P < 0.05$ ). Similarly, Figure 2B shows the effect of oral administration of soy extracts on serum triglyceride levels. A significant effect on serum TGs was observed in week 12 in the 3  $\mu\text{g}$  OS + 3  $\mu\text{g}$  M1 treated group ( $P < 0.03$ ). Oral administration of the soy extracts improved insulin resistance. Figure 2C shows the effect of oral administration of soy extracts on fasting glucose levels. The fasting glucose levels were significantly decreased by the combination of M1 and OS in two doses (0.3 and 3  $\mu\text{g}$  for each extract,  $P < 0.008$ ,  $P < 0.0005$ , respectively). The glucose tolerance test improved in all treatment groups by weeks 4 and 12 (data not shown). Serum insulin was reduced in two of the treated groups (3  $\mu\text{g}$  OS and 3  $\mu\text{g}$  of each OS and M1) but did not reach statistical significance.

Oral administration of the soy extracts improved liver damage in this model. Figure 2D shows the effect of oral administration of soy extracts on liver triglyceride content. A significant reduction in hepatic TGs was observed in all soy extracts, as the combination of 3  $\mu\text{g}$  M1 and 3  $\mu\text{g}$  OS, and the 3  $\mu\text{g}$  M1, were the most effective ones compared with the control group ( $P < 0.003$ ). Figure 2E shows representative sections from liver biopsies performed at the end of the treatment period in all mice in all groups. Alleviation of hepatic steatosis and improved liver cell architecture were noted in mice treated with the combination of 3  $\mu\text{g}$  M1 and 3  $\mu\text{g}$  OS.

The beneficial effects noted in this model were associated with alteration of the immune response. Figure 2F shows the effect of oral administration of soy extracts on TNF- $\alpha$  serum levels. Serum TNF- $\alpha$  was significantly reduced in mice treated with the combination of 3  $\mu\text{g}$  M1 and 3  $\mu\text{g}$  OS ( $P < 0.05$ ). Figure 2G shows the effect of oral administration of soy extracts on NKT and CD8<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> regulatory T lymphocytes. Both combinations of M1 + OS, and the M1 alone treated group showed a significant reduction of these subsets in the spleens.

#### **Oral administration of soy extracts alleviated fatty liver in mice given HFD/MCR diet, in a NASH model**

Similar to the effect on HFD mice, oral M1 and OS soy extracts alleviated liver damage and insulin resistance in the HFD/MCD model without a significant change in body weight. Treatment with 3  $\mu\text{g}$  M1 and 6  $\mu\text{g}$  OS extracts showed a trend in reduction of ALT serum levels (data not shown,  $P = 0.6$ ). Oral administration of soy extracts was associated with a beneficial effect on serum TGs and a significant decrease was noted in 3  $\mu\text{g}$  M1 and 6  $\mu\text{g}$  OS treated mice and for 0.6  $\mu\text{g}$  M1 treated mice at week 12 (data not shown,  $P < 0.02$ ). Figure 3A shows the effect of oral administration of soy extracts on fasting serum glucose levels measured on weeks 0, 7 and 13. Fasting glucose levels were significantly improved in all treated groups compared with the controls ( $P < 0.05$ ). The effect was more

profound in the 3  $\mu\text{g}$  M1 and 6  $\mu\text{g}$  OS, and 9  $\mu\text{g}$  M1 and 18  $\mu\text{g}$  OS treated groups ( $P = 0.003$  vs  $P = 0.008$ , respectively). No significant difference was noted for fasting serum insulin levels.

Figure 3B shows the effect of oral administration of soy extracts on liver triglyceride content. Treatment with the soy extract was associated with a significant reduction of hepatic tissue triglycerides at the end of the study for the 9  $\mu\text{g}$  M1 and 18  $\mu\text{g}$  OS treated group ( $P < 0.05$ ). Figure 3C shows representative sections from liver biopsies performed at the end of the treatment period in all mice in all groups. Alleviation of hepatic steatosis and improved liver cell architecture were noted in mice in all groups, and was mostly profound for mice treated with the combination of 9  $\mu\text{g}$  M1 and 18  $\mu\text{g}$  OS. Figure 3D shows the significant improvement in the pathological NAS score noted in all treated groups.

The beneficial effects noted in this model were associated with alteration of the immune response. Serum TNF- $\alpha$  levels were significantly decreased in the 3  $\mu\text{g}$  OS + 3  $\mu\text{g}$  M1 treated group ( $P < 0.05$ , not shown). Figure 3E shows the effect of oral administration of soy extracts on regulatory lymphocytes. CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> and NKT lymphocyte subsets were significantly increased in the spleen of mice treated with the combination of 9  $\mu\text{g}$  M1 and 18  $\mu\text{g}$  OS ( $P < 0.05$ ).





## DISCUSSION


Oral administration of the combination of OS and M1 soy-derived extracts exerted a systemic immune effect *via* an effect on the gut-immune system, alleviating immune-mediated liver injury in the ConA hepatitis model, hyperlipidemia, insulin resistance and liver damage, in HFD and HFD/MCD mouse models of NASH.

In the ConA hepatitis model, liver damage results from a cytokine storm mediated by kupffer cells and NKT cells. Both IFN- $\gamma$  and TNF- $\alpha$  play a role in mediating the damage<sup>[28]</sup>. Oral administration of the combination of OS and M1 alleviated the immune-mediated liver damage that was manifested by a decrease in liver enzymes. The combination of OS and M1 was superior to either fraction alone in reducing liver enzymes in a dose-dependent manner. The hepatoprotective effect of the two soy extracts examined in the present study was associated with reduction of pro-inflammatory cytokines IFN- $\gamma$  and TNF- $\alpha$ . Similar to the effect on liver enzymes, the anti-inflammatory effect was mostly noted in mice treated with the combination of the two extracts in a dose dependent manner.

Alteration of the immune system, increase in oxidative stress, and pro-inflammatory cytokines are associated with insulin resistance and liver inflammatory response in NASH<sup>[29-31]</sup>. Both the HFD, and the HFD/MCR, models of NASH are characterized by severe depletion of hepatic anti-oxidants<sup>[32]</sup>. The HFD







**Figure 3** Effect of soy extracts on the high-fat diet/MCD mouse model of non-alcoholic steatohepatitis. A: The effect of oral administration of soy extracts on fasting serum glucose levels measured on weeks 0, 7 and 13. Glucose levels improved in all treated groups compared to the controls, and the effect was more profound in the 3 µg M1 and 6 µg OS, and 9 µg M1 and 18 µg OS treated groups (<sup>a</sup>*P* < 0.01, <sup>b</sup>*P* < 0.001 vs control); B: Hepatic triglyceride content was measured in all mice at the end of the study. The combination of 9 µg M1 and 18 µg OS treated groups was associated with significant reduction of hepatic fat content (*P* < 0.05 vs control); C: Representative HE (magnification × 10) sections from liver biopsies performed from all groups at the end of the treatment period; D: The pathological NAS score was performed on liver sections from all groups at the end of treatment. A significant improvement was noted in mice in all treated groups (<sup>a</sup>*P* < 0.001, <sup>b</sup>*P* = 0.0001 vs control); E: FACS analysis was performed on splenocytes for NKT and CD4+CD25+Foxp3+ regulatory T lymphocytes. Both regulatory lymphocyte subsets significantly increased in the spleen of mice treated with the combination of 9 µg M1 and 18 µg OS (<sup>a</sup>*P* < 0.05, <sup>b</sup>*P* < 0.005 vs control).

model is associated with insulin resistance but not always with histological steatohepatitis<sup>[33,34]</sup>. The MCD diet model is associated with histological features of steatohepatitis, but usually is not associated with insulin resistance and obesity<sup>[35]</sup>. The HFD/MCD model provides a model for NASH in which mice manifest both insulin resistance and histological features of NASH<sup>[25]</sup>. TNF- $\alpha$  plays a key role in the development of insulin resistance and NASH in both models, similar to its role in humans with metabolic syndrome<sup>[36,37]</sup>. The data of the present study supports the notion of a systemic immune modulatory anti-inflammatory effect of the two tested soy-extracts, which may target the inflammatory pathways activated in these models.

In the both models of NASH, oral administration of the combination of OS and M1 was associated with alleviation of liver damage and a decrease in plasma levels of cholesterol and/or triglycerides, decrease of hepatic content of triglycerides, improved liver histology and alleviation of insulin resistance. These effects were associated with skewing the immune profile towards an anti-inflammatory paradigm as manifested by a decrease in TNF- $\alpha$  serum levels and alteration in the distribution of different subsets of regulatory lymphocytes.

Tregs are important for the control of the immune response in the setting of immune mediated disorders<sup>[38,39]</sup>. They play a role in protection from liver damage in immune-mediated hepatitis, NAFLD, NASH, insulin resistance and metabolic syndrome<sup>[40-43]</sup>. The results of the present study suggests that the two tested soy extracts altered the distribution of Tregs, further supporting the notion that immune modulatory

therapy may be beneficial in NASH<sup>[11]</sup>.

Previous studies showed that animals fed with soy manifested a reduction in total cholesterol, triglycerides, and blood pressure<sup>[43]</sup>. Soy protein lowered plasma cholesterol concentrations, and body fat accumulation in the animal model of NASH<sup>[24]</sup>. Soy protein intake decreased the hepatic lipid depots of triacylglycerols and cholesterol, and decreased the concentrations of lipid peroxides<sup>[24]</sup>. Rats fed a soy protein diet showed improved anti-oxidative potential due to increases in superoxide dismutase and catalase activities, and a decrease in the protein expression of cytochrome P450 2E1<sup>[24]</sup>. Soy derived products decreased the amount of subcutaneous and mesenteric fat<sup>[24]</sup>. Plasma leptin levels were down-regulated and the HFD-induced increase in liver weight and lipid accumulation in the liver were suppressed<sup>[44]</sup>. Genistein, a soy isoflavone, improves insulin sensitivity<sup>[45]</sup>. Administration of genistein to fructose-fed rats significantly reduced biochemical and histological abnormalities, activating the antioxidant profile, decreasing IL-6 and TNF- $\alpha$  concentrations, ameliorating liver steatosis and insulin-resistance<sup>[45]</sup>. Soy protein reduced liver steatosis in the HFD model *via* induction of PPARalpha-regulated genes<sup>[46]</sup>. An association between the anti-inflammatory effect of isoflavones and the regulation of their immune function was described<sup>[47]</sup>. The effects of soy on *de novo* lipogenic carbohydrate responsive element binding protein (ChREBP) and anti-adipogenic Wnt signaling were recently described<sup>[48]</sup>. Isoflavones suppress ChREBP signaling *via* protein kinase A (PKA) and/or 5'-AMP activated protein kinase (AMPK)-

dependent phosphorylation, which prevents ChREBP from binding to the promoter regions of lipogenic enzymes. Isoflavones stimulate Wnt signaling *via* estrogen receptor-dependent pathways, which in turn inactivate glycogen synthase kinase-3 beta (GSK-3beta), transactivate T-cell factor/lymphoid-enhancer factor (TCF/LEF), degrade adipogenic peroxisome proliferator-activated receptor gamma (PPARgamma), and augment p300/CBP, the transcriptional co-activators of TCF/LEF<sup>[48]</sup>.

Soy derived products are safe and beneficial in humans. Soy is part of the regular diet in many countries. Soybean is a rich source of vegetable protein, complex carbohydrates, polyunsaturated fat, soluble sugars and phytoestrogens<sup>[14-17]</sup>. Some soy proteins are also associated with fatty acids, saponins, isoflavones and phospholipids. Epidemiological studies showed that soy products might decrease the morbidity of NASH<sup>[23]</sup>, lower serum lipids, reduce liver uptake of lipids, improve the anti-oxidant ability, and decrease insulin resistance<sup>[24]</sup>. Soy milk consumption was associated with better blood pressure control among diabetic patients with nephropathy<sup>[49]</sup>. Soy phytoestrogen supplementation reduced insulin resistance, hsCRP, and blood pressure<sup>[50]</sup>. Pasta enriched with soy improved endothelial function and had beneficial effects on cardiovascular risk markers in patients with type 2 diabetes<sup>[51]</sup>. Consumption of soy protein improved serum lipids in adults with type 2 diabetes<sup>[52]</sup>. Soy-protein consumption reduced proteinuria in type 2 diabetes with nephropathy<sup>[53]</sup>. A daily supplement of soy protein prevents the increase in subcutaneous and total abdominal fat in postmenopausal women<sup>[54]</sup>. A significantly greater improvement was observed in CVD risk factors in postmenopausal women on a program incorporating 30 g of soy protein and 4 g of phytosterols per day than with standard therapy<sup>[55]</sup>. In postmenopausal hypercholesterolemic women soy protein improves endothelial function regardless of changes in plasma lipoproteins<sup>[56]</sup>. Consumption of soy protein has been shown to improve blood lipid levels in nondiabetic subjects<sup>[16]</sup>.

Oral immune therapy is an active process that uses the inherent ability of the GI tract's immune system to control unwanted systemic immune responses, by inducing Tregs that suppress the chronic inflammatory state associated with metabolic syndrome<sup>[57]</sup>. This may involve alteration of NKT-DC cell interaction in the gut. The interaction between DC and NKT cells in gut associated lymphoid tissue may contribute to the promotion of Tregs, thereby alleviating immune mediated target organ damage<sup>[12,58]</sup>. One possible mechanism for the effect of these extracts may be *via* the soy-derived glycosphingolipid-mediated alteration of NKT-dendritic cell (DC) cross talk in the gut immune system<sup>[57,59]</sup>. NKT cell function is dependent on the activity of glycosphingolipids<sup>[60]</sup>.  $\beta$ -glucosylceramide, an active ingredient in soy, was shown to exert an

immune modulatory effect in various models and was also tested in clinical trials<sup>[21,26,42,61-63]</sup>. The combination of several soy derived- $\beta$ -glycosphingolipids had a synergistic effect in comparison to each other alone<sup>[64,65]</sup>.

The data in the present study suggests that a combination of the two tested soy extracts exerts a synergistic effect on most of the clinical and immunological parameters tested. While the active molecules in each of the tested extracts remains to be determined, it is postulated that different compositions of the extracts activate different pathways of the immune system that may underline the outcome.

In conclusion, our study suggests that the oral administration of soy derived molecules exerts a hepatoprotective effect alleviating immune-mediated liver injury, hyperlipidemia, insulin resistance, and liver damage, in several models. Being soy-derived safe products, the OS and M1 extracts may become a treatment option for humans with NASH and metabolic syndrome.

## COMMENTS

### Background

Non-alcoholic steatohepatitis (NASH) is a growing health problem and no approved therapy is available. NASH is associated with a pro-inflammatory state. Soy derived molecules were suggested to exert an adjuvant effect *via* activation of innate immune cells in the gut wall, thereby alleviating immune mediated disorders. The aim of the present study was to determine the immune-modulatory and the hepatoprotective effects of oral administration of two soy extracts in immune mediated liver injury and NASH.

### Research frontiers

Two soy extracts, M1 and OS, were orally administered to mice with Concanavalin A (ConA) immune-mediated hepatitis, to high-fat diet (HFD) mice and to methionine and choline reduced diet combined with HFD mice. Oral administration of OS and M1 had an additive effect in alleviating ConA hepatitis manifested by a decrease in ALT and AST serum levels. Oral administration of the OS and M1 soy derived fractions, ameliorated liver injury in the high fat diet model of NASH, manifested by a decrease in hepatic triglyceride levels, improvement in liver histology, decreased serum cholesterol and triglycerides and improved insulin resistance. In the methionine and choline reduced diet combined with the high fat diet model, the authors noted a decrease in hepatic triglycerides and improvement in blood glucose levels and liver histology. The effects were associated with reduced serum TNF- $\alpha$  and alteration of regulatory T cell distribution.

### Innovations and breakthroughs

Oral administration of the combination of OS and M1 soy derived extracts exerted an adjuvant effect in the gut-immune system, altering the distribution of regulatory T cells, and alleviating immune mediated liver injury, hyperlipidemia and insulin resistance.

### Applications

The results of the study suggest that these extracts can be further developed to be used for the treatment of non-alcoholic fatty liver disease and NASH.

### Peer-review

The authors tried to determine the immune-modulatory and the hepatoprotective effects of oral administration of two soy extracts in immune mediated liver injury and NASH. Oral administration of the combination of OS and M1 soy derived extracts exerted an adjuvant effect in the gut-immune system, altering the distribution of regulatory T cells, and alleviating immune mediated liver injury, hyperlipidemia and insulin resistance. They present evidence for the idea that soy extracts are hepatoprotective and immuno-modulatory in several different models for both livers damage (ConA, HFD, MCR) and altered immune response. Overall, the evidence that there is something in the two extracts that ameliorates liver damage and alters the immune response is strong.

REFERENCES

- 1 **Sorbi D**, McGill DB, Thistle JL, Therneau TM, Henry J, Lindor KD. An assessment of the role of liver biopsies in asymptomatic patients with chronic liver test abnormalities. *Am J Gastroenterol* 2000; **95**: 3206-3210 [PMID: 11095343 DOI: 10.1111/j.1572-0241.2000.03293.x]
- 2 **Mathiesen UL**, Franzén LE, Frydén A, Foberg U, Bodemar G. The clinical significance of slightly to moderately increased liver transaminase values in asymptomatic patients. *Scand J Gastroenterol* 1999; **34**: 85-91 [PMID: 10048738 DOI: 10.1080/00365529950172880]
- 3 **Lackner C**. Hepatocellular ballooning in nonalcoholic steatohepatitis: the pathologist's perspective. *Expert Rev Gastroenterol Hepatol* 2011; **5**: 223-231 [PMID: 21476917 DOI: 10.1586/egh.11.8]
- 4 **Neuschwander-Tetri BA**, Caldwell SH. Nonalcoholic steatohepatitis: summary of an AASLD Single Topic Conference. *Hepatology* 2003; **37**: 1202-1219 [PMID: 12717402 DOI: 10.1053/jhep.2003.50193]
- 5 **Anty R**, Lemoine M. Liver fibrogenesis and metabolic factors. *Clin Res Hepatol Gastroenterol* 2011; **35** Suppl 1: S10-S20 [PMID: 21742296 DOI: 10.1016/S2210-7401(11)70003-1]
- 6 **Stanton MC**, Chen SC, Jackson JV, Rojas-Triana A, Kinsley D, Cui L, Fine JS, Greenfeder S, Bober LA, Jenh CH. Inflammatory Signals shift from adipose to liver during high fat feeding and influence the development of steatohepatitis in mice. *J Inflamm (Lond)* 2011; **8**: 8 [PMID: 21410952 DOI: 10.1186/1476-9255-8-8]
- 7 **Robertson G**, Leclercq I, Farrell GC. Nonalcoholic steatosis and steatohepatitis. II. Cytochrome P-450 enzymes and oxidative stress. *Am J Physiol Gastrointest Liver Physiol* 2001; **281**: G1135-G1139 [PMID: 11668021]
- 8 **Porepa L**, Ray JG, Sanchez-Romeu P, Booth GL. Newly diagnosed diabetes mellitus as a risk factor for serious liver disease. *CMAJ* 2010; **182**: E526-E531 [PMID: 20566726 DOI: 10.1503/cmaj.092144]
- 9 **Vignali DA**, Collison LW, Workman CJ. How regulatory T cells work. *Nat Rev Immunol* 2008; **8**: 523-532 [PMID: 18566595 DOI: 10.1038/nri2343]
- 10 **Winer S**, Chan Y, Paltser G, Truong D, Tsui H, Bahrami J, Dorfman R, Wang Y, Zielenski J, Mastronardi F, Maezawa Y, Drucker DJ, Engleman E, Winer D, Dosch HM. Normalization of obesity-associated insulin resistance through immunotherapy. *Nat Med* 2009; **15**: 921-929 [PMID: 19633657 DOI: 10.1038/nm.2001]
- 11 **Ilan Y**. Immune therapy for nonalcoholic steatohepatitis: are we there yet? *J Clin Gastroenterol* 2013; **47**: 298-307 [PMID: 23442833 DOI: 10.1097/MCG.0b013e31827873dc]
- 12 **Ilan Y**, Maron R, Tukpah AM, Maioli TU, Murugaiyan G, Yang K, Wu HY, Weiner HL. Induction of regulatory T cells decreases adipose inflammation and alleviates insulin resistance in ob/ob mice. *Proc Natl Acad Sci USA* 2010; **107**: 9765-9770 [PMID: 20445103 DOI: 10.1073/pnas.0908771107]
- 13 **Mizrahi M**, Shabat Y, Ben Ya'acov A, Lalazar G, Adar T, Wong V, Muller B, Rawlin G, Ilan Y. Alleviation of insulin resistance and liver damage by oral administration of Imm124-E is mediated by increased Tregs and associated with increased serum GLP-1 and adiponectin: results of a phase I/II clinical trial in NASH. *J Inflamm Res* 2012; **5**: 141-150 [PMID: 23293533]
- 14 **Friedman M**, Brandon DL. Nutritional and health benefits of soy proteins. *J Agric Food Chem* 2001; **49**: 1069-1086 [PMID: 11312815 DOI: 10.1021/jf0009246]
- 15 **Erdman JW**. Control of serum lipids with soy protein. *N Engl J Med* 1995; **333**: 313-315 [PMID: 7596379 DOI: 10.1056/NEJM19950803330511]
- 16 **Hermansen K**, Søndergaard M, Høie L, Carstensen M, Brock B. Beneficial effects of a soy-based dietary supplement on lipid levels and cardiovascular risk markers in type 2 diabetic subjects. *Diabetes Care* 2001; **24**: 228-233 [PMID: 11213870 DOI: 10.2337/diacare.24.2.228]
- 17 **Zhang HM**, Chen SW, Zhang LS, Feng XF. The effects of soy isoflavone on insulin sensitivity and adipocytokines in insulin resistant rats administered with high-fat diet. *Nat Prod Res* 2008; **22**: 1637-1649 [PMID: 19085422 DOI: 10.1080/14786410701869598]
- 18 **Ustundag B**, Bahcecioglu IH, Sahin K, Duzgun S, Koca S, Gulcu F, Ozeran IH. Protective effect of soy isoflavones and activity levels of plasma paraoxonase and arylesterase in the experimental nonalcoholic steatohepatitis model. *Dig Dis Sci* 2007; **52**: 2006-2014 [PMID: 17420940 DOI: 10.1007/s10620-006-9251-9]
- 19 **Yanagihara K**, Ito A, Toge T, Numoto M. Antiproliferative effects of isoflavones on human cancer cell lines established from the gastrointestinal tract. *Cancer Res* 1993; **53**: 5815-5821 [PMID: 8242641]
- 20 **Livovsky DM**, Lalazar G, Ben Ya'acov A, Pappo O, Preston S, Zolotaryova L, Ilan Y. Administration of beta-glycolipids overcomes an unfavorable nutritional dependent host milieu: a role for a soy-free diet and natural ligands in intrahepatic CD8+ lymphocyte trapping and NKT cell redistribution. *Int Immunopharmacol* 2008; **8**: 1298-1305 [PMID: 18602077 DOI: 10.1016/j.intimp.2008.05.005]
- 21 **Zigmond E**, Preston S, Pappo O, Lalazar G, Margalit M, Shalev Z, Zolotarov L, Friedman D, Alper R, Ilan Y. Beta-glucosylceramide: a novel method for enhancement of natural killer T lymphocyte plasticity in murine models of immune-mediated disorders. *Gut* 2007; **56**: 82-89 [PMID: 17172586 DOI: 10.1136/gut.2006.095497]
- 22 **Zhao H**, Przybylska M, Wu IH, Zhang J, Maniatis P, Pacheco J, Piepenhagen P, Copeland D, Arbeeny C, Shayman JA, Aerts JM, Jiang C, Cheng SH, Yew NS. Inhibiting glycosphingolipid synthesis ameliorates hepatic steatosis in obese mice. *Hepatology* 2009; **50**: 85-93 [PMID: 19444873 DOI: 10.1002/hep.22970]
- 23 **Nomura H**, Kashiwagi S, Hayashi J, Kajiyama W, Tani S, Goto M. Prevalence of fatty liver in a general population of Okinawa, Japan. *Jpn J Med* 1988; **27**: 142-149 [PMID: 3047469 DOI: 10.2169/internalmedicine1962.27.142]
- 24 **Yang HY**, Tzeng YH, Chai CY, Hsieh AT, Chen JR, Chang LS, Yang SS. Soy protein retards the progression of non-alcoholic steatohepatitis via improvement of insulin resistance and steatosis. *Nutrition* 2011; **27**: 943-948 [PMID: 21333494 DOI: 10.1016/j.nut.2010.09.004]
- 25 **Cong WN**, Tao RY, Tian JY, Liu GT, Ye F. The establishment of a novel non-alcoholic steatohepatitis model accompanied with obesity and insulin resistance in mice. *Life Sci* 2008; **82**: 983-990 [PMID: 18417155 DOI: 10.1016/j.lfs.2008.01.022]
- 26 **Margalit M**, Abu Gazala S, Alper R, Elinav E, Klein A, Doviner V, Sherman Y, Thalenfeld B, Engelhardt D, Rabbani E, Ilan Y. Glucocerebroside treatment ameliorates ConA hepatitis by inhibition of NKT lymphocytes. *Am J Physiol Gastrointest Liver Physiol* 2005; **289**: G917-G925 [PMID: 15976388 DOI: 10.1152/ajpgi.00105.2005]
- 27 **Falcone M**, Facciotti F, Ghidoli N, Monti P, Olivieri S, Zaccagnino L, Bonifacio E, Casorati G, Sanvito F, Sarvetnick N. Up-regulation of CD1d expression restores the immunoregulatory function of NKT cells and prevents autoimmune diabetes in nonobese diabetic mice. *J Immunol* 2004; **172**: 5908-5916 [PMID: 15128771 DOI: 10.4049/jimmunol.172.10.5908]
- 28 **Wang HX**, Liu M, Weng SY, Li JJ, Xie C, He HL, Guan W, Yuan YS, Gao J. Immune mechanisms of Concanavalin A model of autoimmune hepatitis. *World J Gastroenterol* 2012; **18**: 119-125 [PMID: 22253517 DOI: 10.3748/wjg.v18.i2.119]
- 29 **Hotamisligil GS**. Inflammation and metabolic disorders. *Nature* 2006; **444**: 860-867 [PMID: 17167474 DOI: 10.1038/nature05485]
- 30 **Hotamisligil GS**. Inflammation and endoplasmic reticulum stress in obesity and diabetes. *Int J Obes (Lond)* 2008; **32** Suppl 7: S52-S54 [PMID: 19136991 DOI: 10.1038/ijo.2008.238]
- 31 **Hotamisligil GS**. Endoplasmic reticulum stress and the inflammatory basis of metabolic disease. *Cell* 2010; **140**: 900-917 [PMID: 20303879 DOI: 10.1016/j.cell.2010.02.034]
- 32 **Koteish A**, Mae Diehl A. Animal models of steatohepatitis. *Best Pract Res Clin Gastroenterol* 2002; **16**: 679-690 [PMID: 12406439 DOI: 10.1053/bega.2002.0332]
- 33 **Collins JL**. Therapeutic opportunities for liver X receptor

- modulators. *Curr Opin Drug Discov Devel* 2004; **7**: 692-702 [PMID: 15503871]
- 34 **Collins S**, Martin TL, Surwit RS, Robidoux J. Genetic vulnerability to diet-induced obesity in the C57BL/6J mouse: physiological and molecular characteristics. *Physiol Behav* 2004; **81**: 243-248 [PMID: 15159170 DOI: 10.1016/j.physbeh.2004.02.006]
- 35 **Rinella ME**, Green RM. The methionine-choline deficient dietary model of steatohepatitis does not exhibit insulin resistance. *J Hepatol* 2004; **40**: 47-51 [PMID: 14672613 DOI: 10.1016/j.jhep.2003.09.020]
- 36 **Leite NC**, Salles GF, Cardoso CR, Villela-Nogueira CA. Serum biomarkers in type 2 diabetic patients with non-alcoholic steatohepatitis and advanced fibrosis. *Hepatol Res* 2013; **43**: 508-515 [PMID: 23067270 DOI: 10.1111/j.1872-034X.2012.01106.x]
- 37 **Rinella ME**, Siddiqui MS, Gardikiotes K, Gottstein J, Elias M, Green RM. Dysregulation of the unfolded protein response in db/db mice with diet-induced steatohepatitis. *Hepatology* 2011; **54**: 1600-1609 [PMID: 21748768 DOI: 10.1002/hep.24553]
- 38 **Wing K**, Sakaguchi S. Regulatory T cells exert checks and balances on self tolerance and autoimmunity. *Nat Immunol* 2010; **11**: 7-13 [PMID: 20016504 DOI: 10.1038/ni.1818]
- 39 **Tang Q**, Bluestone JA. The Foxp3+ regulatory T cell: a jack of all trades, master of regulation. *Nat Immunol* 2008; **9**: 239-244 [PMID: 18285775 DOI: 10.1038/ni1572]
- 40 **Hotamisligil GS**. Inflammatory pathways and insulin action. *Int J Obes Relat Metab Disord* 2003; **27** Suppl 3: S53-S55 [PMID: 14704746]
- 41 **Numata K**, Kubo M, Watanabe H, Takagi K, Mizuta H, Okada S, Kunkel SL, Ito T, Matsukawa A. Overexpression of suppressor of cytokine signaling-3 in T cells exacerbates acetaminophen-induced hepatotoxicity. *J Immunol* 2007; **178**: 3777-3785 [PMID: 17339476 DOI: 10.4049/jimmunol.178.6.3777]
- 42 **Ilan Y**, Zigmund E, Lalazar G, Dembinsky A, Ben Ya'acov A, Hemed N, Kasis I, Axelrod E, Zolotarov L, Klein A, El Haj M, Gandhi R, Baecher-Allan C, Wu H, Murugaiyan G, Kivisakk P, Farez MF, Quintana FJ, Khoury SJ, Weiner HL. Oral administration of OKT3 monoclonal antibody to human subjects induces a dose-dependent immunologic effect in T cells and dendritic cells. *J Clin Immunol* 2010; **30**: 167-177 [PMID: 19756989 DOI: 10.1007/s10875-009-9323-7]
- 43 **Barrios-Ramos JP**, Garduño-Siciliano L, Loredó M, Chamorro-Cevallos G, Jaramillo-Flores ME. The effect of cocoa, soy, oats and fish oil on metabolic syndrome in rats. *J Sci Food Agric* 2012; **92**: 2349-2357 [PMID: 22430394 DOI: 10.1002/jsfa.5637]
- 44 **Kang JS**, Lee WK, Lee CW, Yoon WK, Kim N, Park SK, Lee HS, Park HK, Han SB, Yun J, Lee K, Lee KH, Park SK, Kim HM. Improvement of high-fat diet-induced obesity by a mixture of red grape extract, soy isoflavone and L-carnitine: implications in cardiovascular and non-alcoholic fatty liver diseases. *Food Chem Toxicol* 2011; **49**: 2453-2458 [PMID: 21745528 DOI: 10.1016/j.fct.2011.06.071]
- 45 **Mohamed Salih S**, Nallasamy P, Muniyandi P, Periyasami V, Carani Venkatraman A. Genistein improves liver function and attenuates non-alcoholic fatty liver disease in a rat model of insulin resistance. *J Diabetes* 2009; **1**: 278-287 [PMID: 20923528 DOI: 10.1111/j.1753-0407.2009.00045.x]
- 46 **Badger TM**, Ronis MJ, Wolff G, Stanley S, Ferguson M, Shankar K, Simpson P, Jo CH. Soy protein isolate reduces hepatosteatosis in yellow Avy/a mice without altering coat color phenotype. *Exp Biol Med* (Maywood) 2008; **233**: 1242-1254 [PMID: 18791133 DOI: 10.3181/0802-RM-60]
- 47 **Akiyama T**, Ishida J, Nakagawa S, Ogawara H, Watanabe S, Itoh N, Shibuya M, Fukami Y. Genistein, a specific inhibitor of tyrosine-specific protein kinases. *J Biol Chem* 1987; **262**: 5592-5595 [PMID: 3106339]
- 48 **Kim MH**, Kang KS. Isoflavones as a smart curer for non-alcoholic fatty liver disease and pathological adiposity via ChREBP and Wnt signaling. *Prev Med* 2012; **54** Suppl: S57-S63 [PMID: 22227283 DOI: 10.1016/j.jpmed.2011.12.018]
- 49 **Miraghajani MS**, Najafabadi MM, Surkan PJ, Esmailzadeh A, Mirlohi M, Azadbakht L. Soy milk consumption and blood pressure among type 2 diabetic patients with nephropathy. *J Ren Nutr* 2013; **23**: 277-282.e1 [PMID: 23498346 DOI: 10.1053/j.jrn.2013.01.017]
- 50 **Sathyapalan T**, Manuchehri AM, Thatcher NJ, Rigby AS, Chapman T, Kilpatrick ES, Atkin SL. The effect of soy phytoestrogen supplementation on thyroid status and cardiovascular risk markers in patients with subclinical hypothyroidism: a randomized, double-blind, crossover study. *J Clin Endocrinol Metab* 2011; **96**: 1442-1449 [PMID: 21325465 DOI: 10.1210/jc.2010-2255]
- 51 **Clerici C**, Nardi E, Battezzati PM, Ascitti S, Castellani D, Corazzi N, Giuliano V, Gizzi S, Perriello G, Di Matteo G, Galli F, Setchell KD. Novel soy germ pasta improves endothelial function, blood pressure, and oxidative stress in patients with type 2 diabetes. *Diabetes Care* 2011; **34**: 1946-1948 [PMID: 21788625 DOI: 10.2337/dc11-0495]
- 52 **Pipe EA**, Gobert CP, Capes SE, Darlington GA, Lampe JW, Duncan AM. Soy protein reduces serum LDL cholesterol and the LDL cholesterol: HDL cholesterol and apolipoprotein B: apolipoprotein A-I ratios in adults with type 2 diabetes. *J Nutr* 2009; **139**: 1700-1706 [PMID: 19605528 DOI: 10.3945/jn.109.109595]
- 53 **Azadbakht L**, Esmailzadeh A. Soy-protein consumption and kidney-related biomarkers among type 2 diabetics: a crossover, randomized clinical trial. *J Ren Nutr* 2009; **19**: 479-486 [PMID: 19758824 DOI: 10.1053/j.jrn.2009.06.002]
- 54 **Sites CK**, Cooper BC, Toth MJ, Gastaldelli A, Arabshahi A, Barnes S. Effect of a daily supplement of soy protein on body composition and insulin secretion in postmenopausal women. *Fertil Steril* 2007; **88**: 1609-1617 [PMID: 17412329 DOI: 10.1016/j.fertnstert.2007.01.061]
- 55 **Lukaczer D**, Liska DJ, Lerman RH, Darland G, Schiltz B, Tripp M, Bland JS. Effect of a low glycemic index diet with soy protein and phytosterols on CVD risk factors in postmenopausal women. *Nutrition* 2006; **22**: 104-113 [PMID: 16459222 DOI: 10.1016/j.jnut.2005.05.007]
- 56 **Cuevas AM**, Irribarra VL, Castillo OA, Yañez MD, Germain AM. Isolated soy protein improves endothelial function in postmenopausal hypercholesterolemic women. *Eur J Clin Nutr* 2003; **57**: 889-894 [PMID: 12879082 DOI: 10.1038/sj.ejcn.1601622]
- 57 **Ilan Y**. Oral tolerance: can we make it work? *Hum Immunol* 2009; **70**: 768-776 [PMID: 19559742 DOI: 10.1016/j.humimm.2009.06.018]
- 58 **Masubuchi Y**, Sugiyama S, Horie T. Th1/Th2 cytokine balance as a determinant of acetaminophen-induced liver injury. *Chem Biol Interact* 2009; **179**: 273-279 [PMID: 19014921 DOI: 10.1016/j.cbi.2008.10.028]
- 59 **Ilan Y**. Alpha versus beta: are we on the way to resolve the mystery as to which is the endogenous ligand for natural killer T cells? *Clin Exp Immunol* 2009; **158**: 300-307 [PMID: 19793337 DOI: 10.1111/j.1365-2249.2009.04030.x]
- 60 **Bendelac A**, Savage PB, Teyton L. The biology of NKT cells. *Annu Rev Immunol* 2007; **25**: 297-336 [PMID: 17150027 DOI: 10.1146/annurev.immunol.25.022106.141711]
- 61 **Margalit M**, Shalev Z, Pappo O, Sklair-Levy M, Alper R, Gomori M, Engelhardt D, Rabbani E, Ilan Y. Glucocorticoid ameliorates the metabolic syndrome in OB/OB mice. *J Pharmacol Exp Ther* 2006; **319**: 105-110 [PMID: 16815870 DOI: 10.1124/jpet.106.104950]
- 62 **Ilan Y**, Elstein D, Zimran A. Glucocorticoid: an evolutionary advantage for patients with Gaucher disease and a new immunomodulatory agent. *Immunol Cell Biol* 2009; **87**: 514-524 [PMID: 19529001 DOI: 10.1038/icb.2009.42]
- 63 **Zigmund E**, Pappo O, Zangen S, Levy Sklair M, Hemed N, Rabbani E, Itamar R, Ilan Y, Margalit M. Treatment of non-alcoholic steatohepatitis by B-glycosylceramide: A phase I/II clinical study. *Hepatology* 2006; **44**: 180A
- 64 **Lalazar G**, Ben Ya'acov A, Livovsky DM, El Haj M, Pappo O, Preston S, Zolotarov L, Ilan Y. Beta-glycosylsphingolipid-induced alterations of the STAT signaling pathways are dependent on CD1d and the lipid raft protein flotillin-2. *Am J Pathol* 2009; **174**:

1390-1399 [PMID: 19246642 DOI: 10.2353/ajpath.2009.080841]  
65 **Zigmond E**, Zangen SW, Pappo O, Sklair-Levy M, Lalazar G, Zolotaryova L, Raz I, Ilan Y. Beta-glycosphingolipids improve

glucose intolerance and hepatic steatosis of the Cohen diabetic rat. *Am J Physiol Endocrinol Metab* 2009; **296**: E72-E78 [PMID: 18940939 DOI: 10.1152/ajpendo.90634.2008]

**P- Reviewer:** Al-Gayyar MMH, Busuttill RW, Grammatikos G  
**S- Editor:** Yu J **L- Editor:** A **E- Editor:** Ma S





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>



ISSN 1007-9327

