

RESEARCH ARTICLE

A Critical Role of the mTOR/eIF2 α Pathway in Hypoxia-Induced Pulmonary Hypertension

Ai-ping Wang^{1,2}, Xiao-hui Li¹, Yong-mei Yang^{1,2}, Wen-qun Li¹, Wang Zhang¹, Chang-ping Hu¹, Zheng Zhang¹, Yuan-jian Li^{1*}

1 Department of Pharmacology, School of Pharmaceutical Sciences, Central South University, Changsha, 410078, China, **2** Department of Anatomy, School of Medicine, University of South China, Hengyang, 421001, China

* yuan_jianli@sina.com



OPEN ACCESS

Citation: Wang A-p, Li X-h, Yang Y-m, Li W-q, Zhang W, Hu C-p, et al. (2015) A Critical Role of the mTOR/eIF2 α Pathway in Hypoxia-Induced Pulmonary Hypertension. PLoS ONE 10(6): e0130806. doi:10.1371/journal.pone.0130806

Editor: Tim Lahm, Indiana University, UNITED STATES

Received: October 15, 2014

Accepted: May 25, 2015

Published: June 29, 2015

Copyright: © 2015 Wang et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was supported by the National Natural Science Foundation of China (No. 81200035 to X-H.L, No. 81273513 to Y-J.L, and No. 81072638 to C-P.H).

Competing Interests: The authors have declared that no competing interests exist.

Abstract

Enhanced proliferation of pulmonary arterial vascular smooth muscle cells (PASMCs) is a key pathological component of vascular remodeling in hypoxia-induced pulmonary hypertension (HPH). Mammalian targeting of rapamycin (mTOR) signaling has been shown to play a role in protein translation and participate in the progression of pulmonary hypertension. Eukaryotic translation initiation factor-2 α (eIF2 α) is a key factor in regulation of cell growth and cell cycle, but its role in mTOR signaling and PASMCs proliferation remains unknown. Pulmonary hypertension (PH) rat model was established by hypoxia. Rapamycin was used to treat rats as an mTOR inhibitor. Proliferation of primarily cultured rat PASMCs was induced by hypoxia, rapamycin and siRNA of mTOR and eIF2 α were used in loss-of-function studies. The expression and activation of eIF2 α , mTOR and c-myc were analyzed. Results showed that mTOR/eIF2 α signaling was significantly activated in pulmonary arteries from hypoxia exposed rats and PASMCs cultured under hypoxia condition. Treatment with mTOR inhibitor for 21 days attenuated vascular remodeling, suppressed mTOR and eIF2 α activation, inhibited c-myc expression in HPH rats. In hypoxia-induced PASMCs, rapamycin and knockdown of mTOR and eIF2 α by siRNA significantly abolished proliferation and increased c-myc expression. These results suggest a critical role of the mTOR/eIF2 α pathway in hypoxic vascular remodeling and PASMCs proliferation of HPH.

Introduction

Pulmonary hypertension (PH) is a multi-factorial disease with poor prognosis [1]. The disease is progressive and characterized by obstructive remodeling of distal pulmonary arteries, causing an increase in pulmonary vascular resistance and subsequent right ventricular hyperplasia, which ultimately leads to right heart failure and death [2]. Dysregulation in proliferation of pulmonary arterial smooth muscle cells (PASMCs) due to aberrant expression of c-myc and cyclin-dependent kinases (CDKs) are recognized as critical cellular event in pulmonary vascular remodeling [3, 4].

Hypoxia-induced PH (HPH) is one of the most common classes of PH. A number of intracellular signaling mechanisms are observed to be involved in PASMCs proliferation and vascular remodeling in HPH, among which hypoxia-inducible factor 1 α (HIF-1 α), transforming growth factor (TGF), Wnt and mitogen-activated protein kinases (MAPKs) are widely studied [5–8]. Unfortunately, effective treatment for HPH is still lacking and the identification of new therapeutic targets remains a significant challenge.

Recent research has indicated that the mammalian target of rapamycin (mTOR) in PH is potentially involved in pathogenesis of PAH [9, 10]. Evidence has shown that mTOR contributes to the proliferation and survival of PASMCs in idiopathic pulmonary arterial hypertension (IPAH) in human and experimental PH as well. Chronic hypoxia can lead to mTOR activation [11], while inhibition of mTOR by rapamycin is beneficial for hypoxia- and monocrotaline-induced PH in animal models [11–13]. These results suggest an important role of mTOR in the development of PH. However, the underlying mechanism remains unknown at large.

mTOR is a highly conserved member of the serine/threonine protein kinase family activated by phosphorylation. There is also evidence shows that mTOR plays an important role in regulation of translation initiation [14, 15]. S6K and eIF4E binding protein (4E-BP) are two major substrates of mTOR and are key regulators of protein synthesis by targeting eIF4B, eIF4E, and eukaryotic translation elongation factor 4 (eEF4) [16, 17]. The formation of Met-tRNA and eIF2 α complex is a critical step for translation initiation, and eIF2 α plays an important role in cell growth and cell cycle regulation through multiple pathways [18–20]. eIF2 α is also reported to be important in regulation of airway smooth muscle cell proliferation, hyperplasia and migration [21]. Interestingly, rapamycin is able to prevent eIF2 α phosphorylation in hepatic cells [22]. Thus, we hypothesized that eIF2 α may play a role in regulation of PASMCs proliferation, and activation of eIF2 α might be modulated by mTOR.

In summary, the purpose of this study was to explore the contribution of mTOR and eIF2 α in vascular remodeling in HPH rat model, and the role of the mTOR/eIF2 α pathway in hypoxia induced PASMCs proliferation and vascular remodeling in PH.

Materials and Methods

Animals

Male Sprague–Dawley rats (weighing 180–220 g) were obtained from the Laboratory Animal Center, Xiangya School of Medicine, Central South University (Changsha, China). All surviving animals were handled in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals. The experimental protocol was approved by the medicine animal welfare committee of Xiangya Medical School, Central South University (Changsha, China).

Animal experiments

Rats were randomly divided into three groups ($n = 10/\text{group}$): (i) normoxia group; (ii) hypoxia group, and (iii) hypoxia plus rapamycin group. Rats in the hypoxia or hypoxia plus rapamycin groups were placed in a chamber and exposed to 10% O₂ continuously for 21 days, and rats in the hypoxia plus rapamycin group were administrated with rapamycin (1 mg/kg/d, oral gavage) from the first day of their exposure to hypoxia for 21 days. On day 22, animals were anesthetized with sodium pentobarbital (30 mg/kg, i.p.), right ventricle systolic pressure (RVSP) and mean pulmonary artery pressure (mPAP) were monitored. After sacrificing the animals, the right and left ventricle (RV, LV) and the interventricular septum (S) were dissected. The tibia length was measured and weighed for calculating the ratio of RV to (LV+S) and tibia length/RV, which are the key indexes used for evaluating RV hyperplasia. The

pulmonary arterial samples were isolated for determination of mRNA expression. Portions of excised lungs were fixed in 4% paraformaldehyde for hematoxylin–eosin and immunohistochemical staining. The remainder of the excised lungs was frozen in liquid nitrogen for immunofluorescent staining.

Histology, immunohistochemistry and immunofluorescent

Vascular remodeling of lung tissues was studied. Lung hematoxylin–eosin staining was performed as previously described [23]. Slices of arteries at a thickness of 3 μ m were stained with hematoxylin–eosin staining. Pulmonary arterial wall thickness (WT) was calculated by the following formula: $WT (\%) = \frac{\text{area}_{\text{ext}} - \text{area}_{\text{int}}}{\text{area}_{\text{ext}}} \times 100$, where area_{ext} and area_{int} are the area bounded by external and internal elastic lamina, respectively. For immunohistochemistry and immunofluorescent staining, sections were stained with anti-eIF2 α antibody (ab131495, 1:100; abcam), anti-p-eIF2 α antibody (#3398, 1:100; Cell Signaling Technology), anti-mTOR antibody (#2972, 1:100; Cell Signaling Technology), anti-p-mTOR antibody (#2974, 1:100; Cell Signaling Technology), anti-c-myc antibody (#5605, 1:100; Cell Signaling Technology), anti- α -SMA(ab7817; 1:200; abcam), developed with DAB, and counterstained with hematoxylin. The immunohistochemistry samples were treated with horseradish peroxidase (HRP)-conjugated goat anti-rabbit secondary antibodies (ZB-2301, 1:500) for 1 h at 37°C. Images were acquired using a Nikon Olympus BX51. The immunofluorescence samples were treated with cy3-conjugated (BA1032, 1:50) and/or fluorescein isothiocyanate (FITC)-conjugated secondary antibodies (BA1101, 1:50) for 1 h at 37°C. Analysis of the intensity in smooth muscle actin (SMA)-positive areas of small muscular PAs was performed using an Olympus BX51.

Cell cultures

PASMCs were isolated from the pulmonary artery of 10-week-old male Sprague-Dawley rats using the method described previously [24]. The cells were cultured at 37°C under 5% CO₂ in Dulbecco's modified Eagle's medium (DMEM) containing 20% fetal bovine serum (FBS). PASMCs were identified by immunohistochemical staining using an antibody against smooth muscle α -actin (ab7817, 1:50; Abcam, Hong Kong, China). PASMCs between passages 3 and 8 were used for the experiments.

Three series of experiments were designed. The first series of experiments investigated the role of hypoxia on PASMCs proliferation. PASMCs were subjected to hypoxia for 12, 24, 48, 60 and 72 h to determine the optimal duration of exposure to hypoxia. The second series of experiments were designed to explore the effect of eIF2 α siRNA on PASMCs proliferation and the involvement of c-myc pathway. PASMCs were treated with eIF2 α siRNA (10, 25, 50 nmol/L) under normal culture conditions for 4 h, then subjected to hypoxia (3%O₂, 5%CO₂ and 92% N₂) for 48 h. Cell proliferation assays were performed. eIF2 α and c-myc expression were analyzed. The 48 h hypoxia duration was determined based on previous pilot study. The three series of experiments were designed to explore the effect of rapamycin (RAPA) on PASMCs proliferation and involvement of the eIF2 α /c-myc pathway. PASMCs were treated with rapamycin (10, 20, 40 and 80 nmol/L), then subjected to hypoxia (3%O₂, 5%CO₂ and 92%N₂) for 48 h. Cell proliferation assays were performed and the expression of eIF2 α , mTOR and c-myc was analyzed.

Small interference RNA transfection

The mTOR and eIF2 α siRNA were purchased from Ribobio (Guangzhou, China). TurboFect transfection reagent (#0531; Thermo scientific, USA) were used for cell transfection. PASMCs grown to 60% to 70% confluence were starved with 2% fetal bovine serum in Dulbecco's

modified Eagle's medium (DMEM) for 24 h and then transfected with Turbofect transfection reagent for 4–6 h, and then fresh medium were replaced according to the manufacturer's instruction. Cells were harvested 4h after for the experiments as mentioned above. Transfection efficiency was evaluated by eIF2 α mRNA and protein expression using real-time PCR and Western blot analysis, respectively. Each experiment was repeated using primarily cultured cells (3–8 passage). The experiments were repeated three times.

DNA synthesis analysis

DNA synthesis analysis was performed using CellTiter 96 AQueous One Solution cell Proliferation Assay (MTS; Promega, USA) according to the manufacturer's instruction. CellTiter 96 AQueousOne Solution Cell Proliferation Assay is a colorimetric method for determination of cell viability in proliferation or cytotoxicity assays. PSMCs were plated at a dense of 6,000 cells/well in 96-well plates. After the cells were attached to the 96-well plates, the medium was replaced with DMEM culture supplemented with 2% fetal bovine serum and incubated for 24 h. The first experiment was designed to determine the effect of hypoxia on the proliferation of PSMCs. Cells were subjected to hypoxia for 12, 24, 48, 60, or 72 h, cultured, and then incubated for 2 h with 20 μ l/well MTS. Absorbance was measured at 490 nm (DTX880; Beckman, Miami, FL). To determine the effect of eIF2 α siRNA on proliferation of PSMCs, cells were exposed to hypoxia for 48 h in the presence or absence of eIF2 α siRNA (10, 25, 50, 100 nmol/l) for 4 h, 20 μ l/well MTS was added followed by incubation for 2 h. Absorbance was measured at 490 nm. All experiments were performed in triplicate and repeated at least three times.

RNA preparation and quantitative reverse transcription PCR analysis

Total RNA was extracted by TRIzol (Invitrogen, Carlsbad, CA). For detection of mRNAs and miRNAs expression, RNA (0.2–0.5 μ g) was subjected to reverse transcription reaction using the PrimeScript reverse transcription reagent Kit (DRR037A; TaKaRa, Dalian, China) according to the manufacturer's instructions. Quantitative analysis of mRNAs and miRNA expression was performed using SYBR Premix Ex Taq (DRR420A; TaKaRa) at ABI 7300 system. PCR cycling conditions were as follows: an initial incubation at 95°C for 15 s, followed by 40 cycles of denaturation at 95°C for 5 s, and annealing at 60°C for 31 s. For detection of mature miRNAs, Bulge-Loop miRNA Primers (Ribobio) were used. The sequences of oligonucleotides used were as follows: Primers for eIF2 α were: 5'- GGACAAATGGAAGTATGGGATG-3' (forward), 5'-CAA-GAGAGAGCCAGTGTAATGC-3' (reverse). Primers for mTOR were: 5'- ATCCAGACCC TGACCCAAAC-3'(forward), 5'- TCCACCCACTTCCTCATCTC -3'(reverse). Primers for c-myc were: 5'- TGTCCGTTCAAGCAGATGAG -3'(forward), 5'- CGGTCAGTTTATGCACC AGA -3'(reverse). Primers for β -actin were: 5'-TGTCACC-AACTGGGACGATA-3'(forward), 5'-ACCCTCATAGATGGGCACAG-3'(reverse). Data analysis was performed by comparative Ct method using the ABI software. U6 and β -actin were used to normalize the expression of mRNA and miRNAs, respectively. The results were repeated at least three times independently using three different pools of templates.

Western blot analysis

Protein was extracted from PSMCs with RIPA buffer (contain 0.1% PMSF), and equal amounts of protein from each sample (20 or 40 μ g) were separated by 12% sodium dodecyl sulfate polyacrylamide gel electrophoresis and transferred to polyvinylidene fluoride membranes. The membranes were then incubated with primary antibodies overnight at 4°C and horseradish peroxidase-coupled goat anti-rat secondary antibody (ZB-2301, 1:5,000; Beijing, China). The chemiluminescence signals were detected with the EasySee Western Blot Kit (Beijing

TransGen Biotech, Beijing, China). The densitometric analysis was conducted with Image J 1.43 (National Institutes of Health). Primary antibodies against eIF2 α (ab131495, 1:500) were purchased from abcam (Hong Kong, China). Primary antibodies against p-eIF2 α (#3398, 1:1000), c-myc (#5605, 1:1000), mTOR (#2972, 1:1000), and p-mTOR (#2974, 1:1000) were obtained from Cell Signaling Technology. Primary antibody against β -actin (BM0627, 1:250) was purchased from Boster.

Data analysis

Data were presented as the mean \pm S.E.M (standard errors). Statistical analysis was performed by unpaired Student's *t*-test for two groups or analysis of variance followed by Newman-Student-Keuls test for multiple groups. Statistical significance was defined as $P < 0.05$.

Results

mTOR was activated in hypoxia-induced PH rats and primary PSMCs

Immunofluorescence analysis of lung tissues from hypoxia-induced PH rats showed a significant increase in p-S2481-mTOR expression level, a marker for mTOR catalytic activity. The expression of mTOR was located in SMA-positive areas in small remodeled pulmonary arteries in lungs of hypoxia-induced PH rats, suggesting a role of mTOR in medial hyperplasia of pulmonary arteries (Fig 1A and 1B). PSMCs cultured under hypoxia condition demonstrated a significant increase in phosphorylation of mTOR and mTOR mRNA expression (Fig 1C and 1D). These data indicated that hypoxia can induce mTOR activation in primarily cultured PSMCs, which may contribute to pulmonary vascular medial hyperplasia.

Activation of eIF2 α in hypoxia-induced PH rats and primarily cultured PSMCs accompanied by up-regulation of c-myc expression

eIF2 α is a critical protein for cell growth and cell cycle regulation. A recent study demonstrated that eIF2 α is involved in the regulation of airway smooth muscle cell proliferation, hyperplasia




Fig 1. Expression and activity of mTOR in pulmonary arteries from HPH rats and PSMCs under hypoxia. (A-B) Dual immunofluorescence analysis with anti-p-mTOR (red), anti-mTOR (red) and anti-SMA antibodies (green) in lung tissue specimens in rats. Images were acquired with an Olympus BX51 microscope, scale bars = 100 μ m. (C) The mRNA expression of mTOR was determined by real-time PCR in primary PSMCs. (D) The expression and phosphorylation of mTOR proteins was determined by western blot in PSMCs. Control: cells were treated with 21% O $_2$ for 48 h. Hypoxia: cells were treated with 3% O $_2$ for 48 h. Data represent fold changes in p-mTOR/ β -actin protein ratios. Data are means \pm S.E.M. ** $P < 0.01$ vs. control; *** $P < 0.001$ vs. control.

doi:10.1371/journal.pone.0130806.g001




Fig 2. Expression and activity of eIF2 α and c-myc expression in pulmonary arteries from HPH rats and PASMCs under hypoxic conditions. (A-B) Dual immunofluorescence analysis with anti-p-eIF2 α (red), anti-eIF2 α (red) and anti-SMA antibodies (green) in lung tissue specimens in rats. Images were acquired with an Olympus BX51 microscope, scale bars = 100 μ m. (C) The mRNA expression of eIF2 α was determined by real-time PCR in PASMCs. (D) The phosphorylation of eIF2 α proteins was determined by western blot in PASMCs. Data represent fold changes in p-eIF2 α /total eIF2 α protein ratios. (E-G) The expression of c-myc protein was determined by immunohistochemistry staining (brown precipitate indicates c-myc positive staining). Images (E, F) were taken under $\times 20$ magnification. G is an enlarged portion of the red box in E. (H) The expression of c-myc mRNA was determined by real-time PCR in pulmonary arteries. (I) The expression of c-myc mRNA was determined by real-time PCR in PASMCs. (J) The expression of c-myc protein was determined by western blot in PASMCs. Control: cells were treated with 21% O $_2$ for 48 h. Hypoxia: cells were treated with 3% O $_2$ for 48 h. Data are means \pm S.E.M. n = 3. *P<0.05 vs. control; **P<0.01 vs. control.

doi:10.1371/journal.pone.0130806.g002

and migration [21]. Thus, we hypothesized that eIF2 α may be involved in the regulation of PASMCs proliferation. Immunofluorescence analysis of lung tissue from hypoxia-induced PH rats showed a marked increase in eIF2 α phosphorylation. The expression of eIF2 α was located in SMA-positive areas in small remodeled pulmonary arteries in hypoxia-induced PH lungs (Fig 2A and 2B) as observed for mTOR, suggesting a possible link between eIF2 α and PASMCs proliferation and medial hyperplasia. And also, primarily cultured PASMCs from SD rats demonstrated a significant elevation of p-eIF2 α under hypoxic conditions, which persisted for up to 48 h (Fig 2C and 2D).

c-myc is an important member of the myc family and is involved in the regulation of cell proliferation and differentiation [25]. Our previous study had shown that c-myc is involved in hypoxia-induced PASMCs proliferation [24]. Results from both immunohistochemical

analysis and western blot showed elevated c-myc expression in pulmonary arteries from HPH rats and in PASMCs cultured under hypoxia condition (Fig 2E–2H, Fig 2I and 2J).

mTOR inhibition attenuated vascular remodeling in HPH rats and inhibited hypoxia-induced PASMCs proliferation

In keeping with a previous study [26], three weeks of exposure to hypoxia induced pulmonary hypertension in rats, as demonstrated by a significant elevation in RVSP and mPAP compared with the control rats (Fig 3A and 3B). Hypoxia also significantly induced hyperplasia of pulmonary arteries, compared with controls (Fig 3C and 3D). The elevation in RVSP and mPAP were decreased by mTOR inhibitor rapamycin (Fig 3A and 3B). Rapamycin treatment also attenuated hyperplasia in the vascular media of small pulmonary arteries (Fig 3C and 3D). Treatment with rapamycin also reversed hypoxia-induced PASMCs proliferation (Fig 5G and 5H). Taken together, these results showed that the mTOR inhibitor rapamycin attenuates vascular remodeling of HPH, most likely through inhibition of PASMCs proliferation.

mTOR inhibition prevents eIF2 α expression and activity

As observed in our study, mTOR and eIF2 α were up-regulated both in the media of pulmonary arteries and in hypoxia-induced cultured PASMCs. We next determined whether mTOR contributes to PASMCs proliferation and pulmonary vascular remodeling via eIF2 α . We performed immunofluorescence staining and immunohistochemical analysis of lung tissues from rats with chronic hypoxia-induced pulmonary vascular remodeling. Rapamycin markedly decreased mTOR, p-eIF2 α and eIF2 α levels (Fig 4A, 4D, 4F and 4G). At the same time, rapamycin decreased p-mTOR, p-eIF2 α and eIF2 α levels, and reduced cell proliferation in PASMCs under hypoxia (Fig 5A and 5B, 5D and 5E, 5G). In addition, p-eIF2 α and total eIF2 α expression was decreased and PASMCs proliferation by hypoxia-induced was inhibited after mTOR knockdown by siRNA (Fig 6A and 6B, 6D and 6E). These results were consistent with those obtained from the use of the mTOR pharmacologic blocker rapamycin, which strongly suggested that mTOR may act as an upstream positive regulator of eIF2 α .

c-myc expression and proliferation of PASMCs is regulated by mTOR and eIF2 α

We next determined whether mTOR and eIF2 α can control c-myc expression and cell proliferation. Our results showed that rapamycin markedly decreased c-myc expression *in vivo* and *in*




Fig 3. Effect of rapamycin on hemodynamic changes and vascular remodeling in HPH rats. (A) Right ventricle systolic pressure (RVSP). (B) Mean pulmonary artery pressure (mPAP). (C) Statistical graph analysis of pulmonary arterial media wall thickness (WT). (D) Hematoxylin-eosin staining in small PAs. Data are means \pm S.E.M. n = 8. RAPA: rapamycin. ** P <0.01 vs. control; *** P <0.001 vs. control; ## P <0.05 vs. hypoxia.

doi:10.1371/journal.pone.0130806.g003




Fig 4. Effect of rapamycin on mTOR, eIF2 α and c-myc expression and activity in HPH rats. (A-D) Dual immunofluorescence analysis with anti-p-mTOR (red), anti-mTOR (red), anti-p-eIF2 α (red), anti-eIF2 α (red), and anti-SMA antibodies (green) in lung tissue specimens in rats. Images were acquired with an Olympus BX51 microscope, scale bars = 100 μ m. (E) The expression of c-myc protein was determined by immunohistochemistry staining (brown precipitate indicates c-myc positive staining) scale bars = 200 μ m. (F-H) The expression of mTOR, eIF2 α and c-myc mRNA was determined by real-time PCR in pulmonary arteries. Data are means \pm S.E.M. n = 8. RAPA: rapamycin. *P < 0.05 vs. control; **P < 0.01 vs. control; #P < 0.05 vs. hypoxia.

doi:10.1371/journal.pone.0130806.g004

vitro (Figs 4E, 4H, 5C and 5F), and mTOR knockdown by siRNA obviously decreased c-myc expression in cultured PSMCs (Fig 6C). Furthermore knockdown of eIF2 α by siRNA was conducted in PSMCs. Firstly, we confirmed the efficiency of siRNA knockdown by western blot (Fig 7A and 7C). Secondly, we found that eIF2 α siRNA treatment suppressed hypoxia-induced increase in c-myc expression and PSMCs proliferation markedly (Fig 7B, 7D and 7E). Taken together, these data demonstrated that activation of eIF2 α and upregulation in c-myc expression are critical downstream events to mTOR that contribute to PSMCs proliferation and vascular remodeling in HPH.

Discussion

Dysregulation in proliferation and migration of PSMCs in small pulmonary arteries are critical components of pulmonary vascular remodeling and lead to medial hyperplasia in PH [27, 28, 9, 11]. Activation in mTOR signaling has been recognized as a hallmark of vascular remodeling in PH. Our present work further confirms the beneficial effect of mTOR/eIF2 α inhibition by rapamycin in HPH rats and provides a novel mechanistic link between mTOR/eIF2 α pathway and PSMCs proliferation and vascular remodeling in HPH.

Vasoconstriction and vascular remodeling are two major pathological characteristics of PH. Current medicines for the treatment of PH such as nitric oxide (NO), endothelin 1 receptor




Fig 5. Rapamycin prevented mTOR, eIF2 α and c-myc expression and PSMCs proliferation induced by hypoxia. The mRNA expression of mTOR (A), eIF2 α (C) and c-myc (E) was determined by real-time PCR. The protein expression of p-mTOR and mTOR (B), p-eIF2 α and eIF2 α (D), c-myc (F) was determined by western blot. (G) Proliferation of PSMCs was measured by MTS assay after incubation with hypoxia for 12, 24, 36, 48, 60, 72h. (H) Cell proliferation was measured by MTS assay. control: cells were treated with 21% O₂ for 48 h. hypoxia: cells were treated with 3% O₂ for 48 h; +RAPA 10–80 nM: cells were treated with RAPA 10–80 nM and then subjected to 3% O₂ for 48 h. Data are means \pm S.E.M. n = 3. RAPA: rapamycin. *P<0.05 vs. con; **P<0.01 vs. con; #P<0.05 vs. hypo; ##P<0.01 vs. hypo; ###P<0.001 vs. hypo.

doi:10.1371/journal.pone.0130806.g005

(ET1R) antagonist, prostacyclin and sildenafil are focused on improving vasodilatation. However, preventing vascular remodeling is supposed to have a greater impact on improving prognosis [1]. Several intracellular signaling proteins including mammalian target of rapamycin (mTOR), hypoxia-inducible factor 1 α (HIF-1 α), transforming growth factor (TGF) have been linked to PSMCs proliferation and vascular remodeling in PH [29, 13, 5, 7]. Research on the efficiency and safety of treatments targeting these signaling pathways has already been carried out in experimental PH models.

A critical role of mTOR signaling in cellular function has been reported mainly in cancer cells [30]. Emerging evidence has shown that mTOR is also involved in the regulation of PSMCs proliferation and survival in human IPAH and experimental PH [31, 11]. Increased expression and activity of mTOR has been observed in pulmonary arteries from monocrotaline and hypoxia-induced PH models [32–34]. Our results showed that three weeks exposure to hypoxia up-regulated mTOR expression and phosphorylation in pulmonary vessels. Further, immunofluorescence histochemistry showed co-expression of mTOR and α -SMA, a marker of smooth muscle cells. *In vitro* experiments showed that chronic hypoxia induced mTOR activation in primarily cultured rat PSMCs. These results suggest that mTOR is activated in PSMCs proliferation and pulmonary vascular remodeling in HPH. To further investigate whether activated mTOR is involved in the progression of PH, the mTOR inhibitor rapamycin was used *in vivo* and *in vitro*. Our findings showed that rats treated with rapamycin were




Fig 6. mTOR siRNA inhibited mTOR, eIF2 α and c-myc expression and PASMCS proliferation induced by hypoxia. The protein expression of mTOR (A), p-eIF2 α and eIF2 α (B), c-myc (C) was determined by western blot, statistical result of total gray value was estimated by the image analysis software. (D) Proliferation of PASMCS was measured by MTS assay. con: cells were treated with 21% O₂ for 48 h; hypo: cells were treated with 3% O₂ for 48 h; +Nc: cells were transfected with negative control 50nM for 4 h before stimulation with hypoxia for 48 h; +Veh: cells were transfected with transfection reagent (30uL/6 well) for 4 h before stimulation with hypoxia for 48 h; +mTOR siRNA 50 nM: cells were transfected with mTOR siRNA 50 nM for 4 h before stimulation with hypoxia for 48 h. Data represent the means \pm S.E.M. n = 3. **P*<0.05 vs. con; ***P*<0.01 vs. con; #*P*<0.05 vs. hypo; ##*P*<0.01 vs. hypo.

doi:10.1371/journal.pone.0130806.g006

resistant to chronic hypoxia, as evidenced by lower right ventricle systolic pressure (RVSP) and mean pulmonary arterial pressure (mPAP) and thinner pulmonary vessel walls. Pretreatment with rapamycin or knockdown of mTOR by siRNA also abolished hypoxia-induced proliferation of PASMCS. Collectively, our results demonstrate that mTOR plays an important role in PASMCS proliferation and vascular remodeling in HPH.

Results from our lab and other scientists [35–38] provide conclusive evidence that mTOR inhibition is beneficial for PH and that mTOR signaling could represent a new therapeutic target for PH treatment. However the exact underlying mechanism is still not very clear. Two forms of the mTOR complex, mTORC1 (mTOR-raptor) and mTORC2 (mTOR-riCTOR), are identified. mTORC1 is sensitive to rapamycin and is thought to mainly function in cell growth and the regulation of proliferation, while mTORC2 usually play a role in cell structure and can also be affected by long-term rapamycin treatment [31, 39]. Recently, Goncharov DA [11] et al found that mTORC2 plays a coordinating role in the metabolism, proliferation and survival of PASMCS in PAH. Krymskaya VP et al have shown that both mTORC1 and mTORC2 are required for PASMCS proliferation induced by chronic hypoxia in vitro and in vivo. Previous studies have revealed that S6K and eIF4E binding protein (4E-BP1) are major downstream




Fig 7. Effect of eIF2 α siRNA on c-myc expression and PASCs proliferation induced by hypoxia. (A) The expression of eIF2 α mRNA was determined by real-time PCR. (B) The expression of c-myc mRNA was determined by real-time PCR. Data represent the means \pm S.E.M. n = 3. * P <0.05 vs. control; # P <0.05 vs. hypoxia; ### P <0.001 vs. hypoxia. (C) The expression of eIF2 α protein was determined by western blot. Statistical result of total gray value was estimated by the image analysis software. (D) The expression of c-myc protein was determined by western blot. Statistical result of total gray value was estimated by the image analysis software. PASCs were transfected with the siRNA of eIF2 α or negative control (Nc) 4 h before stimulation with hypoxia for 48 h. (E) Proliferation of PASCs was measured by MTS assay. con: cells were treated with 21% O $_2$ for 48 h. hypo: cells were treated with 3% O $_2$ for 48 h; +eIF2 α siRNA 10–50 nM: cells were transfected with eIF2 α siRNA 10–50 nM for 4 h before stimulation with hypoxia for 48 h. Data represent the means \pm S.E.M. n = 3. * P <0.05 vs. con; # P <0.05 vs. hypo; ### P <0.01 vs. hypo; ### P <0.001 vs. hypo.

doi:10.1371/journal.pone.0130806.g007

effectors of mTOR. But more details on mechanism of mTOR activation requires further investigation.

The present study suggested that eIF2 α acts as a new mediator of mTOR and an important regulator of PASCs proliferation in HPH. Firstly, under hypoxic conditions, mTOR activation was accompanied by up-regulation of eIF2 α expression and activity (as determined by phosphorylation) in pulmonary arteries and PASCs. Secondly, rapamycin treatment significantly reversed the activation of eIF2 α in vitro and in vivo. Thirdly, knockdown of mTOR with siRNA reversed the increase in eIF2 α expression and phosphorylation in hypoxia-induced PASCs. In addition to eIF2 α , other translation initiation factors are also involved in mTOR signaling. For example, mTOR can bind to eIF3 and lead to release and activation of S6K. 4E-BP1 is one of the main downstream proteins of mTOR [40, 17]. All these results confirmed a critical role of mTOR signaling in protein translation process. Notably, although the present results give a clue that the effects of mTOR inhibitor might be mediated at least in part by eIF2 α down-regulation, studies on the causal relationship of eIF2 α and mTOR is still lacking.

The mechanism by which mTOR was activated and which complex of mTOR activates eIF2 α in PASCs should be fully revealed in future study.

Anyway, a critical role of eIF2 α in PASCs proliferation and vascular remodeling in HPH enlarged our understanding on the role of eIFs in PH. In this study, we observed a marked increase in p-eIF2 α , a marker for eIF2 α catalytic activity, in SMA-positive areas in small muscular remodeled vessels of lungs of HPH rats. Importantly, we found that eIF2 α siRNA prevented hypoxia-induced PASCs proliferation. It suggested that eIF2 α might be crucial for the media hyperplasia in HPH rat through stimulation of PASCs proliferation. We think it is only the first step to reveal the role of eIFs in PH, investigating the effect of other eIFs including eIF3, eIF4 on PASCs proliferation and vascular remodeling would be helpful to decipher the role of eIFs in PH and provide more evidence for potential therapeutic strategy targeting eIFs.

In this study, we found that eIF2 α siRNA can reverse hypoxia-induced PASCs proliferation and c-myc expression. However, much work remains to be done to elucidate the mechanism underlying eIF2 α regulation of PASCs proliferation. Previous studies showed that activation of c-myc is considered to be one of the initiating steps in vascular smooth muscle cell proliferation. An aberrant expression of c-myc is observed in injured vessels after angioplasty [41]. Consistent with previous research, c-myc expression was up-regulated in pulmonary arteries and PASCs exposed to hypoxia and was inhibited by either mTOR or eIF2 α inhibition. Proteins that regulate cell proliferation and differentiation such as c-myc may be important downstream targets of eIF2 α in HPH. However, further studies with eIF2 α knock-down in vivo and c-myc siRNA treatment or overexpression would help decipher the role of eIF2 α in PH.

Conclusions

Our findings suggest a critical role of the mTOR/eIF2 α pathway in hypoxic vascular remodeling and PASCs proliferation in HPH. Inhibition of mTOR/eIF2 α pathway could serve as a new solution for the treatment of PH.

Supporting Information

S1 Fig. Cellular proliferation stimulated by 20% FBS. Cellular proliferation stimulated by 20% FBS (48 hours) was assayed in PASCs treated with siRNA against mTOR or eIF2 α , or in the presence of rapamycin. As shown in MTS experiments, cell proliferation was inhibited by either rapamycin (10, 20, 40, 80 nmol/L) (A), eIF2 α siRNA (25, 50, 100 nmol/L) (B), or mTOR siRNA (25, 50, 100 nmol/L) (C). These results are in agreement with cell proliferation data acquired with hypoxia. Data represent the means \pm S.E.M. n = 3. * P <0.05 vs. con; ** P <0.01 vs. con; *** P <0.001 vs. con; # P <0.05 vs. hypoxia; ## P <0.01 vs. hypoxia. The above experiments were repeated three times with similar results.

(TIF)

S2 Fig. mRNA expression of Ki-67 and PCNA in pulmonary arteries. (A) The expression of Ki-67 mRNA was determined by real-time PCR. (B) The expression of PCNA mRNA was determined by real-time PCR. Data represent the means \pm S.E.M. n = 3. * P <0.05 vs. control; ** P <0.01 vs. control; # P <0.05 vs. hypoxia; ## P <0.01 vs. hypoxia. The above experiments were repeated three times with similar results.

(TIF)

S3 Fig. Schematic representation of proposed function of mTOR/eIF2 α pathway in PASMCs in HPH.

(TIF)

Author Contributions

Conceived and designed the experiments: YJL XHL APW. Performed the experiments: APW WQL WZ. Analyzed the data: YMY CPH ZZ. Contributed reagents/materials/analysis tools: YJL CPH. Wrote the paper: APW XHL YJL.

References

1. Morrell NW, Adnot S, Archer SL, Dupuis J, Jones PL, Maclean MR, et al. (2009) Cellular and Molecular Basis of Pulmonary Arterial Hypertension. *J Am Coll Cardiol* 54(1 Suppl):S20–31. doi: [10.1016/j.jacc.2009.04.018](https://doi.org/10.1016/j.jacc.2009.04.018) PMID: [19555855](https://pubmed.ncbi.nlm.nih.gov/19555855/)
2. Ricard N, Tu L, Le Hress M, Huertas A, Phan C, Thuillet R, et al. (2014) Increased Pericyte Coverage Mediated by Endothelial-Derived Fibroblast Growth Factor-2 and Interleukin-6 Is a Source of Smooth Muscle-Like Cells in Pulmonary Hypertension. *Circulation* 129(15): 1586–97. doi: [10.1161/CIRCULATIONAHA.113.007469](https://doi.org/10.1161/CIRCULATIONAHA.113.007469) PMID: [24481949](https://pubmed.ncbi.nlm.nih.gov/24481949/)
3. Sakao S, Tatsumi K. (2011) Vascular remodeling in pulmonary arterial hypertension: multiple cancer-like pathways and possible treatment modalities. *Int J Cardiol* 147(1): 4–12. doi: [10.1016/j.ijcard.2010.07.003](https://doi.org/10.1016/j.ijcard.2010.07.003) PMID: [20692712](https://pubmed.ncbi.nlm.nih.gov/20692712/)
4. Masri FA, Xu W, Comhair SAA, Asosingh K, Koo M, Vasanthi A, et al. (2007) Hyperproliferative apoptosis-resistant endothelial cells in idiopathic pulmonary arterial hypertension. *Am J Physiol Lung Cell Mol Physiol* 293(3): L548–54. PMID: [17526595](https://pubmed.ncbi.nlm.nih.gov/17526595/)
5. Hubbi ME, Gilkes DM, Hu H, Kshitiz, Ahmed I, Semenza GL. (2014) Cyclin-dependent kinases regulate lysosomal degradation of hypoxia-inducible factor 1 α to promote cell-cycle progression. *Proc Natl Acad Sci USA* 111(32): E3325–34. doi: [10.1073/pnas.1412840111](https://doi.org/10.1073/pnas.1412840111) PMID: [25071185](https://pubmed.ncbi.nlm.nih.gov/25071185/)
6. Zeng XC, Liu FQ, Yan R, Yi HM, Zhang T, Wang GY, et al. (2014) Downregulation of miR-610 promotes proliferation and tumorigenicity and activates Wnt/ β -catenin signaling in human hepatocellular carcinoma. *Mol Cancer* 13: 261. doi: [10.1186/1476-4598-13-261](https://doi.org/10.1186/1476-4598-13-261) PMID: [25491321](https://pubmed.ncbi.nlm.nih.gov/25491321/)
7. Shi X, Guo LW, Seedial SM, Si Y, Wang B, Takayama T, et al. (2014) TGF- β /Smad3 inhibit vascular smooth muscle cell apoptosis through an autocrine signaling mechanism involving VEGF-A. *Cell Death Dis* 5: e1317. doi: [10.1038/cddis.2014.282](https://doi.org/10.1038/cddis.2014.282) PMID: [25010983](https://pubmed.ncbi.nlm.nih.gov/25010983/)
8. Karelina K, Liu Y, Alzate-Correa D, Wheaton KL, Hoyt KR, Arthur JS, et al. (2015) Mitogen and stress-activated kinases 1/2 regulate ischemia-induced hippocampal progenitor cell proliferation and neurogenesis. *Neuroscience* 285: 292–302. doi: [10.1016/j.neuroscience.2014.10.053](https://doi.org/10.1016/j.neuroscience.2014.10.053) PMID: [25451279](https://pubmed.ncbi.nlm.nih.gov/25451279/)
9. Sun LQ, Cairns MJ, Gerlach WL, Witherington C, Wang L, King A, et al. (1999) Suppression of smooth muscle cell proliferation by a c-myc RNA-cleaving deoxyribozyme. *J Biol Chem* 274(24): 17236–41. PMID: [10358082](https://pubmed.ncbi.nlm.nih.gov/10358082/)
10. Goncharova EA, Goncharov DA, Li H, Pimtong W, Lu S, Khavin I, et al. (2011) mTORC2 is required for proliferation and survival of TSC2-Null Cells. *Mol Cell Biol* 31(12): 2484–2498. doi: [10.1128/MCB.01061-10](https://doi.org/10.1128/MCB.01061-10) PMID: [21482669](https://pubmed.ncbi.nlm.nih.gov/21482669/)
11. Goncharov DA, Kudryashova TV, Ziai H, Ihida-Stansbury K, DeLisser H, Krymskaya VP, et al. (2014) Mammalian target of rapamycin complex2(mTORC2) coordinates pulmonary artery smooth muscle cell metabolism, proliferation, and survival in pulmonary arterial hypertension. *Circulation* 129(8): 864–74. doi: [10.1161/CIRCULATIONAHA.113.004581](https://doi.org/10.1161/CIRCULATIONAHA.113.004581) PMID: [24270265](https://pubmed.ncbi.nlm.nih.gov/24270265/)
12. Paddenberg R, Stieger P, von Lilien AL, Faulhammer P, Goldenberg A, Tillmanns HH, et al. (2007) Rapamycin attenuates hypoxia-induced pulmonary vascular remodeling and right ventricular hyperplasia in mice. *Respir Res* 8: 15. PMID: [17319968](https://pubmed.ncbi.nlm.nih.gov/17319968/)
13. Goncharova EA. (2013) mTOR and vascular remodeling in lung diseases: current challenges and therapeutic prospects. *FASEB J* 27(5): 1796–807. doi: [10.1096/fj.12-222224](https://doi.org/10.1096/fj.12-222224) PMID: [23355268](https://pubmed.ncbi.nlm.nih.gov/23355268/)
14. Krymskaya VP, Snow J, Cesarone G, Khavin I, Goncharov DA, Lim PN, et al. (2011) mTOR is required for pulmonary arterial vascular smooth cell proliferation under chronic hypoxia. *FASEB J* 25(6): 1922–33. doi: [10.1096/fj.10-175018](https://doi.org/10.1096/fj.10-175018) PMID: [21368105](https://pubmed.ncbi.nlm.nih.gov/21368105/)
15. Zhang P, Shan T, Liang X, Deng C, Kuang S. (2014) Mammalian target of rapamycin is essential for cardiomyocyte survival and heart development in mice. *Biochem Biophys Res Commun* 452(1): 53–9. doi: [10.1016/j.bbrc.2014.08.046](https://doi.org/10.1016/j.bbrc.2014.08.046) PMID: [25139234](https://pubmed.ncbi.nlm.nih.gov/25139234/)

16. Liu T, Yacoub R, Taliaferro-Smith LD, Sun SY, Graham TR, Dolan R, et al. (2011) Combinatorial effects of lapatinib and rapamycin in triple-negative breast cancer cells. *Mol Cancer Ther* 10(8): 1460–9. doi: [10.1158/1535-7163.MCT-10-0925](https://doi.org/10.1158/1535-7163.MCT-10-0925) PMID: [21690228](https://pubmed.ncbi.nlm.nih.gov/21690228/)
17. Eckerdt F, Beauchamp E, Bell J, Iqbal A, Su B, Fukunaga R, et al. (2014) Regulatory effects of a Mnk2-eIF4E feedback loop during mTORC1 targeting of human medulloblastoma cells. *Oncotarget* 5(18):8442–51. PMID: [25193863](https://pubmed.ncbi.nlm.nih.gov/25193863/)
18. Liu Y, László C, Liu Y, Liu W, Chen X, Evans SC, et al. (2010) Regulation of G(1) arrest and apoptosis in hypoxia by PERK and GCN2-mediated eIF2 α phosphorylation. *Neoplasia* 12(1): 61–8. PMID: [20072654](https://pubmed.ncbi.nlm.nih.gov/20072654/)
19. Liu X, Bennett RL, Cheng X, Byrne M, Reinhard MK, May WS Jr. (2013) PKR regulates proliferation, differentiation, and survival of murine hematopoietic stem/progenitor cells. *Blood* 121(17): 3364–74. doi: [10.1182/blood-2012-09-456400](https://doi.org/10.1182/blood-2012-09-456400) PMID: [23403623](https://pubmed.ncbi.nlm.nih.gov/23403623/)
20. Bennett RL, Pan Y, Christian J, Hui T, May WS Jr. (2012) The RAX/PACT-PKR stress response pathway promotes p53 sumoylation and activation, leading to G₁ arrest. *Cell Cycle* 11(2): 407–17. doi: [10.4161/cc.11.2.18999](https://doi.org/10.4161/cc.11.2.18999) PMID: [22214662](https://pubmed.ncbi.nlm.nih.gov/22214662/)
21. Bentley JK, Hershenson MB. (2008) Airway smooth muscle growth in asthma: proliferation, hyperplasia, and migration. *Proc Am Thorac Soc* 5(1): 89–96. PMID: [18094090](https://pubmed.ncbi.nlm.nih.gov/18094090/)
22. Anand P, Gruppuso PA. (2006) Rapamycin inhibits liver growth during refeeding in rats via control of ribosomal protein translation but not cap-dependent translation initiation. *J Nutr* 136(1): 27–33. PMID: [16365054](https://pubmed.ncbi.nlm.nih.gov/16365054/)
23. Li XW, Hu CP, Wu WH, Zhang WF, Zou XZ, Li YJ. (2012) Inhibitory effect of calcitonin gene-related peptide on hypoxia-induced rat pulmonary artery smooth muscle cells proliferation: role of ERK1/2 and p27. *Eur J Pharmacol* 679(1–3): 117–26. doi: [10.1016/j.ejphar.2012.01.015](https://doi.org/10.1016/j.ejphar.2012.01.015) PMID: [22306243](https://pubmed.ncbi.nlm.nih.gov/22306243/)
24. Li XH, Peng J, Tan N, Wu WH, Li TT, Shi RZ, et al. (2010) Involvement of asymmetric dimethylarginine and Rho kinase in the vascular remodeling in monocrotaline-induced pulmonary hypertension. *Vascul Pharmacol* 53(5–6): 223–229. doi: [10.1016/j.vph.2010.09.002](https://doi.org/10.1016/j.vph.2010.09.002) PMID: [20840872](https://pubmed.ncbi.nlm.nih.gov/20840872/)
25. Bernard S, Eilers M. (2006) Control of cell proliferation and growth by myc proteins. *Results Probl Cell Differ* 42: 329–42. PMID: [16903216](https://pubmed.ncbi.nlm.nih.gov/16903216/)
26. Liu B, Luo XJ, Yang ZB, Zhang JJ, Li TB, Zhang XJ, et al. (2014) Inhibition of NOX/VPO1 Pathway and Inflammatory Reaction by Trimethoxystilbene in Prevention of Cardiovascular Remodeling in Hypoxia-induced Pulmonary Hypertensive Rats. *J Cardiovasc Pharmacol* 63(6): 567–76. doi: [10.1097/FJC.000000000000082](https://doi.org/10.1097/FJC.000000000000082) PMID: [24492474](https://pubmed.ncbi.nlm.nih.gov/24492474/)
27. Chan SY, Loscalzo J. (2008) Pathogenic mechanisms of pulmonary arterial hypertension. *J Mol Cardiol* 44(1): 14–30.
28. Dahal BK, Cornitescu T, Tretny A, Pullamsetti SS, Kosanovic D, Dumitrascu R, et al. (2010) Role of epidermal growth factor inhibition in experimental pulmonary hypertension. *Am J Respir Crit Care Med* 181(2): 158–67. doi: [10.1164/rccm.200811-1682OC](https://doi.org/10.1164/rccm.200811-1682OC) PMID: [19850946](https://pubmed.ncbi.nlm.nih.gov/19850946/)
29. Yang J, Li X, Li Y, Southwood M, Ye L, Long L, et al. (2013) Id proteins are critical downstream effectors of BMP signaling in human pulmonary arterial smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol* 305(4): L312–21. doi: [10.1152/ajplung.00054.2013](https://doi.org/10.1152/ajplung.00054.2013) PMID: [23771884](https://pubmed.ncbi.nlm.nih.gov/23771884/)
30. Evangelisti C, Ricci F, Tazzari P, Tabellini G, Battistelli M, Falcieri E, et al. (2011) Targeted inhibition of mTORC1 and mTORC2 by active-site mTOR inhibitors has cytotoxic effects in T-cell acute lymphoblastic leukemia. *Leukemia* 25: 781–791. doi: [10.1038/leu.2011.20](https://doi.org/10.1038/leu.2011.20) PMID: [21331075](https://pubmed.ncbi.nlm.nih.gov/21331075/)
31. Houssaini A, Abid S, Mouraret N, Wan F, Rideau D, Saker M, et al. (2013) Rapamycin reverses pulmonary artery smooth muscle cell proliferation in pulmonary hypertension. *Am J Respir Cell Mol Biol* 48(5): 568–77. doi: [10.1165/rcmb.2012-0429OC](https://doi.org/10.1165/rcmb.2012-0429OC) PMID: [23470622](https://pubmed.ncbi.nlm.nih.gov/23470622/)
32. Wang W, Liu J, Ma A, Miao R, Jin Y, Zhang H, et al. (2014) mTORC1 Is Involved in Hypoxia-Induced Pulmonary Hypertension through the Activation of Notch3. *J Cell Physiol* 229(12): 2117–25. doi: [10.1002/jcp.24670](https://doi.org/10.1002/jcp.24670) PMID: [24825564](https://pubmed.ncbi.nlm.nih.gov/24825564/)
33. Ogawa A, Firth AL, Yao W, Madani MM, Kerr KM, Auger WR, et al. (2009) Inhibition of mTOR attenuates store-operated Ca²⁺ entry in cells from endarterectomized tissues of patients with chronic thromboembolic pulmonary hypertension. *Am J Physiol Lung Cell Mol Physiol* 297(4): L666–76. doi: [10.1152/ajplung.90548.2008](https://doi.org/10.1152/ajplung.90548.2008) PMID: [19633069](https://pubmed.ncbi.nlm.nih.gov/19633069/)
34. Ogawa A, Firth AL, Ariyasu S, Yamadori I, Matsubara H, Song S, et al. (2013) Thrombin-mediated activation of Akt signaling contributes to pulmonary vascular remodeling in pulmonary hypertension. *Physiol Rep* 1(7): e00190. doi: [10.1002/phy2.190](https://doi.org/10.1002/phy2.190) PMID: [24744867](https://pubmed.ncbi.nlm.nih.gov/24744867/)
35. Humar R, Kiefer FN, Berns H, Resink TJ, Bättegay EJ. (2002) Hypoxia enhances vascular cell proliferation and angiogenesis in vitro via rapamycin (mTOR)-dependent signaling. *FASEB J* 16(8): 771–80. PMID: [12039858](https://pubmed.ncbi.nlm.nih.gov/12039858/)

36. Nührenberg TG, Voisard R, Fahlisch F, Rudelius M, Braun J, Gschwend J, et al. (2005) Rapamycin attenuates vascular wall inflammation and progenitor cell promoters after angioplasty. *FASEB J* 19(2): 246–8. PMID: [15546959](#)
37. Tang Z, Wang Y, Fan Y, Zhu Y, Chien S, Wang N. (2008) Suppression of c-Cbl tyrosine phosphorylation inhibits neointimal formation in balloon-injured rat arteries. *Circulation* 118(7): 764–72. doi: [10.1161/CIRCULATIONAHA.107.761932](#) PMID: [18663086](#)
38. Moss SC, Lightell DJ Jr, Marx SO, Marks AR, Woods TC. (2010) Rapamycin regulates endothelial cell migration through regulation of the cyclin-dependent kinase inhibitor p27Kip1. *J Biol Chem* 285(16): 11991–7. doi: [10.1074/jbc.M109.066621](#) PMID: [20097763](#)
39. Sarbassov DD, Ali SM, Sengupta S, Sheen JH, Hsu PP, Bagley AF, et al. (2006) Prolonged rapamycin treatment inhibits mTORC2 assembly and Akt/PKB. *Mol Cell* 22: 159–168. PMID: [16603397](#)
40. Harris TE, Chi A, Shabanowitz J, Hunt DF, Rhoads RE, Lawrence JC Jr. (2006) mTOR-dependent stimulation of the association of eIF4G and eIF3 by insulin. *EMBO J* 25(8):1659–68. PMID: [16541103](#)
41. Sun LQ, Cairns MJ, Gerlach WL, Witherington C, Wang L, King A, et al. (1999) Suppression of smooth muscle cell proliferation by a c-myc RNA-cleaving deoxyribozyme. *J Biol Chem* 274(24): 17236–41. PMID: [10358082](#)