

CORRECTION

Correction: Prevalence and Transmission of *Trypanosoma cruzi* in People of Rural Communities of the High Jungle of Northern Peru

Karen A. Alroy, Christine Huang, Robert H. Gilman, Victor R. Quispe-Machaca, Morgan A. Marks, Jenny Ancca-Juarez, Miranda Hillyard, Manuela Verastegui, Gerardo Sanchez, Lilia Cabrera, Elisa Vidal, Erica M. W. Billig, Vitaliano A. Cama, César Náquira, Caryn Bern, Michael Z. Levy, Working Group on Chagas Disease in Peru

There are several errors in this article. In [Table 2](#), the age ranges are incorrectly listed under Community in column 1. The age ranges should be listed under Age Category in column 2. Please see the corrected [Table 2](#) here.

There is an error in affiliation 4 for authors Robert H. Gilman, Victor R. Quispe-Machaca, Jenny Ancca-Juarez, Manuela Verastegui, Gerardo Sanchez, Lilia Cabrera, César Náquira, Michael Z. Levy, and Working Group on Chagas Disease in Peru. Affiliation 4 should be: Faculty of Science and Philosophy Alberto Cazorla Talleri, University Peruana Cayetano Heredia, Urbanización Ingeniería, Lima, Peru.

In Table 4, footnote 1 has a typo. *T.cruzi* should be italicized.



OPEN ACCESS

Citation: Alroy KA, Huang C, Gilman RH, Quispe-Machaca VR, Marks MA, Ancca-Juarez J, et al. (2015) Correction: Prevalence and Transmission of *Trypanosoma cruzi* in People of Rural Communities of the High Jungle of Northern Peru. PLoS Negl Trop Dis 9(7): e0003910. doi:10.1371/journal.pntd.0003910

Published: July 6, 2015

Copyright: © 2015 Alroy et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Table 2. Summary of age specific seropositive study participants by community.

Community (years)	Age Category	N	<i>T. cruzi</i> seropositiven (%)	95% Confidence Interval
<i>Pindoc</i>		75	20 (26.7%)	17.1–38.1%
	<10	14	0	0–23.2%
	11–20	19	0	0–17.7%
	21–40	14	4 (28.6%)	8.4–58.1%
	>40	28	16 (57.1%)	37.2–75.5%
<i>Nuevo Guayaquil</i>		63	7 (11.1%)	4.6–21.6%
	<10	11	0	0–28.5%
	11–20	16	1 (6.3%)	0.2–30.2%
	21–40	19	2 (10.5%)	1.3–33.1%
	>40	17	4 (23.5%)	6.8–49.9%
<i>Casa Blanca</i>		130	10 (7.7%)	3.8–13.7%
	<10	24	0	0–14.3%
	11–20	35	2 (5.7%)	0.70–19.2%
	21–40	31	3 (9.7%)	2.0–25.8%
	>40	40	5 (12.5%)	4.2–26.8%
<i>La Esperanza</i>		123	12 (9.8%)	5.1–16.4%
	<10	32	0	0–10.9%
	11–20	32	6 (18.8%)	7.2–36.4%
	21–40	36	3 (8.3%)	1.8–22.5%
	>40	23	3 (13.0%)	2.8–33.6%
<i>Campo Florido</i>		195	40 (20.5%)	15.0–26.9%
	<10	74	7 (9.5%)	3.9–18.5%
	11–20	37	15 (40.5%)	24.8–57.9%
	21–40	42	8 (19.1%)	8.6–34.1%
	>40	42	10 (23.8%)	12.1–39.5%
<i>Rumiaco</i>		25	2 (8.0%)	1.0–26.0%
	<10	4	1 (25.0%)	0.6–80.1%
	11–20	8	0	0–36.9%
	21–40	7	1 (14.3%)	0.4–57.9%
	>40	6	0	0–45.9%

doi:10.1371/journal.pntd.0003910.t001

Reference

1. Alroy KA, Huang C, Gilman RH, Quispe-Machaca VR, Marks MA, Ancca-Juarez J, et al. (2015) Prevalence and Transmission of *Trypanosoma cruzi* in People of Rural Communities of the High Jungle of Northern Peru. *PLoS Negl Trop Dis* 9(5): e0003779. doi:[10.1371/journal.pntd.0003779](https://doi.org/10.1371/journal.pntd.0003779) PMID: [26000770](https://pubmed.ncbi.nlm.nih.gov/26000770/)