

CORRECTION

# Correction: Preoperative Axillary Staging with 3.0-T Breast MRI: Clinical Value of Diffusion Imaging and Apparent Diffusion Coefficient

The *PLOS ONE* Staff

## Notice of Republication

This article was republished on June 22, 2015, to correct Figs 1 and 2, which were darker than intended. The publisher apologizes for the error. Please download this article again to view the corrected version.

## Reference

1. Rautiainen S, Könönen M, Sironen R, Masarwah A, Sudah M, Hakumäki J, et al. (2015) Preoperative Axillary Staging with 3.0-T Breast MRI: Clinical Value of Diffusion Imaging and Apparent Diffusion Coefficient. *PLoS ONE* 10(3): e0122516. doi: [10.1371/journal.pone.0122516](https://doi.org/10.1371/journal.pone.0122516) PMID: [25823016](https://pubmed.ncbi.nlm.nih.gov/25823016/)



## OPEN ACCESS

**Citation:** The *PLOS ONE* Staff (2015) Correction: Preoperative Axillary Staging with 3.0-T Breast MRI: Clinical Value of Diffusion Imaging and Apparent Diffusion Coefficient. *PLoS ONE* 10(7): e0133111. doi:10.1371/journal.pone.0133111

**Published:** July 14, 2015

**Copyright:** © 2015 The PLOS ONE Staff. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.