

Trans-regulation of Syndecan Functions by Hetero-oligomerization*

Received for publication, December 25, 2014, and in revised form, May 14, 2015. Published, JBC Papers in Press, May 15, 2015, DOI 10.1074/jbc.M114.611798

Youngsil Choi[‡], Mi-Jung Kwon[‡], Yangmi Lim[‡], Ji-Hye Yun[§], Weontae Lee[§], and Eok-Soo Oh^{‡1}

From the [‡]Department of Life Sciences, the Research Center for Cellular Homeostasis, Ewha Womans University, Seoul 120-750, Korea and [§]Department of Biochemistry, College of Life Science and Biotechnology, Yonsei University, Seoul 120-749, Korea

Background: Syndecans form non-covalently linked homodimers through their highly conserved transmembrane domains.

Results: Syndecan-2 and -4 exhibit heteromolecular interaction, and this interaction regulates syndecan-mediated cellular functions.

Conclusion: Hetero-oligomerization produces distinct syndecan functions.

Significance: Our finding offers new insights into the underlying signaling mechanisms of syndecans.

Syndecans, a family of transmembrane heparansulfate proteoglycans, are known to interact through their transmembrane domains to form non-covalently linked homodimers, a process essential for their individual functions. Because all syndecan transmembrane domains are highly conserved and thus might mediate interactions between different members of the syndecan family, we investigated syndecan interactions in detail. All recombinant syndecan-2 and -4 protein variants containing the transmembrane domain formed not only sodium dodecyl sulfate (SDS)-resistant homodimers but also SDS-resistant heterodimers. Biochemical and structural data revealed that recombinant syndecan-2 and -4 formed intermolecular interactions *in vitro*, and the GXXXG motif in transmembrane domain mediated this interaction. When exogenously expressed in rat embryonic fibroblasts, syndecan-2 interacted with syndecan-4 and vice versa. Furthermore, bimolecular fluorescence complementation-based assay demonstrated specific hetero-molecular interactions between syndecan-2 and -4, supporting hetero-oligomer formation of syndecans *in vivo*. Interestingly, hetero-oligomerization significantly reduced syndecan-4-mediated cellular processes such as protein kinase C α activation and protein kinase C α -mediated cell adhesion as well as syndecan-2-mediated tumorigenic activities in colon cancer cells such as migration and anchorage-independent growth. Taken together, these data provide evidence that hetero-oligomerization produces distinct syndecan functions and offer insights into the underlying signaling mechanisms of syndecans.

Syndecans are a family of evolutionarily conserved transmembrane heparan sulfate proteoglycans that participate in the regulation of a broad range of adhesion-mediated cell functions (1). Structurally, they are composed of an extracellular domain, a single transmembrane domain, and a short cytoplasmic

domain. The extracellular domain of syndecans interacts with a number of ligands in the extracellular matrix through heparan sulfate chains and interactions that activate syndecans and lead to their regulation of receptor signaling events (1–3). The transmembrane domain is composed of 25 hydrophobic amino acid residues that are responsible for the molecular interaction that causes homo-oligomerization of syndecan core proteins (4, 5). The strictly conserved GXXXG motif is involved in these interactions and induces homodimer formation of syndecans, even in the presence of the strong anionic detergent sodium dodecyl sulfate (SDS), a property known as SDS-resistant dimerization (4). The cytoplasmic domain contains two highly conserved C1 and C2 regions separated by a variable region. The intervening variable region, which is distinct for each syndecan member yet conserved across species, promotes syndecan-specific intracellular functions.

Four members of the syndecan family have been identified in vertebrates (6), and their expression is highly regulated in complex patterns in individual cell types and tissues and at different developmental stages (7). Syndecan-1 is the predominant family member in epithelial cells, syndecan-2 is abundantly expressed in fibroblasts and mesenchymal cells, syndecan-3 abounds in neuronal cells, and syndecan-4 is widely expressed (7). Notably, the expression of syndecans is altered in pathological situations such as wound healing and neoplastic transformation. In particular, altered expression of syndecan family members has been reported in various human tumors, including lung, breast, and colorectal cancers (8, 9).

Transduction of signals through non-covalent dimerization/oligomerization of cell surface receptor proteins containing a single transmembrane domain in response to ligand binding is a well-recognized signaling paradigm (10). As is the case for many other cell surface receptors, homodimerization/oligomerization is the first step in the activation of syndecan signaling, and the syndecan transmembrane domain plays a major role in this process (5). Given the high pairwise sequence identity of syndecan transmembrane domains (48–72%) and the conservation of the GXXXG dimerization motif, all syndecan transmembrane domains might be expected to mediate heteromeric oligomerization as well as homomeric interactions. A previous study using synthetic peptides corresponding to the transmem-

* This work was supported by the National Research Foundation of Korea, funded by Korea government (MSIP) Grants 2012R1A5A1048236 and 2013R1A2A2A01013565. The authors declare that they have no conflicts of interest with the contents of this article.

¹ To whom correspondence should be addressed: Dept. of Life Sciences, Ewha Womans University, Daehyun-dong, Seodaemun-Gu, Seoul 120-750 Korea. Tel.: 82-2-3277-3761; Fax: 82-2-3277-3760; E-mail: OhES@ewha.ac.kr.

Syndecan Hetero-oligomerization

brane domain of each syndecan paralog has measured the association tendencies of the syndecan transmembrane domain (11). This study found that the transmembrane domains of the four syndecan paralogs self-associate to very different degrees and are capable of a distinct set of heteromeric interactions, independent of extracellular and cytoplasmic domains (11). However, whether full-length syndecans form heterodimers and, if so, how this heterodimerization is involved in regulating syndecan functions, have not been investigated. Here, we demonstrate that transmembrane domain-mediated hetero-oligomerization of syndecan-2 with syndecan-4 occurs *in vitro* and in intact cells and regulates syndecan-mediated signal transduction, producing unique functional outcomes.

Experimental Procedures

Antibodies and Materials—Monoclonal antibodies against glutathione *S*-transferase (GST), His, HA, β -actin, RhoA, integrin β 1, extracellular signal-regulated kinase (ERK) and phospho-ERK, and polyclonal antibodies against Myc and protein kinase C α (PKC α) were purchased from Santa Cruz Biotechnology (Santa Cruz, CA). A monoclonal antibody to α -actinin was purchased from Sigma, and a polyclonal antibody to syndecan-4 was purchased from Santa Cruz Biotechnology. Monoclonal antibodies to paxillin and Rac1 were purchased from Upstate Biotechnology, Inc. (Lake Placid, NY), and a monoclonal anti-syndecan-2 antibody was produced by AdipoGen Inc. (Incheon, Korea). Fibronectin was from Upstate Biotechnology, Inc. (Lake Placid, NY). Transient transfection was carried out using Effectene (Qiagen, Hilden, Germany) as described by the provided protocol.

Cell Culture—Rat embryonic fibroblasts (REFs)² were maintained in α -modified Eagle's medium (α -MEM; Gibco) supplemented with 5% (v/v) fetal bovine serum (FBS), penicillin (1000 units/ml), and streptomycin (1 mg/ml). HT29 cells were maintained in McCoy's 5a medium (Welgene, Daegu, Korea); HEK293T and B16F10 cells were maintained in Dulbecco's modified Eagle's medium (Welgene) supplemented with 10% (v/v) FBS, penicillin (100 units/ml) and streptomycin (10 μ g/ml, Gibco). All cell lines were maintained at 37 °C in a humidified 5% CO₂ atmosphere.

Expression and Purification of Recombinant GST-Syndecan Core Proteins—cDNAs of rat syndecan-1, -2, and -4 encoding full-length proteins (SDC1, SDC2, SDC4), the transmembrane domain and cytoplasmic domain (2TC, 4TC), the extracellular domain (2E), the transmembrane domain (2T), the cytoplasmic domain (2C), or the transmembrane domain containing four additional flanking extracellular amino acid residues (KRTE and ERTE) in the extracellular domain and the cytoplasmic domain (2eTC, 4eTC) were synthesized by polymerase chain reaction (PCR) and subcloned into the GST expression vector pGEX-5X-1 (Amersham Biosciences). Mutations were generated by PCR-based site-directed mutagenesis (Stratagene, La Jolla, CA) and confirmed by DNA sequence analysis of the resulting plasmids. These constructs were used to transform

Escherichia coli DH5 α , and the expression of GST fusion proteins was induced by incubation with 1 mM isopropyl- β -D-thiogalactopyranoside for 4 h at 37 °C. The fusion proteins were purified with glutathione-agarose beads (GE Healthcare) as described previously (5).

Purification of Recombinant His-syndecan Proteins by GST-Syndecan-bound Glutathione-agarose Bead Affinity Chromatography—cDNA encoding the entire rat syndecan-2 or -4 core protein was subcloned into the His expression vector pET32a+ (Novagen, Madison, WI), and the expression of fusion proteins in *E. coli* BL21 was induced by incubating with 0.3 mM isopropyl- β -D-thiogalactopyranoside at 25 °C for 16 h. Proteins were released by lysing *E. coli* cells with lysis buffer (20 mM Na₂HPO₄, pH 8.0, 150 mM NaCl, 5 mM β -mercaptoethanol, 1% Triton X-100) and sonicating on ice for 1 m. After removing insoluble material by centrifugation at 13,000 \times *g* for 30 min at 4 °C, the supernatants containing His-syndecan fusion proteins were applied to a glutathione-agarose column containing pre-bound GST-syndecans. The column was washed 3 times, and bound proteins were eluted with elution buffer (50 mM Tris-HCl, pH 8.0, 5 mM reduced glutathione). Fractions were analyzed by SDS-PAGE followed by Coomassie Blue staining and Western blotting using antibodies against GST, His, syndecan-2, and syndecan-4.

Sample Preparation for the Nuclear Magnetic Resonance (NMR) Experiment—cDNA encoding the rat transmembrane domain of syndecan-2 and syndecan-4 were subcloned into the His-thioredoxin expression vector pET32a+, and the enterokinase enzyme recognition site, DDDDK, was inserted between His-thioredoxin tag and target protein. Fusion protein expression was induced in *E. coli* BL21(DE3) cells with 1 mM isopropyl- β -D-thiogalactopyranoside in optical density values of 0.55 at 600 nm and overexpressed at 25 °C for 18 h. Harvested cell pellet was lysed with lysis buffer (50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 5 mM β -mercaptoethanol) and sonicated on ice for 1 m. After centrifugation at 13,000 \times *g* for 30 min, supernatant was removed, and insoluble precipitant was used for the resolubilization step using refolding buffer (50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 5 mM β -mercaptoethanol, 1% *n*-dodecyl β -D-maltoside (Affymetrix). Resolubilized fraction was applied to Ni-NTA affinity column. The column was washed two times using 50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 5 mM β -mercaptoethanol, 20 mM imidazole, and 0.1% *n*-dodecyl β -D-maltoside, and target protein was eluted in 50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 5 mM β -mercaptoethanol, 250 mM imidazole, 0.05% *n*-dodecyl β -D-maltoside. After desalting column work using PD-10 (GE Healthcare), enterokinase enzyme was treated to remove the tag at 37 °C for 12 h. Nonspecific impurity and His-thioredoxin tag was removed using additional Ni-NTA affinity column work. Final target protein was dialyzed in 20 mM ammonium bicarbonate and lyophilized. For the preparation of NMR samples, M9 minimal media and isotopes such as [¹³C]glucose and ¹⁵NH₄Cl were used during cell culture. All NMR samples are prepared in 10 mM Na₂HPO₄, pH 7.0, 4 mM DTT, 1 mM sodium azide, 100 mM dodecylphosphocholine, 10% D₂O.

NMR Experiment and Data Analysis—All NMR spectra were obtained at 318 K on a Bruker DRX 900 MHz spectrometer with

²The abbreviations used are: REF, rat embryonic fibroblast; Ni-NTA, nickel-nitrilotriacetic acid; EGFR, epidermal growth factor receptor; PAK-PBD, p21 binding domain of PAK.

Cryoprobe™ and processed and analyzed using NMRPipe/NMRDraw and the Sparky program. Sequential resonance assignment was executed using ¹H,¹⁵N heteronuclear single quantum coherence, three-dimensional HNCACB, CBCA-CONH, and HNCO. NMR titration experiment was performed using ¹⁵N-labeled syndecan-4 and unlabeled syndecan-2 transmembrane domain using different molar ratios (1:1, 1:2, and 1:3). Chemical shift change values were calculated using the equation $\Delta\delta_{AV} = ((\Delta\delta_{1H})^2 + (\Delta\delta_{15N}/5)^2)^{1/2}$.

Immunoprecipitation—For the immunoprecipitation of HA-tagged syndecans, REFs transfected with the indicated cDNAs were lysed with 1% Nonidet P-40 lysis buffer (50 mM Tris, pH 8.0, 150 mM NaCl, 1% Nonidet P-40, 1 mM EDTA, 2 mM Na₃VO₄, and 10 mM NaF) containing a protease inhibitor mixture. Individual cell lysates were mixed for 30 min on ice, and protein mixtures were incubated with anti-HA affinity Matrix (Roche Applied Science) for 2 h at 4 °C. Immune complexes were collected by centrifugation, washed 3 times with lysis buffer, eluted with 1 M NaCl, and analyzed by slot blotting using the indicated antibodies. For the immunoprecipitation of endogenous syndecan proteins, cells were lysed in 1% Nonidet P-40 lysis buffer containing a protease inhibitor mixture. Each cell lysate was incubated with anti-syndecan antibodies for 2 h at 4 °C followed by incubation with protein G-Sepharose beads for 1 h. Immune complexes were collected by centrifugation, washed 3 times with lysis buffer, and eluted with 1 M NaCl. To remove the IgG of eluted proteins and extract the syndecan antibodies, eluted proteins were incubated with DEAE-Sepharose beads (GE Healthcare) for 2 h at 4 °C in 50 mM Tris, pH 8.0, containing 4 M urea, 200 mM NaCl, 0.1% Tween 20, 5 mM EDTA, and 2 mM PMSF, washed with 30 mM sodium acetate, pH 4.0, containing 4 M urea, 200 mM NaCl, 0.1% Tween 20, 25 mM EDTA, and 2 mM PMSF, eluted with 15 mM sodium acetate buffer, pH 4.0, containing 4 M guanidine HCl and 0.1% Tween-20, and then analyzed by slot blotting using the indicated antibodies. For Western blotting of immunoprecipitates, cells were lysed with 1% radioimmune precipitation assay lysis buffer (50 mM Tris, pH 8.0, 150 mM NaCl, 1% Nonidet P-40, 0.1% SDS, 0.5% sodium deoxycholate, 10 mM NaF, 2 mM Na₃VO₄) containing a protease inhibitor mixture. Cell lysates were incubated with anti-syndecan-4 antibody for 2 h at 4 °C followed by incubation with protein A-Sepharose beads for 1 h. Immune complexes were collected by centrifugation, washed 3 times with lysis buffer, and eluted with 1 M NaCl. Glycosaminoglycan chains were removed by incubating samples with a pH 1.5 nitrous acid solution as previously described (12) and analyzed by Western blotting using an anti-syndecan-2 antibody.

Bimolecular Fluorescence Complementation—The vector constructs encoding fragments of pECFP were kindly provided by Dr. Minsoo Kim (University of Rochester Medical Center). In the current study the sequences encoding pECFP amino acid residues 1–154 (CN) or 155–238 (CC) were fused to the 3' ends of the coding regions of syndecans by conventional restriction enzyme-based cloning; the resulting constructs generated are depicted schematically in Fig. 4E. For bimolecular fluorescence complementation assays, HT29 cells were transiently co-transfected with the indicated cDNAs then fixed for 5 min in 4% paraformaldehyde, washed with PBS, and imaged on a Zeiss

Axiovert 200 M inverted microscope equipped with a Zeiss 510 META confocal head using a C-Apochromat 40×/1.2-W objective lens and a 458-nm argon laser light and Meta detector (462.6–516.1 nm) for pECFP excitation and emission.

In Vitro PKC Assay—For nonradioactive Peptag assays, Peptag C1 peptide (Promega, Madison, WI) was incubated in reaction buffer (20 mM HEPES, pH 7.4, 1 mM DTT, 10 mM MgCl₂, 1 mM ATP) containing phosphatidylinositol 4,5-bisphosphate (50 μM) in a final volume of 25 μl for 30 min at 30 °C. The reactions were stopped by heating to 95 °C for 10 min. The samples were separated on a 0.8% agarose gel at 100 V for 15 min. Phosphorylated peptides migrated toward the cathode (+), whereas non-phosphorylated peptides migrated toward the anode (–). The gel was photographed on a transilluminator.

Centrifugal Detachment Assay—Cell-substratum adhesiveness was quantified using an inverted centrifugal detachment assay. Fibronectin was diluted in serum-free medium (10 μg/ml), added to the plates, and incubated at 25 °C for 1 h. The plates were then washed with PBS and blocked with 0.2% heat-inactivated BSA for 1 h. After washing with PBS, cells were incubated for additional 2 h at 37 °C in 5% CO₂. After removing unattached cells, plates were then filled with serum-free medium, sealed with Parafilm, and centrifuged inverted for 5 min at 300 × g at room temperature using a large capacity tabletop centrifuge (Hanil science industrial, Korea). After the centrifugation, retained cells were counted using a hemocytometer.

Cellular Fractionation—After washing twice with PBS, a hypo-osmotic solution (20 mM Tris/HCl, pH 7.5, 2 mM 2-mercaptoethanol, 5 mM EGTA, 2 mM EDTA) containing a protease inhibitor mixture was added to the culture plates. Cells were subsequently scraped off the plates and homogenized on ice. The homogenate was centrifuged at 13,000 × g for 15 min at 4 °C. The membrane fraction was collected by solubilizing the remaining pellet in radioimmune precipitation assay buffer containing a protease inhibitor mixture and then centrifuged at 13,000 × g for 15 min at 4 °C. Equal amounts of lysates were resolved by SDS-PAGE, transferred onto PVDF membranes, and probed with the indicated antibodies.

Rac and Rho Activity Assay—GST-PAK-PBD binding assays were performed essentially as described previously (13). Briefly, the p21 binding domain of PAK1 (PBD) was expressed in *E. coli* as a GST-PAK-PBD fusion protein, purified using glutathione-agarose beads, and added to cell lysates. Bound proteins were collected by centrifugation and suspended in SDS sample buffer. Proteins were fractionated by SDS-PAGE and transferred to PVDF membranes, and the amount of precipitated Rac1 was estimated by Western blotting with an anti-Rac1 antibody. Rho activity was measured in a pull-down assay using the Rho binding domain from Rhotekin. Equal volumes of lysates were incubated with GST-Rho binding domain beads at 4 °C for 2 h, after which the beads were washed 4 times with lysis buffer, and bound RhoA proteins were detected by Western blotting using a monoclonal antibody against RhoA.

Immunofluorescence—Cells were transfected with the indicated cDNAs, fixed with 3.5% paraformaldehyde in PBS at room temperature for 5 min, permeabilized with 0.1% Triton X-100 in PBS for 10 min, blocked with 0.5% bovine serum albu-

Syndecan Hetero-oligomerization

men in PBS for 1 h, and incubated with the indicated antibodies. The slides were mounted with VECTASHIELD Mounting Medium (Vector Laboratories, Burlingame, CA) and imaged using a fluorescence confocal microscope (Carl Zeiss, Göttingen, Germany).

Monitoring Cell Spreading and Migration—Cell spreading and migration were monitored using the xCELLigence system (Roche Diagnostics). For cell spreading, E-plate 16 assemblies (Roche Diagnostics) were coated with fibronectin (10 $\mu\text{g}/\text{ml}$) and seeded with cells (1.5×10^4 cells/well). Each plate was then assembled on the RTCA DP Analyzer, and data were gathered at 5-min intervals for 5 h at 37 °C, 5% CO_2 . The data were analyzed using the provided RTCA software. For cell migration, a CIM-plate 16 (Roche Diagnostics) was used. The lower chambers were filled with fresh medium (160 $\mu\text{l}/\text{well}$) containing 10 $\mu\text{g}/\text{ml}$ fibroblast growth factor-2 and 10% FBS, and the upper chambers were filled with medium (30 $\mu\text{l}/\text{well}$) containing 5% FBS. Plates were incubated at 37 °C, 5% CO_2 for 1 h, and background was measured using the RTCA DP Analyzer. Transfected cells were added to each well, and the plate was incubated at 25 °C. After 30 min, the CIM-plate was assembled onto the RTCA DP Analyzer, and cell migration was assessed at 15-min intervals for 24 h at 37 °C, 5% CO_2 . The obtained data were analyzed using the provided RTCA software.

Transwell Migration Assay—The lower surface of Transwell inserts (Costar) was coated with gelatin (10 $\mu\text{g}/\text{ml}$), and the membranes were allowed to dry for 1 h at room temperature. The Transwell inserts were assembled into a 24-well plate, and the lower chamber was filled with McCoy's 5a media containing 10% FBS and fibroblast growth factor-2 (10 $\mu\text{g}/\text{ml}$). Cells (5×10^5) were added to each upper chamber, and the plate was incubated at 37 °C in a 5% CO_2 incubator for 30 h. Migrated cells were stained with 0.6% hematoxylin and 0.5% eosin and counted.

Anchorage-independent Growth in Soft Agarose—Each well of a 6-well culture plate was coated with 3 ml of bottom agar mixture (McCoy's 5a, 10% FBS, 0.6% agar). After the bottom layer had solidified, 2 ml of top agar mixture (McCoy's 5a, 10% fetal bovine serum, 0.3% agar) containing 2×10^6 cells was added to each well, and the cultures were incubated at 37 °C in a 5% CO_2 atmosphere. Every 5 days, normal growth medium was gently layered over the cultures. Colony formation was monitored daily with a light microscope. Colonies in soft agar were photographed with a digital camera after incubation for 21 days.

Statistical Analysis—Data are represented as the mean from at least three independent experiments. Statistical analysis was performed using an unpaired Student's *t* test. A *p* value less than 0.01 was considered statistically significant.

Results

Syndecans Form SDS-resistant Heterodimers through Transmembrane Domain Interactions—To examine interactions of syndecan molecules, we first investigated the well known property of SDS-resistant dimer formation. Several recombinant syndecan-2 and -4 constructs were generated and expressed in *E. coli* as GST-fusion proteins (Fig. 1A). Recombinant syndecan proteins were induced with isopropyl- β -D-thiogalactopyrano-

side and purified. wild type syndecans (SDC2 or SDC4) were mixed with deletion mutants of syndecan-2 or -4 containing transmembrane and cytoplasmic domains (2TC and 4TC, respectively). Consistent with a previous report (5), recombinant syndecan-2 and -4 proteins were resolved as SDS-resistant homomeric dimers and monomers (Fig. 1B, left). Interestingly, mixtures of syndecan-2 or -4 proteins with deletion mutants showed only one extra band between the monomer- and dimer-migrating positions, corresponding to a heteromeric molecular mass of ~ 90 kDa (Fig. 1B, right; Tables 1 and 2). Consistently, His-tagged wild type syndecan proteins showed strong SDS-resistant heterodimers (Fig. 1C). Furthermore, His-tagged syndecan-2 formed SDS-resistant dimer with the tag-free recombinant syndecan-4, and the tag-free recombinant syndecan-2 also formed SDS-resistant dimer with His-tagged syndecan-4 (Fig. 1D). In addition, recombinant syndecan-1 formed SDS-resistant heterodimers with syndecan-2 or -4 (data not shown). Collectively, these data indicate that syndecans form strong SDS-resistant heterodimers as well as homodimers *in vitro*.

Because the highly conserved transmembrane domain of syndecan core proteins is responsible for the formation of stable, non-covalent homodimeric complexes, we next investigated whether the transmembrane domain is necessary for heterodimerization. As expected, wild type syndecan-4 formed SDS-resistant heterodimers with the deletion mutant containing the transmembrane domain but not formed with those containing either the cytoplasmic domain (2C) or the extracellular domain (2E) alone (Fig. 1E). To further demonstrate the role of the transmembrane domain in heterodimerization, we used syndecan mutants defective in transmembrane domain-mediated homo-oligomerization (2EPT2C, 4EPT2C, 2GL, or 4GL; see Fig. 1A and Ref. 5). Oligomerization-defective mutations abolished SDS-resistant heterodimer formation by syndecan-2 or -4 (Fig. 1F, bottom), supporting the notion that the transmembrane domain is crucial for SDS-resistant heterodimerization of syndecans.

Transmembrane Domain Mediates the Intermolecular Interaction between Syndecans—To investigate hetero-molecular interactions of the syndecan transmembrane domain, we immobilized GST-syndecan fusion proteins on glutathione beads and used them to purify their respective binding partners among various His-tagged syndecans (Fig. 2A). We found that His-tagged syndecan-2 could be purified from GST-syndecan-2 immobilized on glutathione beads (Fig. 2C), whereas GST protein alone could not (Fig. 2B), supporting the homomeric association of syndecan-2. Notably, His-syndecan-4 fusion protein could also be purified from glutathione bead-immobilized GST-syndecan-2, probably through transmembrane domain-mediated hetero-molecular interactions (Fig. 2D). In the reciprocal experiment, His-syndecan-2 fusion protein was purified from GST-syndecan-4 immobilized on glutathione beads (Fig. 2E). These data indicate that, in addition to well known intermolecular interactions involving syndecans of a single type, hetero-molecular interactions occur between different syndecan members.

We next carried out a set of NMR experiments to examine the intermolecular interaction between syndecan-2 and -4 transmembrane domains in solution. As shown in Fig. 3A, in




FIGURE 1. Syndecans form SDS-resistant heterodimers through their transmembrane domains. *A*, schematic representation of syndecan-2 (SDC2) and syndecan-4 (SDC4) core proteins. The extracellular domain (*E*), the transmembrane domain (*T*), the cytoplasmic domain (*C*), and four amino acid residues in the membrane flanking region (ERTE, KRTE) are shown. Two of the conserved glycine residues (*) replaced with leucine residues are indicated by arrows in 2GL and 4GL. The transmembrane domain of syndecan-2 and -4 was replaced with the single transmembrane domain of PDGF receptor to yield 2EPT2C and 4EPT4C, respectively. *B*, recombinant GST-syndecan core proteins were purified with glutathione-agarose beads as described under “Experimental Procedures” (left). Purified recombinant proteins (SDC2 or SDC4) were mixed with the corresponding mutants (2TC or 4TC) for 30 min on ice, separated by SDS-PAGE, and stained with Coomassie Blue (right). Molecular mass markers in kilodaltons are shown. Migration positions of SDS-resistant syndecan-2 dimers (●●) and monomers (○), syndecan-4 dimers (■) and monomers (□), syndecan-2TC dimers (○○) and monomers (○), and syndecan-4TC dimers (□□) and monomers (□) are indicated. *C*, His-tagged thioredoxin-syndecan proteins were affinity-purified over Ni-NTA columns, and purified recombinant proteins (SDC2 or SDC4) were mixed with the corresponding mutants (2eTC or 4eTC) for 30 min on ice. Mixtures were separated by SDS-PAGE and stained with Coomassie Blue. Migration positions of SDS-resistant syndecan-2 dimers (●●) and monomers (○), syndecan-4 dimers (■) and monomers (□), syndecan-2eTC dimers (○○) and monomers (○), syndecan-4eTC dimers (□□) and monomers (□), heterodimers of syndecan-2 and syndecan-4eTC (●○), and heterodimers of syndecan-4 and syndecan-2eTC (○□) are indicated. *D*, His-tagged thioredoxin-syndecan proteins including the tobacco etch virus cleavage site were incubated with tobacco etch virus protease. Each tag-free recombinant syndecan protein was mixed with His-tagged syndecan proteins, separated by SDS-PAGE, and stained with Coomassie Blue. Migration positions of His-tagged syndecan dimers (■) and monomers (□) and cleaved syndecan dimers (□□) and monomers (□) are indicated. *E*, purified recombinant syndecan-4 protein was mixed with the indicated recombinant proteins, separated by SDS-PAGE, and stained with Coomassie Blue. Migration positions of SDS-resistant wild type syndecan dimers (■) and monomers (□), heterodimers (■○), transmembrane domain mutant dimer (○○), extracellular domain mutant (<) and cytoplasmic domain mutant (◄) are indicated. *F*, recombinant GST-syndecan core proteins (SDC2 and SDC4) and oligomerization-defective mutants (2GL, 2EPT2C, 4GL, and 4EPT4C) were purified using glutathione-agarose beads (top). Purified recombinant proteins (2TC or 4TC) were mixed with the indicated recombinant proteins, and mixtures were separated by SDS-PAGE (bottom).

TABLE 1

The extra band of GST-tagged recombinant protein mixture was identified to be heparan sulfate proteoglycan core protein (syndecan-2 and -4) by mass spectrometry

LC-MS/MS sequencing results of 10 peptides, all of which are the matched peptides that led to sequence coverage of 44%, provided identification of syndecan-2 core protein (*Rattus norvegicus*). Nominal molecular weight (M_r) was 22,135.

Calculated mass	Expected mass	Delta mass	Score	Sequence
M_r	M_r			
978.5134	978.5472	0.0338	27	RKPSSAAYQKA
1209.5401	1209.5760	0.0360	23	KEFEISEAEEKQ
1383.7358	1383.7636	0.0279	111	KGSPDLTTSQLIPRI
1689.8859	1689.9098	0.0240	12	RISLTSAAPEVETMTLKT
1689.8859	1689.9247	0.0388	43	RISLTSAAPEVETMTLKT
1705.8808	1705.8818	0.0011	44	RISLTSAAPEVETMTLKT + Oxidation (Met)
1705.8808	1705.9086	0.0279	2	RISLTSAAPEVETMTLKT + Oxidation (Met)
1705.8808	1705.9092	0.0285	8	RISLTSAAPEVETMTLKT + Oxidation (Met)
3751.5367	3751.6066	0.0699	40	KDMLDSSSIEEASGLYPIDDDYSSASGSGAYEDKG
4511.8969	4512.0257	0.1288	85	RAELTSDKMDMLDSSSIEEASGLYPIDDDYSSASGSGAYEDKG + Oxidation (Met)

the presence of syndecan-2 transmembrane domain, the chemical shift perturbation of glycine residues of syndecan-4 transmembrane domain were observed, indicating the participation of GXXXG motif in the intermolecular interaction. In addition, evident chemical shift perturbations were observed in Ile-155, Gly-157, Gly-158, Val-159, Val-160, Val-166, and Phe-167 residues in the transmembrane domain of syndecan-4 upon syn-

decn-2 interaction (Fig. 3B). Particularly, the most dramatic perturbations were observed in the residues, Gly-157, Val-159, and Val-160 residues (Fig. 3C). The intensity ratio between syndecan-4 homodimer and heterodimer of syndecan-2 and -4 was 1:3 based on peak intensities of the heteronuclear single quantum correlation spectrum (Fig. 3D), implying that the molecular interaction of syndecan-4 transmembrane domain to synde-

Syndecan Hetero-oligomerization

TABLE 2

The extra band of GST-tagged recombinant protein mixture was identified to be heparan sulfate proteoglycan core protein (syndecan-2 and -4) by mass spectrometry

LC-MS/MS sequencing results of 2 peptides, all of which are the matched peptides that led to sequence coverage of 19% provided identification of syndecan-4 core protein (*R. norvegicus*). Nominal mass (M_r) is 21,948.

Calculated mass	Expected mass	Delta mass	Score	Sequence
M_r 1612.7944	M_r 1612.8196	0.0252	38	RETEVIDPQDLLEGRY
2752.4548	2752.4986	0.0438	20	RTFFEVISPLVPLDNHIPPENAPQGIKRV




FIGURE 2. Syndecans form both homomeric and heteromeric associations with partner syndecans. *A*, scheme for purifying His-syndecan proteins using GST-syndecans immobilized on glutathione beads. #1, cell lysates from *E. coli* expressing GST-syndecan fusion proteins were applied to a column of glutathione-agarose beads, and beads were washed with lysis buffer containing 1% Triton X-100. #2, cell lysates from *E. coli* expressing His-syndecan fusion proteins were applied to a column of glutathione-agarose beads containing bound GST-syndecan fusion proteins, and beads were washed with His lysis buffer containing 1% Triton X-100. #3, bound materials were eluted with elution buffer containing reduced glutathione. *B* and *C*, total lysates of cells expressing His-tagged syndecan-2 were passed over a column of glutathione-agarose bound to GST (*B*) or GST-syndecan-2 (*C*), washed three times, and eluted. Each fraction was analyzed by SDS-PAGE followed by Coomassie Blue staining or Western blotting using the indicated antibodies. *D* and *E*, total lysates of cells expressing His-tagged syndecan-4 (*D*) or His-tagged syndecan-2 (*E*) were passed over a column of glutathione-agarose bound to GST-syndecan-2 (*D*) or GST-syndecan-4 (*E*), washed three times, and eluted. Each fraction was analyzed by SDS-PAGE followed by Western blotting using the indicated antibodies.

can-2 is stronger than that of syndecan-4. Taken together, these data confirm that syndecan-2 and -4 form heteromeric interactions *in vitro* and that the transmembrane domain mediates the intermolecular interaction between syndecans.

Syndecan-2 Interacts with Syndecan-4—Because syndecans are modified with highly negatively charged glycosaminoglycan chains that may disrupt their interactions through charge repulsion, the hetero-oligomerization characteristics of endogenously expressed syndecans may differ from those of recombinant proteins. To investigate the intermolecular interactions of syndecans at the cellular level, we separately expressed HA- or Myc-tagged syndecan-2 or -4 in rat embryonic fibroblasts, and equal amounts of total cell lysate from REF-HA-syndecan-2 or REF-HA-syndecan-4 cells were mixed with those from REF-Myc-syndecan-2 or REF-Myc-syndecan-4 cells. Each mixture of total cell lysates was then immunoprecipitated with an anti-HA antibody, and the immunoprecipitate was analyzed by slot blotting using an anti-Myc antibody. As shown in Fig. 4A,

HA-tagged syndecan-2 was co-immunoprecipitated with Myc-tagged syndecan-2 (Fig. 4A, bottom, left). A similar interaction of HA-tagged syndecan-4 and Myc-tagged syndecan-4 was also observed (Fig. 4A, bottom, right), supporting homo-oligomerization of both syndecan-2 and -4. Notably, HA immunocomplexes from the mixed lysates from REF-HA-syndecan-2 and REF-Myc-syndecan-4 cells or REF-HA-syndecan-4 and REF-Myc-syndecan-2 cells were immunoreactive to anti-Myc antibodies, indicating the hetero-oligomeric interactions of syndecan-2 with syndecan-4 (Fig. 4B, bottom). In addition, both slot blotting (Fig. 4C) and Western blotting (Fig. 4D) with appropriate anti-syndecan antibodies showed co-immunoprecipitation of endogenous syndecan-2 and -4, suggesting that syndecan-2 and -4 form both homomeric and heteromeric interactions in intact cells.

To further demonstrate the interaction of syndecan-2 with syndecan-4 at the cellular level, we employed a bimolecular fluorescence complementation assay. This protein-protein interaction technique relies on the ability of molecular fragments of a fluorescent protein, in this case fragments of pECFP (enhanced cyan fluorescent protein) fused to syndecan-2 and -4 (Fig. 4E), to re-associate and refold into a fluorescent structure. Cells were co-transfected with CN and CC fusion constructs of pECFP with syndecan-2 (2CN and 2CC), syndecan-4 (4CN and 4CC), and subsequently analyzed by laser-scanning confocal microscopy for fluorescence complementation. Under each co-transfection condition, fluorescence complementation of both 2CC and 2CN and of 4CC and 4CN was detected at the cell surface (Fig. 4F). Similarly, fluorescence complementation of both 2CC and 4CN and of 4CC and 2CN was detected (Fig. 4F), indicating homomeric association of both syndecan-2 and syndecan-4 and heteromeric interactions of syndecan-2 and -4. However, we failed to detect complementation of fluorescence between wild type syndecan and oligomerization-defective mutants (Fig. 4F). Collectively, these data strongly suggest that, when expressed in the same cell, syndecan-2 and -4 are capable of both homo-oligomerization and hetero-oligomerization.

Hetero-oligomerization Inhibits Syndecan-regulated Molecular Events That Depend on Homo-oligomerization—Because homo-oligomerization is crucial for syndecan functions, hetero-oligomer formation, which inevitably decreases homo-oligomerization, might result in inhibition of known syndecan functions. Previously, we and others have shown that oligomerization of the syndecan-4 cytoplasmic domain is critical for the interaction with and activation of PKC α (5, 14, 15). Solutions of purified wild type syndecan-2, purified 4eTC (a syndecan-4 deletion mutant containing the transmembrane domain with a 4-amino acid extracellular region and the cytoplasmic domain), or a mixture of 4eTC and wild type syndecan-2 were resolved by




FIGURE 3. **Syndecan-4 transmembrane domain interacts with syndecan-2 transmembrane domain.** A and B, ¹⁵N-labeled syndecan-4 transmembrane domain was titrated with non-labeled syndecan-2 transmembrane domain up to 1:3 molar ratio, and syndecan-4 and syndecan-4-syndecan-2 complex were displayed as red and green. C, NMR chemical shift perturbation map of the syndecan-4 transmembrane domain with syndecan-2 transmembrane domain titration. The bar diagram reveals the chemical shift changes at syndecan-4 transmembrane domain: syndecan-2 transmembrane domain molar ratio of 1:3. Chemical shift change values were calculated using the equation $\Delta\delta_{AV} = ((\Delta\delta_{1H})^2 + (\Delta\delta_{15N}/5)^2)^{1/2}$. D, heterodimerization level of syndecan-2 with syndecan-4 was calculated as the intensity ratio between the syndecan-4 that interacted with syndecan-2 and the syndecan-4 that remained according to the intensity in NMR titration.




FIGURE 4. **Syndecan-2 interacts with syndecan-4 in intact cells.** A and B, REFs transiently transfected with HA- or Myc-tagged syndecans were lysed and analyzed by slot blotting (top). Individual cell lysates were mixed for 30 min on ice and protein mixtures were immunoprecipitated (IP) with HA-conjugated agarose beads. The levels of protein in each immunoprecipitate were determined by slot blotting with either anti-Myc or anti-HA antibodies (bottom). C, REF cell lysates were immunoprecipitated with the indicated antibodies as described under "Experimental Procedures." Immunocomplexes were slot-blotted with the indicated antibodies. GST-SDC2 and -SDC4 were used as positive controls for anti-syndecan antibodies. D, REF cell extracts were immunoprecipitated with anti-syndecan-2 or -4 antibodies, and immunocomplexes were treated with nitrous acid to degrade glycosaminoglycan chains. The levels of protein in each immunoprecipitate were determined by immunoblotting with anti-syndecan-4 or -2 antibodies. E, schematic representation of bimolecular fluorescence complementation plasmid constructs. E, extracellular domain; T, the transmembrane domain; C, the cytoplasmic domain. F, HT29 cells were transiently co-transfected with the indicated cDNAs expressing SDC-CC and SDC-CN constructs. At 24 h post-transfection, the cells were fixed, and images were captured by confocal microscopy using a filter set specific for pECFP.

Syndecan Hetero-oligomerization




FIGURE 5. Disruption of homo-oligomerization inhibits the function of individual syndecans. *A* and *B*, purified recombinant syndecan-2 proteins were mixed with the syndecan-4 deletion mutant 4eTC for 30 min on ice and resolved by SDS-PAGE. Half of the gel was stained with Coomassie Blue (*left*). Proteins in the remaining gel half were transferred to PVDF membranes (*right*), after which membranes were incubated with B16F10 melanoma cell lysates, and interacting proteins were analyzed by Western blotting. Molecular mass markers, in kilodaltons, are shown. Migration positions of SDS-resistant syndecan-2 dimers (●●), -4eTC dimers (□□), and heteromeric syndecan (□●) are indicated. *C*, purified recombinant syndecan-2 protein was mixed with the 4eTC, and PKC activity assays were performed (*top*). Molecular mass markers, in kilodaltons, are shown. Migration positions of SDS-resistant syndecan-2 dimers (●●), -4eTC dimers (□□), and heteromeric syndecan (□●) are indicated. *D*, purified recombinant proteins were mixed as indicated, and gel overlay assays were performed as described in *A* using REF cell lysates. *E*, schematic representation of syndecan chimeras (PE2TPC) and deletion mutants (4ET, *top*). HEK293T cells were transiently co-transfected with the indicated cDNAs. After 24 h, cells were lysed, and MAPK activation was assessed using an antibody specific for phospho-ERK (pERK) followed by stripping and reprobing with an anti-ERK antibody.

SDS-PAGE. Half the gel was stained with Coomassie Blue to measure the amounts of homomeric (Fig. 5*A*, *left*, *first* and *second* lanes) and heteromeric (Fig. 5*A*, *left*, *third* lane) syndecans. Proteins in the other gel half were transferred to PVDF membranes (Fig. 5*A*, *right*); membranes were then incubated with B16F10 melanoma cell lysates, and interacting proteins were analyzed by Western blotting. Only homodimeric syndecan-4 mutants containing the cytoplasmic domain bound to PKC α (Fig. 5*A*, *right*); syndecan-2 homodimers and heterodimers of syndecan-2 and -4 did not. The increase in heterodimerization induced a significant decrease in syndecan-4 homodimers and led to a reduction in syndecan-4-PKC α interaction (Fig. 5*B*) and diminished PKC α activation (Fig. 5*C*, *top*), indicating an inhibitory effect of hetero-oligomerization on syndecan-4 function. Consistent with this, increased heterodimerization reduced the interaction of syndecan-4 with α -actinin, which is known to directly interact with the cytoplasmic domain of syndecan-4 (Fig. 5*D*).

Previous studies have shown that dimerization of PDGF receptor subsequently activates mitogen-activated protein kinase (MAPK) (16, 17). Moreover, a chimeric protein contain-

ing the transmembrane domain of syndecan fused to the extracellular and cytoplasmic domain of β -PDGF receptor is sufficient to induced MAPK activation through chimera oligomerization (5). Accordingly, we explored the potential effect of transmembrane domain-mediated hetero-oligomerization on chimera-induced MAPK activation. HEK293T cells were transiently co-transfected with the chimeras PE2TPC and syndecan mutants containing extracellular and transmembrane domains of syndecan-4 (4ET), and chimera-induced MAPK activity was analyzed by Western blotting with an anti-phospho-ERK antibody. Consistent with previous data (5), phosphorylation of ERK was increased in chimera-transfected cells compared with that in vector-transfected cells (Fig. 5*E*). Co-transfection of the chimera with greater amounts of syndecan mutants caused a reduction in MAPK activity, implying the possibility of hetero-oligomer formation through the syndecan transmembrane domain and subsequent regulation of MAPK activation. Taken together, these results suggest that disruption of homo-oligomerization inhibits the function of individual syndecans.

Hetero-oligomerization Negatively Regulates Syndecan-mediated Cellular Processes—Because syndecan-4 binding to the high affinity heparin binding domain of fibronectin drives focal adhesion through PKC α and RhoA activation during cell adhesion (Fig. 6*A*; Refs. 18–20), we examined whether the formation of hetero-oligomers affected syndecan-4-mediated signaling. Consistent with prior studies (21), a centrifugal detachment assay showed greater cell adhesion to fibronectin by REFs overexpressing vectors encoding wild type syndecan-4 compared with those expressing the empty vector (Fig. 6*B*). Whereas adhesion to fibronectin increased in cells expressing individual syndecans, co-expression of syndecan-4 and -2 inhibited syndecan-4-mediated cell attachment to fibronectin. Furthermore, co-expression of syndecan-4 and -2 reduced syndecan-4-mediated PKC α membrane localization (Fig. 6*C*, *top*) and RhoA activation (Fig. 6*D*). In contrast, co-expression of wild type syndecan-4 and syndecan-2 dimerization-defective mutants had no effect on syndecan-4-mediated PKC α localization (Fig. 6*C*, *bottom*). As expected, induction of hetero-oligomerization reduced syndecan-4-mediated focal adhesion formation (Fig. 6*E*) and cell adhesion (Fig. 6*F*). These data suggest that hetero-oligomerization negatively regulates syndecan-4-mediated cellular processes, including PKC α activation and PKC α -mediated signaling during cell adhesion.

Because syndecan-2 regulates tumorigenic activity in an oligomerization-dependent manner (Fig. 7*A*; Refs. 22–24), we investigated whether hetero-oligomerization of syndecans might affect syndecan-2-mediated signal transduction involved in the regulation of tumorigenic activity. Co-localization of syndecan-2 and -4 was observed in HT29 colon adenocarcinoma cells transiently expressing syndecan-2 and -4 (Fig. 7*B*), suggesting that the two syndecans undergo hetero-molecular interactions in cancer cells. Similar to the observed effects on syndecan-4 signaling, our results showed that induction of hetero-oligomerization through co-overexpression of syndecan-2 and -4 significantly reduced the membrane localization of Tiam1 (T-cell lymphoma invasion and metastasis 1) (Fig. 7*C*) and activation of Rac (Fig. 7*D*), crucial regulators in syndecan-2-mediated signal transduction in colon cancer cells, compared




FIGURE 6. Hetero-oligomerization regulates syndecan-4-mediated functions. *A*, scheme showing syndecan-4-mediated molecular mechanisms involved in adhesion signaling. *B*, cell-substrate adhesiveness was quantified using an inverted centrifugal detachment assay as described under “Experimental Procedures.” The results are expressed as the mean percentages of retained cells per field \pm S.E. from three independent experiments. *, $p < 0.01$. *C*, REFs were co-transfected with the indicated cDNAs, and the amount of PKC α in the membrane fraction was determined by immunoblotting with the indicated antibodies. An antibody against integrin $\beta 1$ (α -Int $\beta 1$) was used as a loading control. *D*, REFs were co-transfected with indicated cDNAs. Cell lysates were subsequently prepared and incubated with purified GST-Rhotekin-Rho binding domain proteins for 2 h. Bound proteins were collected, and GTP-bound RhoA was detected by immunoblotting with anti-RhoA antibody. *E*, transfected cells were stained with anti-paxillin antibody (*top*). *Bottom*, percentages of cells showing focal adhesions (mean \pm S.E.; vector transfectants set as 100%). The data shown are representative of three independent experiments. *, $p < 0.01$. *F*, REFs co-transfected with syndecan-4 and either vector or syndecan-2 were plated on E-plates pre-coated with fibronectin (FN). After the indicated times cell spreading was monitored and plotted using the xCELLigence system. Representative results from three independent experiments are shown.

with cells transfected with syndecan-2. Consistent with this, both syndecan-2-mediated cell migration and anchorage-independent growth were reduced in HT29 cells co-expressing syndecan-2 and -4 compared with those expressing syndecan-2 alone (Fig. 7, *E–G*), suggesting that hetero-oligomer formation inhibits syndecan-2 functions in colon cancer. Taken together, our findings indicate that hetero-oligomerization negatively regulates fundamental syndecan-mediated cellular processes.

Discussion

Syndecan core proteins are known to have the propensity to form non-covalently linked homodimers through interactions of a strictly conserved GXXXG motif in the transmembrane domain (4, 5, 25), but the potential of different syndecans to interact to form heterodimers has not been studied. Our investigation of the detailed interactions of syndecans reported here revealed that in addition to homodimer formation, all exogenously expressed syndecan-2 and -4 protein variants containing the transmembrane domain showed strong SDS-resistant heterodimer formation (Fig. 1). Consistent with this, endogenous syndecan-2 interacted with syndecan-4 (Fig. 4) notwithstanding potential repulsive forces contributed by heparan sulfate chains and steric hindrance by the large syndecan extracellular domains, both of which would tend to prevent transmembrane domain association. We also found that syndecan-1, which has a low tendency to self-associate (11), showed SDS-resistant heterodimer formation with syndecan-2 and -4.

Therefore, syndecan transmembrane domains appear to be involved in both homomeric interactions of a single syndecan type and heteromeric interactions between syndecan paralogs.

At least one syndecan is expressed in virtually all normal cells, but the expression pattern is distinct in each cell type. For example, epithelial cells express syndecan-1 but not syndecan-2. Thus, opportunities for formation of syndecan hetero-oligomers may be rare in normal cells; a notable exception is vascular smooth muscle cells, which express all four syndecan family members (26). However, under pathological conditions, syndecan expression is commonly altered. During carcinogenesis, cancer cells may undergo “receptor exchange,” in which anti-tumorigenic receptors are down-regulated with the concomitant up-regulation of pro-tumorigenic receptors. For instance, the expression of syndecan-2, which is not expressed in normal epithelial cells, is increased during colorectal carcinogenesis (22). Altered syndecan-2 expression not only indicates that receptor exchange is sufficient for induction of colon tumorigenesis, it also suggests that the altered expression of syndecan-2 subsequently increases the likelihood of syndecan hetero-oligomerization. Syndecan hetero-oligomerization creates a new paradigm for integrating regulatory mechanism of syndecans *in vivo*. For example, syndecan-2 function has been studied independently from that of syndecan-4. However, hetero-oligomer formation between syndecan-2 and -4 raises the possibility of three cooperative regulatory mechanisms with

Syndecan Hetero-oligomerization




FIGURE 7. Hetero-oligomerization regulates syndecan-2-mediated functions in colon cancer cells. *A*, overall scheme of syndecan-2-mediated signal transduction in colorectal tumorigenesis. *B*, HT29 cells were co-transfected with the indicated cDNAs and stained with antibodies against syndecan-2 and -4. *C*, HT29 cells were co-transfected with the indicated cDNAs, and the amount of Tiam1 in the membrane fraction was determined by immunoblotting with anti-Tiam1 antibody. An antibody against integrin $\beta 1$ (*αIntβ1*) was used as a loading control. *D*, purified GST-PAK-PBD protein was incubated with lysates from transfected cells for 2 h. Bound proteins were collected, and GTP-bound Rac was detected by immunoblotting with anti-Rac1 antibody. *E*, HT29 cells were transfected with the indicated cDNAs, and Transwell migration assays were performed as described under "Experimental Procedures." The data shown are representative of three independent experiments. *, $p < 0.01$. *F*, transfected cells were seeded in duplicate in the upper chamber of CIM-plates. The lower chambers were filled with medium containing 10% FBS and fibroblast growth factor-2 (10 $\mu\text{g}/\text{ml}$), and migration was monitored and plotted using the xCELLigence system. Representative results from three independent experiments are shown. *G*, HT29 cells were seeded in soft agar and allowed to grow for 21 days, and the number of viable colonies was counted. Representative results from three independent experiments are shown. *, $p < 0.01$.

in cells: those involving homo-oligomers of syndecan-2, those involving homo-oligomers of syndecan-4, and those involving hetero-oligomers of syndecan-2 and -4. We hypothesized that syndecan hetero-oligomerization might reorganize the formation of functional complexes depending on the paralogs expressed and their relative abundance. Because transmembrane domain-induced homodimerization is important for the regulation of syndecan functions, the transition to heterodimers, which would necessarily constrain the formation of homo-oligomers, would be expected to disrupt syndecan functions. Consistent with this, our results revealed that hetero-oligomer formation between syndecan-2 and -4 inhibited both syndecan-4-mediated cell adhesion functions (Fig. 6) and syndecan-2-mediated tumorigenic activity (Fig. 7). These data suggest that increased hetero-oligomerization negatively regulates syndecan-mediated cellular signaling through restriction of homo-oligomerization. The oligomeric status of the syndecan-4 cytoplasmic domain is particularly important for syndecan-4 functions, including interactions with effector molecules. Hetero-oligomerization prevents homo-association of syndecan-4 cytoplasmic domains, thereby negatively regulating syndecan-4-mediated functions. These hetero-interactions may also contribute to novel signaling specificities because, in hetero-oligomers, the identity and geometry of ligand pairs differ from that of homo-oligomers. Distinct sets of syndecans formed through hetero-oligomerization could be regulated by different ligands, resulting in differential coupling to signaling

effectors and biological responses. Dissecting the cooperative signal transduction pathways of syndecan homo- and hetero-oligomers would be an interesting avenue for future research.

Transmembrane domain interactions have been shown to play a functional role in other receptors, including growth factor receptor signal transduction. Binding of ligands to the extracellular domain of the epidermal growth factor receptor (EGFR or Erb) induces the formation of receptor homo- or heterodimers and subsequent activation of the intrinsic tyrosine kinase domain (27). A hierarchy of homomeric and heteromeric association has been identified for ErbB transmembrane domain peptides (28). These interactions may contribute to unique signaling specificities because kinase/substrate pairs in a trans-phosphorylation reaction within a heterodimer differ from those of a homodimer. In NIH3T3 cells ectopically expressing EGFR and ErbB3, neuregulin stimulates anchorage-independent growth, whereas EGF does not despite the fact that EGF stimulates higher levels of receptor phosphorylation than does neuregulin (29). Similarly, EGFR is activated by HB-EGF in a breast tumor cell line, leading to tyrosine phosphorylation of the downstream signaling effector Cbl and subsequent Cbl-EGFR interactions, whereas in EGFR-expressing cells in which EGFR is activated by neuregulin through transmodulation by ErbB3 or ErbB4, Cbl is not tyrosine phosphorylated and does not complex with the EGFR (30). These data suggest that heterodimerization mediates phosphorylation of distinct sets of receptor tyrosine residues in response to differ-

ent ligands, creating an extracellular regulatory mechanism. Therefore, hetero-oligomerization through transmembrane domains might potentiate the diversification of biological functions.

This work clearly showed that syndecans form transmembrane domain-mediated heterodimers, and this dimerization regulates syndecan-mediated cellular functions. However, the effect of hetero-oligomerization might differ in various cells and tissues, perhaps due to tissue-specific differences in expression patterns and syndecan regulatory mechanisms. Further studies will be required to clarify the roles of hetero-oligomerization in specific syndecan regulatory mechanisms.

References

1. Tkachenko, E., Rhodes, J. M., and Simons, M. (2005) Syndecans: new kids on the signaling block. *Circ. Res.* **96**, 488–500
2. Rapraeger, A. C. (2000) Syndecan-regulated receptor signaling. *J. Cell Biol.* **149**, 995–998
3. Lambaerts, K., Wilcox-Adelman, S. A., and Zimmermann, P. (2009) The signaling mechanisms of syndecan heparan sulfate proteoglycans. *Curr. Opin. Cell Biol.* **21**, 662–669
4. Asundi, V. K., and Carey, D. J. (1995) Self-association of N-syndecan (syndecan-3) core protein is mediated by a novel structural motif in the transmembrane domain and ectodomain flanking region. *J. Biol. Chem.* **270**, 26404–26410
5. Choi, S., Lee, E., Kwon, S., Park, H., Yi, J. Y., Kim, S., Han, I. O., Yun, Y., and Oh, E. S. (2005) Transmembrane domain-induced oligomerization is crucial for the functions of syndecan-2 and syndecan-4. *J. Biol. Chem.* **280**, 42573–42579
6. Carey, D. J. (1997) Syndecans: multifunctional cell-surface co-receptors. *Biochem. J.* **327**, 1–16
7. Kim, C. W., Goldberger, O. A., Gallo, R. L., and Bernfield, M. (1994) Members of the syndecan family of heparan sulfate proteoglycans are expressed in distinct cell-, tissue-, and development-specific patterns. *Mol. Biol. Cell* **5**, 797–805
8. Fears, C. Y., and Woods, A. (2006) The role of syndecans in disease and wound healing. *Matrix Biol.* **25**, 443–456
9. Han, I., Park, H., and Oh, E. S. (2004) New insights into syndecan-2 expression and tumorigenic activity in colon carcinoma cells. *J. Mol. Histol.* **35**, 319–326
10. Schlessinger, J. (2000) Cell signaling by receptor tyrosine kinases. *Cell* **103**, 211–225
11. Dews, I. C., and Mackenzie, K. R. (2007) Transmembrane domains of the syndecan family of growth factor coreceptors display a hierarchy of homotypic and heterotypic interactions. *Proc. Natl. Acad. Sci. U.S.A.* **104**, 20782–20787
12. Shively, J. E., and Conrad, H. E. (1976) Formation of anhydrosugars in the chemical depolymerization of heparin. *Biochemistry* **15**, 3932–3942
13. Hecker, T. P., Grammer, J. R., Gillespie, G. Y., Stewart, J. Jr., and Gladson, C. L. (2002) Focal adhesion kinase enhances signaling through the Shc/extracellular signal-regulated kinase pathway in anaplastic astrocytoma tumor biopsy samples. *Cancer Res.* **62**, 2699–2707
14. Oh, E. S., Woods, A., and Couchman, J. R. (1997) Multimerization of the cytoplasmic domain of syndecan-4 is required for its ability to activate protein kinase C. *J. Biol. Chem.* **272**, 11805–11811
15. Horowitz, A., and Simons, M. (1998) Regulation of syndecan-4 phosphorylation *in vivo*. *J. Biol. Chem.* **273**, 10914–10918
16. Schlessinger, T. K., Demali, K. A., Johnson, G. L., and Kazlauskas, A. (1999) Platelet-derived growth factor-dependent association of the GTPase-activating protein of Ras and Src. *Biochem. J.* **344**, 519–526
17. DeMali, K. A., Balciunaite, E., and Kazlauskas, A. (1999) Integrins enhance platelet-derived growth factor (PDGF)-dependent responses by altering the signal relay enzymes that are recruited to the PDGF β receptor. *J. Biol. Chem.* **274**, 19551–19558
18. Saoncella, S., Echtermeyer, F., Denhez, F., Nowlen, J. K., Mosher, D. F., Robinson, S. D., Hynes, R. O., and Goetinck, P. F. (1999) Syndecan-4 signals cooperatively with integrins in a Rho-dependent manner in the assembly of focal adhesions and actin stress fibers. *Proc. Natl. Acad. Sci. U.S.A.* **96**, 2805–2810
19. Woods, A., and Couchman, J. R. (2001) Syndecan-4 and focal adhesion function. *Curr. Opin. Cell Biol.* **13**, 578–583
20. Dovas, A., Choi, Y., Yoneda, A., Multhaupt, H. A., Kwon, S. H., Kang, D., Oh, E. S., and Couchman, J. R. (2010) Serine 34 phosphorylation of rho guanine dissociation inhibitor (RhoGDI α) links signaling from conventional protein kinase C to RhoGTPase in cell adhesion. *J. Biol. Chem.* **285**, 23296–23308
21. Choi, Y., Kang, D., Han, I. O., and Oh, E. S. (2012) Hierarchy between the transmembrane and cytoplasmic domains in the regulation of syndecan-4 functions. *Cell Signal.* **24**, 1522–1530
22. Park, H., Kim, Y., Lim, Y., Han, I., and Oh, E. S. (2002) Syndecan-2 mediates adhesion and proliferation of colon carcinoma cells. *J. Biol. Chem.* **277**, 29730–29736
23. Choi, Y., Kim, H., Chung, H., Hwang, J. S., Shin, J. A., Han, I. O., and Oh, E. S. (2010) Syndecan-2 regulates cell migration in colon cancer cells through Tiam1-mediated Rac activation. *Biochem. Biophys. Res. Commun.* **391**, 921–925
24. Lee, H., Kim, Y., Choi, Y., Choi, S., Hong, E., and Oh, E. S. (2011) Syndecan-2 cytoplasmic domain regulates colon cancer cell migration via interaction with syntenin-1. *Biochem. Biophys. Res. Commun.* **409**, 148–153
25. Rapraeger, A. C., and Ott, V. L. (1998) Molecular interactions of the syndecan core proteins. *Curr. Opin. Cell Biol.* **10**, 620–628
26. Cizmeci-Smith, G., Langan, E., Youkey, J., Showalter, L. J., and Carey, D. J. (1997) Syndecan-4 is a primary-response gene induced by basic fibroblast growth factor and arterial injury in vascular smooth muscle cells. *Arterioscler. Thromb. Vasc. Biol.* **17**, 172–180
27. Olayioye, M. A., Neve, R. M., Lane, H. A., and Hynes, N. E. (2000) The ErbB signaling network: receptor heterodimerization in development and cancer. *EMBO J.* **19**, 3159–3167
28. Duneau, J. P., Vegh, A. P., and Sturgis, J. N. (2007) A dimerization hierarchy in the transmembrane domains of the HER receptor family. *Biochemistry* **46**, 2010–2019
29. Cohen, B. D., Green, J. M., Foy, L., and Fell, H. P. (1996) HER4-mediated biological and biochemical properties in NIH 3T3 cells: evidence for HER1-HER4 heterodimers. *J. Biol. Chem.* **271**, 4813–4818
30. Graus-Porta, D., Beerli, R. R., Daly, J. M., and Hynes, N. E. (1997) ErbB-2, the preferred heterodimerization partner of all ErbB receptors, is a mediator of lateral signaling. *EMBO J.* **16**, 1647–1655