



HHS Public Access

Author manuscript

JAMA Otolaryngol Head Neck Surg. Author manuscript; available in PMC 2016 July 01.

Published in final edited form as:

JAMA Otolaryngol Head Neck Surg. 2015 July 1; 141(7): 641–648. doi:10.1001/jamaoto.2015.0889.

Hearing Impairment Prevalence and Associated Risk Factors in the Hispanic Community Health Study/Study of Latinos (HCHS/SOL)

Karen J Cruickshanks, Ph.D.^{1,2}, Sumitrajit Dhar, Ph.D.³, Elizabeth Dinces, M.D., M.S.⁴, Robert C. Fifer, Ph.D.⁵, Franklyn Gonzalez II, M.S.⁶, Gerardo Heiss, M.D., Ph.D.⁶, Howard J. Hoffman, M.A.⁷, David J. Lee, Ph.D.⁸, Marilyn Newhoff, Ph.D.^{9,10}, Laura Tocci, Au.D.⁴, Peter Torre III, Ph.D., M.S.¹⁰, and Ted S. Tweed, M.A.¹

¹Department of Ophthalmology and Visual Sciences, University of Wisconsin School of Medicine and Public Health, Madison, WI 53705

²Department of Population Health Sciences, University of Wisconsin School of Medicine and Public Health, Madison, WI 53705

³Roxelyn and Richard Pepper Department of Communication Sciences and Disorders, Northwestern University, Evanston, IL 60208

⁴Department of Otorhinolaryngology- Head and Neck Surgery, Montefiore Medical Center, The University Hospital for Albert Einstein College of Medicine, Bronx, NY 10467

⁵Mailman Center for Child Development, Department of Pediatrics, University of Miami Miller School of Medicine, Fifer - Miami, FL 33136

⁶Collaborative Studies Coordinating Center, Department of Biostatistics, University of North Carolina Gillings School of Global Public Health, Chapel Hill, NC, 27514

Corresponding Author: Karen J. Cruickshanks, PhD, Dept. of Ophthalmology and Visual Sciences and Dept. of Population Health Sciences, School of Medicine and Public Health, University of Wisconsin, 610 Walnut Street, 10th Floor WARF, Madison, WI 53705, (608) 262-4032, FAX (608) 265-2148, cruickshanks@episense.wisc.edu.

Author contributions: Dr. Cruickshanks and Mr. Gonzalez had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. Drs. Cruickshanks, Dhar, Dinces, Fifer, Heiss, Lee, Tocci, and Torre and Mr. Hoffman, Mr. Gonzalez, and Mr. Tweed had responsibility for the conception and design of the study. All authors made substantial contributions to the acquisition, analysis or interpretation of the data. Dr. Cruickshanks drafted the manuscript and all authors reviewed the draft and contributed to the critical revision of the manuscript for important intellectual content. All authors contributed to obtaining funding for the study. Mr. Gonzalez had primary responsibility for conducting the statistical analyses. All authors contributed administrative, technical or material support and were responsible for supervision of aspects of the study.

Role of the Sponsors: Under the cooperative agreement the funding agencies were involved in the design and conduct of the study, reviewed and approved the manuscript, and participated in the decision to submit the manuscript for publication.

Conflict of Interest Disclosures: All authors declare that they have no conflicts of interest.

FINANCIAL DISCLOSURE: The Hispanic Community Health Study/Study of Latinos was carried out as a collaborative study supported by contracts from the National Heart, Lung, and Blood Institute (NHLBI) to the University of North Carolina (N01-HC65233), University of Miami (N01-HC65234), Albert Einstein College of Medicine (N01-HC65235), Northwestern University (N01-HC65236), and San Diego State University (N01-HC65237). The hearing protocol and Hearing Reading Center were sponsored and funded as an adjunct to HCSH/SOL through Intra-Agency Agreement with the National Institute on Deafness and Other Communication Disorders (NIDCD), NIH. In addition, the following Institutes/Centers/Offices contributed to specific components in the protocol and, also, to the overall collection of data in the HCHS/SOL through a transfer of funds to the NHLBI: National Center on Minority Health and Health Disparities, the National Institute on Deafness and Other Communications Disorders, the National Institute of Dental and Craniofacial Research, the National Institute of Diabetes and Digestive and Kidney Diseases, the National Institute of Neurological Disorders and Stroke, and the Office of Dietary Supplements.

⁷Epidemiology and Statistics Program, National Institute on Deafness and Other Communication Disorders (NIDCD), National Institutes of Health (NIH), Bethesda, MD 20892

⁸Department of Public Health Sciences, University of Miami Miller School of Medicine Miami, FL 33101

⁹College of Health and Human Services, San Diego State University, San Diego, CA 92182-1518

¹⁰School of Speech, Language, & Hearing Sciences, San Diego State University, San Diego, CA 92182-1518

Abstract

Importance—Hearing impairment (HI) is a common problem in adults but there have been few studies of hearing in the U.S. Hispanic/Latino population. Little is known about factors associated with HI among Hispanics/Latinos.

Objective—To determine the prevalence of HI among U.S. Hispanic/Latino adults of diverse backgrounds and determine associations with sociodemographic factors, noise exposure, diabetes, smoking, cardiovascular disease, and other potential risk factors.

Design and Setting—The Hispanic Community Health Study/Study of Latinos (HCHS/SOL) is a population-based sample of Hispanics/Latinos in four U.S. communities (Bronx, NY, Chicago, IL, Miami, FL, and San Diego, CA). Examinations were conducted in 2008–2011.

Participants—The HCHS/SOL examined 16,415 self-identified Hispanic/Latino persons aged 18 to 74 years recruited from randomly selected households using a stratified 2-stage area probability sample design based on census block groups and households within block groups.

Intervention(s)—None

Main Outcome(s) and Measures—Hearing thresholds were measured by pure-tone audiometry. HI was defined as a pure-tone average (PTA) of thresholds at 0.5, 1, 2, 4 kHz >25 dB HL. Bilateral hearing impairment (BHI) required a PTA >25 dB HL in both ears. Multivariable analyses included adjustments for sociodemographic and lifestyle variables, body mass index, and medical conditions.

Results—The prevalence of HI was 15.1% overall and 8.2% had BHI. The prevalence of HI was higher among people aged 45 and older, ranging by Hispanic/Latino background from 29–41% among men and 18–31% among women. The multivariable-adjusted odds of HI was greater for participants of Puerto Rican background compared to Mexican background (OR = 1.57, 95% CI = 1.10, 2.25). The odds of HI were lower with more education and higher income. People with noise exposure were about 30% more likely to have HI. Diabetes (OR = 1.57, 95% CI = 1.27, 1.94) and pre-diabetes (OR = 1.37, 95% CI = 1.12, 1.67) were associated with higher odds of HI.

Conclusions and Relevance—HI is a common problem for older Hispanics/Latinos in these communities and is associated with socioeconomic factors, noise exposure and abnormal glucose metabolism. Longitudinal studies are needed to determine if these factors are involved in the etiology of HI and to identify ways to prevent or delay age-related changes in hearing.

Keywords

Hearing impairment; presbycusis; Hispanic; prevalence; risk factors; diabetes

INTRODUCTION

Hearing Impairment (HI) is one of the most common chronic conditions affecting adults but data specific to the Hispanic/Latino population have been limited.¹⁻³ In the Hispanic Health and Nutrition Examination Survey (HHANES) conducted between 1982 and 1984, the prevalence of HI among men aged 55–74 years varied by Hispanic/Latino background, ranging from 46% among Mexican Americans and 48% among Cuban Americans to only 23% among Puerto Rican participants.¹ Among women the rates varied slightly with 35%, 43% and 33% of Mexican American, Cuban American, and Puerto Rican participants having HI, respectively.¹ More recently reports from the National Health and Nutrition Examination Survey (NHANES), have suggested that HI is less common among Mexican Americans than Non-Hispanic whites (NHW) but did not provide data about other Hispanic/Latino backgrounds.² Other large U.S. cohort studies⁴⁻⁷ include few Hispanic/Latino participants, leaving large gaps in knowledge about the prevalence, impact, unmet health care needs, and factors associated with HI among Hispanic/Latino populations.

There is growing evidence that age-related HI may be associated with lower socioeconomic status (SES), noisy jobs, diabetes, smoking, and obesity, risk factors that may be more common among the Hispanic/Latino population.^{3,6,8-13} HI may lead to lower quality of life and be associated with increased risk of dementia.¹⁴⁻²⁴ Most people with HI are undiagnosed and untreated, an unmet need for health care which may be exacerbated in groups typically underserved for health care.²⁵⁻²⁷ The purpose of this study was to determine the prevalence of HI among Hispanic/Latino adults from diverse backgrounds and to identify factors associated with HI among Hispanics/Latinos.

METHODS

The HCHS/SOL is a population-based cohort study of Hispanics/Latinos in four U.S. cities (Bronx, New York; Chicago, Illinois; Miami, Florida; San Diego, California) which was designed to examine risk and protective factors for chronic diseases by Hispanic/Latino background. Detailed descriptions of the sampling methods and baseline examination (2008–2011) have been presented previously.²⁸⁻³⁰ The HCHS/SOL examined 16,415 self-identified Hispanic/Latino persons aged 18 to 74 years recruited from randomly selected households. The largest background groups were Central American (n=1,730), Cuban (n=2,348), Dominican (n=1,460), Mexican (n=6,471), Puerto Rican (n=2,728), and South American (n=1,068). Households were selected using a stratified 2-stage area probability sample design based on census block groups and households within each sampled block group. Oversampling occurred at each stage, with block groups in areas of Hispanic/Latino concentration, households associated with a Hispanic/Latino surname, and persons aged 45 to 74 years selected at higher rates than their counterparts. All reported values (means and prevalence rates) were weighted to account for the disproportionate selection of the sample

and to at least partially adjust for any bias effects due to differential nonresponse in the selected sample at the household and person levels. The adjusted weights were also trimmed to limit precision losses due to the variability of the adjusted weights, and calibrated to the 2010 Census characteristics by age, sex and Hispanic background in each field site's target population. All analyses also accounted for cluster sampling and the use of stratification in sample selection. The study was approved by institutional review boards at each participating institution; written informed consent was obtained from all participants.

Hearing Examination Methods

All technicians were centrally trained and certified by the EpiSense Audiometry Reading (EAR) Center at the University of Wisconsin. Detailed protocols are available on line at <http://www.csc.unc.edu/hchs/>.

Ambient sound levels were measured daily at each test site.³¹ GSI 61 clinical audiometers (Grason-Stadler, Inc., Madison, WI) equipped with TDH-50P and insert earphones (E-A-Rtone 3A, Cabot Safety Corp., Indianapolis, IN) were calibrated annually according to ANSI standards.³² A brief otoscopic examination and screening tympanogram (Earscan®, Micro Audiometrics, Murphy, North Carolina) were obtained to assist in identifying obstructing cerumen and middle ear disease. Hearing thresholds were measured in sound-treated booths by pure tone air (0.5, 1, 2, 3, 4, 6, 8 kHz) and bone conduction (0.5, 2, 4 kHz) audiometry using a modified Hughson-Westlake procedure according to the guidelines of the American Speech-Language-Hearing Association.³³ Masking was used as necessary. The presence of hearing impairment was defined as a pure-tone average (PTA) of thresholds at 0.5, 1, 2, 4 kHz greater than 25 decibels Hearing Level (dB HL) in either ear. A second definition of PTA > 25 dB HL in the better ear was also used to capture bilateral hearing impairment. This four frequency average has long been used as a measure of hearing impairment in epidemiologic studies.³⁴

Sociodemographic (education and income), lifestyle and medical history data were obtained by questionnaires administered in English or Spanish. Participants were asked about smoking (never, former and current) and alcohol consumption (never, former, current). History of cardiovascular disease (CVD) including myocardial infarction, angina, stroke or transient ischemic attacks was ascertained. History of noise exposure was considered positive if the participant reported using firearms, military service or exposure to loud noises during leisure time. Occupational noise exposure was defined as a current or former job which was noisy (had to speak in a raised voice or louder to be heard two feet away). The longest held job was categorized as professional/managerial/technical, service, or production/manufacturing/labor.

Hypertension was defined as a systolic blood pressure 140 mm Hg or greater, diastolic blood pressure 90 mm Hg or greater, or receiving antihypertensive medication.^{30,35} Diabetes mellitus was a fasting plasma glucose 126 mg/dL or greater, 2-hour-postload plasma glucose 200 mg/dL or greater, an HbA1c 6.5% or greater, self-report of doctor-diagnosed diabetes (excluding gestational diabetes) or use of antihyperglycemic medications.^{30,36} Pre-diabetes was a fasting plasma glucose between 100–125 mg/dL, 2-hour-postload plasma glucose between 140–199 mg/dL or an HbA1c between 5.7–6.4%. Height was measured to the

nearest centimeter and weight to the nearest 0.1 kg. Body mass index (BMI) was calculated as weight in kilograms divided by height in meters squared.

Statistical Analyses

All reported values (means, prevalence, and odds ratios [ORs]) were weighted to adjust for sampling probability and nonresponse. Age-adjusted prevalence estimates for the target population of Hispanics/Latinos in the four HCHS/SOL communities were calculated using survey-specific logistic regression procedures adjusting each subgroup to the age distribution of the target population. Survey-specific logistic regression analyses were used to examine associations of risk factors with HI. Initial models adjusted for age, sex, center, and Hispanic background. Multivariable models adjusted for all covariates shown. Odds ratios with 95% CIs were computed using survey-specific logistic regression procedures to account for the 2-stage sampling design, stratification, and clustering. Age groups were collapsed to ages 18–44 and 45+ to ensure good precision of prevalence estimates; the cut-point reflects the original study design which oversampled the 45+ age group. All analyses were performed using SAS version 9.3 (SAS Institute, Cary, NC) and SUDAAN Release 10.0.0 (RTI, Research Triangle Park, NC).

RESULTS

The descriptive characteristics of the study participants (unweighted) and the sample population (weighted) are shown in Table 1. In the weighted sample, 52% were female and 40.5% were age 45 years or older, the age range when age-related hearing loss would present. Only 0.85% reported a history of ear surgery or disease. Overall, 15.06% (Standard error (SE) = 0.44) had hearing impairment, and about half of these (8.24%, SE=0.33) had bilateral hearing impairment (Table 2). Only a small proportion of those with hearing impairment (either ear or bilateral) had some evidence of conductive hearing problems as 3.12% (SE=0.19) of left ears and 2.92% (SE=0.17) of right ears demonstrated an air-bone gap of 15 dB or more.

In general, the prevalence of hearing impairment was higher among men and older age groups (Table 2). Among men 18–44 years of age, the prevalence of hearing impairment in either ear ranged from 2.68% among men reporting South American background to 7.14% among men of Puerto Rican background (Table 2). Similarly, among men 45 years of age or older, the prevalence of hearing impairment (in either ear) was highest among men reporting Puerto Rican background (41.20%) and varied among the other groups with a low of 29.35% among men with Dominican background.

The prevalence of HI also varied among women by Hispanic/Latino background (Table 2). For women ages 18–44 years, the prevalence ranged from a high of 8.14% among participants reporting Puerto Rican background to a low of 2.72% among those reporting South American background. Older women (age 45+ years) who reported mixed/other background had a prevalence of 32.11% and the lowest prevalence was among the women of Mexican background (17.89%). As expected, the prevalence of bilateral hearing impairment was substantially lower in all groups of men and women, but the patterns were

similar, with higher prevalence among participants reporting Puerto Rican background and lower prevalence among persons of South American background.

Adjusting for age, gender, center, and Hispanic/Latino background, indicators of higher socioeconomic status were associated with lower prevalence (Table 3). Greater education was associated with a 30% lower prevalence of HI (Odds Ratio (OR) = 0.71 for a high school diploma or equivalent vs none, 95% Confidence Interval (CI) = 0.61, 0.83). Those with the highest income level were significantly less likely to have HI than people with the lowest income (OR=0.56, 95% CI = 0.29, 0.73 for \$75,000 vs <\$10,000). People with diabetes had a 50% increased odds of HI. Obesity, current smoking, history of cardiovascular disease, service sector occupations, former exposure to occupational noise, and noise exposure history were associated with greater odds of hearing impairment (Table 3). Associations with bilateral hearing impairment (BHI) were very similar (Table 3), although statistical significance varied slightly.

In the multivariable-adjusted model (Table 4), age was strongly associated with HI; people 65 years of age or older had an 18.5-fold (95%CI = 12.40, 27.68) greater prevalence of HI compared to younger adults (ages 18–44 years). Greater education and higher income remained protective with effect sizes similar to the simpler models. Compared to Mexican background, Puerto Rican background remained associated with increased odds of HI (OR=1.57, 95%CI=1.10, 2.25); there were no other Hispanic/Latino background groups with higher or lower odds of HI. Exposure to noise from firearms and leisure activities as well as former exposure to occupational noise were associated with increased odds of HI. Having diabetes, and pre-diabetes, was associated with significantly increased odds of HI. CVD history, hypertension, BMI, smoking, and alcohol were retained in the model, although not significantly associated with HI. The bilateral hearing impairment model showed similar associations; in addition, hypertension and former but not current alcohol consumption were independently associated with increased odds of bilateral hearing impairments.

DISCUSSION

In this study of U.S. Hispanics/Latinos from diverse backgrounds, HI was common among adults and the prevalence increased sharply after age 45 years. This finding is consistent with results from studies in other populations and with the national data from the NHANES.^{1–7} As in other studies men were more likely to have HI than women. Unlike the early HHANES¹ in the 1980s, Puerto Rican background was associated with higher prevalence of HI than Mexican background. The hearing testing procedures were similar between these studies. It is possible the different pattern by background reflects temporal changes as the children with Puerto Rican background in this early study had worse hearing than Mexican American children; those children would now be middle-aged adults.³⁷ Alternatively, sample differences may account for this difference as the typical male excess of hearing impairment was not seen among participants with Puerto Rican backgrounds in the HHANES, suggesting that participation or selection bias in HHANES may have led to underestimating the prevalence of HI in this group.¹ Results from the NHANES have suggested that Hispanics/Latinos have lower rates of hearing impairment than NHW.^{3,38} However, in that study Hispanics/Latinos were primarily from Mexican backgrounds, which

may have led to over-estimating the ethnic differences in HI. Clearly, it is important for future studies to include people from diverse backgrounds to accurately determine the public health burden of hearing impairment.

Hearing impairment is associated with lower overall quality of life, and has been associated with depression and risk of dementia.^{14–24} It is common for hearing impairment to go undetected and untreated for many years.^{25–27} In this cohort, some of the participants with hearing impairment had conductive components which may be amenable to medical treatment. Health professionals and the public need to be aware that hearing impairment is a common and important health disorder affecting Hispanics/Latinos in the U.S. It is important to develop culturally appropriate effective intervention strategies to meet the communication needs of the Hispanic/Latino community.

Education and higher income have been associated with lower rates of hearing impairment in other cross-sectional studies and in longitudinal studies of NHW participants.^{3,6,9,10,12} Similarly, in the HCHS/SOL cohort, people with higher educational levels and higher incomes were less likely to have HI than others. Socioeconomic factors are often associated with disease prevalence, including risk of cardiovascular disease and other age-related conditions, and occupational noise exposure which is a well-recognized risk factor for hearing loss. In this study, histories of noise exposure at work and outside of work (firearms and leisure sources) were associated with higher prevalence of HI. This finding is consistent with other cross-sectional studies of hearing.^{3,5–7,38} The education/income gradient remained in the multivariable model adjusting for noise exposure suggesting that unidentified factors which vary by socioeconomic status factors may contribute to the loss of hearing with aging.

Cardiovascular disease and its risk factors have often been thought to contribute to aging changes in hearing, at least partially through its effects on the stria vascularis which is important for maintaining normal signal transduction in the inner ear.^{40–42} Vascular factors also may damage central signal processing and contribute to deterioration in hearing with aging.³⁹ The higher prevalence of HI among HCHS/SOL participants with Puerto Rican background is consistent with a vascular hypothesis as they also were more likely than other groups to have multiple cardiovascular disease risk factors.³⁰ In this study, smoking, obesity, history of CVD, and diabetes also were associated with HI in simpler models adjusting for demographic factors; only diabetes and pre-diabetes remained significantly associated with HI in the final multivariable model. These findings are consistent with other studies.^{3,6,7,8,9,11,12, 39,40} Diabetes is associated with generalized inflammation, basement membrane thickening, vascular disease and neural changes all of which also may damage the auditory system and result in HI. However, to date, diabetes has not been associated with the incidence of hearing impairment in large longitudinal studies, so the etiologic significance of this association remains uncertain. Inflammatory markers have been associated with HI in other studies.^{43–44}

The strengths of the study are the population-based sampling strategy, large numbers of Hispanics/Latinos from diverse backgrounds and use of standardized, sensitive assessments of hearing and diabetes, as well as the availability of extensive lifestyle and health data.

There are, of course, limitations in drawing causal inferences due to the cross-sectional design. The smaller numbers of participants within some of the background groups may have resulted in lower power to detect differences. In addition, participants are representative of the four communities in United States included in the study but may not be representative of residents in other parts of the United States or of the countries from which they or their ancestors emigrated. The groups reporting Central or South American backgrounds are likely to be very heterogeneous because of the large number of countries and geographic regions encompassed. Measures of auditory function were limited as time constraints in this study of numerous health outcomes did not permit the inclusion of tests of speech understanding or central auditory processing. Self-reports of health conditions, noise exposure, and smoking, etc. may be subject to recall or reporting bias which may have led to under- or over-estimates of effect sizes. Nonetheless, this is the largest study of hearing sensitivity among people of diverse Hispanic/Latino backgrounds in the United States and suggests that the prevalence of hearing impairment varies within the U.S. Hispanic/Latino population in these communities.

Conclusion

This large multicenter study has demonstrated that HI is common among Hispanic/Latino adults from all backgrounds, as 15% of all adults had HI. Diabetes, less education, lower income, and more noise exposure were associated with higher odds of hearing impairments. Health care professionals need to be aware that Hispanic/Latino patients, especially those with abnormal glucose metabolism, may have hearing impairments and need screening and appropriate referrals. Future longitudinal studies of Hispanics/Latinos from diverse backgrounds could strengthen the determination of the risks associated with hearing loss. This longitudinal information is needed to identify modifiable risk factors to slow the progression of hearing loss with aging and to develop culturally appropriate effective intervention strategies to meet the communication needs of the Hispanic/Latino community.

Acknowledgments

The authors thank the staff and participants of HCHS/SOL for their important contributions. A complete list of staff and investigators has been provided by Sorlie P., et al. in *Ann Epidemiol.* 2010 Aug; 20: 642–649 and is also available on the study website <http://www.csc.unc.edu/hchs/>

FUNDING

The Hispanic Community Health Study/Study of Latinos was carried out as a collaborative study supported by contracts from the National Heart, Lung, and Blood Institute (NHLBI) to the University of North Carolina (N01-HC65233), University of Miami (N01-HC65234), Albert Einstein College of Medicine (N01-HC65235), Northwestern University (N01-HC65236), and San Diego State University (N01-HC65237). The hearing protocol and Hearing Reading Center were sponsored and funded as an adjunct to HCHS/SOL through Intra-Agency Agreement with the National Institute on Deafness and Other Communication Disorders (NIDCD), NIH. In addition, the following Institutes/Centers/Offices contributed to specific components in the protocol and, also, to the overall collection of data in the HCHS/SOL through a transfer of funds to the NHLBI: National Center on Minority Health and Health Disparities, the National Institute on Deafness and Other Communications Disorders, the National Institute of Dental and Craniofacial Research, the National Institute of Diabetes and Digestive and Kidney Diseases, the National Institute of Neurological Disorders and Stroke, and the Office of Dietary Supplements.

References

1. Lee DJ, Carlson DL, Lee HM, Ray LA, Markides KS. Hearing loss and hearing aid use in Hispanic adults: results from the Hispanic Health and Nutrition Examination Survey. *Am J Public Health*. 1991 Nov; 81(11):1471–4. [PubMed: 1951805]
2. Davanipour Z, Lu NM, Lichtenstein M, Markides KS. Hearing problems in Mexican American elderly. *Am J Otol*. 2000 Mar; 21(2):168–72. [PubMed: 10733179]
3. Agrawal Y, Platz EA, Niparko JK. Risk factors for hearing loss in US adults. Data from the National Health and Nutrition Examination Survey, 1999 to 2002. *Otol Neurotol*. 2009 Feb; 30(2): 139–45. [PubMed: 19092714]
4. Mo cicki EK, Elkins EF, Baum HM, McNamara PM. Hearing Loss in the Elderly: An Epidemiologic Study of the Framingham Heart Study Cohort. *Ear Hear*. 1985; 6:184–190. [PubMed: 4043571]
5. Cruickshanks KJ, Wiley TL, Tweed TS, Klein BEK, Klein R, Mares-Perlman JA, et al. Prevalence of hearing loss in older adults in Beaver Dam, Wisconsin: The Epidemiology of Hearing Loss Study. *American Journal of Epidemiology*. 1998; 148(9):879–886. [PubMed: 9801018]
6. Nash SD, Cruickshanks KJ, Klein R, Klein BEK, Nieto JF, Huang GH, et al. The prevalence of hearing impairment and associated risk factors: The Beaver Dam Offspring Study. *Archives of Otolaryngology Head & Neck Surgery*. 2011 May; 137(5):432–439. [PubMed: 21339392]
7. Helzner EP, Cauley JA, Pratt SR, Wisniewski SR, Zmuda JM, Talbott EO, et al. Race and sex differences in age-related hearing loss; The Health, Aging, and Body Composition Study. *J Am Geriatr Soc*. 2005; 53:2119–2127. [PubMed: 16398896]
8. Bainbridge KE, Hoffman HJ, Cowie CC. Diabetes and hearing impairment in the United States: Audiometric evidence from the National Health and Nutrition Exam Survey, 1999 to 2004. *Ann Intern Med*. 2008; 149(1):1–10. [PubMed: 18559825]
9. Zhan W, Cruickshanks KJ, Klein BEK, Klein R, Huang GH, Pankow JS, et al. Modifiable determinants of hearing impairment in adults. *Preventive Medicine*. 2011 Oct; 53(4–5):338–342. [PubMed: 21871479]
10. Cruickshanks KJ, Nondahl DM, Tweed TS, Wiley TL, Klein BEK, Klein RK, et al. Education, occupation, noise exposure history and the 10-yr cumulative incidence of hearing impairment in older adults. *Hearing Research*. 2010; 264(1–2):3–9. [PubMed: 19853647]
11. Cruickshanks KJ, Klein R, Klein BEK, Wiley TL, Nondahl DM, Tweed TS. Cigarette smoking and hearing loss: The Epidemiology of Hearing Loss Study. *JAMA*. 1998; 279(21):1715–1719. [PubMed: 9624024]
12. Curhan SG, Eavey R, Wang M, Stampfer MJ, Curhan GC. Body mass index, waist circumference, physical activity, and risk of hearing loss in women. *Am J Med*. 2013 Dec; 126(12):1142.e1–8. Epub 2013 Oct 11. 10.1016/j.amjmed.2013.04.026 [PubMed: 24125639]
13. Lee DJ, Gomez-Martin O, Lee HM. Sociodemographic correlates of hearing loss and hearing aid use in Hispanic adults. *Epidemiology*. 1996 Jul; 7(4):443–6. [PubMed: 8793375]
14. Lee DJ, Gomez-Marin O. Major depressive disorder, depressive symptoms, and bilateral hearing loss in Hispanic adults. *J Affect Disord*. 1997 Jul; 44(2–3):189–95. [PubMed: 9241579]
15. Salomon JA, Vos T, Hogan DR, Gagnon M, Naghavi M, Mokdad A, et al. Common values in assessing health outcomes from disease and injury: disability weights measurement study for the Global Burden of Disease Study 2010. *Lancet*. 2012; 380(9859):2129–2143. [PubMed: 23245605]
16. Murray CJL, Vos T, Lozano R, Naghavi M, Flaxman AD, Michaud C, et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012; 380(9859):2197–2223. [PubMed: 23245608]
17. Dalton DS, Cruickshanks KJ, Klein BEK, Klein R, Wiley TL, Nondahl DM. The impact of hearing loss on quality of life in older adults. *Gerontologist*. 2003; 43(5):661–668. [PubMed: 14570962]
18. Fischer ME, Cruickshanks KJ, Klein BEK, Klein R, Schubert CR, Wiley TL. Multiple sensory impairment and quality of life. *Ophthalmic Epidemiol*. 2009; 16(6):346–353. [PubMed: 19995199]

19. Strawbridge WJ, Wallhagen MI, Sherma SJ, Kaplan GA. Negative consequences of hearing impairment in old age: a longitudinal analysis. *Gerontologist*. 2000; 40(3):320–26. [PubMed: 10853526]
20. Nachtegaal J, Smit JH, Smits C, Bezemer PD, Van Beek JH, Festen JM, et al. The association between hearing status and psychosocial health before the age of 70 years: results from an Internet-based national survey on hearing. *Ear Hear*. 2009; 30(3):302–312. [PubMed: 19322094]
21. Uhlmann RF, Larson EB, Rees TS, Koepsell TD, Duckert LG. Relationship of hearing impairment to dementia and cognitive dysfunction in older adults. *JAMA*. 1989; 261(13):1916–1919. [PubMed: 2926927]
22. Lin FR, Yaffe K, Xia J, Xue QL, Harris TB, Purchase-Helzner E, et al. Hearing loss and cognitive decline in older adults. *JAMA Internal Medicine*. 2013; 173(4):292–299.
23. Lin FR, Metter EJ, O'Brien RJ, Resnick SM, Zonderman AB, Ferrucci L. Hearing loss and incident dementia. *Archives of Neurology*. 2011; 68(2):214–220.
24. Li C-M, Zhang X, Hoffman HJ, Cotch MF, Themann CL, Wilson MR. Hearing impairment associated with depression in US adults, National Health and Nutrition Examination Survey 2005–2010. *JAMA Otolaryngol Head Neck Surgery*. Published online March 6, 2014. 10.1001/jamaoto.2014.42
25. Popelka MM, Cruickshanks KJ, Wiley TL, Tweed TS, Klein BEK, Klein R. Low prevalence of hearing aid use among older adults with hearing loss: the Epidemiology of Hearing Loss Study. *J Am Geriatr Soc*. 1998; 46(9):1075–1078. [PubMed: 9736098]
26. Donahue A, Dubno JR, Beck L. Guest editorial: accessible and affordable hearing health care for adults with mild to moderate hearing loss. *Ear Hear*. 2010; 31(1):2–6. [PubMed: 20040828]
27. Nash SD, Cruickshanks KJ, Huang GH, Klein BEK, Klein R, Nieto J, et al. Unmet hearing health care needs: The Beaver Dam Offspring Study. *Am J Public Health*. 2013 Jun; 103(6):1134–1139. [PubMed: 23597370]
28. Sorlie PD, Avilés-Santa LM, Wassertheil-Smoller S, Kaplan RC, Daviglius ML, Giachello AL, et al. G. Design and implementation of the Hispanic Community Health Study/Study of Latinos. *Ann Epidemiol*. 2010 Aug; 20(8):629–41. [PubMed: 20609343]
29. LaVange LM, Kalsbeek WD, Sorlie PD, Aviles-Santa LM, Kaplan RC, Barnhart J, et al. Sample Design and Cohort Selection in the Hispanic Community Health Study/Study of Latinos. *Ann Epidemiol*. 2010; 20(8):642–49. [PubMed: 20609344]
30. Daviglius ML, Talavera GA, Avilés-Santa ML, Allison M, Cai J, Criqui MH, et al. Prevalence of major cardiovascular risk factors and cardiovascular diseases among Hispanic/Latino individuals of diverse backgrounds in the United States. *JAMA*. 2012 Nov 7; 308(17):1775–84.10.1001/jama.2012.14517 [PubMed: 23117778]
31. American National Standards Institute. Maximum permissible ambient noise levels for audiometric test rooms. New York: ANSI; 2008. p. S3.1
32. American National Standards Institute. Specifications for audiometers. New York: ANSI; 2010. p. S3.6
33. American Speech and Language-Hearing Association. Guidelines for manual pure tone audiometry. *AHSA*. 1987; 20:287–301.
34. Davis AC. The prevalence of hearing impairment and reported hearing disability among adults in Great Britain. *Int J Epidemiol*. 1989; 18(4):911–917. [PubMed: 2621028]
35. Sorlie PD, Allison MA, Aviles-Santa M, Cai J, Daviglius ML, Howard AG, et al. Prevalence of hypertension, awareness, treatment, and control in the Hispanic Community Health Study/Study of Latinos. *Am J Hypertens*. 2014 Jun; 27(6):793–800. [PubMed: 24627442]
36. Schneiderman N, Llabre M, Cowie CC, Barnhart J, Carnethon M, Gallo LC, et al. Prevalence of diabetes among Hispanics/Latians from diverse backgrounds: The Hispanic Community Health Study/Study of Latinos (HCHS/SOL). *Diabetes Care*. 2014 Aug; 37:2233–2239. [PubMed: 25061138]
37. Lee DJ, Gomez-Martin O, Lee HM. Prevalence of childhood hearing loss. *Am J Epidemiol*. 1996 Sep 1; 144(5):442–9. [PubMed: 8781458]

38. Lin FR, Maas P, Chien W, Carey JP, Ferrucci L, Thorpe R. Association of skin color, race/ethnicity, and hearing loss among adults in the USA. *J Assoc Res Otolaryngol*. 2012 Feb; 13(1): 109–17. [PubMed: 22124888]
39. Fransen E, Topsakal V, Hendrickx JJ, Van Laer L, Huyghe JR, Van Eyken E, et al. Occupational noise, smoking, and a high body mass index are risk factors for age-related hearing impairment and moderate alcohol consumption is protective: A European population- multicenter study. *J Assn Res Otolaryngol*. 2008; 9:264–276.
40. Gates GA, Cobb JL, D'Agostino RB, Wolf P. The relation of hearing in the elderly to the presence of cardiovascular disease and cardiovascular risk factors. *Arch Otolaryngol Head Neck Surg*. 1993; 119:156–161. [PubMed: 8427676]
41. Johnsson LG, Hawkins JE. Vascular changes in the human inner ear associated with aging. *Ann Otol*. 1972; 81:364–76.
42. Trune DR, Nguyen-Huynh A. Vascular pathophysiology in hearing disorders. *Seminars in Hearing*. 2012; 33(3):242–250. [PubMed: 25346568]
43. Verschuur CA, Dowell A, Syddall HE, Simmonds SS, Baylis D, Gale CR, et al. Markers of inflammatory status are associated with hearing threshold in older people: findings from the Hertfordshire Ageing Study. *Age and Ageing*. 2012; 41(1):92–97. [PubMed: 22086966]
44. Nash SD, Cruickshanks KJ, Zhan W, Tsai MY, Klein R, Chappell R, et al. Long-term Assessment of Systemic Inflammation and the Cumulative Incidence of Age-related Hearing Impairment in the Epidemiology of Hearing Loss Study. *J Gerontol A Biol Sci Med Sci*. 2014 Feb; 69(2):207–14. [PubMed: 23739996]

Table 1

Demographic Characteristics (unweighted and weighted for sampling design)

	Unweighted		Weighted	
	N	%	Mean or PCT	SE
Age				
18–24 years	1590	10.12	16.81	0.56
25–34 years	1969	12.53	21.50	0.61
35–44 years	2829	18.00	21.20	0.57
45–54 years	4729	30.09	19.00	0.48
55–64 years	3336	21.23	12.97	0.41
65+ years	1263	8.04	8.53	0.38
Gender				
Female	9415	59.91	52.03	0.57
Male	6301	40.09	47.97	0.57
Hispanic/Latino Background				
Central American	1674	10.65	7.39	0.55
Cuban	2301	14.64	20.51	1.72
Dominican	1340	8.53	9.48	0.71
Mexican	6354	40.43	38.21	1.68
Puerto Rican	2525	16.07	15.37	0.77
South American	1030	6.55	4.88	0.31
Other or Mixed	492	3.13	4.16	0.29
Center				
Bronx	3690	23.48		
Chicago	3985	25.36		
Miami	3996	25.43		
San Diego	4045	25.74		
Education				
No High school diploma or GED			32.37	0.73
At least High school diploma or GED			67.63	0.73

Table 2
Prevalence of Hearing Impairment by Hispanic/Latino Background (weighted for sampling design)

Gender	Age (yrs)	Hearing Impairment (Worse ear)															
		Central American	Cuban	Dominican	Mexican	Puerto Rican	South American	Other/Mixed	All	%	SE						
Bilateral Hearing Impairment																	
Males	18-44	5.05	1.16	6.46	1.22	6.32	1.99	4.33	0.74	7.14	1.34	2.68	1.15	3.25	1.48	5.20	0.49
	45+	37.65	3.72	39.71	2.32	29.35	3.58	33.39	2.14	41.20	3.50	33.44	4.13	34.42	6.99	36.76	1.25
Females	18-44	3.63	0.91	4.73	1.00	7.12	2.96	4.13	0.78	8.14	1.62	2.72	1.03	3.15	1.42	4.89	0.59
	45+	22.05	2.29	26.59	1.99	23.62	2.54	17.89	1.41	31.36	2.76	23.49	3.42	32.11	12.17	23.97	1.06
All		12.88	1.02	20.65	1.08	14.11	1.47	10.94	0.64	21.25	1.32	13.81	1.58	10.06	2.28	15.06	0.44
Males	18-44	3.10	1.04	2.45	0.79	0.55	0.40	1.33	0.40	3.19	0.94	0.16	0.16	0.25	0.25	1.77	0.29
	45+	24.99	3.27	25.68	2.05	15.8	2.91	21.76	1.78	26.77	3.61	14.78	3.26	20.47	5.72	23.22	1.17
Females	18-44	2.17	0.71	0.62	0.35	1.37	0.66	1.71	0.61	4.60	1.35	0.28	0.28	0.79	0.57	1.78	0.34
	45+	9.00	1.55	15.40	1.67	14.22	2.51	10.29	1.11	16.60	2.03	14.28	2.73	8.97	3.40	13.18	0.74
All		7.16	0.77	11.94	0.90	6.23	0.75	6.06	0.46	12.25	1.15	6.64	1.04	3.67	0.82	8.24	0.33

Table 3Adjusted¹ Odds Ratios of Hearing Impairment by Demographic and Other Factors

	<u>Hearing Impairment</u>		<u>Bilateral Hearing Impairment</u>	
	OR	95%CI	OR	95%CI
<u>Education</u>				
No High school diploma or GED	1.0		1.0	
At least High school diploma or GED	0.71	0.61, 0.83	0.64	0.53, 0.77
<u>Income</u>				
Less than \$10,000	1.0		1.0	
\$10,001 – \$20,000	0.88	0.70, 1.10	0.67	0.52, 0.87
\$20,001 – \$40,000	0.77	0.60, 1.00	0.55	0.42, 0.74
\$40,001 – \$75,000	0.53	0.40, 0.70	0.50	0.36, 0.69
More than \$75,000	0.56	0.29, 0.73	0.29	0.16, 0.54
<u>Diabetes</u>				
No diabetes	1.0		1.0	
Pre-diabetes	1.14	0.95, 1.36	1.20	0.93, 1.56
Diabetes	1.50	1.24, 1.82	1.58	1.21, 2.08
<u>Body Mass Index (Kg/m²)</u>				
Normal/Underweight (0–24.9)	1.0		1.0	
Overweight (25–29.9)	0.98	0.81, 1.18	1.01	0.80, 1.28
Obese (30+)	1.23	1.03, 1.48	1.28	1.00, 1.63
<u>Smoking Status</u>				
Never	1.0		1.0	
Former	1.09	0.93, 1.28	1.14	0.93, 1.41
Current	1.26	1.06, 1.50	1.33	1.06, 1.67
<u>Alcohol Drinking Status</u>				
Never	1.0		1.0	
Former	1.14	0.93, 1.39	1.17	0.89, 1.53
Current	0.87	0.72, 1.05	0.97	0.74, 1.26
<u>History of CVD, Stroke, TIA</u>				
No	1.0		1.0	
Yes	1.32	1.07, 1.62	1.06	0.84, 1.33
<u>Hypertension</u>				
No	1.0		1.0	
Yes	1.07	0.91, 1.26	1.08	0.89, 1.32
<u>Longest Held Job Categories</u>				
Professional/Managerial	1.0		1.0	
Service	1.30	1.04, 1.61	1.34	0.96, 1.86
Production/Manufacturing/Labor	1.18	0.96, 1.44	1.32	0.98, 1.78
<u>History of Noise Exposure²</u>				
No	1.0		1.0	
Yes	1.39	1.16, 1.66	1.31	1.05, 1.64

	Hearing Impairment		Bilateral Hearing Impairment	
	OR	95%CI	OR	95%CI
<u>Occupational Noise Exposure</u>				
Never	1.0		1.0	
Former	1.38	1.18, 1.60	1.41	1.16, 1.72
Current	1.16	0.95, 1.41	1.09	0.83, 1.45

¹ Adjusted for age, gender, center and Hispanic/Latino background.

² Defined as use of firearms, military service or exposure to loud noises during leisure time.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 4

Multivariate Adjusted Odds Ratios (OR) for Hearing Impairment

	<u>Hearing Impairment</u>		<u>Bilateral Hearing Impairment</u>	
	OR	95% CI	OR	95% CI
Age				
18–44 years	1.0		1.0	
45–64 years	4.84	3.83, 6.12	6.97	4.92, 9.89
65+years	18.52	12.40, 27.68	27.14	17.32, 42.52
Gender				
Female	1.0		1.0	
Male	1.66	1.33, 2.07	1.73	1.32, 2.25
Hispanic/Latino Background				
Dominican	1.42	0.84, 2.41	1.19	0.73, 1.94
Central American	1.02	0.70, 1.49	1.17	0.75, 1.81
Cuban	1.03	0.67, 1.57	0.92	0.57, 1.49
Mexican	1.0		1.0	
Puerto Rican	1.57	1.10, 2.25	1.38	0.94, 2.03
South American	1.01	0.68, 1.51	0.85	0.50, 1.44
Other or mixed	0.76	0.45, 1.29	0.69	0.38, 1.27
Education				
No High school diploma or GED	1.0		1.0	
At least High school diploma or GED	0.71	0.59, 0.86	0.63	0.50, 0.80
Income				
Less than \$10,000	1.0		1.0	
\$10,001 – \$20,000	0.83	0.64, 1.07	0.62	0.47, 0.83
\$20,001 – \$40,000	0.74	0.56, 0.96	0.52	0.38, 0.71
\$40,001 – \$75,000	0.58	0.42, 0.80	0.59	0.39, 0.87
More than \$75,000	0.58	0.36, 0.92	0.34	0.18, 0.64
Diabetes				
No diabetes	1.0		1.0	
Pre-diabetes	1.37	1.12, 1.67	1.44	1.07, 1.94
Diabetes	1.57	1.27, 1.94	1.61	1.22, 2.13
Body Mass Index (Kg/m ²)				
Normal/Underweight (0–24.9)	1.0		1.0	
Overweight (25–29.9)	1.24	0.98, 1.56	1.32	0.98, 1.79
Obese (30+)	1.18	0.94, 1.49	1.29	0.92, 1.80
Smoking Status				
Never	1.0		1.0	
Former	1.16	0.95, 1.40	1.13	0.90, 1.42
Current	1.19	0.97, 1.46	1.17	0.88, 1.56
Alcohol Drinking Status				
Never	1.0		1.0	

	Hearing Impairment		Bilateral Hearing Impairment	
	OR	95% CI	OR	95% CI
Former	1.21	0.94, 1.56	1.54	1.15, 2.06
Current	0.86	0.69, 1.08	1.17	0.86, 1.59
History of CVD, Stroke, TIA				
No	1.0		1.0	
Yes	1.11	0.88, 1.41	1.01	0.76, 1.34
Hypertension				
No	1.0		1.0	
Yes	1.15	0.94, 1.41	1.29	1.02, 1.62
Longest Held Job Categories				
Professional/Managerial	1.0		1.0	
Service Jobs	1.01	0.79, 1.29	1.02	0.70, 1.47
Production/Manufacturing/Labor	0.85	0.67, 1.08	0.89	0.63, 1.27
History of Noise Exposure ¹				
No	1.0		1.0	
Yes	1.35	1.07, 1.70	1.27	0.99, 1.63
Occupational Noise Exposure				
Never	1.0		1.0	
Former	1.27	1.03, 1.57	1.38	1.07, 1.79
Current	1.27	1.00, 1.63	1.23	0.92, 1.65

¹ Defined as use of firearms, military service or exposure to loud noises during leisure time.