

CORRECTION

# Correction: Population Genetic Structure Within and among Seasonal Site Types in the Little Brown Bat (*Myotis lucifugus*) and the Northern Long-Eared Bat (*M. septentrionalis*)

The *PLOS ONE* Staff

There are multiple errors in the fifth sentence of the Abstract. The publisher apologizes for the errors. The correct sentence is: Both species exhibited moderate degrees of mitochondrial DNA differentiation (little brown bat:  $F_{ST(SUMMER)} = 0.093$ ,  $F_{ST(SWARMING)} = 0.052$ ; northern long-eared bat:  $F_{ST(SUMMER)} = 0.117$ ,  $F_{ST(SWARMING)} = 0.043$ ) and little microsatellite DNA differentiation among summering and among swarming sites.

## Reference

1. Johnson LNL, McLeod BA, Burns LE, Arseneault K, Frasier TR, Broders HG (2015) Population Genetic Structure Within and among Seasonal Site Types in the Little Brown Bat (*Myotis lucifugus*) and the Northern Long-Eared Bat (*M. septentrionalis*). PLoS ONE 10(5): e0126309. doi:[10.1371/journal.pone.0126309](https://doi.org/10.1371/journal.pone.0126309) PMID: [25942425](https://pubmed.ncbi.nlm.nih.gov/25942425/)



## OPEN ACCESS

**Citation:** The *PLOS ONE* Staff (2015) Correction: Population Genetic Structure Within and among Seasonal Site Types in the Little Brown Bat (*Myotis lucifugus*) and the Northern Long-Eared Bat (*M. septentrionalis*). PLoS ONE 10(7): e0133457. doi:[10.1371/journal.pone.0133457](https://doi.org/10.1371/journal.pone.0133457)

**Published:** July 20, 2015

**Copyright:** © 2015 The PLOS ONE Staff. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.