



# HHS Public Access

Author manuscript

*Curr HIV/AIDS Rep.* Author manuscript; available in PMC 2015 July 22.

Published in final edited form as:

*Curr HIV/AIDS Rep.* 2011 December ; 8(4): 269–276. doi:10.1007/s11904-011-0092-6.

## Secondary HIV Prevention: Novel Intervention Approaches to Impact Populations Most at Risk

Jennifer L. Brown and Ralph J. DiClemente

Rollins School of Public Health, Emory University, 1518 Clifton Road NE, Room 426, Atlanta, GA 030322, USA

Jennifer L. Brown: jennifer.brown@emory.edu; Ralph J. DiClemente: rdiclem@emory.edu

### Abstract

This paper reviews recent secondary prevention interventions designed to reduce sexual risk behaviors among people living with HIV/AIDS (PLWHA). A summary of findings from previous meta-analyses and narrative reviews of interventions is provided. Next, novel HIV prevention approaches for PLWHA are reviewed. The review reports on the efficacy of interventions delivered in primary care settings or by technology-formats, interventions that also address mental health difficulties, and programs to address particular at-risk populations (eg, men who have sex with men). A critique of recent interventions for people living with HIV/AIDS is provided as well as suggestions for future research.

### Keywords

HIV/AIDS; Public health; Research synthesis; Review; Behavioral interventions; Secondary prevention; Sexual transmission of HIV

### Introduction

The Centers for Disease Control and Prevention estimates that there are now more than 1 million people living with HIV/AIDS (PLWHA) in the United States (US) currently, with approximately 53,600 new HIV infections diagnosed annually [1, 2]. Highly active antiretroviral therapy (HAART) has improved the long-term health outlook of PLWHA and has markedly decreased HIV-related mortality [3, 4]. While an HIV diagnosis is initially accompanied by decreases in sexual risk behavior [5], improvements in PLWHA's health status are associated with increased risky sexual practices [6]. Indeed, one study noted that between 20% and 30% of PLWHA reported unprotected sexual encounters a year after being diagnosed with HIV [7]. Unprotected sexual encounters pose the risk of HIV transmission to uninfected partners and increased possibility of negative health consequences for PLWHA through acquisition of other sexually transmitted infections (STIs) or drug-resistant HIV strains [8–11]. Thus, the development of effective sexual risk reduction interventions for PLWHA is critical from a broader societal perspective, to reduce

Correspondence to: Jennifer L. Brown, jennifer.brown@emory.edu.

**Disclosure** No potential conflicts of interest relevant to this article were reported.

the risk of new infections among HIV-negative sex partners, and from an individual perspective, to reduce the adverse health consequences associated with exposure to STIs and drug-resistant strains of HIV.

Early in the HIV epidemic, prevention efforts focused primarily on *primary prevention* to decrease HIV risk behaviors among uninfected individuals. In contrast, *secondary prevention* refers to interventions designed to reduce sexual risk behaviors among PLWHA. Secondary prevention efforts are essential since only HIV-infected individuals can transmit HIV [12, 13]. Additionally, PLWHA who engage in unprotected sex may acquire syphilis, gonorrhea, chlamydia, or other STIs [8]. In addition to deleterious STI symptoms, comorbid STIs among PLWHA can increase HIV transmission [14, 15] and accelerate HIV disease progression [16]. PLWHA may also be exposed to more virulent multidrug-resistant strains of HIV [9] or HIV super-infection [10, 11]. Given the health risks posed by unprotected sex to PLWHA and their uninfected partners, the World Health Organization identified secondary prevention as an urgent public health priority [17]. Existing reviews of secondary prevention trials for PLWHA suggest promising intervention effects, with reductions in sexual risk practices similar to studies of uninfected populations [18•, 19•].

In this review, we provide a summary of key findings from previous narrative reviews and meta-analyses that have examined the efficacy of secondary prevention interventions for PLWHA [18•, 19•, 20•, 21]. Next, we update the existing reviews by highlighting novel secondary prevention interventions designed to reduce sexual risk behaviors among PLWHA. Specifically, we describe core components of recent intervention approaches, review intervention efficacy findings, summarize intervention strengths and weaknesses, and offer guidance for future research.

## Literature Search Method and Criteria for Inclusion

Database searches of PsycINFO, PsycARTICLES, Medline, and PubMed were conducted to identify published articles in peer-reviewed journals that evaluated sexual risk reduction interventions for PLWHA. Combinations of the following search terms were used to identify relevant articles: HIV prevention, behavioral intervention, secondary prevention, sexual risk reduction, intervention, HIV, HIV-positive, HIV-infected, and AIDS. A search of references cited in relevant studies was conducted to identify additional articles. English language articles published in peer-reviewed journals were screened for inclusion. Studies were included if they met all of the following criteria: 1) the intervention included content to reduce sexual risk behaviors (eg, increase condom use, decrease number of sexual partners); 2) the study's sample was composed of only PLWHA; and 3) the intervention did not focus exclusively on perinatal HIV transmission. For this review, we excluded studies that were not conducted with US samples. Studies included in previous reviews were not separately reviewed; instead a summary of previous meta-analyses and review papers is provided.

## Summary of Previous Secondary Prevention Reviews

To date, two meta-analyses [18•, 19•] and two narrative reviews [20•, 21] have reviewed the secondary prevention literature. In what follows, we provide a succinct summary of findings from each of the reviews. For the meta-analyses, we summarize the format of the

interventions, describe study sample characteristics, report on the efficacy of reviewed interventions, and note intervention characteristics associated with differential intervention efficacy.

Gordon and colleagues [21] first summarized the state of secondary prevention interventions in 2005. This review provided an overview of 13 initial secondary prevention interventions that were either completed or currently in progress. Among the six completed trials [22–27], five were rigorous randomized controlled trials (RCTs) that demonstrated efficacy in modifying sexual risk behaviors. Additionally, of trials currently underway as this review was published, there was preliminary evidence for intervention feasibility and efficacy for programs targeting particular populations at increased HIV risk (eg, serodiscordant couples). This review noted that existing interventions were conducted with individuals or among small groups using variations of social-cognitive theory to guide the intervention framework.

Crepaz and colleagues [18•] conducted a meta-analysis of 12 secondary prevention interventions implemented between 1998 and 2004 [22–33]. The majority of interventions (67%) utilized a small group intervention format led by health care workers (75%) or peers (41%). Most studies recruited primarily males, with only one intervention trial conducted with women. Four of the 12 studies had samples comprised of more than two-thirds men who have sex with men (MSM); three studies enrolled injecting drug users (IDU) or other PLWHA who abuse other substances. Across studies, the interventions were efficacious in reducing the frequency of unprotected sex and STIs. However, no significant effects for reduced needle sharing were observed. Interventions that were based on a behavior change theory and incorporated content to change specific HIV risk behaviors by fostering skill development were most efficacious. In addition, intensive interventions that were delivered by health care workers where PLWHA receive medical or related services demonstrated increased efficacy. Lastly, interventions that also addressed related concerns of PLWHA including mental health problems or difficulty maintaining medication adherence showed particular promise.

In a second meta-analysis, Johnson and colleagues [19•] reviewed 15 secondary prevention interventions published between 1993 and 2004 [22–24, 26–29, 32–41]. Eleven studies were conducted with only HIV-infected individuals, while four studies also included at-risk HIV-uninfected participants. Most studies recruited primarily male (64%) and African American (49%) individuals. Results indicated that the reviewed interventions increased condom use, but there were no differences in the reported number of sexual partners between the intervention and control conditions. Interventions demonstrated increased efficacy when content included both motivational and behavioral skills training. When examining sample characteristics associated with differential efficacy, increased efficacy was noted among younger PLWHA, with interventions conducted with MSM demonstrating limited efficacy in reducing sexual risk behaviors.

In 2009, Fisher and colleagues [20•] reviewed 11 secondary prevention studies conducted with US and international PLWHA [42–52]. Unlike earlier interventions for PLWHA that typically utilized a small group format, studies chronicled in this review utilized a diverse

range of intervention approaches to reduce sexual risk behavior. For example, three studies utilized Motivational Interviewing delivered by health care workers [42, 43] or via computer [44]. Other intervention approaches focused on addressing co-occurring challenges including mental health difficulties (eg, trauma) [48], substance use [49–51], and initiation of antiretroviral medication [45]. Across studies, interventions were successful in reducing sexual risk behaviors, including fewer unprotected sexual encounters and sexual partners.

In summary, secondary prevention interventions demonstrate great promise for reducing a range of sexual risk behaviors. However, there is less support for secondary prevention interventions to decrease needle sharing or the sexual risk behaviors of MSM and older adults. Efficacious interventions were guided by theory (eg, social-cognitive theory) and incorporated content to increase PLWHA's motivation and behavioral skills to reduce sexual risk behaviors. While the earliest secondary prevention interventions were typically delivered in a small group format with most content geared toward reducing sexual risk behaviors, more recent interventions have been implemented within HIV medical settings and address additional challenges faced by PLWHA.

## **Novel Secondary Prevention Intervention Approaches**

The goal of this segment of the review is to highlight novel secondary prevention interventions that have either been published since Fisher and colleagues' review paper [20] or were not included in previous reviews. Interventions fall into one of four categories: 1) interventions delivered in primary care settings; 2) technology-delivered interventions; 3) interventions that seek to reduce sexual risk behaviors and psychological distress; and 4) interventions that target specific at-risk populations including MSM, serodiscordant couples, older individuals, and women.

### **Secondary Prevention in Primary Care Settings**

Given the complex medical and related health care needs of PLWHA there is an emphasis on providing comprehensive services through funding programs such as the Ryan White Act [53]. Such care centers offer the potential to provide sexual risk reduction interventions to PLWHA within the context of their routine medical care. To address this secondary prevention need, the Health Resources and Services Administration funded an initiative to develop and evaluate secondary prevention interventions delivered in the context of primary care settings [54]. While a number of papers have reported on the development and implementation of such intervention programs [55–58], two recent studies report outcomes from secondary prevention interventions delivered by medical providers in primary HIV care [59, 60].

In one study, Gardner and colleagues [60] gave medical providers a 4-hour training session intervention to improve their communication skills and ability to provide behavioral sexual risk reduction counseling. The study measured patients' behaviors over 12 months and observed that patients' frequency of unprotected sexual encounters decreased; more frequent discussions with providers about sexual risk reduction were associated with incremental decreases in sexual risk behavior [60]. In a second study, Rose and colleagues [59] randomized HIV medical care providers to either a 4-hour training to improve their ability to

conduct sexual risk behavior assessments and deliver tailored, individualized risk reduction messages or a standard care condition. At a 6-month follow-up, patients whose providers completed the training reported increased discussions surrounding safer sex and more frequent assessment of risk behaviors by their providers; patients also reported having fewer sexual partners [59].

### **Technology-Delivered Secondary Prevention Programs**

Secondary prevention interventions have typically been administered in a group format led by a health care provider or an experienced health educator. These treatment programs often involved multiple sessions and a significant time commitment. Although group-delivered training provides the added benefit of fostering social support, some PLWHA may be reticent to attend groups due to confidentiality concerns or dislike of group meetings. Thus, the usefulness of group sexual risk reduction interventions may be limited to the subset of patients who are willing to participate in a group-based program. Group interventions also require significant financial resources to implement and may not be feasible in some resource-limited clinic settings. To address these concerns, a promising approach is the use of technology-delivered approaches to secondary prevention.

One such intervention utilized a web-based sexual risk reduction intervention for adolescent PLWHA that was accessed by youth while waiting for their scheduled medical appointments [61]. In a pilot-test of this intervention, adolescents and clinic staff found the program to be acceptable and feasible within the constraints of usual clinic flow [61]. Additionally, the intervention demonstrated efficacy for increasing antecedent constructs of sexual behavior, including condom use self-efficacy and increased desire to wait before having sex [61]. A second intervention utilized a computerized assessment within a primary care setting to conduct a sexual behavior risk assessment and measure an individual's readiness for changing their sexual behavior [62]. The results of this computerized assessment generated a provider advice sheet and a prescription to reduce the reported risk behaviors [62]. In turn, providers utilized the risk assessment and behavior change prescription to discuss sexual risk reduction with their patients [62]. In the initial process evaluation of the intervention, HIV medical service providers reported increased confidence in discussing sexual risk strategies with their patients [62]. However, this study also found that providers' attitudes toward HIV prevention did not improve as the result of the intervention.

### **Secondary Prevention and Mental Health Interventions**

Evidence suggests that PLWHA experience high rates of mental health difficulties and substance abuse problems [63, 64]. Recent reviews suggest that cognitive-behavioral interventions tailored to the unique concerns of PLWHA can result in significant decreases in depressive symptoms, anxiety levels, and overall stress [65–67]. A number of studies also highlight the efficacy of substance use interventions for PLWHA to reduce the use of alcohol [68, 69], opiates [70], injection drugs [71], and crack cocaine [72]. While there are efficacious interventions to address the psychological needs of PLWHA, PLWHA with mental illness often have limited access to psychiatric treatment and do not receive adequate mental health services [73]. Unfortunately, untreated mental illness may be associated with increased health risk behaviors by PLWHA, including unprotected sexual encounters [74,

75]. Additionally, a minority of secondary prevention interventions incorporates mental health treatment [76, 77].

Despite the clear need for both mental health treatment and secondary prevention, few interventions have addressed both of these needs. In one such intervention trial, Wyatt and colleagues [78] developed and evaluated an intervention for HIV-infected women with histories of child abuse. The intervention consisted of 11 cognitive-behavioral sessions designed to reduce sexual risk behaviors, improve medication adherence, and address psychological symptoms [78]. This intervention had a particular focus on addressing the association between childhood sexual abuse and the individual's current cognitive, affective, and behavioral patterns [78]. Results indicate that women assigned to the treatment group reduced their sexual risk behaviors and reported improved medication adherence, particularly among participants attending more of the group sessions [78]. Similarly, Sikkema and colleagues [48] evaluated a 15-session intervention for HIV-infected women and men with abuse histories and found that participants in the treatment condition reported decreased frequency of unprotected sex relative to the support group control condition.

### Secondary Prevention Tailored to Specific At-Risk Populations

**MSM**—MSM continue to experience elevated rates of HIV infection [79], but previous interventions with MSM have demonstrated limited efficacy in reducing sexual risk behaviors [19•]. Two recent studies have examined the efficacy of secondary prevention interventions for MSM [80, 81]. Rosser and colleagues [81] recruited 675 HIV-infected MSM who were randomized to one of three conditions: 1) a weekend, 14-hour to 16-hour, large group-delivered intervention to reduce sexual risk behaviors with a focus *specific to HIV-infected MSM*; 2) a weekend, 14-hour to 16-hour, large group-delivered intervention to reduce sexual risk behaviors with a focus on *MSM more broadly*; or 3) a 3-hour group session utilizing HIV prevention videos for MSM. Those in the two more intensive intervention conditions reported greater intentions to reduce sexual risk behaviors; however, there were equivalent reductions in unprotected anal sexual encounters with serodiscordant partners across conditions [81]. In a second trial, Lapinski and colleagues [80] compared the efficacy of an intervention combining group and individual-counseling to an individual-counseling only condition to improve communication skills related to sexual risk reduction. HIV-infected MSM in the combined treatment group demonstrated improvements in desire to not engage in sexual encounters when under the influence of alcohol or drugs, improved communication skills, and increased likelihood of disclosing their HIV serostatus to partners [80]. However, there were no differences in perceived norms regarding sexual behaviors or HIV knowledge [80].

**Serodiscordant Couples**—In the US, African Americans have experienced disproportionately elevated rates of HIV/AIDS relative to other racial groups [1]. Among African American women, the most common form of HIV transmission is via heterosexual contact, and this is the second most common transmission mode among African American men [1, 82]. To reduce rates of HIV transmission, the multisite Eban intervention was conducted to examine the efficacy of couples-based secondary prevention among couples with one HIV-infected partner and one uninfected partner [83]. In this large randomized

controlled trial, 535 couples were randomly assigned to either an intervention consisting of eight weekly 2-hour sexual risk reduction sessions led by male and female co-facilitators or a time-equivalent health promotion comparison condition [83]. Couples in the treatment condition reported increased condom-protected encounters, consistent condom use, and fewer unprotected sexual encounters at the 12-month follow-up [83]. However, there were no differences in STI rates or rates of concurrent partners between the intervention and control conditions over the follow-up period [83].

**Older Adults**—In the meta-analysis conducted by Johnson and colleagues [19•] they found that secondary prevention interventions demonstrated greater efficacy for sexual risk reduction among younger individuals. Given the improved long-term prognosis of PLWHA with HAART medication, there is a clear need for secondary prevention programs to sustain engagement in safer sexual practices for older individuals. However, only one intervention has been designed for older PLWHA [84]. The intervention consisted of four sessions based on the Information-Motivation-Behavior skills model of HIV risk behavior tailored to the prevention needs of PLWHA who were 45 years or older [84]. Relative to a comparison condition consisting of an educational HIV prevention brochure, those in the treatment group reported decreased unprotected sexual encounters with all partners, but particularly with HIV-negative or unknown serostatus partners [84]. However, participants in both conditions demonstrated overall reductions in sexual risk behaviors [84].

**Women**—Rates of HIV infection are on the rise among several subgroups in the US, including low-income, minority women [82]. However, relatively few secondary prevention programs have been developed to decrease sexual risk behaviors among HIV-infected women. As described in previous reviews, Wingood and colleagues [27] developed an efficacious four-session gender and culturally tailored intervention for African American HIV-infected women that resulted in decreased unprotected sexual encounters, fewer STIs, and improved related HIV risk reduction constructs (eg, condom use self-efficacy, HIV knowledge). In a newly developed intervention, Teti and colleagues [85] described the development of a multifaceted intervention for HIV-infected women that included prevention messages delivered by PLWHA's clinicians, a group-based intervention led by a health educator, and a peer-led support group. While a multicomponent intervention shows promise, the authors identified a number of feasibility concerns including staff burden for administering the intervention and challenges enrolling and retaining women in the program that resulted in poor intervention participation [85].

## Conclusions

The first generation of secondary prevention interventions demonstrates promise for reducing sexual risk behaviors among PLWHA [18•, 19•, 20•]. While previous intervention approaches typically relied on multiple group-delivered intervention sessions, more recent secondary prevention interventions have been integrated within routine medical care for HIV using formats that have the potential to be more cost effective to implement (eg, brief provider-delivered approaches). In particular, interventions to improve HIV medical providers' ability to incorporate sexual risk reduction within routine care show potential for delivering tailored, individualized prevention messages to PLWHA on an ongoing basis.

Additionally, technology-based risk assessments may serve as an effective adjunct to secondary prevention within primary care. To advance such interventions, strategies to bolster the long-term efficacy of provider-delivered interventions are needed. Additional research to understand characteristics of PLWHA for whom briefer, provider-delivered prevention is most effective is needed. In turn, the use of an adaptive intervention approach [86] could facilitate ongoing assessment of sexual risk behaviors and guide the selection of an appropriate intervention approach for a particular individual.

Newer interventions that provide a combined focus on sexual risk reduction and mental health treatment [76, 78] may be more acceptable to PLWHA [87] and offer the potential to reduce both psychological distress and sexual risk behaviors. Indeed, both of the reviewed interventions were successful in reducing sexual risk behaviors, while also addressing psychological distress associated with a history of childhood sexual abuse [76, 78]. Unfortunately, secondary prevention interventions to address sexual risk reduction for PLWHA with more severe mental illness are lacking [65, 77]. First, improving strategies to identify PLWHA with mental health concerns as part of routine HIV care is needed [73]. Psychiatric treatment provided within the context of comprehensive HIV care may be an optimal forum to integrate secondary prevention. Similar to the primary care model of secondary prevention delivery, mental health professionals treating PLWHA may benefit from additional training to address HIV-specific concerns and incorporate prevention messages as a part of mental health treatment.

With the improved long-term prognosis following an HIV diagnosis, secondary prevention intervention approaches to target subpopulations at increased risk for either transmitting HIV or experiencing negative health consequences are needed. Interventions such as Eban that provided a couples-delivered intervention demonstrated efficacy to address the complicated sexual behavior dynamics within serodiscordant couples [83]. Interventions that move beyond individual-level approaches to behavior change should be examined to address sexual risk behavior reduction within the broader socio-ecological context. For example, secondary prevention interventions could be bolstered by complementary structural interventions [21]. In addition, efficacious interventions are needed for specific subpopulations. Thus far, secondary prevention interventions have demonstrated limited efficacy for MSM. Targeting other high-risk groups, such as incarcerated individuals or those with comorbid substance abuse or mental illness, is also urgently needed.

As we enter the third decade of the HIV epidemic, continued secondary prevention intervention efforts are needed to reduce the number of new HIV infections and promote PLWHA's sexual health. While initial prevention efforts were directed toward primary prevention, the growing body of secondary prevention literature highlights a number of efficacious intervention approaches. Moving forward, strategies to disseminate efficacious interventions to reach a larger number of PLWHA are needed. Programs such as the CDC's Dissemination of Evidenced-Based Interventions offer one approach to promote widespread use of secondary prevention interventions. Ultimately, intervention approaches that can be effectively delivered in resource-limited settings and result in reduced sexual risk behaviors offer great potential to improve the health of PLWHA and also prevent the spread of HIV to uninfected partners.



## Acknowledgments

This research was supported by a grant from the National Institute of Mental Health (5R01 MH070537) to Ralph J. DiClemente (PI). Additional support was provided by the Emory Center for AIDS Research (P30 AI050409). Jennifer L. Brown was supported by K12 GM000680 from the National Institute of General Medical Sciences.

## References

Papers of particular interest, published recently, have been highlighted as:

- Of importance

1. CDC. HIV prevalence estimates, United States, 2006. *Morb Mortal Wkly Rep.* 2008; 57:1073–6.
2. CDC. HIV Surveillance Report, 2008. Atlanta: 2010.
3. Schackman BR, Gebo KA, Walensky RP, Losina E, Muccio T, Sax PE, et al. The lifetime cost of current human immunodeficiency virus care in the United States. *Med Care.* 2006; 44:990–7. [PubMed: 17063130]
4. Palella FJ Jr, Delaney KM, Moorman AC, Loveless MO, Fuhrer J, Satten GA, et al. Declining morbidity and mortality among patients with advanced human immunodeficiency virus infection. HIV Outpatient Study Investigators. *N Engl J Med.* 1998; 338:853–60. [PubMed: 9516219]
5. Marks G, Crepaz N, Senterfitt JW, Janssen RS. Meta-analysis of high-risk sexual behavior in persons aware and unaware they are infected with HIV in the United States: implications for HIV prevention programs. *J Acquir Immune Defic Syndr (1999).* 2005; 39:446–53.
6. Eaton LA, Kalichman SC. Changes in transmission risk behaviors across stages of HIV disease among people living with HIV. *J Assoc Nurses AIDS Care: JANAC.* 2009; 20:39–49. [PubMed: 19118770]
7. Weinhardt, LS. HIV diagnosis and risk behavior. In: Kalichman, SC., editor. *Positive prevention: reducing HIV transmission among people living with HIV/AIDS.* New York: Springer; 2005.
8. Kalichman SC, Pellowski J, Turner C. Prevalence of sexually transmitted co-infections in people living with HIV/AIDS: systematic review with implications for using HIV treatments for prevention. *Sex Transm Infect.* 2011; 87:183–90. [PubMed: 21330572]
9. Taiwo B. Understanding transmitted HIV resistance through the experience in the USA. *Int J Infect Dis: IJID: Official Publication of the International Society for Infectious Diseases.* 2009; 13:552–9.
10. Campbell MS, Gottlieb GS, Hawes SE, Nickle DC, Wong KG, Deng W, et al. HIV-1 superinfection in the antiretroviral therapy era: are seroconcordant sexual partners at risk? *Plos One.* 2009; 4:e5690. [PubMed: 19479055]
11. Brenner B, Routy J-P, Quan Y, Moisi D, Oliveira M, Turner D, et al. Persistence of multidrug-resistant HIV-1 in primary infection leading to superinfection. *AIDS (London, England).* 2004; 18:1653–60.
12. DiClemente RJ, Wingood GM, Del Rio C, Crosby RA. Prevention interventions for HIV positive individuals. *Sex Transm Infect.* 2002; 78:393–5. [PubMed: 12473796]
13. DiClemente, RJ. Looking forward: future directions for HIV prevention research. In: Peterson, JL.; DiClemente, RJ., editors. *Handbook of HIV prevention.* New York: Kluwer/Plenum; 2000. p. 311-24.
14. Ward H, Rönn M. Contribution of sexually transmitted infections to the sexual transmission of HIV. *Curr Opin HIV AIDS.* 2010; 5:305–10. [PubMed: 20543605]
15. Royce RA, Seña A, Cates W Jr, Cohen MS. Sexual transmission of HIV. *N Engl J Med.* 1997; 336:1072–8. [PubMed: 9091805]
16. White MK, Gorrill TS, Khalili K. Reciprocal transactivation between HIV-1 and other human viruses. *Virology.* 2006; 352:1–13. [PubMed: 16725168]
17. Bunnell R, Mermin J, De Cock KM. HIV prevention for a threatened continent: implementing positive prevention in Africa. *JAMA: J Am Med Assoc.* 2006; 296:855–8.
- 18•. Crepaz N, Lyles CM, Wolitski RJ, Passin WF, Rama SM, Herbst JH, et al. Do prevention interventions reduce HIV risk behaviours among people living with HIV? A meta-analytic

- review of controlled trials. *AIDS*. 2006; 20:143–57. This paper provides a meta-analysis of the efficacy of secondary prevention interventions for PLWHA. [PubMed: 16511407]
19. Johnson BT, Carey MP, Chaudoir SR, Reid AE. Sexual risk reduction for persons living with HIV research synthesis of randomized controlled trials, 1993 to 2004. *JAIDS: J Acquir Immune Defic Syndr*. 2006; 41:642–50. This meta-analysis of secondary prevention interventions highlights key intervention features associated with differential intervention efficacy. [PubMed: 16652039]
  20. Fisher JD, Smith L. Secondary prevention of HIV infection: the current state of prevention for positives. *Curr Opin HIV AIDS*. 2009; 4:279–87. This paper provides an excellent review of secondary prevention interventions for PLWHA conducted both domestically in the US and also internationally. [PubMed: 19532065]
  21. Gordon CM, Forsyth AD, Stall R, Cheever LW. Prevention interventions with persons living with HIV/AIDS: state of the science and future directions. *AIDS Educ Prev*. 2005; 17:6–20. [PubMed: 15843114]
  22. Kalichman SC, Rompa D, Cage M, DiFonzo K, Simpson D, Austin J, et al. Effectiveness of an intervention to reduce HIV transmission risks in HIV-positive people. *Am J Prev Med*. 2001; 21:84–92. [PubMed: 11457627]
  23. Rotheram-Borus MJ, Lee MB, Murphy DA, Futterman D, Duan N, Birnbaum JM, et al. Efficacy of a preventive intervention for youths living with HIV. *Am J Public Health*. 2001; 91:400–5. [PubMed: 11236404]
  24. Margolin A, Avants SK, Warburton LA, Hawkins KA, Shi JA. A randomized clinical trial of a manual-guided risk reduction intervention for HIV-positive injection drug users. *Heal Psychol*. 2003; 22:223–8.
  25. Patterson TL, Shaw WS, Semple SJ. Reducing the sexual risk behaviors of HIV+ individuals: outcome of a randomized controlled trial. *Annals Behav Med*. 2003; 25:137–45.
  26. Richardson JL, Milam J, McCutchan A, Stoyanoff S, Bolan R, Weiss J, et al. Effect of brief safer-sex counseling by medical providers to HIV-1 seropositive patients: a multi-clinic assessment. *AIDS*. 2004; 18:1179–86. [PubMed: 15166533]
  27. Wingood GM, DiClemente RJ, Mikhail I, Lang DL, Hubbard McCree D, Davies SL, et al. A randomized controlled trial to reduce HIV transmission risk behaviors and sexually transmitted diseases among women living with HIV: the willow program. *J Acquir Immune Defic Syndr*. 2004; 37:S58–67. [PubMed: 15385901]
  28. Rotheram-Borus MJ, Swendeman D, Comulada WS, Weiss RE, Lee M, Lightfoot M. Prevention for substance-using HIV-positive young people: telephone and in-person delivery. *J Acquir Immune Defic Syndr* (1999). 2004; 37 (Suppl 2):S68–77.
  29. Cleary PD, Van Devanter N, Steilen M, Stuart A, Shipton-Levy R, McMullen W, et al. A randomized trial of an education and support program for HIV-infected individuals. *AIDS* (London, England). 1995; 9:1271–8.
  30. Grinstead O, Zack B, Faigeles B. Reducing postrelease risk behavior among HIV seropositive prison inmates: the health promotion program. *AIDS Educ Prev: Official Publication Of The International Society For AIDS Education*. 2001; 13:109–19.
  31. Sorensen JL, Dille J, London J, Okin RL, Delucchi KL, Phibbs CS. Case management for substance abusers with HIV/AIDS: a randomized clinical trial. *Am J Drug Alcohol Abuse*. 2003; 29:133–50. [PubMed: 12731685]
  32. Kelly JA, Murphy DA, Bahr GR, Kalichman SC, Morgan MG, Stevenson LY, et al. Outcome of cognitive-behavioral and support group brief therapies for depressed, HIV-infected persons. *Am J Psychiatr*. 1993; 150:1679–86. [PubMed: 8214177]
  33. Wolitski RJ, Gomez CA, Parsons JT. Effects of a peer-led behavioral intervention to reduce HIV transmission and promote serostatus disclosure among HIV-seropositive gay and bisexual men. *AIDS*. 2005; 19:S99–S109. [PubMed: 15838199]
  34. Roffman RA, Picciano JF, Ryan R, Beadnell B, Fisher D, Downey L, et al. HIV-prevention group counseling delivered by telephone: an efficacy trial with gay and bisexual men. *AIDS Behav*. 1997; 1:137–54.

35. El-Bassel N, Witte SS, Gilbert L, Wu E, Chang M, Hill J, et al. The efficacy of a relationship-based HIV/STD prevention program for heterosexual couples. *Am J Public Health*. 2003; 93:963–9. [PubMed: 12773363]
36. Fogarty LA, Heilig CM, Armstrong K, Cabral R, Galavotti C, Gielen AC, et al. Long-term effectiveness of a peer-based intervention to promote condom and contraceptive use among HIV-positive and at-risk women. *Public Health Reports (Washington, DC: 1974)*. 2001; 116 (Suppl 1): 103–19.
37. Wolitski RJ, Parsons JT, Gomez CA. Prevention with HIV-seropositive men who have sex with men: lessons from the Seropositive Urban Men’s Study (SUMS) and the Seropositive Urban Men’s Intervention Trial (SUMIT). *J Acquir Immune Defic Syndr*. 2004; 37:S101–9. [PubMed: 15385906]
38. Dushay RA, Singer M, Weeks MR, Rohena L, Gruber R. Lowering HIV risk among ethnic minority drug users: comparing culturally targeted intervention to a standard intervention. *Am J Drug Alcohol Abuse*. 2001; 27:501–24. [PubMed: 11506265]
39. Gielen AC, Fogarty LA, Armstrong K, Green BM, Cabral R, Milstein B, et al. Promoting condom use with main partners: a behavioral intervention trial for women. *AIDS Behav*. 2001; 5:193–204.
40. MacNeil JM, Mberesero F, Kilonzo G. Is care and support associated with preventive behaviour among people with HIV? *AIDS Care*. 1999; 11:537–46. [PubMed: 10755029]
41. The National Institute of Mental Health (NIMH) Multisite HIV Prevention Trial Group. . The NIMH Multisite HIV Prevention Trial: reducing HIV sexual risk behavior. *Science (New York, NY)*. 1998; 280:1889–94.
42. Fisher JD, Fisher WA, Cornman DH, Amico RK, Bryan A, Friedland GH. Clinician-delivered intervention during routine clinical care reduces unprotected sexual behavior among HIV-infected patients. *J Acquir Immune Defic Syndr (1999)*. 2006; 41:44–52.
43. Cornman DH, Kiene SM, Christie S, Fisher WA, Shuper PA, Pillay S, et al. Clinic-based intervention reduces unprotected sexual behavior among HIV-infected patients in KwaZulu-Natal, South Africa: results of a pilot study. *J Acquir Immune Defic Syndr (1999)*. 2008; 48:553–60.
44. Gilbert P, Ciccarone D, Gansky SA, Bangsberg DR, Clanon K, McPhee SJ, et al. Interactive “Video Doctor” counseling reduces drug and sexual risk behaviors among HIV-positive patients in diverse outpatient settings. *Plos One*. 2008; 3:e1988. [PubMed: 18431475]
45. Bonell C, Hargreaves J, Strange V, Pronyk P, Porter J. Should structural interventions be evaluated using RCTs? The case of HIV prevention. *Soc Sci Med*. 2006; 63:1135–42. [PubMed: 16697512]
46. Jones DL, Weiss SM, Bhat GJ, Bwalya V. Influencing sexual practices among HIV-positive Zambian women. *AIDS Care*. 2006; 18:629–34. [PubMed: 16831792]
47. Sikkema KJ, Kalichman SC, Hoffmann R, Koob JJ, Kelly JA, Heckman TG. Coping strategies and emotional wellbeing among HIV-infected men and women experiencing AIDS-related bereavement. *AIDS Care*. 2000; 12:613–24. [PubMed: 11218547]
48. Sikkema KJ, Wilson PA, Hansen NB, Kochman A, Neufeld S, Ghebremichael MS, et al. Effects of a coping intervention on transmission risk behavior among people living with HIV/AIDS and a history of childhood sexual abuse. *J Acquir Immune Defic Syndr (1999)*. 2008; 47:506–13.
49. Jones TS, Vlahov D. What we can learn from the INSPIRE Study about improving prevention and clinical care for injection drug users living with HIV. *J Acquir Immune Defic Syndr (1999)*. 2007; 46 (Suppl 2):S31–4.
50. Mausbach BT, Semple SJ, Strathdee SA, Zians J, Patterson TL. Efficacy of a behavioral intervention for increasing safer sex behaviors in HIV-positive MSM methamphetamine users: results from the EDGE study. *Drug Alcohol Depend*. 2007; 87:249–57. [PubMed: 17182196]
51. Margolin A, Schuman-Olivier Z, Beitel M, Arnold RM, Fulwiler CE, Avants SK. A preliminary study of spiritual self-schema (3-S (+)) therapy for reducing impulsivity in HIV-positive drug users. *J Clin Psychol*. 2007; 63:979–99. [PubMed: 17828761]
52. Effects of a behavioral intervention to reduce risk of transmission among people living with HIV: the healthy living project randomized controlled study. *J Acquir Immune Defic Syndr*. 2007; 44:213–21. [PubMed: 17146375]
53. Ryan White HIV/AIDS Treatment Extension Act of 2009. 2009.

54. Malitz FE, Eldred L. Evolution of the special projects of national significance prevention with HIV-infected persons seen in primary care settings initiative. *AIDS Behav.* 2007; 11:S1–5. [PubMed: 17546495]
55. Callahan EJ, Flynn NM, Kuenneth CA, Enders SR. Strategies to reduce HIV risk behavior in HIV primary care clinics: brief provider messages and specialist intervention. *AIDS Behav.* 2007; 11:S48–57. [PubMed: 17205389]
56. Nollen C, Drainoni M-L, Sharp V. Designing and delivering a prevention project within an HIV treatment setting: lessons learned from a specialist model. *AIDS Behav.* 2007; 11:S84–94. [PubMed: 17577654]
57. Grimley DM, Bachmann LH, Jenckes MW, Erbeling EJ. Provider-delivered, theory-based, individualized prevention interventions for HIV positive adults receiving HIV comprehensive care. *AIDS Behav.* 2007; 11:S39–47. [PubMed: 17149671]
58. Koester KA, Maiorana A, Vernon K, Myers J, Rose CD, Morin S. Implementation of HIV prevention interventions with people living with HIV/AIDS in clinical settings: challenges and lessons learned. *AIDS Behav.* 2007; 11:S17–29. [PubMed: 17436072]
59. Rose CD, Courtenay-Quirk C, Knight K, Shade SB, Vittinghoff E, Gomez C, et al. HIV intervention for providers study: a randomized controlled trial of a clinician-delivered HIV risk-reduction intervention for HIV-positive people. *JAIDS J Acquir Immune Defic Syndr.* 2010; 55:572–81. [PubMed: 20827218]
60. Gardner LI, Marks G, O'Daniels CM, Wilson TE, Golin C, Wright J, et al. Implementation and evaluation of a clinic-based behavioral intervention: positive steps for patients with HIV. *AIDS Patient Care STDs.* 2008; 22:627–35. [PubMed: 18627280]
61. Markham CM, Shegog R, Leonard AD, Bui TC, Paul ME. +CLICK: harnessing web-based training to reduce secondary transmission among HIV-positive youth. *AIDS Care.* 2009; 21:622–31. [PubMed: 19444671]
62. Chen HT, Grimley DM, Waithaka Y, Aban IB, Hu J, Bachmann LH. A process evaluation of the implementation of a computer-based, health provider-delivered HIV-prevention intervention for HIV-positive men who have sex with men in the primary care setting. *AIDS Care.* 2008; 20:51–60. [PubMed: 18278615]
63. Klinkenberg WD, Sacks S. Mental disorders and drug abuse in persons living with HIV/AIDS. *AIDS Care.* 2004; 16:S22–42. [PubMed: 15736820]
64. Galvan FH, Bing EG, Fleishman JA, London AS, Caetano R, Burnam MA, et al. The prevalence of alcohol consumption and heavy drinking among people with HIV in the United States: results from the HIV cost and services utilization study. *J Stud Alcohol.* 2002; 63:179–86. [PubMed: 12033694]
65. Brown JL, Venable PA. Cognitive-behavioral stress management interventions for persons living with HIV: a review and critique of the literature. *Annals Behav Med.* 2008; 35:26–40.
66. Crepaz N, Passin WF, Herbst JH, Rama SM, Malow RM, Purcell DW, et al. Meta-analysis of cognitive-behavioral interventions on HIV+ persons' mental health and immune functioning. *Heal Psychol.* 2008; 27:4–14.
67. Scott-Sheldon LAJ, Kalichman SC, Carey MP, Fielder RL. Stress management interventions for HIV+ adults: a meta-analysis of randomized controlled trials, 1989 to 2006. *Heal Psychol.* 2008; 27:129–39.
68. Morgenstern J, Irwin TW, Wainberg ML, Parsons JT, Muench F, Bux DA Jr, et al. A randomized controlled trial of goal choice interventions for alcohol use disorders among men who have sex with men. *J Consult Clin Psychol.* 2007; 75:72–84. [PubMed: 17295566]
69. Velasquez MM, von Sternberg K, Johnson DH, Green C, Carbonari JP, Parsons JT. Reducing sexual risk behaviors and alcohol use among HIV-positive men who have sex with men: a randomized clinical trial. *J Consult Clin Psychol.* 2009; 77:657–67. [PubMed: 19634959]
70. Marsch LA. The efficacy of methadone maintenance interventions in reducing illicit opiate use, HIV risk behavior and criminality: a meta-analysis. *Addiction.* 1998; 93:515–32. [PubMed: 9684390]

71. Rotheram-Borus MJ, Rhodes F, Desmond K, Weiss RE. Reducing HIV risks among active injection drug and crack users: the safety counts program. *AIDS Behav.* 2010; 14:658–68. [PubMed: 19757019]
72. Weeks MR, Li J, Dickson-Gomez J, Convey M, Martinez M, Radda K, et al. Outcomes of a peer HIV prevention program with injection drug and crack users: the risk avoidance partnership. *Subst Use Misuse.* 2009; 44:253–81. [PubMed: 19142824]
73. Stepleman LM, Hann G, Santos M, House AS. Reaching under-served HIV-positive individuals by using patient-centered psychological consultation. *Prof Psychol Res Pract.* 2006; 37:75–82.
74. Gore-Felton C, Koopman C, Turner-Cobb JM, Duran R, Israelski D, Spiegel D. The influence of social support, coping and mood on sexual risk behavior among HIV+ men and women. *J Heal Psychol.* 2002; 7:713–22.
75. Morin SF, Steward WT, Charlebois ED, Remien RH, Pinkerton SD, Johnson MO, et al. Predicting HIV transmission risk among HIV-infected men who have sex with men: findings from the healthy living project. *JAIDS J Acquir Immune Defic Syndr.* 2005; 40:226–35. [PubMed: 16186742]
76. Sikkema KJ, Watt MH, Drabkin AS, Meade CS, Hansen NB, Pence BW. Mental health treatment to reduce HIV transmission risk behavior: a positive prevention model. *AIDS Behav.* 2010; 14:252–62. [PubMed: 20013043]
77. Grossman CI, Gordon CM. Mental health considerations in secondary HIV prevention. *AIDS Behav.* 2010; 14:263–71. [PubMed: 19057989]
78. Wyatt GE, Longshore D, Chin D, Carmona JV, Loeb TB, Myers HF, et al. The efficacy of an integrated risk reduction intervention for HIV-positive women with child sexual abuse histories. *AIDS Behav.* 2004; 8:453–62. [PubMed: 15690118]
79. CDC. High-risk sexual behavior by HIV-positive men who have sex with men: 16 sites, United States 2000–2002. *Morb Mortal Wkly Rep.* 2004; 53:891–910.
80. Lapinski MK, Randall LM, Peterson M, Peterson A, Klein KA. Prevention options for positives: the effects of a health communication intervention for men who have sex with men living with HIV/AIDS. *Heal Commun.* 2009; 24:562–71.
81. Rosser BR, Oakes JM, Konstan J, Hooper S, Horvath KJ, Danilenko GP, et al. Reducing HIV risk behavior of men who have sex with men through persuasive computing: results of the Men's INternet Study-II. *AIDS.* 2010; 24:2099–107. [PubMed: 20601853]
82. CDC. HIV/AIDS among women: fact sheet. Atlanta: 2006.
83. El-Bassel N, Jemmott JB, Landis JR, Pequegnat W, Wingood GM, Wyatt GE, et al. National Institute of Mental Health Multisite Eban HIV/STD Prevention Intervention for African American HIV Serodiscordant Couples: a cluster randomized trial. *Arch Int Med.* 2010; 170:1594–601. [PubMed: 20625011]
84. Illa L, Echenique M, Jean GS, Bustamante-Avellaneda V, Metsch L, Mendez-Mulet L, et al. Project ROADMAP: reeducating older adults in maintaining AIDS prevention: a secondary intervention for older HIV-positive adults. *AIDS Educ Prev: Official Publication Of The International Society For AIDS Education.* 2010; 22:138–47.
85. Teti M, Rubinstein S, Lloyd L, Aaron E, Merron-Brainerd J, Spencer S, et al. The Protect and Respect program: a sexual risk reduction intervention for women living with HIV/AIDS. *AIDS Behav.* 2007; 11:S106–16. [PubMed: 17605102]
86. Luce BR, Kramer JM, Goodman SN, Connor JT, Tunis S, Whicher D, et al. Rethinking randomized clinical trials for comparative effectiveness research: the need for transformational change. *Annals Intern Med.* 2009; 151:206–9.
87. Vanable, PA.; Carey, MP.; Brown, JL.; Littlewood, RA.; Bostwick, RA.; Blair, DC. What HIV+ MSM want from sexual risk reduction interventions: findings from a qualitative study. under review