

Crystal structure of 1-[(2,4,6-triisopropylphenyl)sulfonyl]aziridine

Lena Knauer, Christopher Golz and Carsten Strohmann*

Fakultät für Chemie und Chemische Biologie, Technische Universität Dortmund, Otto-Hahn-Strasse 6, 44227 Dortmund, Germany. *Correspondence e-mail: carsten.strohmann@tu-dortmund.de

Received 30 April 2015; accepted 27 May 2015

Edited by U. Flörke, University of Paderborn, Germany

The title compound, $C_{17}H_{27}NO_2S$, exhibits a distorted geometry of the aromatic ring with elongated bonds at the *ipso*-C atom. The S atom deviates from the aromatic ring plane by 0.393 (4) Å. Similar to this, the adjacent isopropyl groups are bent out of the aromatic ring plane by -0.125 (4) and -0.154 (4) Å. Even the distant isopropyl group in *para*-position to the sulfonyl moiety shows a slight deviation from the ring plane of 0.111 (5) Å. These distortions, which are caused by the bulky substituents, can also be observed in related sulfonylaziridine structures.

Keywords: crystal structure; aziridine; triisopropylbenzenesulfonyl; consecutive ring-opening reactions.

CCDC reference: 1403377

1. Related literature

For the crystal structure of a related phenyl-substituted compound, see: Golz *et al.* (2014). For a discussion of the geometry of the triisopropylbenzenesulfonyl moiety, see: Sandrock *et al.* (2004). For a discussion of the pyramidalized geometry of *N*-sulfonylamides, see: Ohwada *et al.* (1998). By regioselective ring opening reactions, countless nitrogen-containing compounds are accessible, see: Stamm (1999); Schneider (2009). For consecutive ring-opening reactions of aziridines by triethylamine, see: Golz & Strohmann (2015). In some cases, the three-membered aziridine ring is further activated by electron-withdrawing groups (Hu, 2004) to increase its reactivity.

2. Experimental

2.1. Crystal data

$C_{17}H_{27}NO_2S$	$V = 1771.5$ (3) Å ³
$M_r = 309.45$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 6.2679$ (8) Å	$\mu = 0.19$ mm ⁻¹
$b = 17.5289$ (18) Å	$T = 173$ K
$c = 16.3890$ (13) Å	$0.33 \times 0.25 \times 0.01$ mm
$\beta = 100.331$ (10)°	

2.2. Data collection

Agilent Xcalibur Sapphire3 diffractometer	9521 measured reflections
Absorption correction: multi-scan (<i>CrysAlis PRO</i> ; Agilent, 2013)	3449 independent reflections
$T_{\min} = 0.782$, $T_{\max} = 1.000$	2479 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.049$

2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.053$	196 parameters
$wR(F^2) = 0.141$	H-atom parameters constrained
$S = 1.07$	$\Delta\rho_{\text{max}} = 0.33$ e Å ⁻³
3449 reflections	$\Delta\rho_{\text{min}} = -0.58$ e Å ⁻³

Data collection: *CrysAlis PRO* (Agilent, 2013); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009); software used to prepare material for publication: *OLEX2*.

Acknowledgements

We are grateful to the Deutsche Forschungsgemeinschaft (DFG) for financial support.

Supporting information for this paper is available from the IUCr electronic archives (Reference: FK2088).

References

- Agilent (2013). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, Oxfordshire, England.
- Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
- Golz, C., Preut, H. & Strohmann, C. (2014). *Acta Cryst. E* **70**, o153.
- Golz, C. & Strohmann, C. (2015). *Acta Cryst. E* **71**, 564–566.
- Hu, X. E. (2004). *Tetrahedron*, **60**, 2701–2743.

- Ohwada, T., Okamoto, I., Shudo, K. & Yamaguchi, K. (1998). *Tetrahedron Lett.* **39**, 7877–7880.
- Sandrock, P. B., Meyers, C. Y., Rath, N. P. & Robinson, P. D. (2004). *Acta Cryst. E* **60**, o544–o546.
- Schneider, C. (2009). *Angew. Chem. Int. Ed.* **48**, 2082–2084.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst. C* **71**, 3–8.
- Stamm, H. (1999). *J. Prakt. Chem.* **341**, 319–331.

supporting information

Acta Cryst. (2015). E71, o438–o439 [doi:10.1107/S2056989015010221]

Crystal structure of 1-[(2,4,6-triisopropylphenyl)sulfonyl]aziridine

Lena Knauer, Christopher Golz and Carsten Strohmann

S1. Structural commentary


Aziridines are interesting and versatile building blocks in synthetic chemistry due to their high ring strain. By regioselective ring opening reactions, countless nitrogen-containing compounds are accessible (Stamm, 1999; Schneider, 2009). For example, the aziridine ring can be opened by triethylamine (Golz & Strohmann, 2015). In some cases, the three-membered aziridine ring is further activated by electron-withdrawing groups (Hu, 2004) to increase its reactivity.

The title compound, $C_{17}H_{27}NO_2S$, is a representative of the class of activated aziridines, as it contains a triisopropylbenzene substituted sulfonyl ester attached to the nitrogen atom. In the aromatic ring, the bulky substituents lead to a distortion of its geometry. This is expressed by the increased bond lengths and out-of-plane bent substituents around the benzene ring. At the *ipso*-carbon, the bonds C10–C11 and C10–C6 are slightly elongated to 1.410 (3) Å. In contrast, the other bonds of the aromatic ring exhibit usual lengths [C4–C15 1.374 (3) Å, C5–C6 1.380 (3) Å, C11–C15 1.388 (3) Å, C4–C5 1.389 (3) Å]. The sulfonyl group as well as the adjacent isopropyl groups bend out of the aromatic plane. This is caused by steric repulsion between the isopropyl groups and the sulfonyl oxygen atoms. The sulfur atom deviates from the mean aromatic ring plane by 0.393 (4) Å. The carbon atoms C7 and C12 show distances of -0.125 (4) Å and -0.154 (4) Å, respectively (see Table 5). A similar distortion can also be observed at the isopropyl group in *para* position to the sulfonyl moiety. Here, C1 has a distance to the aromatic ring plane of 0.111 (5) Å, thus being distorted in the same direction as the sulfur atom. This is caused by steric repulsion between the C1 isopropyl group and the adjacent isopropyl groups in *ortho* position in respect of the sulfonyl moiety.

S2. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 1.

Hydrogen atoms were located from difference Fourier maps, refined at idealized positions riding on the carbon atoms with isotropic displacement parameters $U_{iso}(H) = 1.2U_{eq}(C)$ or $1.5U_{eq}(-CH_3)$ and aromatic C–H = 0.95 Å, primary C–H = 0.98 Å, secondary C–H = 0.99 Å, tertiary C–H = 1.00 Å. All CH_3 hydrogen atoms were allowed to rotate but not to tip. Aziridine protons could be located from difference Fourier maps, but were refined as idealized CH_2 groups.

**Figure 1**

Molecular structure of the title compound with anisotropic displacement ellipsoids drawn at 50% probability level.

(I)

Crystal data

$C_{17}H_{27}NO_2S$

$M_r = 309.45$

Monoclinic, $P2_1/c$

$a = 6.2679 (8) \text{ \AA}$

$b = 17.5289 (18) \text{ \AA}$

$c = 16.3890 (13) \text{ \AA}$

$\beta = 100.331 (10)^\circ$

$V = 1771.5 (3) \text{ \AA}^3$

$Z = 4$

$F(000) = 672$

$D_x = 1.160 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 2277 reflections

$\theta = 3.3\text{--}29.2^\circ$

$\mu = 0.19 \text{ mm}^{-1}$

$T = 173 \text{ K}$

Plate, colourless

$0.33 \times 0.25 \times 0.01 \text{ mm}$

Data collection

Agilent Xcalibur Sapphire3
diffractometer

Radiation source: Enhance (Mo) X-ray Source

Graphite monochromator

Detector resolution: 16.0560 pixels mm^{-1}

ω scans

Absorption correction: multi-scan
(*CrysAlis PRO*; Agilent, 2013)

$T_{\min} = 0.782$, $T_{\max} = 1.000$

9521 measured reflections

3449 independent reflections

2479 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.049$

$\theta_{\max} = 26.0^\circ$, $\theta_{\min} = 2.5^\circ$

$h = -7 \rightarrow 7$

$k = -15 \rightarrow 21$

$l = -20 \rightarrow 20$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.141$
 $S = 1.07$
 3449 reflections
 196 parameters
 0 restraints

Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0473P)^2 + 0.8936P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.33 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.58 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. Absorption correction: CrysAlisPro, Agilent Technologies, Version 1.171.36.28 (release 01-02-2013 CrysAlis171 .NET) (compiled Feb 1 2013,16:14:44) Empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.71473 (10)	0.20619 (4)	0.37952 (4)	0.0245 (2)
C10	0.7527 (4)	0.26810 (13)	0.29748 (14)	0.0205 (5)
C15	0.8227 (4)	0.28863 (14)	0.16069 (15)	0.0272 (6)
H15	0.8229	0.2705	0.1061	0.033*
C6	0.8105 (4)	0.34434 (14)	0.31802 (14)	0.0246 (6)
C5	0.8842 (4)	0.38836 (15)	0.25903 (15)	0.0297 (6)
H5	0.9273	0.4394	0.2726	0.036*
C11	0.7462 (4)	0.24050 (14)	0.21624 (15)	0.0243 (6)
C4	0.8981 (4)	0.36110 (15)	0.18058 (15)	0.0279 (6)
O1	0.7219 (3)	0.12858 (10)	0.35547 (11)	0.0370 (5)
O2	0.8625 (3)	0.22687 (11)	0.45315 (10)	0.0360 (5)
N1	0.4729 (4)	0.23217 (13)	0.39562 (14)	0.0345 (6)
C17	0.3491 (5)	0.17365 (19)	0.4317 (2)	0.0500 (9)
H17A	0.4185	0.1234	0.4451	0.060*
H17B	0.2536	0.1908	0.4702	0.060*
C16	0.2886 (6)	0.1945 (3)	0.3447 (2)	0.0701 (12)
H16A	0.1543	0.2246	0.3280	0.084*
H16B	0.3190	0.1572	0.3029	0.084*
C12	0.6577 (5)	0.16388 (14)	0.18281 (15)	0.0312 (6)
H12	0.5872	0.1391	0.2261	0.037*
C7	0.7975 (4)	0.38301 (14)	0.40041 (15)	0.0290 (6)
H7	0.7291	0.3463	0.4347	0.035*
C1	0.9900 (5)	0.41027 (15)	0.11913 (15)	0.0331 (7)
H1	0.9766	0.3811	0.0660	0.040*
C13	0.4860 (6)	0.17400 (18)	0.10515 (18)	0.0489 (9)
H13A	0.3751	0.2099	0.1166	0.073*
H13B	0.4184	0.1246	0.0887	0.073*

H13C	0.5539	0.1940	0.0602	0.073*
C3	0.8614 (5)	0.48402 (17)	0.10107 (18)	0.0457 (8)
H3A	0.7095	0.4719	0.0787	0.068*
H3B	0.9223	0.5142	0.0605	0.068*
H3C	0.8695	0.5133	0.1525	0.068*
C14	0.8399 (5)	0.11176 (17)	0.1661 (2)	0.0471 (8)
H14A	0.9167	0.1361	0.1259	0.071*
H14B	0.7787	0.0631	0.1436	0.071*
H14C	0.9415	0.1026	0.2180	0.071*
C2	1.2285 (5)	0.42692 (19)	0.1493 (2)	0.0496 (9)
H2A	1.2459	0.4567	0.2006	0.074*
H2B	1.2856	0.4560	0.1068	0.074*
H2C	1.3083	0.3788	0.1597	0.074*
C8	1.0244 (5)	0.40142 (17)	0.44750 (16)	0.0407 (8)
H8A	1.1099	0.3544	0.4564	0.061*
H8B	1.0137	0.4243	0.5012	0.061*
H8C	1.0951	0.4374	0.4151	0.061*
C9	0.6542 (6)	0.45350 (17)	0.38647 (18)	0.0476 (8)
H9A	0.7228	0.4919	0.3562	0.071*
H9B	0.6351	0.4744	0.4401	0.071*
H9C	0.5125	0.4396	0.3541	0.071*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0289 (4)	0.0256 (4)	0.0205 (3)	0.0015 (3)	0.0085 (3)	0.0032 (3)
C10	0.0213 (13)	0.0238 (13)	0.0168 (11)	0.0008 (10)	0.0048 (10)	0.0025 (10)
C15	0.0372 (16)	0.0275 (14)	0.0178 (12)	-0.0032 (12)	0.0077 (11)	-0.0036 (10)
C6	0.0284 (14)	0.0279 (14)	0.0174 (12)	-0.0012 (11)	0.0039 (11)	-0.0011 (10)
C5	0.0401 (17)	0.0269 (14)	0.0228 (13)	-0.0082 (12)	0.0077 (12)	-0.0035 (11)
C11	0.0283 (14)	0.0233 (13)	0.0212 (12)	0.0003 (11)	0.0043 (11)	-0.0005 (10)
C4	0.0358 (16)	0.0297 (15)	0.0190 (12)	-0.0034 (12)	0.0068 (11)	0.0003 (11)
O1	0.0590 (14)	0.0250 (10)	0.0292 (10)	0.0063 (9)	0.0137 (10)	0.0051 (8)
O2	0.0431 (13)	0.0460 (12)	0.0184 (9)	-0.0037 (9)	0.0036 (9)	0.0068 (8)
N1	0.0286 (13)	0.0448 (14)	0.0326 (12)	-0.0001 (11)	0.0128 (11)	0.0007 (11)
C17	0.0423 (19)	0.053 (2)	0.063 (2)	-0.0197 (15)	0.0294 (17)	-0.0063 (17)
C16	0.0294 (19)	0.123 (4)	0.059 (2)	-0.014 (2)	0.0122 (17)	-0.030 (2)
C12	0.0473 (18)	0.0251 (14)	0.0221 (13)	-0.0042 (12)	0.0088 (12)	-0.0015 (11)
C7	0.0442 (17)	0.0251 (14)	0.0197 (12)	-0.0084 (12)	0.0115 (12)	-0.0048 (11)
C1	0.0504 (18)	0.0296 (15)	0.0224 (13)	-0.0063 (13)	0.0152 (13)	-0.0008 (11)
C13	0.060 (2)	0.0442 (19)	0.0360 (16)	-0.0173 (16)	-0.0084 (16)	-0.0013 (14)
C3	0.060 (2)	0.0423 (19)	0.0371 (16)	-0.0001 (16)	0.0160 (16)	0.0155 (14)
C14	0.059 (2)	0.0336 (17)	0.0517 (19)	-0.0011 (15)	0.0181 (17)	-0.0145 (15)
C2	0.047 (2)	0.055 (2)	0.0497 (19)	-0.0071 (16)	0.0182 (16)	0.0124 (16)
C8	0.056 (2)	0.0432 (18)	0.0215 (13)	-0.0145 (15)	0.0028 (14)	-0.0060 (13)
C9	0.070 (2)	0.0402 (18)	0.0353 (16)	0.0090 (16)	0.0172 (16)	-0.0109 (14)

Geometric parameters (\AA , $\text{^{\circ}}$)

S1—C10	1.777 (2)	C7—H7	1.0000
S1—O1	1.4193 (19)	C7—C8	1.526 (4)
S1—O2	1.4300 (19)	C7—C9	1.520 (4)
S1—N1	1.649 (2)	C1—H1	1.0000
C10—C6	1.410 (3)	C1—C3	1.524 (4)
C10—C11	1.410 (3)	C1—C2	1.517 (4)
C15—H15	0.9500	C13—H13A	0.9800
C15—C11	1.388 (3)	C13—H13B	0.9800
C15—C4	1.374 (4)	C13—H13C	0.9800
C6—C5	1.380 (3)	C3—H3A	0.9800
C6—C7	1.526 (3)	C3—H3B	0.9800
C5—H5	0.9500	C3—H3C	0.9800
C5—C4	1.389 (3)	C14—H14A	0.9800
C11—C12	1.518 (4)	C14—H14B	0.9800
C4—C1	1.516 (3)	C14—H14C	0.9800
N1—C17	1.473 (3)	C2—H2A	0.9800
N1—C16	1.456 (4)	C2—H2B	0.9800
C17—H17A	0.9900	C2—H2C	0.9800
C17—H17B	0.9900	C8—H8A	0.9800
C17—C16	1.455 (5)	C8—H8B	0.9800
C16—H16A	0.9900	C8—H8C	0.9800
C16—H16B	0.9900	C9—H9A	0.9800
C12—H12	1.0000	C9—H9B	0.9800
C12—C13	1.523 (4)	C9—H9C	0.9800
C12—C14	1.526 (4)		
O1—S1—C10	111.10 (11)	C8—C7—H7	107.9
O1—S1—O2	115.43 (12)	C9—C7—C6	110.6 (2)
O1—S1—N1	112.65 (12)	C9—C7—H7	107.9
O2—S1—C10	109.23 (11)	C9—C7—C8	112.1 (2)
O2—S1—N1	105.67 (12)	C4—C1—H1	107.8
N1—S1—C10	101.77 (11)	C4—C1—C3	111.1 (2)
C6—C10—S1	117.51 (17)	C4—C1—C2	111.3 (2)
C6—C10—C11	120.9 (2)	C3—C1—H1	107.8
C11—C10—S1	121.32 (19)	C2—C1—H1	107.8
C11—C15—H15	118.3	C2—C1—C3	110.9 (2)
C4—C15—H15	118.3	C12—C13—H13A	109.5
C4—C15—C11	123.4 (2)	C12—C13—H13B	109.5
C10—C6—C7	125.5 (2)	C12—C13—H13C	109.5
C5—C6—C10	117.8 (2)	H13A—C13—H13B	109.5
C5—C6—C7	116.7 (2)	H13A—C13—H13C	109.5
C6—C5—H5	118.6	H13B—C13—H13C	109.5
C6—C5—C4	122.7 (2)	C1—C3—H3A	109.5
C4—C5—H5	118.6	C1—C3—H3B	109.5
C10—C11—C12	126.2 (2)	C1—C3—H3C	109.5
C15—C11—C10	117.1 (2)	H3A—C3—H3B	109.5

C15—C11—C12	116.6 (2)	H3A—C3—H3C	109.5
C15—C4—C5	117.5 (2)	H3B—C3—H3C	109.5
C15—C4—C1	121.6 (2)	C12—C14—H14A	109.5
C5—C4—C1	120.9 (2)	C12—C14—H14B	109.5
C17—N1—S1	116.0 (2)	C12—C14—H14C	109.5
C16—N1—S1	116.1 (2)	H14A—C14—H14B	109.5
C16—N1—C17	59.5 (2)	H14A—C14—H14C	109.5
N1—C17—H17A	117.8	H14B—C14—H14C	109.5
N1—C17—H17B	117.8	C1—C2—H2A	109.5
H17A—C17—H17B	114.9	C1—C2—H2B	109.5
C16—C17—N1	59.7 (2)	C1—C2—H2C	109.5
C16—C17—H17A	117.8	H2A—C2—H2B	109.5
C16—C17—H17B	117.8	H2A—C2—H2C	109.5
N1—C16—H16A	117.7	H2B—C2—H2C	109.5
N1—C16—H16B	117.7	C7—C8—H8A	109.5
C17—C16—N1	60.8 (2)	C7—C8—H8B	109.5
C17—C16—H16A	117.7	C7—C8—H8C	109.5
C17—C16—H16B	117.7	H8A—C8—H8B	109.5
H16A—C16—H16B	114.8	H8A—C8—H8C	109.5
C11—C12—H12	107.9	H8B—C8—H8C	109.5
C11—C12—C13	110.9 (2)	C7—C9—H9A	109.5
C11—C12—C14	111.0 (2)	C7—C9—H9B	109.5
C13—C12—H12	107.9	C7—C9—H9C	109.5
C13—C12—C14	111.1 (2)	H9A—C9—H9B	109.5
C14—C12—H12	107.9	H9A—C9—H9C	109.5
C6—C7—H7	107.9	H9B—C9—H9C	109.5
C8—C7—C6	110.4 (2)		
S1—C10—C6—C5	166.7 (2)	C5—C6—C7—C8	−67.6 (3)
S1—C10—C6—C7	−13.6 (3)	C5—C6—C7—C9	57.1 (3)
S1—C10—C11—C15	−166.5 (2)	C5—C4—C1—C3	−58.7 (4)
S1—C10—C11—C12	15.3 (4)	C5—C4—C1—C2	65.4 (3)
S1—N1—C17—C16	−106.4 (3)	C11—C10—C6—C5	−7.3 (4)
S1—N1—C16—C17	106.1 (2)	C11—C10—C6—C7	172.4 (3)
C10—S1—N1—C17	154.1 (2)	C11—C15—C4—C5	−3.9 (4)
C10—S1—N1—C16	87.0 (2)	C11—C15—C4—C1	177.0 (3)
C10—C6—C5—C4	1.6 (4)	C4—C15—C11—C10	−1.5 (4)
C10—C6—C7—C8	112.8 (3)	C4—C15—C11—C12	176.8 (2)
C10—C6—C7—C9	−122.6 (3)	O1—S1—C10—C6	−162.51 (19)
C10—C11—C12—C13	125.2 (3)	O1—S1—C10—C11	11.4 (3)
C10—C11—C12—C14	−110.8 (3)	O1—S1—N1—C17	35.1 (2)
C15—C11—C12—C13	−52.9 (3)	O1—S1—N1—C16	−32.0 (3)
C15—C11—C12—C14	71.0 (3)	O2—S1—C10—C6	−34.0 (2)
C15—C4—C1—C3	120.4 (3)	O2—S1—C10—C11	139.9 (2)
C15—C4—C1—C2	−115.5 (3)	O2—S1—N1—C17	−91.8 (2)
C6—C10—C11—C15	7.2 (4)	O2—S1—N1—C16	−158.9 (2)
C6—C10—C11—C12	−170.9 (2)	N1—S1—C10—C6	77.4 (2)
C6—C5—C4—C15	3.8 (4)	N1—S1—C10—C11	−108.7 (2)

supporting information

C6—C5—C4—C1

−177.1 (3)

C7—C6—C5—C4

−178.1 (3)
