

Improve survival from prolonged mechanical ventilation: beginning with first step

Chun Pan, Haibo Qiu

Department of Critical Care Medicine, Zhongda Hospital, School of Medicine, Southeast University, Nanjing 210009, China

Correspondence to: Haibo Qiu, MD, PhD. Department of Critical Care Medicine, Zhongda Hospital, School of Medicine, Southeast University, Nanjing 210009, China. Email: haiboq2000@163.com.

Submitted Jun 15, 2015. Accepted for publication Jul 06, 2015.

doi: 10.3978/j.issn.2072-1439.2015.07.13

View this article at: <http://dx.doi.org/10.3978/j.issn.2072-1439.2015.07.13>

Although conventional invasive mechanical ventilation is a life-saving procedure for critically ill patients, it is the most important risk factor for ventilator-associated events (VAEs) (1-3). Patients with VAEs often need prolonged mechanical ventilation (PMV), which leads to increased family and financial burden (1) and worse prognosis (2). However, long-term outcomes associated with PMV are not fully understood (3). A recent meta-analysis reporting long-term survival of critically ill patients treated with PMV further highlights the worse prognosis posed by the patients (3). Pooled mortality associated with PMV at hospital discharge, 1 year, and beyond 1 year was 29% (95% CI: 26-32), 62% (95% CI: 57-67), and 66% (95% CI: 61-70), respectively. Moreover, only 19% [16-24] were discharged to home and only 50% [47-53] were successfully liberated from mechanical ventilation (MV). These results alarm the physicians that PMV increases not intensive care unit (ICU) and hospital mortality, but long-term mortality. Considering the worse short-term and long-term outcomes of PMV, timely and effective weaning from PMV is an important priority in primary care for critically ill patients. However, difficult or prolonged weaning is common and effective management weaning remains unclear. If physicians and intensivists focus on the problems, a first step to improve the outcomes of patients with PMV is reducing the incidence of PMV.

Predictors of difficult or prolonged weaning

Peñuelas screened the weaning patients, and clarified simple, difficult and prolonged weaning (2). About 69% patients can be extubated at the first weaning attempt (simple

weaning), while the remaining 31% need more than one spontaneous breathing trial (SBT) (difficult plus prolonged weaning) (4). Difficult weaning includes patients who required up to 7 days being extubated from the first attempt of withdrawal from mechanical ventilation. Prolonged weaning refers to failure at least three weaning attempts or required >7 days of weaning after the first SBT. Predictors for difficult or prolonged weaning include high simplified acute physiology score II score, duration of MV before simple weaning, chronic pulmonary disease, pneumonia, high PEEP before simple weaning (2). However, airway and lung dysfunction, brain dysfunction, cardiac dysfunction, diaphragm/respiratory muscle function and endocrine and metabolic dysfunction also could induced difficult or prolonged weaning.

Reducing incidence of difficult or prolonged weaning

Focusing on the primary disease is the first line treatment for reducing the incidence of prolonged weaning. Besides the primary diseases control, there are some the do's and don'ts needing attention.

Screen the reasons of prolonged weaning

After intubating the patients, the first purpose of the treatment is weaning from the ventilator. Heunks classified the reasons of a difficult to wean patients: airway or lung dysfunction, brain dysfunction, cardiac dysfunction, diaphragm dysfunction, and endocrine dysfunction (5). Critically ill patients in ICU may have continuing organ

failures leading to protracted periods of organ support (6). Therefore, it is a key step of weaning that primary disease and the complications are controlled.

Optimal mechanical ventilation settings

Improper settings could induce lung injury, especially for acute respiratory distress syndrome (ARDS) patients. Although hospital mortality of ARDS patients is approximately 40%, 1-year mortality decreases to 14% and 3-year mortality is only 15% (7,8). Even the patients treated with extracorporeal membrane oxygenation (ECMO), the 261-day survival is 86% (9), and the survivors had minor lung disabilities with diminished diffusion capacities across the blood-gas barrier (10). If acute lung injury is controlled, it may not influence long-term outcomes of these patients. Therefore, it is essential to establish optimal ventilation strategy to protect lung and improve hospital mortality. The ventilation strategy include: low tidal volume (6 mL/kg predict body weight), optimal PEEP level, plateau pressure <30 cm H₂O, and prone position in ARDS patients with PO₂/FiO₂ <150 mmHg (11). Moreover, Amato in the retrospective study showed that mortality of ARDS patients was associated with driving pressure (12), but the threshold value of the pressure needs to be further clarified.

Early transfer to non-invasive ventilation (NIV)

NIV has been used in weaning protocol for chronic obstructive pulmonary disease (COPD) patients. It could reduce intubation rates and improve ICU mortality (13-15). NIV also could be used in weaning of acute respiratory failure patients, and NIV after early extubation could decrease invasive-ventilation-days and reintubation rates within 48 hours (16,17). However, further studies are needed to explain the characteristics of the patients and the protocol.

Avoiding ICU-acquired weakness

ICU-acquired weakness is an important reason which directly leads to prolonged weaning. The pathophysiological mechanisms of ICU-acquired weakness are multifactorial. Both of critical illness polyneuropathy and critical illness myopathy could induce ICU-acquired weakness. Moreover, sepsis, persistent systemic inflammation, multiorgan system failure, long duration of mechanical ventilation, immobility, hyperglycemia, and the use of glucocorticoids or neuromuscular blocking agents are important risk factors.

Ventilator induced diaphragm dysfunction (VIDD) is the important clinical manifestation of ICU-acquired weakness, and it could induce prolonged weaning. There are many risks contributing to VIDD, e.g., sepsis, mechanical ventilation, corticosteroids using and inflammatory factors (18). Some treatments should be administrated for the patients with mechanical ventilation. Firstly, system inflammatory response syndrome (SIRS) and sepsis should be controlled. Secondly, measures how much benefits the patients can get from corticosteroids, and some antioxidant supplementation (alpha-tocopherol and ascorbic acid) may be benefit for VIDD (19). Thirdly, keeping spontaneous breath and using assisted ventilation mode could improve diaphragm movement, neurally adjusted ventilatory assist ventilation should be the better mode for the difficult and prolonged patients weaning (20,21). Fourthly, sufficient nutrition could help the patients recovery from VIDD (22).

Early progressive mobilization

The strategy for whole-body rehabilitation could improve mechanical patients' functional outcomes and ventilator-free days, decrease incidence of delirium, length of ICU and hospital days (23). Many studies have proved early mobilization is safe and effect for the critically ill patients (24,25), but when and how to start early mobilization protocol is still debated. However, an expert consensus and recommendations could be used for safety criteria. Mobilizing patients in ICU may be considered according to a traffic-light system of low risk of an adverse event (green), potential risk of an adverse event is outweighed by the benefit of early mobilization (yellow) and significant potential risk of an adverse event requiring consultation with senior ICU staff (red) (26), and awakening and breathing coordination, delirium monitoring/management, early exercise/mobility (ABCDE) bundle could help to the treatment of implementation (25).

Light sedation and delirium controlled

Light sedation is associated with improved clinical outcomes (e.g., shorter duration of mechanical ventilation and length of ICU stay). Therefore, the guidelines for the management of pain, agitation, and delirium recommend maintaining light levels of sedation in adult ICU patients (27). Delirium is associated with prolonged length of ICU and hospital stay, increased mortality, and long-term cognitive impairment in ICU patients (28). Early mobilization of the

patients and avoiding benzodiazepine usage could prevent delirium (27).

Improve endocrine and metabolic dysfunctions

Difficult weaning induced by endocrine and metabolic dysfunctions is usually ignored. Adrenal insufficiency, hypothyroidism, and electrolyte disturbances (e.g., very low plasma levels of phosphate and magnesium) are associated with difficult and prolonged weaning (5). These diseases should be taken into consideration when treating difficult weaning patients.

In conclusion, physicians and intensivists should pay more attention to critically ill patients receiving PMV. How to improve long-term outcomes of these patients needs the further study. If we could reduce the risks of PMV at the first step, these patients will start on the journey towards better outcomes.

Acknowledgements

Funding: This work was sponsored by grants (81300043 and 81370180) from the National Natural Science Foundation of China, Jiangsu Clinical Science and Technology Projects (BL2013030) and grants (BK20131302) from Jiangsu Natural Science Foundation.

Footnote

Provenance: This is a Guest Editorial commissioned by the section editor Wan-Jie Gu (Department of Anesthesiology, Affiliated Drum Tower Hospital, Medical College of Nanjing University, Nanjing, China).

Conflicts of Interest: The authors have no conflicts of interest to declare.

References

1. Kahn JM, Le T, Angus DC, et al. The epidemiology of chronic critical illness in the United States*. *Crit Care Med* 2015;43:282-7.
2. Peñuelas O, Frutos-Vivar F, Fernández C, et al. Characteristics and outcomes of ventilated patients according to time to liberation from mechanical ventilation. *Am J Respir Crit Care Med* 2011;184:430-7.
3. Damuth E, Mitchell JA, Bartock JL, et al. Long-term survival of critically ill patients treated with prolonged mechanical ventilation: a systematic review and meta-analysis. *Lancet Respir Med* 2015;3:544-53.
4. Sellares J, Ferrer M, Cano E, et al. Predictors of prolonged weaning and survival during ventilator weaning in a respiratory ICU. *Intensive Care Med* 2011;37:775-84.
5. Heunks LM, van der Hoeven JG. Clinical review: the ABC of weaning failure--a structured approach. *Crit Care* 2010;14:245.
6. Nelson JE, Cox CE, Hope AA, et al. Chronic critical illness. *Am J Respir Crit Care Med* 2010;182:446-54.
7. Herridge MS, Cheung AM, Tansey CM, et al. One-year outcomes in survivors of the acute respiratory distress syndrome. *N Engl J Med* 2003;348:683-93.
8. Khandelwal N, Hough CL, Bansal A, et al. Long-term survival in patients with severe acute respiratory distress syndrome and rescue therapies for refractory hypoxemia*. *Crit Care Med* 2014;42:1610-8.
9. Hodgson CL, Hayes K, Everard T, et al. Long-term quality of life in patients with acute respiratory distress syndrome requiring extracorporeal membrane oxygenation for refractory hypoxaemia. *Crit Care* 2012;16:R202.
10. Luyt CE, Combes A, Becquemin MH, et al. Long-term outcomes of pandemic 2009 influenza A(H1N1)-associated severe ARDS. *Chest* 2012;142:583-92.
11. Esteban A, Frutos-Vivar F, Muriel A, et al. Evolution of mortality over time in patients receiving mechanical ventilation. *Am J Respir Crit Care Med* 2013;188:220-30.
12. Amato MB, Meade MO, Slutsky AS, et al. Driving pressure and survival in the acute respiratory distress syndrome. *N Engl J Med* 2015;372:747-55.
13. Ferrer M, Valencia M, Nicolas JM, et al. Early noninvasive ventilation averts extubation failure in patients at risk: a randomized trial. *Am J Respir Crit Care Med* 2006;173:164-70.
14. Nava S, Gregoretti C, Fanfulla F, et al. Noninvasive ventilation to prevent respiratory failure after extubation in high-risk patients. *Crit Care Med* 2005;33:2465-70.
15. Burns KE, Meade MO, Premji A, et al. Noninvasive positive-pressure ventilation as a weaning strategy for intubated adults with respiratory failure. *Cochrane Database Syst Rev* 2013;12:CD004127.
16. Vaschetto R, Turucz E, Dellapiazza F, et al. Noninvasive ventilation after early extubation in patients recovering from hypoxemic acute respiratory failure: a single-centre feasibility study. *Intensive Care Med* 2012;38:1599-606.
17. Ornicco SR, Lobo SM, Sanches HS, et al. Noninvasive ventilation immediately after extubation improves weaning outcome after acute respiratory failure: a randomized controlled trial. *Crit Care* 2013;17:R39.

18. Picard M, Jung B, Liang F, et al. Mitochondrial dysfunction and lipid accumulation in the human diaphragm during mechanical ventilation. *Am J Respir Crit Care Med* 2012;186:1140-9.
19. Nathens AB, Neff MJ, Jurkovich GJ, et al. Randomized, prospective trial of antioxidant supplementation in critically ill surgical patients. *Ann Surg* 2002;236:814-22.
20. Huang DY, Liu J, Wu XY, et al. Effects of neurally adjusted ventilatory assist on prevention of ventilator-induced diaphragmatic dysfunction in acute respiratory distress syndrome rabbits. *Zhonghua Jie He He Hu Xi Za Zhi* 2011;34:288-93.
21. Liu L, Liu H, Yang Y, et al. Neuroventilatory efficiency and extubation readiness in critically ill patients. *Crit Care* 2012;16:R143.
22. Rahman A, Wilund K, Fitschen PJ, et al. Elderly persons with ICU-acquired weakness: the potential role for β -hydroxy- β -methylbutyrate (HMB) supplementation? *JPEN J Parenter Enteral Nutr* 2014;38:567-75.
23. Stiller K. Physiotherapy in intensive care: an updated systematic review. *Chest* 2013;144:825-47.
24. Schweickert WD, Pohlman MC, Pohlman AS, et al. Early physical and occupational therapy in mechanically ventilated, critically ill patients: a randomised controlled trial. *Lancet* 2009;373:1874-82.
25. Balas MC, Vasilevskis EE, Olsen KM, et al. Effectiveness and safety of the awakening and breathing coordination, delirium monitoring/management, and early exercise/mobility bundle. *Crit Care Med* 2014;42:1024-36.
26. Hodgson CL, Stiller K, Needham DM, et al. Expert consensus and recommendations on safety criteria for active mobilization of mechanically ventilated critically ill adults. *Crit Care* 2014;18:658.
27. Barr J, Fraser GL, Puntillo K, et al. Clinical practice guidelines for the management of pain, agitation, and delirium in adult patients in the intensive care unit. *Crit Care Med* 2013;41:263-306.
28. Salluh JI, Wang H, Schneider EB, et al. Outcome of delirium in critically ill patients: systematic review and meta-analysis. *BMJ* 2015;350:h2538.

Cite this article as: Pan C, Qiu H. Improve survival from prolonged mechanical ventilation: beginning with first step. *J Thorac Dis* 2015;7(7):1076-1079. doi: 10.3978/j.issn.2072-1439.2015.07.13