

RESEARCH ARTICLE

Tumor-Targeting *Salmonella typhimurium* A1-R Arrests a Chemo-Resistant Patient Soft-Tissue Sarcoma in Nude Mice

Yukihiko Hiroshima^{1,2,3}, Ming Zhao¹, Yong Zhang¹, Nan Zhang¹, Ali Maawy², Takashi Murakami^{1,2,3}, Sumiyuki Mii^{1,2}, Fuminari Uehara^{1,2}, Mako Yamamoto^{1,2}, Shinji Miwa^{1,2}, Shuya Yano^{1,2}, Masashi Momiyama³, Ryutaro Mori³, Ryusei Matsuyama³, Takashi Chishima³, Kuniya Tanaka³, Yasushi Ichikawa³, Michael Bouvet², Itaru Endo³, Robert M. Hoffman^{1,2*}

1 AntiCancer, Inc., San Diego, California, United States of America, **2** Department of Surgery, University of California San Diego, San Diego, California, United States of America, **3** Department of Gastroenterological Surgery, Yokohama City University Graduate School of Medicine, Yokohama, Japan

* all@anticancer.com



OPEN ACCESS

Citation: Hiroshima Y, Zhao M, Zhang Y, Zhang N, Maawy A, Murakami T, et al. (2015) Tumor-Targeting *Salmonella typhimurium* A1-R Arrests a Chemo-Resistant Patient Soft-Tissue Sarcoma in Nude Mice. PLoS ONE 10(8): e0134324. doi:10.1371/journal.pone.0134324

Editor: Pankaj K Singh, University of Nebraska Medical Center, UNITED STATES

Received: October 1, 2014

Accepted: July 8, 2015

Published: August 3, 2015

Copyright: © 2015 Hiroshima et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All data underlying the findings in our study are freely available in the manuscript.

Funding: This study was supported in part by National Cancer Institute grant numbers CA183280 and JSPS KAKENHI grant numbers 26830081 to YH, 26462070 to IE and 24592009 to KT. AntiCancer, Inc., provided support in the form of salaries for authors [MJ, YZ, NZ]. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Abstract

A patient-derived nude-mouse model of soft-tissue sarcoma has been established and treated in the following groups: (1) untreated controls; (2) gemcitabine (GEM) (80 mg/kg, ip, weekly, 3 weeks); (3) Pazopanib (100 mg/kg, orally, daily, 3 weeks) and (4) *Salmonella typhimurium* A1-R (5×10^7 CFU/body, ip, weekly, 3 weeks). The sarcoma was resistant to GEM ($p = 0.879$). Pazopanib tended to reduce the tumor volume compared to the untreated mice, but there was no significant difference ($p = 0.115$). *S. typhimurium* A1-R significantly inhibited tumor growth compared to the untreated mice ($p = 0.001$). *S. typhimurium* A1-R was the only effective treatment for the soft-tissue sarcoma nude mouse model among all treatments including a newly approved multiple tyrosine kinase inhibitor; Pazopanib. These results suggest tumor-targeting *S. typhimurium* A1-R is a promising treatment for chemo-resistant soft-tissue sarcoma.

Introduction

Soft-tissue sarcomas are rare mesenchymal cancers comprising approximately 50 histological types [1]. The yearly incidence of soft-tissue sarcomas in the USA is 11,280 cases, with 3,900 deaths. Patients with metastatic soft-tissue sarcomas have a median overall survival of about 12 months.

Pazopanib (Votrient) is a multiple tyrosine kinase inhibitor approved for the treatment of advanced (unresectable and/or metastatic) soft-tissue sarcomas in patients who have received prior chemotherapy. Pazopanib improved progression-free survival in a Phase III clinical trial [2].

An attenuated strain of *Clostridium novyi* (*C. novyi*-NT) was used to treat a human patient who had an advanced leiomyosarcoma. Treatment was intratumor (i.t.) injection of *C. novyi*-NT spores. *C. novyi*-NT regressed the tumor within and surrounding the bone [3].

Competing Interests: Ming Zhao, Yong Zhang and Nan Zhang are employees of AntiCancer Inc. Yukihiko Hiroshima, Takashi Murakami, Mako Yamamoto, Shinji Miwa, Shuya Yano, and Robert M. Hoffman are unpaid affiliates of AntiCancer Inc. Sumiyuki Mii, Fuminari Uehara, Masashi Momiyama and Takashi Chishima were former affiliates of AntiCancer Inc. AntiCancer Inc. markets animal models of cancer. There are no other competing interests. There are no patents, products in development or marketed products to declare. This does not alter the authors' adherence to all the PLOS ONE policies on sharing data and materials, as detailed online in the guide for authors.

Our laboratory has previously developed a genetically-modified bacteria strain, *Salmonella typhimurium* A1-R, selected for tumor-targeting in vivo. *S. typhimurium* A1-R is auxotrophic for leu and arg [4]. The strain targets and grows in tumors. In contrast, normal tissue is cleared of these bacteria even in immunodeficient athymic mice. *S. typhimurium* A1-R is effective against prostate cancer [5], breast cancer [6, 7], pancreatic cancer [8–11], glioma [12, 13], lung cancer [14], fibrosarcoma [15] and osteosarcoma [16].

In the present study, we report the efficacy of *S. typhimurium* A1-R against a human patient chemo-resistant sarcoma growing in nude mice. Tumors growing in nude mice can faithfully replicate important features of the patient's cancer, such as tumor markers even after passage [17, 18].

Materials and Methods

Animals

Male athymic *nu/nu* nude mice (AntiCancer Inc., San Diego, CA), 4–6 weeks old, were used in this study. Mice were kept in a barrier facility under HEPA filtration. Mice were fed with autoclaved laboratory rodent diet. All mouse surgical procedures and imaging were performed with the animals anesthetized by intramuscular injection of 50% ketamine, 38% xylazine, and 12% acepromazine maleate (0.02 ml solution). All animal studies were conducted with an AntiCancer Institutional Animal Care and Use Committee (IACUC)-protocol specifically approved for this study and in accordance with the principals and procedures outlined in the National Institute of Health Guide for the Care and Use of Animals under Assurance Number A3873-1.

Specimen collection

The patient provided informed written consent and samples were procured and the study was conducted under the approval of the Institutional Review Board of the UC San Diego Medical Center.

Establishment of patient nude mouse model of sarcoma

Tumor tissues were obtained from the patient with a metastatic soft-tissue sarcoma of the retroperitoneum at biopsy and cut into fragments (3-mm³) and transplanted subcutaneously in nude mice. Tumors in the present study were in their second passage.

Preparation of bacteria

S. typhimurium A1-R was grown overnight on LB medium and then diluted 1:10 in LB medium. Bacteria were harvested at late-log phase, washed with PBS, and then diluted in PBS. Bacteria were then used for experiments [6].

Treatment of soft-tissue sarcoma in nude mice

The patient sarcoma established in nude mice was passaged subcutaneously to 20 nude mice to determine the efficacy of various treatments. Four weeks after implantation, the mice in each model were randomized to 6 mice in the untreated control and 4 groups of 5 mice in each treatment group and treated as follows: (1) untreated control; (2) gemcitabine (GEM, Eli Lilly and Company, Indianapolis, IN, USA) (80 mg/kg, ip, weekly, 3 weeks); (3) Pazopanib (Selleck Chemicals, Houston, TX, USA, 100 mg/kg, orally, daily, 3 weeks) and (4) *S. typhimurium* A1-R (5×10^7 CFU/body, ip, weekly, 3 weeks). Tumor size was evaluated every 3 to 4 days by caliper measurements. The approximate volume of the mass was calculated using the formula $\frac{4}{3}\pi \cdot (d/2)^2 \cdot D/2$, where *d* is the minor tumor axis and *D* is the major tumor axis. Body weight of the

mice was measured on a balance once a week. Relative tumor volume and body weight were calculated by comparison to Day 1. The endpoint of the experiment was when tumor size in the untreated control mice became approximately 2 cm. The method of euthanasia was CO₂ inhalation.

Tissue histology

Tumor samples were removed with surrounding normal tissues at the time of resection. Fresh tissue samples were fixed in 10% formalin and embedded in paraffin before sectioning and staining. Tissue sections (5 μm) were deparaffinized in xylene and rehydrated in an ethanol series. Hematoxylin and eosin (H & E) staining was performed according to standard protocols. The sections were examined using a BH-2 microscope (Olympus, Tokyo, Japan) equipped with a INFINITY1 2.0 megapixel CMOS digital camera (Lumenera Corporation, Ottawa, Canada). All images were acquired using INFINITY ANALYZE software (Lumenera Corporation) without post-acquisition processing.

Evaluation of histopathological response to treatment

Histopathological response to treatment was defined according to Evans's grading scheme: Grade I, little (<10%) or no cancer cell destruction is evident; Grade IIa, destruction of 10%-50% of cancer cells; Grade IIb, destruction of 51%-90% of cancer cells; Grade III, few (<10%) viable-appearing cancer cells are present; Grade IV, no viable cancer cells are present [11, 19].

Confocal imaging of *S. typhimurium* A1-R-GFP in sarcoma tissue

Resected sarcoma specimens from mice treated with *S. typhimurium* A1-R were embedded with optimal cutting temperature (OCT) compound (Tissue-Tek; Sakura Finetek Europe BV, Zoeterwude, Netherlands) and preserved in liquid nitrogen. Frozen sections of 7–10 μm thickness were prepared with a CM1850 cryostat (Leica, Wetzlar, Germany). The frozen sections were directly observed with a confocal microscope (Fluoview FV1000, Olympus, Tokyo, Japan). Excitation sources were semiconductor lasers at 473 nm for GFP excitation. After confocal imaging, frozen sections were fixed in 10% formalin and H & E staining was performed.

Culture of GFP-labeled *S. typhimurium* A1-R bacteria from tumors and organs

Subcutaneous tumors and normal nude mouse organs (blood and liver) were removed at the termination of the treatment experiments. Bacteria were isolated from the tumors and organs and cultured in LB agar for 24 hours, and imaged with the OV100 small animal imaging system (Olympus, Tokyo, Japan) [20].

Statistical analysis

PASW Statistics 18.0 (SPSS, Inc) was used for all statistical analyses. The Student's t-test was used to compare continuous variables between two groups. A p value of 0.05 was considered statistically significant for all comparisons.




Fig 1. Tumor histology. **A)** Histology of the original patient sarcoma. **B)** Histology of the mouse grown sarcoma.

doi:10.1371/journal.pone.0134324.g001

Results and Discussion

Soft-tissue patient sarcoma grown in nude mice recapitulates the histology of the original tumor

The majority of the original-tumor section was comprised of sarcomatous high grade spindle cells of varying sizes, demonstrating abundant, finely granular cytoplasm and atypical, pleomorphic, round-to-elongated nuclei with irregular nuclear membranes, an open chromatin pattern and prominent nucleoli (Fig 1A). The mouse-grown tumors had histological structures similar to the original tumor (Fig 1B).

S. typhimurium A1-R is highly effective on the patient soft-tissue sarcoma in nude mice

The relative tumor volume on day 22, compared to day 1, of each group was as follows: (1) untreated control: 6.61 ± 2.37 ; (2) GEM: 6.39 ± 2.26 ; (3) Pazopanib: 3.94 ± 2.71 ; (4) *S. typhimurium* A1-R: 1.58 ± 0.37 (Fig 2). The sarcoma did not significantly respond to GEM ($p = 0.879$). Pazopanib tended to reduce the tumor volume compared to the untreated mice, but there was not a significant difference ($p = 0.115$). *S. typhimurium* A1-R significantly inhibited the sarcoma tumor growth compared to the untreated control mice (Fig 2) ($p = 0.001$). Sizes of all tumors are listed in the S1 Table. No body weight loss was found in any treatment groups.

Histological response to treatment

Histopathological response to each treatment was defined according to Evans's grading scheme. In the control (no treatment) and GEM-treated sections, the tissue sections from the tumor were occupied by viable cancer cells (Fig 3A and 3B). Approximately 40% of cancer cells were destroyed and replaced by stromal cells in the tumor sections treated with Pazopanib (Fig 3E). Tumors treated with *S. typhimurium* A1-R consisted of 2 components; one was a viable-like component and the other one was a necrotic component. The viable-like component was occupied by cancer cells (Fig 3D). In contrast, no viable cancer cell were detected in a necrotic component of the *S. typhimurium* A1-R-treated tumor (Fig 3E). Although the viable cancer




Fig 2. Drug-response of soft tissue sarcoma in nude mice. A representative image of nude mouse with the subcutaneous sarcoma in (A) the untreated mice; (B) GEM-treated; (C) Pazopanib-treated or (D) *Salmonella typhimurium* A1-R-treated. Scale bars: 10 mm. (E) Growth curves of the subcutaneous sarcoma's treated with various drugs as described above. The values are mean relative tumor volume \pm S.D. (bars) of five different tumors. ** $p < 0.01$, compared to no treatment group. (F-G) Representative cross-sections of excised subcutaneous tumors from the control and treatment groups with type of treatment indicated.

doi:10.1371/journal.pone.0134324.g002

cells were found, they did not form a tumor as can be seen from Fig 2. The untreated control and GEM were judged as grade I; Pazopanib as IIa; *S. typhimurium* A1-R of a viable-like component as grade I and the necrotic component as IV. The necrotic component was not detected in any treatment group except for *S. typhimurium* A1-R (Fig 3A–3E). Tumor heterogeneity may be a factor in the observed chemoresistance of the soft tissue sarcoma that was overcome by *S. typhimurium* A1-R.

GFP-labeled *S. typhimurium* A1-R was isolated from the tumor (Fig 4), but not from blood and only minimally from liver, indicating the tumor was effectively targeted by *S. typhimurium* A1-R.

S. typhimurium A1-R was the only effective treatment for the soft-tissue patient sarcoma growing in nude mice including GEM and a newly approved multiple tyrosine kinase inhibitor; Pazopanib. One factor in chemoresistance of solid tumors is that the majority of the cancer cells within the tumor are in a chemoresistant G_0/G_1 quiescent cell-cycle phase [21]. We have recently shown that quiescent cancer cells are sensitive to *S. typhimurium* A1-R [22]. *S. typhimurium* A1-R, unlike *C. novyi*-NT, is a facultative anaerobe and can be administered systemically such as in the present study (i.p.), whereas *C. novyi*-NT seems to require i.t. administration which makes it different to target metastasis. The results of the present study indicate tumor-targeting *S. typhimurium* A1-R is a promising treatment for soft-tissue sarcomas. In future experiments, *S. typhimurium* A1-R will be tested in a series of patient soft tissue sarcoma grown in mice as a bridge to the clinic, where we also intend to focus on sarcoma.




Fig 3. Effect of treatment on histology in nude mice. Histopathological responses to treatments were defined according to Evans's grading scheme. (A) Treatment effect of control / no treatment was judged as grade I; (B) GEM treatment as grade I; (C) Pazopanib treatment as grade IIa; (D) *S. typhimurium* A1-R of a viable-like area as grade I and (E) *S. typhimurium* A1-R of a necrotic area as grade IV. Scale bars: 200 μ m.

doi:10.1371/journal.pone.0134324.g003




Fig 4. Tumor-targeting of *S. typhimurium* A1-R. Distribution of GFP-labeled *S. typhimurium* A1-R in tumors and organs. Representative images of GFP-labeled *S. typhimurium* A1-R bacteria culture isolated from the tumor and the normal organs (blood and liver) of the mice treated with *S. typhimurium* A1-R. GFP-labeled *S. typhimurium* A1-R were clearly detected only in the tumor. Only a few GFP-labeled *S. typhimurium* A1-R were detected in the liver and no GFP-labeled *S. typhimurium* A1-R was detected in blood. Scale bars: 10 mm.

doi:10.1371/journal.pone.0134324.g004

Evaluation of biomarker expansion under various treatment regimens will be evaluated in future experiments.

Supporting Information

S1 Table. Tumor sizes of all mice on Day-22.
(DOCX)

Acknowledgments

This study was supported in part by National Cancer Institute grant numbers CA183280 and JSPS KAKENHI grant numbers 26830081 to YH, 26462070 to IE and 24592009 to KT. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Dedication

This paper is dedicated to the memory of A. R. Moossa, M.D. and to the memory of James (Barney) Berglund, Jr., who inspired us to find improved therapy of soft-tissue sarcoma.

Author Contributions

Conceived and designed the experiments: YH RMH. Performed the experiments: YH MZ YZ NZ AM. Analyzed the data: YH MZ YZ NZ AM TM SuM FU MY ShM SY MM RMo RMa TC KT YI MB IE RMH. Contributed reagents/materials/analysis tools: RMH. Wrote the paper: YH RMH.

References

1. Clark MA, Fisher C, Judson I, Thomas JM. (2005) Soft-tissue sarcomas in adults. *N Engl J Med* 353:701–711. PMID: [16107623](#)
2. van der Graaf WT, Blay JY, Chawla SP, Kim DW, Bui-Nguyen B, Casali PG, et al. (2012) Pazopanib for metastatic soft-tissue sarcoma (PALETTE): a randomised, double-blind, placebo-controlled phase 3 trial. *Lancet* 379:1879–1886. doi: [10.1016/S0140-6736\(12\)60651-5](#) PMID: [22595799](#)
3. Roberts NJ, Zhang L, Janku F, Collins A, Bai RY, Staedtke V, et al. (2014) Intratumoral injection of *Clostridium novyi*-NT spores induces antitumor responses. *Sci Transl Med* 6:249ra111. doi: [10.1126/scitranslmed.3008982](#) PMID: [25122639](#)
4. Zhao M, Yang M, Li XM, Jiang P, Baranov E, Li S, et al. (2005) Tumor-targeting bacterial therapy with amino acid auxotrophs of GFP-expressing *Salmonella typhimurium*. *Proc Natl Acad Sci USA* 102:755–760. PMID: [15644448](#)
5. Zhao M, Geller J, Ma H, Yang M, Penman S, Hoffman RM. (2007) Monotherapy with a tumor-targeting mutant of *Salmonella typhimurium* cures orthotopic metastatic mouse models of human prostate cancer. *Proc Natl Acad Sci USA* 104:10170–10174. PMID: [17548809](#)
6. Zhao M, Yang M, Ma H, Li X, Tan X, Li S, et al. (2006) Targeted therapy with a *Salmonella typhimurium* leucine-arginine auxotroph cures orthotopic human breast tumors in nude mice. *Cancer Res* 66:7647–7652. PMID: [16885365](#)
7. Zhang Y, Tome Y, Suetsugu A, Zhang L, Zhang N, Hoffman RM, et al. (2012) Determination of the optimal route of administration of *Salmonella typhimurium* A1-R to target breast cancer in nude mice. *Anticancer Res* 32:2501–2508. PMID: [22753706](#)
8. Nagakura C, Hayashi K, Zhao M, Yamauchi K, Yamamoto N, Tsuchiya H, et al. (2009) Efficacy of a genetically-modified *Salmonella typhimurium* in an orthotopic human pancreatic cancer in nude mice. *Anticancer Res* 29:1873–1878. PMID: [19528442](#)
9. Yam C, Zhao M, Hayashi K, Ma H, Kishimoto H, McElroy M, et al. (2010) Monotherapy with a tumor-targeting mutant of *S. typhimurium* inhibits liver metastasis in a mouse model of pancreatic cancer. *J Surg Res* 164:248–255. doi: [10.1016/j.jss.2009.02.023](#) PMID: [19766244](#)
10. Hiroshima Y, Zhao M, Zhang Y, Maawy A, Hassanein MK, Uehara F, et al. (2013) Comparison of efficacy of *Salmonella typhimurium* A1-R and chemotherapy on stem-like and non-stem human pancreatic cancer cells. *Cell Cycle* 12:2774–2780. doi: [10.4161/cc.25872](#) PMID: [23966167](#)
11. Hiroshima Y, Zhao M, Maawy A, Zhang Y, Katz MH, Fleming JB, et al. (2014) Efficacy of *Salmonella typhimurium* A1-R versus chemotherapy on a pancreatic cancer patient-derived orthotopic xenograft (PDOX). *J Cell Biochem* 115:1254–1261. doi: [10.1002/jcb.24769](#) PMID: [24435915](#)
12. Kimura H, Zhang L, Zhao M, Hayashi K, Tsuchiya H, Tomita K, et al. (2010) Targeted therapy of spinal cord glioma with a genetically-modified *Salmonella typhimurium*. *Cell Prolif* 43:41–48. doi: [10.1111/j.1365-2184.2009.00652.x](#) PMID: [19922490](#)
13. Momiyama M, Zhao M, Kimura H, Tran B, Chishima T, Bouvet M, et al. (2012) Inhibition and eradication of human glioma with tumor-targeting *Salmonella typhimurium* in an orthotopic nude-mouse model. *Cell Cycle* 11:628–632. doi: [10.4161/cc.11.3.19116](#) PMID: [22274398](#)
14. Zhao M, Suetsugu A, Ma H, Zhang L, Liu F, Zhang Y, et al. (2012) Efficacy against lung metastasis with a tumor-targeting mutant of *Salmonella typhimurium* in immunocompetent mice. *Cell Cycle* 11:187–193. doi: [10.4161/cc.11.1.18667](#) PMID: [22186786](#)
15. Hayashi K, Zhao M, Yamauchi K, Yamamoto N, Tsuchiya H, Tomita K, et al. (2009) Cancer metastasis directly eradicated by targeted therapy with a modified *Salmonella typhimurium*. *J Cell Biochem* 106:992–998. doi: [10.1002/jcb.22078](#) PMID: [19199339](#)
16. Hayashi K, Zhao M, Yamauchi K, Yamamoto N, Tsuchiya H, Tomita K, et al. (2009) Systemic targeting of primary bone tumor and lung metastasis of high-grade osteosarcoma in nude mice with a tumor-selective strain of *Salmonella typhimurium*. *Cell Cycle* 8:870–875. PMID: [19221501](#)
17. Fu X, Guadagni F, Hoffman RM. (1992) A metastatic nude-mouse model of human pancreatic cancer constructed orthotopically from histologically intact patient specimens. *Proc Natl Acad Sci USA* 89:5645–5649. PMID: [1608975](#)
18. Hiroshima Y, Zhang Y, Zhang M, Maawy A, Mii S, Yamamoto M, et al. (2015) Establishment of a patient-derived orthotopic xenograph (PDOX) model of HER-2-positive cervical cancer expressing the clinical metastatic pattern. *PLOS ONE* 10:e0117417. doi: [10.1371/journal.pone.0117417](#) PMID: [25689852](#)
19. Evans DB, Rich TA, Byrd DR, Cleary KR, Connelly JH, Levin B, et al. (1992) Preoperative chemoradiation and pancreaticoduodenectomy for adenocarcinoma of the pancreas. *Arch Surg* 127:1335–1339. PMID: [1359851](#)

20. Yamauchi K, Yang M, Jiang P, Xu M, Yamamoto N, Tsuchiya H, et al. (2006) Development of real-time subcellular dynamic multicolor imaging of cancer cell trafficking in live mice with a variable-magnification whole-mouse imaging system. *Cancer Res* 66:4208–4214. PMID: [16618743](#)
21. Yano S, Zhang Y, Miwa S, Tome Y, Hiroshima Y, Uehara F, et al. (2014) Spatial-temporal FUCCI imaging of each cell in a tumor demonstrates locational dependence of cell cycle dynamics and chemoresponsiveness. *Cell Cycle* 13:2110–2119. doi: [10.4161/cc.29156](#) PMID: [24811200](#)
22. Yano S, Zhang Y, Zhao M, Hiroshima Y, Miwa S, Uehara F, et al. (2014) Tumor-targeting *Salmonella typhimurium* A1-R decoys quiescent cancer cells to cycle as visualized by FUCCI imaging and become sensitive to chemotherapy. *Cell Cycle* 13:3958–3963. doi: [10.4161/15384101.2014.964115](#) PMID: [25483077](#)