



# HHS Public Access

Author manuscript

*J Holist Nurs.* Author manuscript; available in PMC 2016 September 01.

Published in final edited form as:

*J Holist Nurs.* 2015 September ; 33(3): 247–259. doi:10.1177/0898010115569349.

## Mindfulness-Based Stress Reduction in Advanced Nursing

### Practice:

**A Nonpharmacologic Approach to Health Promotion, Chronic Disease Management, and Symptom Control**

**Hants Williams, BSN, RN, PHN, Leigh Ann Simmons, PhD, MFT, and Paula Tanabe, PhD, MSN, MPH, RN, FAEN, FAAN**

Duke University

### Abstract

The aim of this article is to discuss how advanced practice nurses (APNs) can incorporate mindfulness-based stress reduction (MBSR) as a nonpharmacologic clinical tool in their practice. Over the last 30 years, patients and providers have increasingly used complementary and holistic therapies for the nonpharmacologic management of acute and chronic diseases. Mindfulness-based interventions, specifically MBSR, have been tested and applied within a variety of patient populations. There is strong evidence to support that the use of MBSR can improve a range of biological and psychological outcomes in a variety of medical illnesses, including acute and chronic pain, hypertension, and disease prevention. This article will review the many ways APNs can incorporate MBSR approaches for health promotion and disease/symptom management into their practice. We conclude with a discussion of how nurses can obtain training and certification in MBSR. Given the significant and growing literature supporting the use of MBSR in the prevention and treatment of chronic disease, increased attention on how APNs can incorporate MBSR into clinical practice is necessary.

### Keywords

psychosocial/mental health; health promotion; meditation/mindfulness; alternative therapies; chronic disease; holistic nursing; mind–body techniques

### Introduction

Due to changes resulting from implementation of the Affordable Care Act, advanced practice nurses (APNs) will assume a growing proportion of the responsibility for patient care over the next decade and beyond (H.R. 3590, Patient Protection and Affordable Care Act, 2010). Thus, it is critical that APNs continue to advance their knowledge and practices in the management of patient problems, including both pharmacologic and nonpharmacologic approaches.

---

Please address correspondence to Hants Williams, BSN, RN, PHN, Duke University, 301 Swift Avenue, #25 Durham, NC 27705; hants.williams@dm.duke.edu.

Over the past 30 years, both patients and providers have increasingly used complementary and holistic therapies for the nonpharmacologic management of acute and chronic diseases (Harris & Rees, 2000). Studies have shown that up to 40% of patients in the United States, and up to 44% in England, use one or more nontraditional therapies, some of which are prescribed by providers and some not (Eisenberg et al., 1998; Hunt et al., 2010). Mind–body practices such as acupuncture, chiropractic and spinal manipulation, massage therapy, yoga, tai chi, and mindfulness are among the most commonly used.

While there is a growing body of research establishing the benefits of many of these practices, mindfulness meditation—specifically mindfulness-based stress reduction (MBSR)—has robust evidence for its effectiveness in family, adult-gerontology, pediatric, women’s health, and psychiatric-mental health populations (Ludwig & Kabat-Zinn, 2008). Moreover, research is beginning to demonstrate how health care providers can use MBSR to reduce stress, burnout, and medical errors.

The purpose of this article is to discuss (1) the theory and mechanisms underlying MBSR; (2) the use of MBSR for disease prevention, chronic disease management, and symptom management; (3) how APNs can use MBSR to promote personal well-being; and (4) how nurses can refer patients for these interventions, and if desired, become MBSR certified to incorporate mindfulness into their own clinical care of patients.

## Background

### Overview of MBSR

MBSR is a relatively recent addition to the nonpharmacological approaches for disease management. Originally developed by John Kabat-Zinn in 1979 and first published in 1982, MBSR is based on the Buddhist principle of mindfulness. Mindfulness, defined as an awareness that emerges with purposeful, nonjudgmental attention within the present moment (Kabat-Zinn, 2003), is a fundamental concept of most meditation practices. Individuals learn mindfulness techniques, such as breath awareness (focus on breath and observing thoughts without being caught up in them) and body scan (promoting awareness and acceptance of sensations in different parts of the body), to help them understand that painful sensations and associated negative emotions do not need to be fought, suppressed, or they need not inhibit them from living a meaningful life and accomplishing goals.

### Theory and Concepts

Since the seminal 1982 publication, hundreds of studies have demonstrated the effectiveness of MBSR on health outcomes (Hölzel et al., 2011; MacCoon et al., 2012). However, the mechanism(s) through which MBSR works are still under investigation. Several psychological-based theories of mindfulness have been proposed, such as the Intentionally (I) Attending (A) with openness and nonjudgmental awareness (A) theory (IAA theory; Brown, Ryan, & Creswell, 2007; Shapiro, Carlson, Astin, & Freedman, 2006). In addition to psychological-driven theories of mindfulness, theories have also been based on therapeutic contexts, such as pain management (Day, Jensen, Ehde, & Thorn, 2014).

But at the core of mindfulness theories are four fundamental concepts that work and occur simultaneously during mindfulness practice: awareness, attention, acceptance, and re-perceiving. As described in a recent concept analysis of mindfulness (White, 2014), awareness is being deeply attentive to oneself in moment-to-moment experiences; attention is the act of staying with the moment-to-moment experience; and acceptance is receiving what arises without judgment, resistance, or avoidance.

Re-perceiving can be understood as a shift in perspective similar to the act of detachment (Bohart, 1983) or decentering (Safran & Segal, 1990). Instead of becoming immersed with a bodily sensation or particular situation, the individual attends to the immediate experience, without judgment, leading to a shift in perspective about the situation. For example, consider a person with sickle cell disease during a vaso-occlusive crisis. Common thoughts may include the following: "I'm having a pain crisis, which means I may need to go to the emergency department for treatment, potentially get admitted as an inpatient, and be put on new medications, and I'm really upset about this happening." In this case, the focus is on "what may happen" and the negative feelings associated with fears of future possibilities. Conversely, re-perceiving keeps the focus on the present moment. The individual may scan the body to identify the location and intensity of the pain and the physical response (e.g., clenching fists, shallow breathing). Awareness of breath techniques and yoga help maintain focus on the present moment as opposed to thoughts of future scenarios, which may or may not actually occur. This interoceptive (internally focused) attention has been shown to improve overall well-being (Farb, Segal, & Anderson, 2013).

### **Mechanisms**

Recent neuroimaging studies have started to identify how mindfulness practices influence neuro-cognitive pathways (Vago & Silbersweig, 2012) and brain morphology (Hölzel et al., 2011). Structural and functional magnetic resonance imaging studies have shown that mindfulness practice is associated with changes in the prefrontal cortex, the posterior and anterior cingulate cortex, the insular and amygdalae, and hippocampus. Specifically, after 8 weeks of MBSR, observed morphologic changes include increased cortical thickness and grey matter density in the posterior and anterior cingulate cortex, hippocampus, and insula (Farb et al., 2007; Hölzel et al., 2007; Hölzel et al., 2008; Hölzel et al., 2011; Lazar et al., 2005). Functional studies have demonstrated increased activation in these areas as well (Lutz et al., 2014). Conversely, studies have demonstrated decreases in both size and activation in the amygdala, which is primarily responsible for activation of the stress response (Desbordes et al., 2012).

Given that the anterior cingulate cortex and insula are involved with attention and body awareness (van Veen & Carter, 2002), and that areas of the prefrontal cortex are involved with acceptance and re-perceiving (in addition to other higher level cognitive and emotional processing, e.g., emotion regulation, perspective taking, and compassion), these findings support the proposed theoretical concepts of awareness, attention, acceptance, and re-perceiving as possible mechanisms of MBSR (Ochsner & Gross, 2005; Schmitz & Johnson, 2007). In addition to these mechanisms, many more are being explored to better understand and answer what areas of the brain are involved with mindfulness and self-

regulation exercises like attention regulation, body awareness, emotion regulation, and perspective taking of self (Hölzel et al., 2011).

### Data Sources

To explore the potential uses of MBSR for chronic disease management, symptom management, and how APNs may use MBSR practices to promote personal well-being for improved clinical care, an electronic database search was performed. The search was limited to English and published articles from 1982 (first publication of MBSR) through 2013. CINAHL and PubMed, PsychInfo, PsychArticles, Google Scholar, and EBSCO online databases were searched using keywords and subject headings. The main key phrases included were mindfulness-based stress reduction, MBSR, mindfulness, meditation, and mindfulness-based intervention. The search strategy yielded a list of approximately 500 citations. Because of the large number of findings, only the seminal and most recent works that discussed biological health outcomes for patient populations that are relevant for APNs were included. Limited psychological outcomes are reported, because most discussions of MBSR to date have focused on psychological outcomes, and numerous reviews have documented the efficacy of MBSR for improving psychological health and quality of life (see Keng, Smoski, & Robins, 2011, for most recent literature review). Therefore, the discussion below is not exhaustive but a brief overview of specific applications of MBSR that may be most useful for APNs. Based on the search results, the MBSR literature was classified by clinical applications: (1) preventive health and health enhancement, (2) chronic disease management, and (3) symptom management. Also included is a brief summary of the role of MBSR in supporting clinician well-being as it relates to enhanced clinical care.

### MBSR for Health Promotion and Prevention

Chronic diseases are among the most prevalent and preventable of all health problems (Centers for Disease Control and Prevention, 2014b). Reductions or cessation of risky behaviors such as smoking, drinking, sedentary lifestyle, and unhealthy eating could prevent onset of common cancers (Flanders, Lally, Zhu, Henley, & Thun, 2003), liver disease (Mann, Smart, & Govoni, 2003), obesity (World Health Organization, n.d.-c), cardiovascular disease, and type 2 diabetes. Because APNs are usually the first to see patients presenting with new signs of illness in the outpatient setting, they are at the front lines and best suited to provide patients at risk for chronic diseases with preventive health education and recommend interventions (Centers for Disease Control and Prevention, 2011b).

#### Smoking

Approximately one billion people in the world smoke tobacco-related products, resulting in nearly 6 million deaths annually (World Health Organization, n.d.-b). Estimates show that fewer than 10% of U.S. individuals who attempt to quit smoking will be successful, in large part due to tobacco's addictive nature (Centers for Disease Control and Prevention, 2011c). Emerging evidence suggests MBSR may be used in conjunction with standard therapies (e.g., medications, counseling, and combination therapy) to reduce tobacco consumption and cravings, and improve smoking cessation rates. In a pilot study of 18 tobacco-smoking

outpatients who averaged 19 cigarettes per day for 26 years, 56% (10/18) remained biologically smoke free 6 weeks after completing an 8-week MBSR course (Davis, Fleming, Bonus, & Baker, 2007). A similar study of 158 long-term smokers also found that increased mindfulness was associated with decreased nicotine dependence and nicotine withdrawal and increased sense of control over smoking (Vidrine et al., 2009). Two recent literature reviews on MBSR's for smoking cessation (Carim-Todd, Mitchell, & Oken, 2013; Chiesa & Serretti, 2014) have similarly concluded that preliminary evidence suggests mindfulness-based programs like MBSR can reduce addictive behaviors and substance abuse and consumption, but larger randomized controlled trials (RCT) are warranted.

### Dietary Behaviors

Despite recommendations to reduce consumption of unhealthy fats, there has been little change in the amount of fat intake over the past 20 years in the United States or the United Kingdom (Centers for Disease Control and Prevention, 2014c; National Health Service, n.d.). Moreover, 24% of men and 26% of women in the United Kingdom are overweight or obese (Health and Societal Care Information Centre, 2013), and 35.7% of all American adults *are obese* (Centers for Disease Control and Prevention, 2014a).

As briefly described in the chronic disease management section of this article (diabetes section), MBSR may help patients modify their dietary behaviors. Specifically, MBSR programs that have been modified to address specific dietary concerns have shown promising results. A pilot study that tested a 6-week modified MBSR program for 10 obese adults found statistically significant decreases in body mass index (BMI; mean loss of 1.2 kg over 12 weeks;  $p < .01$ ), large decreases in reported hunger at 6 weeks ( $\mu$  [ $SD$ ]: 7.6 [3.9] to 4.6 [3.5],  $p = .02$ ), and improved cognitive restraint at 12 weeks ( $\mu$  [ $SD$ ]: 9.5 [4.6] to 4.5 [2.5],  $p = .02$ ; Dalen et al., 2010). Conversely, a larger RCT of 47 obese women enrolled in a modified 9-week MBSR program found no statistically significant improvements in BMI or weight compared with a wait-listed control group (Daubenmier et al., 2011). However, the wait-listed control group gained weight over the course of the intervention and at follow-up, while the MBSR participants had no increases in weight or BMI.

There also is evidence to suggest that patients with eating disorders, such as binge eating, may benefit from MBSR. A feasibility study of 25 participants found that after completing a modified MBSR course centered on topics specific for binge eaters, there were statistically significant pre- to post-MBSR declines in binge eating severity, depressive symptoms, and state anxiety, as well as improved levels of self-acceptance (Smith, Shelley, Leahigh, & Vanleit, 2006).

### MBSR for Chronic Disease Management

Patients with chronic conditions must make daily choices about how to best self-manage their illness. MBSR is potentially useful for these patients, because the mindfulness practices can enhance self-efficacy in behavior and lifestyle management (Chang et al., 2004; Semple, Lee, Rosa, & Miller, 2010), in addition to improving physical and psychological outcomes associated with having a chronic condition. MBSR has been applied in the management of numerous chronic diseases, but for this section, we focus on how

MBSR may be useful in the management of chronic diseases with high worldwide prevalence: (1) hypertension, (2) type 2 diabetes, and (3) the human immunodeficiency virus (HIV)/acquired immunodeficiency syndrome (AIDS).

### **Hypertension**

Approximately 20% of the worldwide population has high blood pressure (BP; BP > 140/90), and in 2010, one out of three adults (67 million) in the United States had either hypertension or prehypertension (Egan, Zhao, & Axon, 2010; World Health Organization, 2002). In addition to prescribing anti-hypertension medication, current treatment for both prehypertension and hypertension includes lifestyle modifications like exercise, weight loss, dietary changes, and stress reduction. MBSR may be an additional behavioral approach to help patients manage their BP. Studies have demonstrated significant reductions in BP and BMI, in addition to anxiety and depression related to hypertension (Curiati et al., 2005; Griffiths, Camic, & Hutton, 2009; Parswani, Sharma, & Iyengar, 2013). A recent study of prehypertensive, unmedicated outpatients found statistically and clinically significant reductions in both BP (135/83 to 124/81) and BMI (25.2 to 23) after taking an 8-week MBSR course, and these improvements were maintained at 3 months postintervention (Hughes et al., 2013).

### **Type 2 Diabetes**

Since 1980, diabetes has increased by 133%, and the World Health Organization predicts that by 2030 the number of diabetic-related deaths will double (Centers for Disease Control and Prevention, 2011a; World Health Organization, n.d.-a). Evidence is growing that MBSR may help stabilize glycemic levels. A prospective observational study by Rosenzweig et al. (2007) found that after completing an 8-week MBSR program, type 2 diabetic patients had significant reductions in HbA(1c), mean arterial pressure, and general improvements in quality of life. From baseline (Week 0) to postintervention follow-up (Week 12), HbA(1c) decreased from a mean of 7.5 to 7.02 ( $p = .03$ ;  $d = .88$ ) and mean arterial pressure from 100 mmHg to 94 mmHg ( $p = .009$ ;  $d = .48$ ). Similarly, symptoms of depression dropped by 43% ( $p = .03$ ,  $d = .086$ ), anxiety by 37% ( $p = .33$ ,  $d = .43$ ), and general distress by 35% ( $p = .07$ ,  $d = .60$ ). In addition to glycemic control, MBSR may also help with dietary modification. After completing an 8-week MBSR program, diabetic outpatients reported significant improvements in cognitive control over eating behaviors and self-efficacy of diabetes management, had greater reductions in sugar and trans fat intake, and increased their consumption of fiber (Miller, Kristellar, Headings, & Nagaraja, 2014).

### **HIV/AIDS**

Worldwide, approximately 35 million people now live with HIV/AIDS, and the number of newly affected in 2011 exceeded 2.2 million people (Joint United Nations Programme on HIV/AIDS report, n.d.). For persons infected with HIV/AIDS and receiving antiretroviral therapy (ART), MBSR may provide some relief from ART side effects. Compared with wait-listed controls, HIV+ patients who participated in an 8-week MBSR program experienced a 39.1% reduction in ART symptoms at 3 months postintervention, and 46.2% reduction in ART symptoms 6 months postintervention (Duncan et al., 2012). MBSR has

also led to improvements in T and NK cells. In a nonrandomized, pre–post comparison study by Jam et al. (2010), HIV+ patients showed significant improvements in their CD4 counts after participating in an 8-week MBSR program. Creswell, Myers, Cole, and Irwin (2009) also have demonstrated comparable results; MBSR buffered CD4+ T lymphocyte decline in HIV+ adults who participated in an 8-week MBSR program. Last, in a quasi-experimental study by Robinson, Matthews, and Witek-Janusek (2003), HIV+ adults assigned to MBSR had clinically significant increases and activity of NK cells compared with a comparison group. Taken together, MBSR is a potentially useful tool for reducing the number of side effects related to taking antiretroviral medication and buffering against further declines in immune function.

### **Clinical Depression**

General medical inpatients in both North America and Europe experience levels of depressive symptoms that exceed population averages (Creed et al., 2002). Approximately 5% to 10% of inpatients will present with signs and symptoms of depression (Himelhoch, Weller, Wu, Anderson, & Cooper, 2004), while rates of depression may exceed 20% for patients with chronic illnesses (e.g., coronary heart disease, diabetes, and chronic pain; Anderson, Freedland, Clouse, & Lustman, 2001; Thombs et al., 2006). MBSR has shown great success in reducing depressive symptoms. A recent meta-analysis of 39 MBSR studies conducted between 1992 to 2009 found large effect sizes for depressive symptom reduction in those diagnosed with major depressive disorder (Hedge's  $g = 0.95$ ; 95% confidence interval [CI] = 0.71–1.18,  $p < .01$ ), and moderately strong effect sizes for those without major depressive disorder but with elevated levels of depressive symptoms (Hedge's  $g = 0.59$ ; 95% CI = 0.51–0.66,  $p < .01$ ; Hofmann, Sawyer, Witt, & Oh, 2010). These findings suggest that MBSR is applicable for a variety of patients in reducing depressive symptoms, regardless of primary diagnosis, and for patients both with and without major depressive disorder. Therefore, patients who are experiencing acute depressive symptoms (e.g., related to prolonged hospitalization, poor prognosis, terminal illness, and slow recovery time) or have major depressive disorder may greatly benefit from MBSR.

### **MBSR for Symptom Management**

#### **Stress**

Defined as “a particular relationship between the person and the environment that is appraised by the person as taxing or exceeding his or her resources and endangering his or her wellbeing” (Lazarus, 1999, p. 19), stress is common among patients and closely associated with immune functioning. Regardless of diagnosis, the majority of inpatients will experience some degree of stress related to their hospitalization. Patients admitted through the emergency department may become stressed due to long waiting times, and once admitted, they may experience stress before or during a nursing procedure (e.g., catheter or tube placement, intravenous line insertion), or while in a preoperative suite awaiting surgery. In general, patients usually experience both physiologic stress due to their illness and psychological stress due to being in the hospital and the meaning of the illness in their life.





Von Korff, Simon, & Gater, 1998). Additionally, reports of chronic pain in pediatric admissions have increased by over 800% from 2004 to 2010 (Coffelt, Bauer, & Carroll, 2013). When compared with other chronic conditions such as diabetes, coronary heart disease, stroke, and cancer, more people suffer from chronic pain (116 million in the United States and 10 million in the United Kingdom) than these conditions combined (GfK NOP British Pain Society, n.d.; Institute of Medicine, 2011). Financially, treatment costs between \$11 and \$20 billion annually for just chronic back pain in the United Kingdom (Maniadakis & Gray, 2000), and in the United States, the amount of money lost in productive time exceeds \$61.2 billion (Stewart, Ricci, Chee, & Morganstein, 2003).

The first study that demonstrated efficacy of MBSR in reducing chronic pain (33% reduction in mean total of a pain rating index) was with a diverse group of chronic pain patients 30 years ago (Kabat-Zinn, 1984), and since that first publication, numerous studies have underscored these seminal findings. MBSR has been shown to reduce pain intensity (Grossman, Tiefenthaler-Gilmer, Raysz, & Kesper, 2007; Rosenzweig et al., 2010), increase gray matter density in areas of the brain that regulate pain perception (Carrasquillo & Gereau, 2007; Hölzel et al., 2010; Hölzel et al., 2011), improve acceptance of chronic pain and continual engagement of daily living activities (Morone, Greco, & Weiner, 2008), and decrease depression and anxiety related to chronic pain (Astin, Shapiro, Eisenberg, & Forsys, 2003; Gardner-Nix, Backman, Barbati, & Grummitt, 2008; Pradhan et al., 2007; Sagula & Rice, 2004; Sephton et al., 2007). What is most important about these findings is that they are long-lasting. Sustained benefits of improved coping, decreased somatic complaints and depression, and reduced pain intensity have been found up to 3 years after MBSR program completion in chronic pain patients (Grossman et al., 2007). Thus, chronic pain patients who experience related psychological and/or psychological symptoms also may greatly benefit by participating in an MBSR program.

## Sleep

Getting a full night of undisturbed sleep is a common problem for hospitalized patients. Compared with nonhospitalized persons, inpatients are more fatigued on awaking, report worse sleep quality and quality of life, and show more circadian rhythm disruption (Davidson, MacLean, Brundage, & Schulze, 2002; Fernandes, Stone, Andrews, Morgan, & Sharma, 2006). MBSR can improve a number of factors related to sleep. Various studies have shown MBSR can improve reported quality of sleep (Carlson & Garland, 2005), decrease the number of sleep disturbances (Gross et al., 2004), and increase sleep duration (Garland et al., 2014). In addition, a literature review of 38 articles on MBSR and sleep concluded that there is evidence to support MBSR is associated with improved sleep and with decreased cognitive processes (e.g., worry) that may interfere with sleep (Winbush, Gross, & Kreitzer, 2007). These data suggest that MBSR may be a potentially powerful tool to help hospitalized patients suffering from poor sleep.

## Provider Development and Self-Care

Growing demands for health care combined with provider shortages, long work hours, and worsening population health contribute to increased stress and burnout among health care workers, including nurses. Highest estimates report that 40% of nurses and 20% of

physicians will experience burnout at some point during their career (Bruce, Conaglen, & Conaglen, 2005; Vahey, Aiken, Sloane, Clarke, & Vargas, 2004). Moreover, studies have shown that provider burnout is associated with higher frequency of medical errors, higher rates of depression, decreased quality of life, lower personal accomplishment, and higher turnover rates (West et al., 2006; West, Tan, Habermann, Sloan, & Shanafelt, 2009). Patient satisfaction with quality of care is also lower among providers reporting higher levels of burnout (Aiken et al., 2012). An increasing number of studies have demonstrated that MBSR may improve physical and emotional health as well as reduce work-related distress among health care providers. Multiple RCTs have shown that compared to wait-listed control groups, nurses and physicians who participate in four to eight sessions of MBSR report significantly lower levels of stress, fewer symptoms of burnout (emotional exhaustion), and increased life satisfaction, empathy, and self-compassion (Cohen-Katz et al., 2005; Mackenzie, Poulin, & Siedman-Carlson, 2006; Shapiro, Astin, Bishop, & Cordova, 2005). Based on these findings, some experts have suggested that mindfulness may be a powerful tool to enhance the patient-provider relationship, improve cognitive processes involved in decision making and diagnosis, and promote resilience (Groopman, 2007; Ludwig & Kabat-Zinn, 2008).

## How to Incorporate MBSR in Clinical Practice

### Referrals

Based on the existing advanced practice nursing regulatory model (which includes the essential elements of licensure, accreditation, certification, and education), nurses are in an ideal situation to make patient referrals to MBSR programs (Advanced Practice Registered Nurses Consensus Work Group, 2008). In the existing advanced practice nursing regulatory model, the major population foci are family, adult-gerontology, pediatric, neonatology, women's health/gender-related, and psychiatric-mental health populations. As described, MBSR has already been implemented and tested within family, adult-gerontology, pediatric, women's health/gender-related, and psychiatric-mental health populations to help treat a variety of symptoms across both acute and chronic illnesses with great success (Ludwig & Kabat-Zinn, 2008). APNs who practice within these areas are thus in an ideal situation to help identify patients who may benefit from referral to an MBSR program.

### Informal Patient Teaching and Education

Advanced practice registered nurses who practice in the hospital or community can also incorporate mindfulness informally into their own practice. Nurses can enroll in MBSR programs to acquire basic knowledge and exercises, which can then be disseminated to their patients. For example, a nurse can guide an anxious patient through a breathing or body-scan exercise. Most major hospitals and universities offer MBSR programs that nurses can enroll in. Within the United States alone, there are almost 1,000 certified MBSR instructors and mindfulness research centers that offer mindfulness-based programs in all 50 states (University of Massachusetts; <http://w3.umassmed.edu/MBSR/>).

## Formal Training and Certification

Last, for nurses that want to become certified MBSR instructors and conduct their own MBSR classes or workshops, they must attend the Center of Mindfulness and Stress Reduction clinic at the University of Massachusetts. Currently, the University of Massachusetts has the only program in the United States that grants MBSR certification. Run by John Kabat-Zinn and associates, training occurs over the course of 2 years, is composed of both online and in-person classes, and runs upwards of \$3,000 to complete all training certification requirements. Although the process of becoming certified does not require an additional degree, certification cannot simply be done in just a few weeks or months. And even after one becomes certified, it can take years of additional practice to become comfortable enough to instruct patients that have severe symptoms or distressing thoughts.

## Conclusion

There is strong evidence to support that MBSR can improve a range of biologic and psychological outcomes in a variety of medical illnesses; therefore, current and future APNs should be familiar with the benefits of MBSR for both their patients and themselves. Advanced practice and registered nurses have important relationships with their patients in both inpatient and outpatient settings that make them ideal candidates to make referrals to MBSR, as well as receive training and certification in MBSR so they can share the practice with their patients in the context of individual and group medical appointments. Future studies should investigate implementation of MBSR into advanced clinical nursing practice to develop best practices.

## References

- Advanced Practice Registered Nurses Consensus Work Group. Consensus model for APRN regulation: Licensure, accreditation, certification, & education. 2008. Retrieved from [https://www.bon.texas.gov/pdfs/forms\\_pdfs/applications\\_pdfs/aprn\\_pdfs/aprnmodel.pdf](https://www.bon.texas.gov/pdfs/forms_pdfs/applications_pdfs/aprn_pdfs/aprnmodel.pdf)
- Aiken LH, Sermeus WS, Van den Heede K, Sloane DM, Busse R, McKee M, Kutney L. Patient safety, satisfaction, and quality of hospital care: Cross sectional surveys of nurses and patients in 12 countries in Europe and the United States. *British Medical Journal*. 2012; 344:e1717.10.1136/bmj.e1717 [PubMed: 22434089]
- Anderson RJ, Freedland KE, Clouse RE, Lustman PJ. The prevalence of comorbid depression in adults with diabetes. *Diabetes Care*. 2001; 24:1069–1078. [PubMed: 11375373]
- Astin JA, Shapiro SL, Eisenberg DM, Forsys KL. Mind-body medicine: State of the science, implications for practice. *Journal of the American Board of Family Practice*. 2003; 16:131–147. [PubMed: 12665179]
- Birnie K, Garland SN, Carlson LE. Psychological benefits for cancer patients and their partners participating in mindfulness-based stress reduction (MBSR). *Psycho-Oncology*. 2010; 19:1004–1009. [PubMed: 19918956]
- Bohart, A. Detachment: A variable common to many psychotherapies?. Paper presented at the 63rd Annual Convention of the Western Psychological Association; San Francisco, CA. 1983 Apr.
- Brown KW, Ryan RM, Creswell JD. Mindfulness: Theoretical foundations and evidence for its salutary effects. *Psychological Inquiry*. 2007; 18:211–237.
- Bruce SM, Conaglen HM, Conaglen JV. Burnout in physicians: A case report for peer-support. *Internal Medicine Journal*. 2005; 35:272–278. [PubMed: 15845108]

- Carim-Todd L, Mitchell SH, Oken BS. Mind-body practices: An alternative, drug-free treatment for smoking cessation? A systematic review of the literature. *Drug and Alcohol Dependence*. 2013; 132:399–410. [PubMed: 23664122]
- Carlson LE, Garland S. Impact of mindfulness-based stress reduction (MBSR) on sleep quality in cancer patients. *International Journal of Behavioral Medicine*. 2005; 12:278–285. [PubMed: 16262547]
- Carrasquillo Y, Gereau RW. Activation of the extracellular signal regulated kinase in the amygdala modulates pain perception. *Journal of Neuroscience*. 2007; 27:1543–1551. [PubMed: 17301163]
- Centers for Disease Control and Prevention. National diabetes fact sheet, 2011. 2011a. Retrieved from [http://www.cdc.gov/diabetes/pubs/pdf/ndfs\\_2011.pdf](http://www.cdc.gov/diabetes/pubs/pdf/ndfs_2011.pdf)
- Centers for Disease Control and Prevention. Physician assistant and advance practice nurse care in hospital outpatient departments: United States, 2008–2009. 2011b. (NCHS data brief). Retrieved from <http://www.cdc.gov/nchs/data/databriefs/db77.htm>
- Centers for Disease Control and Prevention. Quitting smoking among adults: United States, 2001–2010. *Morbidity and Mortality Weekly Report*. 2011c; 60:1513–1519. Retrieved from <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6044a2.htm>. [PubMed: 22071589]
- Centers for Disease Control and Prevention. Adult obesity facts. 2014a. Retrieved from <http://www.cdc.gov/obesity/data/adult.html>
- Centers for Disease Control and Prevention. Chronic diseases and health promotion. 2014b. Retrieved from <http://www.cdc.gov/chronicdisease/overview/>
- Centers for Disease Control and Prevention. Trans fat. 2014c. Retrieved from <http://www.cdc.gov/nutrition/everyone/basics/fat/transfat.html>
- Chang VY, Palesh O, Caldwell R, Glasgow N, Abramson M, Luskin F, Koopman C. The effects of a mindfulness based stress reduction program on stress, mindfulness self-efficacy, and positive states of mind. *Stress & Health*. 2004; 20:141–147.
- Chapman CR, Turner JA. Psychological control of acute pain in medical settings. *Journal of Pain and Symptom Management*. 1986; 1:9–20. [PubMed: 3633305]
- Chiesa A, Serretti A. Are mindfulness-based interventions effective for substance use disorders? A systematic review of the evidence. *Substance Use & Misuse*. 2014; 49:492–512. [PubMed: 23461667]
- Coffelt TA, Bauer BD, Caroll AE. Inpatient characteristics of the child admitted with chronic pain. *Pediatrics*. 2013; 132:422–429.
- Cohen-Katz J, Wileys SD, Capuano T, Bakers DM, Kimmel S, Shapiro S. The effects of mindfulness-based stress reduction on nurse stress and burnout, Part II: A quantitative and qualitative study. *Holistic Nursing Practice*. 2005; 19:26–35. [PubMed: 15736727]
- Cordell WC, Keene KK, Giles BK, Jones JB, Jones JH, Brizendine EJ. The high prevalence of pain in emergency medical care. *American Journal of Emergency Medicine*. 2002; 20:165–169. [PubMed: 11992334]
- Creed F, Morgan R, Fiddler M, Marshall S, Guthrie E, House A. Depression and anxiety impair health-related quality of life and are associated with increased costs in general medical inpatients. *Psychosomatics*. 2002; 43:302–309. [PubMed: 12189256]
- Creswell JD, Myers HF, Cole SW, Irwin MR. Mindfulness meditation training effects on CD4+ T lymphocytes in HIV-1 infected adults: A small randomized controlled trial. *Brain, Behavior, and Immunity*. 2009; 23:184–188.
- Curiati JA, Boochi E, Freire JO, Arantes AC, Braga M, Garcia Y, Fo WJ. Meditation reduces sympathetic activation and improves the quality of life in elderly patients with optimally treated heart failure: A prospective randomized study. *Journal of Alternative and Complementary Medicine*. 2005; 11:465–472. [PubMed: 15992231]
- Dalen J, Smith B, Shelley BM, Sloan A, Leahigh L, Begay D. Pilot study: Mindful eating and living (MEAL): Weight, eating behavior, and psychological outcomes associated with a mindfulness-based intervention for people with obesity. *Complementary Therapies in Medicine*. 2010; 18:260–264. [PubMed: 21130363]
- Daubenmier J, Kristeller J, Hect FM, Maninger N, Kuwata M, Jhaveri K, Epel E. Mindfulness intervention for stress eating to reduce cortisol and abdominal fat among overweight and obese

- women: An exploratory randomized controlled study. *Journal of Obesity*, 2011. 2011:Article ID 651936.10.1155/2011/651936
- Davidson JR, MacLean AW, Brundage MD, Schulze K. Sleep disturbance in cancer patients. *Social Science & Medicine*. 2002; 54:1309–1321. [PubMed: 12058848]
- Davis JM, Fleming MF, Bonus KA, Baker TB. A pilot study on mindfulness based stress reduction for smokers. *BMC Complementary and Alternative Medicine*. 2007; 7:2. [PubMed: 17254362]
- Day M, Jensen MP, Ehde DM, Thorn BE. Toward a theoretical model for mindfulness-based pain management. *Journal of Pain*. 2014; 15:691–703. [PubMed: 24985483]
- Desbordes G, Negi LT, Pace TW, Wallace BA, Raison CL, Schwartz EL. Effects of mindful-attention and compassion meditation training on amygdala response to emotional stimuli in an ordinary, non-meditative state. *Frontiers in Human Neuroscience*. 2012; 6:292. [PubMed: 23125828]
- Duncan LG, Moskowitz JT, Neilands TB, Dilworth SE, Hecht FM, Johnson MO. Mindfulness-based stress reduction for HIV treatment side effects: A randomized, wait-list controlled trial. *Journal of Pain and Symptom Management*. 2012; 43:161–171. [PubMed: 21925831]
- Egan BM, Zhao Y, Axon RN. US trends in prevalence, awareness, treatment, and control of hypertension, 1988–2008. *Journal of the American Medical Association*. 2010; 303:2043–2050. [PubMed: 20501926]
- Eisenberg DM, Davis RB, Ettner SL, Appel S, Wilkey S, Van Rompay M, Kessler RC. Trends in alternative medicine use in the United States, 1990–1997: Results of a follow-up national survey. *Journal of the American Medical Association*. 1998; 280:1569–1575. [PubMed: 9820257]
- Farb NAS, Segal ZV, Anderson AK. Mindfulness meditation training alters cortical representations of interoceptive attention. *Social Cognitive and Affective Neuroscience*. 2013; 8:15–26. [PubMed: 22689216]
- Farb NAS, Segal ZV, Mayberg H, Bean J, McKeon D, Fatima Z, Anderson AK. Attending to the present: Mindfulness meditation reveals distinct neural modes of self-reference. *Social Cognitive and Affective Neuroscience*. 2007; 2:313–322. [PubMed: 18985137]
- Feldner MT, Hekmat H, Zvolensky MJ, Vowles KE, Secrist Z, Leen-Feldner EW. The role of experiential avoidance in acute pain tolerance: A laboratory test. *Journal of Behavior Therapy and Experimental Psychiatry*. 2006; 37:146–158. [PubMed: 15882839]
- Fernandes R, Stone P, Andrews P, Morgan R, Sharma S. Comparison between fatigue, sleep disturbance, and circadian rhythm in cancer inpatients and healthy volunteers: Evaluation of diagnostic criteria for cancer-related fatigue. *Journal of Pain and Symptom Management*. 2006; 32:245–254. [PubMed: 16939849]
- Flanders WD, Lally CA, Zhu BP, Henley SJ, Thun MJ. Lung cancer mortality in relation to age, duration of smoking, and daily cigarette consumption: Results from Cancer Prevention Study II. *Cancer Research*. 2003; 63:6556–6562. [PubMed: 14559851]
- Gardner-Nix J, Backman S, Barbati J, Grummitt J. Evaluating distance education of a mindfulness-based meditation program for chronic pain management. *Journal of Telemedicine and Telecare*. 2008; 14:88–92. [PubMed: 18348755]
- Garland SN, Carlson LE, Stephens AJ, Antle MC, Samuels C, Campbell TS. Mindfulness-based stress reduction compared with cognitive behavioral therapy for the treatment of insomnia comorbid with cancer: A randomized, partially blinded, noninferiority trial. *Journal of Clinical Oncology*. 2014; 32:449–457.10.1200/JCO.2012.47.7265 [PubMed: 24395850]
- Gaylord SA, Palsson OS, Garland EL, Faurot KR, Coble RS, Mann JD, Whitehead WE. Mindfulness training reduces the severity of irritable bowel syndrome in women: Results of a randomized controlled trial. *American Journal of Gastroenterology*. 2011; 106:1678–1688. [PubMed: 21691341]
- GfK NOP British Pain Society. 2005 surveys & reports pain survey. n.d. Retrieved from <http://www.britishpainsociety.org/media-resources/#surveys-and-reports>
- Gureje O, Von Korff M, Simon GE, Gater R. Persistent pain and well-being: A World Health Organization study in primary care. *Journal of the American Medical Association*. 1998; 280:147–151. [PubMed: 9669787]

- Griffiths K, Camic PM, Hutton JM. Participant experiences of a mindfulness-based cognitive therapy group for cardiac rehabilitation. *Journal of Health Psychology*. 2009; 14:675–681. [PubMed: 19515682]
- Groopman, J. *How doctors think*. Boston, MA: Houghton Mifflin; 2007.
- Gross CR, Kreitzer MJ, Russas V, Treesak C, Frazier PA, Hertz MI. Mindfulness meditation to reduce symptoms after organ transplant: A pilot study. *Alternative Therapies in Health and Medicine*. 2004; 10(3):58–66. [PubMed: 15154154]
- Grossman P, Niemann L, Schmidt S, Walach H. Mindfulness-based stress reduction and health benefits: A meta-analysis. *Journal of Psychosomatic Research*. 2004; 57:35–43. [PubMed: 15256293]
- Grossman P, Tiefenthaler-Gilmer U, Raysz A, Kesper U. Mindfulness training as an intervention for fibromyalgia: Evidence of postintervention and 3-year follow-up benefits in well-being. *Psychotherapy and Psychosomatics*. 2007; 76:226–233. [PubMed: 17570961]
- Harris P, Rees R. The prevalence of complementary and alternative medicine use among the general population: A systematic review of the literature. *Complementary Therapies in Medicine*. 2000; 8(2):88–96. [PubMed: 10859601]
- Health and Societal Care Information Centre. *Statistics on obesity, physical activity and diet*: England, 2013 [NS]. 2013. Retrieved from <http://www.hscic.gov.uk/catalogue/PUB10364>
- Himelhoch S, Weller WE, Wu AW, Anderson GF, Cooper LA. Chronic medical illness, depression, and use of acute medical services among Medicare beneficiaries. *Medical Care*. 2004; 42:512–521. [PubMed: 15167319]
- Hofmann S, Sawyer AT, Witt AA, Oh D. The effect of mindfulness-based therapy on anxiety and depression: A meta-analytic review. *Journal of Consulting and Clinical Psychology*. 2010; 78:169–183. [PubMed: 20350028]
- Hölzel BK, Carmody J, Evans KC, Hoge EA, Dusek JA, Morgan L, Lazar SW. Stress reduction correlates with structural changes in the amygdala. *Social Cognitive and Affective Neuroscience*. 2010; 5:11–17. [PubMed: 19776221]
- Hölzel BK, Carmody J, Vangel M, Congleton C, Yerramsetti SM, Gard T, Lazar SW. Mindfulness practice leads to increases in regional brain gray matter density. *Psychiatry Research*. 2011; 191:36–43. [PubMed: 21071182]
- Hölzel BK, Ott U, Gard T, Hempel H, Weygandt M, Morgen K, Vaitl D. Investigation of mindfulness meditation practitioners with voxel-based morphometry. *Social Cognitive and Affective Neuroscience*. 2008; 3:55–61. [PubMed: 19015095]
- Hölzel BK, Ott U, Hempel H, Hackl A, Wolf K, Stark R, Vaitl D. Differential engagement of anterior cingulate and adjacent medial frontal cortex in adept meditators and non-meditators. *Neuroscience Letters*. 2007; 421:16–21. [PubMed: 17548160]
- Hunt KJ, Coelho HF, Wider B, Perry R, Hung SK, Ernst T. Complementary and alternative medicine use in England: Results from a national survey. *International Journal of Clinical Practice*. 2010; 64:1496–1502. [PubMed: 20698902]
- Hughes JW, Fresco DM, Myerscough R, van Dulmen MHM, Carlson LE, Josephson R. Randomized controlled trial of mindfulness-based stress reduction for prehypertension. *Psychosomatic Medicine*. 2013; 75:8.
- Institute of Medicine. *Relieving pain in America: A blueprint for transforming prevention, care, education, and research*. 2011. Retrieved from <http://www.iom.edu/Reports/2011/Relieving-Pain-in-America-A-Blueprint-for-transforming-Prevention-Care-Education-Research.aspx>
- Jam S, Imani AH, Foroughi M, Alinaghi S, Koochack HE, Mohraz M. The effects of mindfulness-based stress reduction (MBSR) program in Iranian HIV/AIDS patients: A pilot study. *Acta Medica Iranica*. 2010; 48:101–106. [PubMed: 21133002]
- Joint United Nations Programme on HIV/AIDS. 2013 fact sheet. n.d. Retrieved from <http://www.unaids.org/en/resources/campaigns/globalreport2013/factsheet/>
- Kabat-Zinn J. An outpatient program in behavioral medicine for chronic pain patients based on the practice of mindfulness meditation: Theoretical considerations and preliminary results. *General Hospital Psychiatry*. 1982; 4:33–47. [PubMed: 7042457]

- Kabat-Zinn J. An outpatient program in behavioral medicine for chronic pain patients based on the practice of mindfulness meditation: Theoretical considerations and preliminary results. *REVISION*. 1984; 7:71–72.
- Kabat-Zinn J. Mindfulness-based interventions in context: Past, present, and future. *Clinical Psychology: Science and Practice*. 2003; 10:144–156.
- Kabat-Zinn J, Wheeler E, Light T, Skillings A, Scharf MJ, Cropley TG, Bernhard JD. Influence of a mindfulness meditation-based stress reduction intervention on rates of skin clearing in patients with moderate to severe psoriasis undergoing phototherapy (UVB) and photochemotherapy (PUVA). *Psychosomatic Medicine*. 1998; 60:625–632. [PubMed: 9773769]
- Keng SL, Smoski MJ, Robins CJ. Effects of mindfulness on psychological health: A review of empirical studies. *Clinical Psychology Review*. 2011; 31:1041–1056. [PubMed: 21802619]
- Lazar SW, Kerr CE, Wasserman RH, Gray JR, Greve DN, Treadway MT, Fischi B. Medication experience is associated with increased cortical thickness. *Neuroreport*. 2005; 16:1893–1897. [PubMed: 16272874]
- Lazarus, RS. *Stress and emotion: A new synthesis*. New York, NY: Springer; 1999.
- Liebkeind JC. Pain can kill. *Pain*. 1991; 44:3–4. [PubMed: 2038486]
- Ludwig DS, Kabat-Zinn J. Mindfulness in medicine. *Journal of the American Medical Association*. 2008; 300:1350–1352. [PubMed: 18799450]
- Lutz J, Herwig U, Opialla S, Hittmeyer A, Janacke L, Rufer M, Bruhl AB. Mindfulness and emotion regulation: An fMRI study. *Social Cognitive and Affective Neuroscience*. 2014; 9:776–785. [PubMed: 23563850]
- MacCoun DG, Imel ZE, Rosenkranz MA, Sheftel JG, Weng HY, Sullivan JC, Lutz A. The validation of an active control intervention for mindfulness based stress reduction (MBSR). *Behaviour Research and Therapy*. 2012; 50:3–12. [PubMed: 22137364]
- Mackenzie CS, Poulin PA, Siedman-Carlson R. A brief mindfulness-based stress reduction intervention for nurses and nurse aides. *Applied Nursing Research*. 2006; 19:105–109. [PubMed: 16728295]
- Maniadakis N, Gray A. The economic burden of back pain in the UK. *Pain*. 2000; 84:95–103. [PubMed: 10601677]
- Mann RE, Smart RG, Govoni R. The epidemiology of alcoholic liver disease. *Alcohol Research & Health*. 2003; 27:209–219. [PubMed: 15535449]
- Miller CK, Kristellar JL, Headings A, Nagaraja H. Comparison of a mindful eating intervention to a diabetes self-management intervention among adults with type 2 diabetes: A randomized controlled trial. *Health Education & Behavior*. 2014; 41:145–154. [PubMed: 23855018]
- Minor HG, Carlson LE, Mackenzie MJ, Zernicke K, Jones L. Evaluation of mindfulness-based stress reduction (MBSR) program for caregivers of children with chronic conditions. *Social Work in Health Care*. 2006; 43(1):91–109. [PubMed: 16723337]
- Molitor, N.; Keck, A.; Nordal, K. *Coping with chronic pain*. Washington, DC: American Psychological Association; 2004. Retrieved from <http://www.apa.org/helpcenter/chronic-pain.aspx>
- Morone NE, Greco CM, Weiner DK. Mindfulness meditation for the treatment of chronic low back pain in adults: A randomized controlled pilot study. *Pain*. 2008; 134:310–319. [PubMed: 17544212]
- National Health Service. *Fat: The facts*. n.d. Retrieved from <http://www.nhs.uk/Livewell/Goodfood/Pages/Fat.aspx>
- Oschner KN, Gross JJ. The cognitive control of emotion. *Trends in Cognitive Sciences*. 2005; 9:242–249. [PubMed: 15866151]
- Parswani MJ, Sharma MP, Iyengar SS. Mindfulness-based stress reduction program in coronary heart disease: A randomized control trial. *International Journal of Yoga*. 2013; 6:111–117. [PubMed: 23930029]
- Patient Protection and Affordable Care Act of 2010, H.R. 3590, 111th Cong. Retrieved from <https://www.govtrack.us/congress/bills/111/hr3590/text>
- Pradhan EK, Baumgarten M, Langenberg P, Handwerger B, Gilpin AK, Magyari T, Berman BM. Effect of mindfulness-based stress reduction in rheumatoid arthritis patients. *Arthritis and Rheumatism*. 2007; 57:1134–1142. [PubMed: 17907231]

- Robinson FP, Matthews HL, Witek-Janusek L. Psycho-endocrine-immune response to mindfulness-based stress reduction in individuals infected with the human immunodeficiency virus: A quasi-experimental study. *Journal of Alternative and Complementary Medicine*. 2003; 9:683–694. [PubMed: 14629846]
- Rosenzweig S, Greeson JM, Reibel DK, Green JS, Jasser SA, Beasley D. Mindfulness-based stress reduction for chronic pain conditions: Variation in treatment outcomes and role of home meditation practice. *Journal of Psychosomatic Research*. 2010; 68:29–36. [PubMed: 20004298]
- Rosenzweig S, Reibel DK, Greeson JM, Edman JS, Jasser SA, McMearty KD, Goldstein BJ. Mindfulness-based stress reduction is associated with improved glycemic control in type 2 diabetes mellitus: A pilot study. *Alternative Therapies in Health and Medicine*. 2007; 13:36–38. [PubMed: 17900040]
- Safran, JD.; Segal, ZV. *Interpersonal process in cognitive therapy*. New York, NY: Basic Books; 1990.
- Sagula D, Rice KG. The effectiveness of mindfulness training on the grieving process and emotional well-being of chronic pain patients. *Journal of Clinical Psychology in Medical Settings*. 2004; 11:333–342.
- Schmitz TW, Johnson SC. Relevance to self: A brief review and framework of neural systems underlying appraisal. *Neuroscience and Biobehavioral Review*. 2007; 31:585–596.
- Semple RJ, Lee J, Rosa D, Miller LF. A randomized trial of mindfulness-based cognitive therapy for children: Promoting mindful attention to enhance social-emotional resiliency in children. *Journal of Child and Family Studies*. 2010; 19:218–229.
- Septon SE, Salmon P, Weissbecker I, Ulmer C, Floyd A, Hoover K, Studts JL. Mindfulness meditation alleviates depressive symptoms in women with fibromyalgia: Results of a randomized clinical trial. *Arthritis and Rheumatism*. 2007; 57:77–85. [PubMed: 17266067]
- Shapiro SL, Astin JA, Bishop SR, Cordova M. Mindfulness-based stress reduction for health care professionals: Results from a randomized trial. *International Journal of Stress Management*. 2005; 12:164–176.
- Shapiro SL, Carlson LE, Astin JA, Freedman B. Mechanisms of mindfulness. *Journal of Clinical Psychology*. 2006; 62:373–386. [PubMed: 16385481]
- Smith BW, Shelley BM, Leahigh L, Vanleit B. A preliminary study of the effects of a modified mindfulness intervention on binge eating. *Complementary Health Practice Review*. 2006; 11:133–143.
- Sommer M, de Rijke JM, van Kleef M, Kessels AG, Peters ML, Geurts JW, Marcus MA. The prevalence of postoperative pain in a sample of 1490 surgical inpatients. *European Journal of Anaesthesiology*. 2008; 25:267–274. [PubMed: 18053314]
- Stewart WF, Ricci JA, Chee E, Morganstein D. Lost productive work time costs from health conditions in the United States: Results from the American productivity audit. *Journal of Occupational and Environmental Medicine*. 2003; 45:1234–1246. [PubMed: 14665809]
- Thombs BD, Bass EB, Ford DE, Stewart KJ, Tsilidis KK, Patel U, Ziegelstein RC. Prevalence of depression in survivors of acute myocardial infarction. *Journal of General Internal Medicine*. 2006; 21:30–38. [PubMed: 16423120]
- Vago DR, Silbersweig DA. Self-awareness, self-regulation, and self-transcendence (S-ART): A framework for understanding the neurobiological mechanisms of mindfulness. *Frontiers in Human Neuroscience*. 2012; 6:296.10.3389/fnhum.2012.00296 [PubMed: 23112770]
- Vahey DC, Aiken LH, Sloane DM, Clarke SP, Vargas D. Nurse burnout and patient satisfaction. *Medical Care*. 2004; 42(Suppl 2):II57–II66. [PubMed: 14734943]
- van Veen V, Carter CS. The timing of action-monitoring processes in the anterior cingulate cortex. *Journal of Cognitive Neuroscience*. 2002; 14:593–602. [PubMed: 12126500]
- Vidrine JI, Vidrine DJ, Costello TJ, Mazas C, Cofta-Woerpel L, Mejia LM, Wetter DW. The Smoking Consequences Questionnaire: Factor structure and predictive validity among Spanish-speaking Latino smokers in the United States. *Nicotine & Tobacco Research*. 2009; 11:1280–1288. [PubMed: 19696309]
- West CP, Huschka MM, Novotny PJ, Sloan JA, Kolars JC, Habermann TM, Shanafelt TD. Association of perceived medical errors with resident distress and empathy: A prospective



longitudinal study. *Journal of the American Medical Association*. 2006; 296:1071–1078. [PubMed: 16954486]

West CP, Tan AD, Habermann TM, Sloan JA, Shanafelt TD. Association of resident fatigue and distress with perceived medical errors. *Journal of the American Medical Association*. 2009; 302:1294–1300. [PubMed: 19773564]

White L. Mindfulness in nursing: An evolutionary concept analysis. *Journal of Advanced Nursing*. 2014; 70:282–294. [PubMed: 23772683]

Winbush NY, Gross CR, Kreitzer MJ. The effects of mindfulness-based stress reduction on sleep disturbance: A systematic review. *Explore*. 2007; 3:585–591. [PubMed: 18005910]

World Health Organization. 2013 fact sheet diabetes. n.d.-a. Retrieved from <http://www.who.int/mediacentre/factsheets/fs312/en/>

World Health Organization. 2013 tobacco fact sheet. n.d.-b. Retrieved from <http://www.who.int/mediacentre/factsheets/fs339/en/>

World Health Organization. Fact sheet 2004: Obesity and overweight. n.d.-c. Retrieved from <http://www.who.int/mediacentre/factsheets/fs311/en/>

World Health Organization. *The World Health Report 2002: Reducing risks, promoting healthy life*. Geneva, Switzerland: Author; 2002. Retrieved from <http://www.who.int/whr/2002/en/>

Zeidan F, Gordon NS, Merchant J, Goolkasian P. The effects of brief meditation training on experimentally induced pain perception. *Journal of Pain*. 2009; 11:199–209. [PubMed: 19853530]

Zernicke KA, Campbell TS, Specia M, McCabe-Ruff K, Flowers S, Carlson LE. A randomized wait-list controlled trial of feasibility and efficacy of an online mindfulness-based cancer recovery program for underserved distressed cancer survivors: The eTherapy for cancer applying mindfulness trial. *Psychosomatic Medicine*. 2013; 76:257–267. [PubMed: 24804884]

## Biographies

**Hants Williams**, RN, BSN is currently a 4th year PhD student at Duke University School of Nursing, with research interests in non-pharmacological therapies for persons with acute and chronic pain.

**Leigh Ann Simmons**, PhD, MFT is a Associate Professor in the School of Nursing at Duke University, with research and public policy interests in women’s health disparities, and personalized approaches to health care in underserved populations.

**Paula Tanabe**, PhD, RN is a Associate Professor in the Schools of Nursing and Medicine at Duke University, with research interests in improving pain management in emergency departments, with a strong focus on patients with Sickle Cell Disease.