

ARTICLE

Targeted next-generation sequencing in steroid-resistant nephrotic syndrome: mutations in multiple glomerular genes may influence disease severity

Gemma Bullich^{1,2}, Daniel Trujillano^{3,4,5,6}, Sheila Santín¹, Stephan Ossowski^{4,7}, Santiago Mendizábal⁸, Gloria Fraga⁹, Álvaro Madrid¹⁰, Gema Ariceta¹⁰, José Ballarín², Roser Torra², Xavier Estivill^{3,4,5,6} and Elisabet Ars^{*,1,2}

Genetic diagnosis of steroid-resistant nephrotic syndrome (SRNS) using Sanger sequencing is complicated by the high genetic heterogeneity and phenotypic variability of this disease. We aimed to improve the genetic diagnosis of SRNS by simultaneously sequencing 26 glomerular genes using massive parallel sequencing and to study whether mutations in multiple genes increase disease severity. High-throughput mutation analysis was performed in 50 SRNS and/or focal segmental glomerulosclerosis (FSGS) patients, a validation cohort of 25 patients with known pathogenic mutations, and a discovery cohort of 25 uncharacterized patients with probable genetic etiology. In the validation cohort, we identified the 42 previously known pathogenic mutations across *NPHS1*, *NPHS2*, *WT1*, *TRPC6*, and *INF2* genes. In the discovery cohort, disease-causing mutations in SRNS/FSGS genes were found in nine patients. We detected three patients with mutations in an SRNS/FSGS gene and *COL4A3*. Two of them were familial cases and presented a more severe phenotype than family members with mutation in only one gene. In conclusion, our results show that massive parallel sequencing is feasible and robust for genetic diagnosis of SRNS/FSGS. Our results indicate that patients carrying mutations in an SRNS/FSGS gene and also in *COL4A3* gene have increased disease severity.

European Journal of Human Genetics (2015) **23**, 1192–1199; doi:10.1038/ejhg.2014.252; published online 19 November 2014

INTRODUCTION

Nephrotic syndrome (NS) is characterized by heavy proteinuria, hypoalbuminemia, edema, and dyslipidemia. Although most patients are steroid-sensitive NS (SSNS), about 20% of children and 40% of adults are steroid-resistant NS (SRNS) and progress to end-stage renal disease (ESRD). In these cases, renal histology typically shows focal segmental glomerulosclerosis (FSGS).^{1–3}

Inherited structural defects in the glomerular filtration barrier proteins are responsible for a significant proportion of SRNS.^{4,5} Patients with SRNS of genetic origin have poor renal survival but low rate of disease recurrence after renal transplantation.⁶ Genetic forms of SRNS can be inherited as an autosomal recessive (AR) or autosomal dominant (AD) condition and can be isolated or syndromic.⁵ Mutations in nephrin (*NPHS1*)⁷ and podocin (*NPHS2*),⁸ with an AR inheritance, are the major cause of congenital and childhood onset NS, respectively. However, mutations in other genes have also been reported.^{5,9} Mutations in inverted formin-2 (*INF2*),¹⁰ transient receptor potential channel 6 (*TRPC6*),¹¹ and rarely, in α -actinin-4 (*ACTN4*)¹² and CD2-associated protein (*CD2AP*)¹³ genes

cause juvenile or adult onset FSGS with AD inheritance. In rare cases, recessive mutations in *NPHS2* are associated with adult onset FSGS.¹⁴ *De novo* heterozygous mutations in exons 8 and 9 of Wilms tumor (*WT1*) gene can cause both syndromic¹⁵ and isolated childhood onset SRNS.¹⁶ The study of the relative frequency of mutations in the most commonly altered genes in patients with SRNS and/or FSGS allowed the development of genetic testing algorithms based on age at onset, family history, or renal histology.^{17–20} However, the genetic heterogeneity and significant phenotypic variability of SRNS make genetic testing using standard Sanger methods costly and time consuming, even if the analysis is restricted to the most frequently mutated genes.

Massive parallel next-generation sequencing (NGS) technology has dramatically increased the throughput and reduced the cost per nucleotide sequenced compared with traditional Sanger methods, enabling cost-effective sequencing of multiple genes simultaneously. Over the past 3 years, whole-exome sequencing has revealed new genes associated with SRNS in a few cases, expanding the genetic heterogeneity of the disease.^{21–25} Based on this scenario, targeted NGS

¹Molecular Biology Laboratory, Fundació Puigvert, Instituto de Investigaciones Biomédicas Sant Pau (IIB-Sant Pau), Universitat Autònoma de Barcelona, REDinREN, Instituto de Investigación Carlos III, Barcelona, Catalonia, Spain; ²Nephrology Department, Fundació Puigvert, Instituto de Investigaciones Biomédicas Sant Pau (IIB-Sant Pau), Universitat Autònoma de Barcelona, REDinREN, Instituto de Investigación Carlos III, Barcelona, Catalonia, Spain; ³Genomics and Disease Group, Bioinformatics and Genomics Programme, Centre for Genomic Regulation (CRG), Barcelona, Catalonia, Spain; ⁴Department of Experimental and Health Sciences, Universitat Pompeu Fabra, Barcelona, Catalonia, Spain; ⁵Hospital del Mar Medical Research Institute (IMIM), Barcelona, Catalonia, Spain; ⁶CIBER in Epidemiology and Public Health (CIBERESP), Barcelona, Catalonia, Spain; ⁷Genomic and Epigenomic Variation in Disease Group, Centre for Genomic Regulation (CRG), Barcelona, Catalonia, Spain; ⁸Pediatric Nephrology Department, Hospital Universitario La Fe, Valencia, Spain; ⁹Pediatric Nephrology Department, Hospital de la Santa Creu i Sant Pau, Barcelona, Catalonia, Spain; ¹⁰Pediatric Nephrology Department, Hospital Vall d'Hebron, Barcelona, Spain

*Correspondence: Dr E Ars, Molecular Biology Laboratory, Fundació Puigvert, Instituto de Investigaciones Biomédicas Sant Pau (IIB-Sant Pau), Universitat Autònoma de Barcelona, REDinREN, Instituto de Investigación Carlos III, Cartagena 340-350, Barcelona, Catalonia 08025, Spain. Tel: +34 93 416 97 00; Fax: +34 93 416 97 30; E-mail: ears@fundacio-puigvert.es

Received 27 May 2014; revised 14 October 2014; accepted 16 October 2014; published online 19 November 2014

of a broad panel of NS-related genes has emerged as a cost-effective strategy to screen the multiple genes involved in SRNS/FSGS,²⁶ but optimal sensitivity and specificity must be demonstrated for each gene in the panel.

In this study, we used targeted NGS to simultaneously sequence 26 genes associated with inherited glomerular diseases in a heterogeneous cohort of 50 SRNS/FSGS patients and 5 control individuals. We aimed to develop a glomerular disease gene panel for SRNS/FSGS and to study the influence of mutations in multiple genes on phenotype variability.

MATERIALS AND METHODS

Patients

A total of 50 Spanish patients with idiopathic SRNS/FSGS were included. Patients developing steroid resistance at a later stage of the disease or with recurrence after kidney transplantation were excluded as we considered that they likely had an immunological cause. Biopsy findings included FSGS, minimal change disease (MCD) or diffuse mesangial sclerosis. The validation cohort consisted of 25 patients with known pathogenic mutations in the five most commonly mutated SRNS/FSGS genes that had been previously identified by Sanger sequencing.¹⁸ The discovery cohort consisted of 25 patients with diagnosis of SRNS/FSGS, 21 genetically uncharacterized, and 4 incompletely characterized. All 25 had a probable genetic etiology, based on early onset of the disease ($n=10$), familial history of SRNS/FSGS ($n=11$), or consanguinity ($n=4$). Four of these patients had been analyzed by Sanger sequencing for the most frequently mutated SRNS/FSGS genes in our previous study, and only one recessive pathogenic mutation was identified.¹⁸ We also included five control individuals without nephropathy who had been previously genome-wide genotyped with a HumanOmni 2.5–8 BeadChip (Illumina, Inc., San Diego, CA, USA) to test the performance of the assay across the whole panel. Blood samples were obtained from other family members if they were available. All the samples were codified, and data analysis was performed blindly. The study was approved by the Institutional Review Board, and all participants gave their signed informed consent.

Sequencing and data analyses

We selected 26 genes associated with hereditary glomerular diseases based on published literature (Table 1). The complete genomic sequence (plus 1 kb of 5' and 3' flanking genomic regions) of *NPHS1*, *NPHS2*, *WT1*, *TRPC6*, *INF2*, *LAMB2*, *COLAA3*, *COLAA4*, *COLAA5*, and *GLA* genes and all exons and intron boundaries (plus 100 bp at each end) of the remaining genes were captured using a custom NimbleGen SeqCap EZ Choice Library (Roche NimbleGen, Madison, WI, USA). After removal of repetitive sequences, 83.6% of the targeted bases were covered with capture baits ranging from 68 to 6689 bp (average 1062 bp), for a final targeted region of 0.9 Mb.

Genomic DNA was isolated from peripheral blood using the salting-out method. Libraries were prepared with the TruSeq DNA Sample Preparation Kit (Illumina, Inc.) according to the manufacturer's instructions. In familial cases, only the proband was analyzed by NGS. Pools of 24 individuals were prepared, hybridized to the custom NimbleGen SeqCap EZ Choice Library (Roche NimbleGen) for 72 h, stringently washed, amplified 17 PCR cycles, and run in a HiSeq2000 instrument (Illumina Inc.).

Data analysis was performed blindly with an in-house developed pipeline previously described.²⁷ All candidate variants were required on both sequenced DNA strands and to account for $\geq 20\%$ of total reads at that site. Common polymorphisms ($\geq 5\%$ in the general population) were discarded by comparison with dbSNP 138, the 1000G (<http://www.1000genomes.org>), the Exome Variant Server (<http://evs.gs.washington.edu>), and an in-house exome variant database to filter out both common benign variants and recurrent artifact variant calls. To identify large structural variants, we used Pindel,²⁸ Conifer,²⁹ and PeSV-Fisher (<http://gd.crg.eu/tools>).

Evaluation of the pathogenicity of the variants

Nonsense, frameshift, and canonical splice site variants were classified as definitely pathogenic mutations (mutation group (MG)=A). Missense variants

were considered *a priori* unclassified sequence variants (UCV), and their potential pathogenicity was evaluated using an *in silico* scoring system developed for the *PKD1* and *PKD2* genes.³⁰ This scoring system with some minor modifications was tested using previously described pathogenic mutations, for which functional studies had been performed, as positive controls, and known neutral variants or polymorphisms as negative controls.^{31–33} This scoring system takes into consideration the biophysical and biochemical difference between wild type and mutant amino acid, the evolutionary conservation of the amino-acid residue in orthologs,³⁴ a number of *in silico* predictors (Sift, Polyphen, Mutation taster, and Condel), and population data. All candidate pathogenic variants not previously identified were validated by conventional PCR amplification and Sanger sequencing and were not detected in 284 control chromosomes. Segregation of these changes with the disease was assessed for all the available family members. We scored each of these factors, and their sum resulted in an overall variant score (VS). The UCV were classified into four MGs: highly likely pathogenic ($VS \geq 11$, MG=B), likely pathogenic ($5 \leq VS \leq 10$, MG=C), indeterminate ($0 \leq VS \leq 4$, MG=I), and highly likely neutral ($VS \leq -1$, MG=N). To evaluate the pathogenicity of non-canonical splice site variants, RNA analysis was performed by RT-PCR and Sanger sequencing. If no RNA was available, these variants were analyzed using Alamut version 2.3 (Interactive Biosoftware, Rouen, France), a software package that uses different splice site prediction programs to compare the normal and variant sequences for differences in potential regulatory signals.³⁵

We designated pathogenic mutations to be: (i) those sequence variants predicted to result in a truncated protein (MG=A), (ii) canonical and non-canonical splice site variants showed to alter splicing patterns (MG=A), and (iii) those amino-acid substitutions expected to severely alter the protein sequence using *in silico* predictors (MG=B). Missense substitutions classified as MG=C or MG=I were considered as mild mutations in *NPHS1*³² or variants of unknown clinical significance. All the variants were entered in the Leiden Open Variation Database (<http://databases.lovd.nl/shared/genes>).

RESULTS

Validation of the technology

Sequencing of the 26 glomerular disease gene panel (Table 1) in 50 patients with SRNS/FSGS and 5 control individuals generated a mean of 14.3 million reads per patient. On average, 99.1% of these reads mapped to the reference genome. A mean depth of coverage of $466 \times$ was achieved for the 26 targeted genes across all individuals, with 99.6% of targeted bases covered by at least 20 reads (Supplementary Table S1).

The validation cohort included 25 SRNS/FSGS patients who carried a total of 42 known pathogenic mutations in *NPHS1*, *NPHS2*, *WT1*, *TRPC6*, or *INF2* genes and with different phenotypic characteristics (Table 2). We identified all known pathogenic mutations (33 different) in their correct heterozygous/homozygous state, specifically: 22 missense, 3 nonsense, 2 splice site, 4 small deletions, 1 small insertion, and 1 deletion/insertion (Indel) (data not shown). No spurious pathogenic mutations were found in any of these samples. Prior Sanger sequencing of these patients had revealed a total of 285 variants in these genes, 281 of which were also detected by NGS, resulting in 98.6% accuracy.

To assess the sensitivity and specificity of our assay across all 26 genes included in the panel, we evaluated 5 control individuals without nephropathy who had been previously genome-wide genotyped. Sensitivity of detecting homozygous and heterozygous polymorphisms across the 26 genes was 95.6% (1315/1375), and specificity of detecting non-variant sites from the reference genome was 99.9% (3387/3391). No spurious pathogenic mutations were found in any of these samples. Detailed quality control parameters are provided in Supplementary Table S2.

Sequence variants in NS genes in the discovery cohort

We identified disease-causing mutations in NS genes in 9 out of the 25 SRNS/FSGS patients in the discovery cohort (Table 3).

Table 1 Panel of genes involved in inherited glomerular diseases

Gene	Disease association	Inheritance	Target	Accession no.	Chromosome
<i>NPHS1</i>	CNS, SRNS	AR	Whole gene	NM_004646.2	19
<i>NPHS2</i>	CNS, SRNS	AR	Whole gene	NM_014625.2	1
<i>WT1</i>	SRNS, Denys–Drash syndrome	AD	Whole gene	NM_000378.4	11
<i>INF2</i>	SRNS, FSGS	AD	Whole gene	NM_001031714.3	14
<i>TRPC6</i>	SRNS, FSGS	AD	Whole gene	NM_004621.5	11
<i>LAMB2</i>	SRNS, Pierson syndrome	AR	Whole gene	NM_002292.3	3
<i>COL4A5</i>	Collagen type IV nephropathy	XL	Whole gene	NM_000495.4	X
<i>COL4A3</i>	Collagen type IV nephropathy	AD/AR	Whole gene	NM_000091.4	2
<i>COL4A4</i>	Collagen type IV nephropathy	AD/AR	Whole gene	NM_000092.4	2
<i>GLA</i>	Fabry disease	XL	Whole gene	NM_000169.2	X
<i>PLCE1</i>	CNS, SRNS	AR	Exons	NM_016341.3	10
<i>ACTN4</i>	SRNS, FSGS	AD	Exons	NM_004924.4	19
<i>CD2AP</i>	SRNS	AD/AR	Exons	NM_012120.2	6
<i>MYO1E</i>	SRNS	AR	Exons	NM_004998.3	15
<i>ARHGAP24</i>	NS, FSGS	AD	Exons	NM_001025616.2	4
<i>CUBN</i>	NS	AR	Exons	NM_001081.3	10
<i>CFH</i>	NS	AR	Exons	NM_000186.3	1
<i>COQ2</i>	NS	AR	Exons	NM_015697.7	4
<i>COQ6</i>	NS	AR	Exons	NM_182476.2	14
<i>ITGA3</i>	NS	AR	Exons	NM_002204.2	17
<i>LMX1B</i>	NS, FSGS	AR	Exons	NM_001174146.1	9
<i>NEIL1</i>	NS	AR	Exons	NM_001256552.1	15
<i>PDSS2</i>	NS	AR	Exons	NM_020381.3	6
<i>PTPRO</i>	SRNS	AR	Exons	NM_030667.2	12
<i>SCARB2</i>	NS	AR	Exons	NM_005506.3	4
<i>SMARCAL1</i>	NS	AR	Exons	NM_001127207.1	2

Abbreviations: AD autosomal dominant; AR, autosomal recessive; CNS, congenital nephrotic syndrome; FSGS, focal segmental glomerulosclerosis; NS, nephrotic syndrome; SRNS, steroid-resistant nephrotic syndrome; XL, X-linked.

Table 2 Overview of genotypic data obtained by next-generation sequencing

	Total	Familial	Sporadic	Congenital onset	Early or late childhood onset	Adolescent or adult onset
<i>Validation cohort</i>	25	10	15	10	9	6
Patients with pathogenic mutations in an SRNS/FSGS gene	23	9	14	9	8	6
Patients with mutations in an SRNS/FSGS gene and <i>COL4A3</i>	2	1	1	1	1	0
Patients with no pathogenic mutations found	0	0	0	0	0	0
<i>Discovery cohort</i>	25	15	10	5	12	8
Patients with pathogenic mutations in an SRNS/FSGS gene	9	4	5	5	2	2
Patients with mutations in an SRNS/FSGS gene and <i>COL4A3</i>	1	1	0	0	0	1
Patients with no pathogenic mutations found	15	10	5	0	10	5

Abbreviations: FSGS, focal segmental glomerulosclerosis; SRNS, steroid-resistant nephrotic syndrome. Onset was classified as follows: congenital, 0–3 months; early childhood, 4 months to 5 years; late childhood, 6–12 years; adolescent, 13–18 years; adult, > 18 years.

The distribution of mutations in SRNS/FSGS genes differed depending on the age at onset. The mutation detection rate decreased as the age at onset of NS increased. In congenital onset patients (from 0 to 3 months), all the five patients (100%) carried mutations in *NPHS1* ($n=3$) and *NPHS2* ($n=2$) genes. In the early-childhood onset cohort (from 4 months to 5 years), two out of the nine patients (22%) had mutations in *NPHS1* ($n=1$) and *WT1* ($n=1$). No disease-causing mutations were found in any of the three patients with late-childhood onset NS (from 6 to 12 years). In patients with adult onset of NS or FSGS (> 18 years), two out of the eight patients (25%) carried mutations in *INF2* ($n=1$) and *TRPC6* ($n=1$) (Table 2). A detailed scoring matrix for the missense variants is provided in Supplementary Table S3.

In the discovery cohort, we included four cases (one familial and three sporadic), with only one recessive pathogenic mutation previously identified by Sanger sequencing. The NGS approach detected variants predicted to alter the non-canonical splice site sequences by the Alamut software but with uncertain clinical significance in three patients.

Phenotypic effect of mutations in multiple glomerular genes

We found four patients belonging to the validation cohort with three mutated alleles in two recessive SRNS/FSGS genes (Supplementary Table S4). Phenotype modification of the third mutated allele could not be assessed in these patients as three of them were sporadic cases,

Table 3 Clinical and genetic data of patients in the discovery cohort with disease-causing mutations in an SRNS/FSGS gene and COL4A3

Patient	Gender	Familial/sporadic	Age at onset (years)	Features at presentation	Renal biopsy	Immunosuppressive therapy	Evolution	Gene	Mutation 1 (MG)	Mutation 2 (MG)	Gene	Mutation (MG)
<i>Patients in the discovery cohort with disease-causing mutations in an SRNS/FSGS gene</i>												
319	M	Sp	0	CNS	NP	—	Dead at 1 year	<i>NPHS1</i>	c.468C>G p.(V156*) (A)	c.3478C>T p.(R1160*) (A)	<i>COL4A3</i>	c.3829G>A p.(G1277S) (B)
336	M	Sp	0	CNS	CNF	—	ESRD at 2 months	<i>NPHS1</i>	c.1655C>A p.(A552D) (B)	c.1655C>A p.(A552D) (B)	<i>COL4A3</i>	c.4504T>C p.(F1502L) (C)
299	F	Sp	0.1	CNS	NP	—	ESRD at 8 months	<i>NPHS1</i>	c.3250dup p.(V1084Gfs*12) (A)	c.3250dup p.(V1084Gfs*12) (A)	<i>COL4A3</i>	c.4028-3C>A p.(V1344_G1385del) (A)
324	M	Sp	0.3	NS without edema	DMS	Cs, CP, CsA, MMF	Normal Cr at 19 years	<i>NPHS1</i>	c.1930+5G>A p.(V634Tfs*13) (A)	c.1930+5G>A p.(V634Tfs*13) (A)	<i>COL4A3</i>	c.4028-3C>A p.(V1344_G1385del) (A)
363	F	Sp	0	CNS	NP	—	Cr 0.37 mg/dl at 1 month	<i>NPHS2</i>	c.413G>A p.(R138P) (B)	c.413G>A p.(R138P) (B)	<i>COL4A3</i>	c.4028-3C>A p.(V1344_G1385del) (A)
330	F	Fam ^a	0.2	Nephrotic proteinuria, MAL	DMS	—	Normal Cr at 4 months	<i>NPHS2</i>	c.320T>C p.(L107P) (B)	c.320T>C p.(L107P) (B)	<i>COL4A3</i>	c.4028-3C>A p.(V1344_G1385del) (A)
320	F	Sp	3	Denys-Drash syndrome	FSGS	—	ESRD at 4 years	<i>WT1</i>	c.1419T>A p.(H473Q) (B)	—	<i>COL4A3</i>	—
347-1	M	Fam ^b	19	Non-nephrotic proteinuria	FSGS	—	CKD stage II at 20 years	<i>INF2</i>	c.658G>A p.(E220K) (B)	—	<i>COL4A3</i>	—
347-2	F	Fam ^c	20	Nephrotic proteinuria	FSGS	Cs, CsA	ESRD at 29 years	<i>INF2</i>	p.(E220K) (B)	—	<i>COL4A3</i>	—
384-1	F	Fam ^b	27	Non-nephrotic proteinuria, MH, MAL	NP	—	Normal Cr at 32 years	<i>TRPC6</i>	c.2656G>A p.(E886K) (B)	—	<i>COL4A3</i>	c.4028-3C>A p.(V1344_G1385del) (A)
384-2	M	Fam ^d	30	MAL	NP	—	Normal Cr at 35 years	<i>TRPC6</i>	c.2656G>A p.(E886K) (B)	—	<i>COL4A3</i>	c.4028-3C>A p.(V1344_G1385del) (A)
384-3	F	Fam ^e	55	Non-nephrotic proteinuria, MH	NP	—	Normal Cr at 63 years	<i>TRPC6</i>	c.2656G>A p.(E886K) (B)	—	<i>COL4A3</i>	c.4028-3C>A p.(V1344_G1385del) (A)
<i>Patients with mutations in an SRNS/FSGS gene and COL4A3</i>												
266 ^f	F	Sp	0	CNS, MH	NP	—	ESRD at 1 year	<i>NPHS1</i> [†]	c.514_516del p.(T172del)	c.3250dup p.(V1084Gfs*12) (A)	<i>COL4A3</i>	c.3829G>A p.(G1277S) (B)
10-1 ^e	M	Fam ^b	4	NS, MH	FSGS	Cs, CsA	ESRD at 12 years	<i>NPHS2</i> [‡]	c.274G>T p.(G92C) (B)	c.506T>C p.(L169P) (B)	<i>COL4A3</i>	c.4504T>C p.(F1502L) (C)
10-2	F	Fam ^g	2	NS	MCD	Cs, CsA ±	Normal Cr at 18 years	<i>NPHS2</i> [‡]	c.274G>T p.(G92C) (B)	c.506T>C p.(L169P) (B)	<i>COL4A3</i>	—
253-1	F	Fam ^b	32	NS, MH	FSGS*	Cs, CsA	ESRD at 33 years	<i>INF2</i>	c.2065C>T p.(R689W) (I)	—	<i>COL4A3</i>	c.4028-3C>A p.(V1344_G1385del) (A)
253-2	M	Fam ^h	39	Non-nephrotic proteinuria, MH	FSGS	—	ESRD at 51 years	<i>INF2</i>	—	—	<i>COL4A3</i>	c.4028-3C>A p.(V1344_G1385del) (A)
253-3	M	Fam ⁱ	—	—	NP	—	Normal Cr at 61 years	<i>INF2</i>	—	—	<i>COL4A3</i>	c.4028-3C>A p.(V1344_G1385del) (A)
253-4	F	Fam ^j	U	MH	NP	—	Normal Cr at 52 years	<i>INF2</i>	—	—	<i>COL4A3</i>	c.4028-3C>A p.(V1344_G1385del) (A)
253-5	F	Fam ^k	—	—	NP	—	Normal Cr at 63 years	<i>INF2</i>	c.2065C>T p.(R689W) (I)	—	<i>COL4A3</i>	—

Abbreviations: CKD, chronic kidney disease; CNF, congenital nephrotic syndrome of Finnish type; CNS, congenital nephrotic syndrome; CP, cyclophosphamide; Cr, creatinine; Cs, corticosteroids; CSA, cyclosporin A; DMS, diffuse mesangial sclerosis; ESRD, end-stage renal disease; F, female; Fam, familial case; FSGS, focal segmental glomerulosclerosis; FSGS*, mesangio proliferative lesions with FSGS; M, male; MAL, microalbuminuria; MCD, minimal change disease; MG, mutation group; MH, microhematuria; MMF, mycophenolate mofetil; NP, not performed; NS, nephrotic syndrome; Sp, sporadic case; U, unknown.
 †Patients of the validation cohort.
 ‡Mutations in these genes were previously known in Sanger sequencing.
 §Proband's sister.
 ¶Proband's father.
 ††Proband's uncle.
 ‡‡Proband's aunt.
 §§Only child of consanguineous parents.
 ¶¶Proband.
 ¶¶¶Proband's mother.
 ¶¶¶¶Proband's brother.
 ¶¶¶¶¶Patients of the validation cohort.
 ¶¶¶¶¶¶Mutations in these genes were previously known in Sanger sequencing.
 ¶¶¶¶¶¶¶Proband's sister.
 ¶¶¶¶¶¶¶¶Proband's father.
 ¶¶¶¶¶¶¶¶¶Proband's uncle.
 ¶¶¶¶¶¶¶¶¶¶Proband's aunt.

and only two siblings, both carrying the three mutated alleles, were identified.

We identified three patients carrying mutations in an SRNS/FSGS gene and also in *COL4A3* (Table 3). Patient 266 carried two *NPHS1* pathogenic mutations, an in-frame deletion and a frameshift, together with a heterozygous missense mutation in *COL4A3*, previously reported by Heidet *et al.*³⁶ She had a congenital NS presenting with microhematuria and no family history of NS. Patient 10-1 and his affected sister (10-2) both carried compound heterozygous missense pathogenic mutations in *NPHS2* gene, but only the proband 10-1 harbored a heterozygous missense variant in *COL4A3* predicted to be likely pathogenic. Both siblings had early childhood onset of SRNS. Patient 10-1 presented with nephrotic range proteinuria and microhematuria. His renal biopsy revealed FSGS, and he developed ESRD at 12 years. His sibling 10-2 presented with borderline nephrotic range proteinuria but no evidence of microhematuria, renal biopsy showed MCD and she presented normal renal function by the age of 18 years (Figure 1a). Patient 253-1 carried a heterozygous splicing mutation in *COL4A3*, demonstrated to produce exon 46 skipping by RNA analysis and predicted to result in a protein lacking 42 amino acids, in combination with a missense variant in the exon 12 of *INF2*. This novel non-conservative substitution, p.R689W, is located at a highly conservative domain (FH2) in the *INF2* protein and scored as highly likely pathogenic, using mutation prediction programs. The arginine in the position 689 is totally conserved in mammals and a basic amino acid in all the species. She presented with SRNS and microhematuria at 32 years, and her renal biopsy showed mesangioproliferative lesions with FSGS. Her renal function rapidly deteriorated, reaching ESRD at 33 years. The *COL4A3* mutation was inherited from her affected father (253-2) who presented with non-nephrotic range proteinuria and hematuria at 39 years. His renal biopsy showed FSGS, and he reached ESRD at 51 years. The *INF2* variant was inherited from her asymptomatic mother (253-5). Two of the proband's uncles carried the *COL4A3* mutation, but they only presented microhematuria at 61 (253-3) and 56 years (253-4) (Figure 1b).

DISCUSSION

In this study, we show that the simultaneous analysis of 26 genes causative of inherited glomerular diseases allows a more complete and efficient characterization of patients with SRNS/FSGS than traditional Sanger sequencing. In addition, we identified three patients carrying combined mutations in an SRNS/FSGS gene and *COL4A3*, suggesting that mutations in different genes that converge in the glomerular filtration barrier influence disease severity.

In the past years, several genetic testing algorithms for SRNS/FSGS have been developed to help in establishing a prioritization of the genes to be sequenced by Sanger. However, the genetic heterogeneity and phenotypic variability of this disease make this approach expensive and time consuming.^{17–20} Recently, two studies used NGS technology to analyze the exons and intron boundaries of 24 genes²⁶ and 21 genes³⁷ associated with SRNS. Our gene panel included not only genes related with SRNS/FSGS but also genes involved in other glomerular diseases, as we hypothesized that disease severity could be influenced by mutations in multiple glomerular genes. The identification of all previously known pathogenic mutations and no spurious pathogenic mutations in our validation cohort, as well as the high sensitivity and specificity obtained with the analysis of the previously genotyped controls, demonstrate the suitability of this approach for genetic diagnosis of SRNS/FSGS.

In the discovery cohort, we identified disease-causing mutations in NS genes in 9 out of the 25 patients. All patients carried pathogenic

mutations in the most likely mutated NS gene according to their age at disease onset.¹⁸ Interestingly, patient 324 had a congenital onset of the disease but still normal renal function at the age of 19 years. He carried a homozygous splicing mutation (c.1930+5G>A) in *NPHS1* found to produce the deletion of the 31 last nucleotides of exon 14 in the mRNA, which is predicted to result in a truncated protein. The mild phenotype of this patient could be explained, because splicing mutations that do not affect the canonical GT/AG splice sites could allow the coexistence of a certain proportion of wild-type *NPHS1* mRNA with the altered mRNA, as previously suggested.³⁸ Although mRNA analysis from patient's blood did not confirm this hypothesis, we cannot discard the occurrence of this phenomenon in kidneys (Supplementary Figure S1).

We also included four patients with only one recessive candidate pathogenic mutation in an SRNS gene identified by Sanger sequencing. We hypothesized that these patients would carry a large insertion or deletion or a deep intronic splicing mutation as a second pathogenic mutation. Thus we included the whole genomic sequence of the most frequently mutated genes in glomerular diseases in our NGS gene panel and analyzed the data using specific algorithms to search for structural variants. No clear pathogenic mutation was detected, but only variants in non-canonical splice sites were found in three patients. However, RNA from these patients was not available, and the pathogenicity of these variants could not be assessed.

The phenotypic variability observed in SRNS/FSGS patients bearing mutations in the same gene suggests that modifier genes and environmental factors may have a significant role in the renal presentation and outcome.⁴ Evidence of oligogenic inheritance with mutations in genes encoding proteins that converge in common pathomechanistic pathways has been reported in Bardet–Biedl syndrome.³⁹ In addition, the p.R229Q variant in *NPHS2* gene has been suggested to contribute to proteinuria and ESRD in thin basement membrane nephropathy.^{40,41} Recently, modifier genes have been proposed to explain early and severe polycystic kidney disease.⁴² McCarthy *et al.*²⁶ described two patients carrying a homozygous mutation in *NPHS1* and a possibly pathogenic variant in *WT1*, who developed a more aggressive disease than a third patient carrying the same mutation in *NPHS1* but without the *WT1* variant. To study the putative role of mutations in multiple glomerular genes on SRNS/FSGS clinical variability, disease severity should ideally be compared among various family members with different genotype combinations. Here, four patients carrying three mutated alleles in two SRNS/FSGS genes were found. Unfortunately, three of them were sporadic cases, and only two affected siblings—both carrying the three mutated alleles—were identified. Therefore, the putative effect of the third variant on disease severity could not be assessed.

We identified three patients carrying mutations in an SRNS/FSGS gene in combination with a heterozygous mutation in *COL4A3* gene. Heterozygous mutations in *COL4A3* and *COL4A4* genes cause the mildest phenotype of collagen type IV ($\alpha3\alpha4$) nephropathy, also named thin basement membrane nephropathy. This nephropathy is characterized by hematuria and low proteinuria,^{43,44} and progression to ESRD has recently been described in 30% of cases.⁴⁵ The clinical phenotype of the three patients with combined mutations in an SRNS/FSGS gene and *COL4A3* stands out for the coexistence of NS and microhematuria at presentation. Interestingly, in two of these three cases, several family members with different genotype combinations were available (Figure 1). In both families, patients with mutations in an SRNS/FSGS gene and *COL4A3* had a more severe phenotype than their family members carrying mutations in only one gene. Variable disease penetrance in *INF2*-mutated patients has been reported⁴⁶ likely




Figure 1 Pedigrees of two families with mutations in an SRNS/FSGS gene and *COL4A3*. (a) In family 10, both siblings had compound heterozygous pathogenic mutations in *NPHS2* gene and the more severely affected individual (10-1) carried an additional likely pathogenic variant in *COL4A3* gene. (b) In family 253, individuals 253-1 to -4 carried a pathogenic mutation in *COL4A3* gene demonstrated to produce exon 46 skipping by reverse transcriptase-PCR and Sanger sequencing and predicted to result in a protein lacking 42 amino acids. Patient 253-1 carried an additional variant in *INF2* gene inherited from her mother and developed a more aggressive phenotype than the other affected family members. Cr, creatinine; wt, wild type. The arrows indicate probands. Squares denote males, circles denote females. Filled symbols indicate affected status. Quarter solid symbols indicate microhematuria.

explaining that, in family 253, the proband's mother (253-5) remained asymptomatic. These findings suggest that mutations in multiple glomerular disease genes explain some of the phenotypic variability in nephropathies. Another possible explanation for clinical intrafamilial variability could arise in families carrying a splicing mutation that

does not affect the canonical splice sites, such as the mutation in *COL4A3* gene detected in family 253. This mutation could lead to variable amounts of the correctly spliced transcript and could explain the phenotypic variability among the three siblings carrying this splicing mutation.³⁸

Despite the broad panel of genes analyzed, we could not find pathogenic mutations in 15 of the patients in the discovery cohort, 8 of whom were familial cases. The fact that some SRNS/FSGS patients present with recurrence after kidney transplantation indicates that some of these cases may be due to an immunological cause, although no evidence of immunological bases was observed in our cohort. In the familial cases, it is highly likely that an SRNS/FSGS gene, as yet non-identified, is responsible for the disease. The next step should therefore be to sequence the whole exome in the 8 familial cases to identify new candidate genes.

The results obtained in the validation cohort demonstrate that our approach is suitable for genetic diagnosis of SRNS/FSGS but, based on the discovery cohort findings, we propose some modifications: (1) to sequence a gene panel with only the six most frequently mutated genes in SRNS/FSGS (*NPHS1*, *NPHS2*, *PLCE1*, *WT1*, *INF2*, *TRPC6*). The *COL4A3*, *COL4A4* and *COL4A5* genes, associated with collagen type IV ($\alpha3\alpha4$) nephropathy, could also be included as they may influence disease severity. If no pathogenic mutations are identified, a more extensive glomerular gene panel or exome sequencing could be performed; and (2) to restrict the targeted sequence to exons and intron boundaries as the assessment of the pathogenicity of deep intronic variants is challenging and their involvement in the disease speculative. In terms of the cost, NGS will allow the simultaneous analysis of around 250 exons for approximately the same cost of consumables than sequencing 40 exons by Sanger, with three times saving in hands-on time. Identifying pathogenic mutations in SRNS is important for many reasons. It can help to avoid the adverse effects of steroid therapy, modify the intensity and duration of immunosuppressive therapies, encourage living donor kidney transplantation, provide prognostic information regarding the gene and type of mutations, and enable genetic counseling. Sequencing a panel of genes involved in glomerular inherited diseases will also help to elucidate cases with atypical renal phenotypes and/or with high clinical intrafamilial variability. Based on our findings, such cases could be more prevalent than previously expected.

In conclusion, this study shows the feasibility and robustness of targeted NGS for genetic diagnosis of SRNS/FSGS, allowing a more complete characterization of patients with SRNS/FSGS. Our results indicate that patients carrying mutations in an SRNS/FSGS gene and also in *COL4A3* gene have increased disease severity.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We thank the patients for taking part in this study and the referring physicians who participated in this study, especially Dr Camino (Hospital San Agustín, Avilés). We thank Justo Gonzalez, Patricia Ruiz and Estefania Eugui for technical support; Cristian Tornador and Georgia Escaramis for bioinformatics assistance; Carolyn Newey for English language editing; and IIB Sant Pau-Fundació Puigvert Biobank for kindly providing some of the NS samples, ReTBioH (Spanish Biobank Network) RD09/76/00064. This project was funded by the Spanish Plan Nacional SAF2008-00357 (NOVADIS); the Instituto de Salud Carlos III (FIS/FEDER PI11/00733); the European Commission 7th Framework Program, Project N. 261123 (GEUVADIS) and Project N. 262055 (ESGI); the Spanish Healthy Ministry (FIS 12/01523 and FIS PI13/01731), ISCIII-RETIC REDinREN/RD06/0016 and RD012/0021 FEDER funds; the Catalan Government (AGAUR 2009/SGR-1116); and the Fundació Renal Iñigo Álvarez de Toledo (FRIAT) in Spain. Dr Roser Torra is supported by the Intensification Program of Research Activity ISCIII/Generalitat de Catalunya (program I3SN).

- Short versus standard prednisone therapy for initial treatment of idiopathic nephrotic syndrome in children. Arbeitsgemeinschaft für Padiatrische. *Lancet* 1988; **1**: 380–383.
- Troyanov S, Wall CA, Miller JA, Scholey JW, Cattran DC: Focal and segmental glomerulosclerosis: definition and relevance of a partial remission. *J Am Soc Nephrol* 2005; **16**: 1061–1068.
- Antignac C: Genetic models: clues for understanding the pathogenesis of idiopathic nephrotic syndrome. *J Clin Invest* 2002; **109**: 447–449.
- Benoit G, Machuca E, Heidet L, Antignac C: Hereditary kidney diseases: highlighting the importance of classical Mendelian phenotypes. *Ann NY Acad Sci* 2010; **1214**: 83–98.
- Gbadegesin RA, Winn MP, Smoyer WE: Genetic testing in nephrotic syndrome—challenges and opportunities. *Nat Rev Nephrol* 2013; **9**: 179–184.
- Conlon PJ, Lynn K, Winn MP *et al*: Spectrum of disease in familial focal and segmental glomerulosclerosis. *Kidney Int* 1999; **56**: 1863–1871.
- Keistilä M, Lenkkeri U, Männikkö M *et al*: Positionally cloned gene for a novel glomerular protein—nephrin—is mutated in congenital nephrotic syndrome. *Mol Cell* 1998; **1**: 575–582.
- Boute N, Gribouval O, Roselli S *et al*: NPHS2, encoding the glomerular protein podocin, is mutated in autosomal recessive steroid-resistant nephrotic syndrome. *Nat Genet* 2000; **24**: 349–354.
- Hinkes BG, Mucha B, Vlangos CN *et al*: Nephrotic syndrome in the first year of life: two thirds of cases are caused by mutations in 4 genes (NPHS1, NPHS2, WT1, and LAMB2). *Pediatrics* 2007; **119**: e907–e919.
- Brown EJ, Schlöndorff JS, Becker DJ *et al*: Mutations in the formin gene *INF2* cause focal segmental glomerulosclerosis. *Nat Genet* 2010; **42**: 72–76.
- Winn MP, Conlon PJ, Lynn KL *et al*: A mutation in the *TRPC6* cation channel causes familial focal segmental glomerulosclerosis. *Science* 2005; **308**: 1801–1804.
- Kaplan JM, Kim SH, North KN *et al*: Mutations in *ACTN4*, encoding alpha-actinin-4, cause familial focal segmental glomerulosclerosis. *Nat Genet* 2000; **24**: 251–256.
- Kim JM, Wu H, Green G *et al*: CD2-associated protein haploinsufficiency is linked to glomerular disease susceptibility. *Science* 2003; **300**: 1298–1300.
- Machuca E, Hummel A, Nevo F *et al*: Clinical and epidemiological assessment of steroid-resistant nephrotic syndrome associated with the NPHS2 R229Q variant. *Kidney Int* 2009; **75**: 727–735.
- Pelletier J, Bruening W, Li FP, Haber DA, Glaser T, Housman DE: WT1 mutations contribute to abnormal genital system development and hereditary Wilms' tumour. *Nature* 1991; **353**: 431–434.
- Jeanpierre C, Denamur E, Henry I *et al*: Identification of constitutional WT1 mutations, in patients with isolated diffuse mesangial sclerosis, and analysis of genotype/phenotype correlations by use of a computerized mutation database. *Am J Hum Genet* 1998; **62**: 824–833.
- Benoit G, Machuca E, Antignac C: Hereditary nephrotic syndrome: a systematic approach for genetic testing and a review of associated podocyte gene mutations. *Pediatr Nephrol* 2010; **25**: 1621–1632.
- Santín S, Bullich G, Tazón-Vega B *et al*: Clinical utility of genetic testing in children and adults with steroid-resistant nephrotic syndrome. *Clin J Am Soc Nephrol* 2011; **6**: 1139–1348.
- Rood IM, Deegens JK, Wetzels JF: Genetic causes of focal segmental glomerulosclerosis: implications for clinical practice. *Nephrol Dial Transplant* 2012; **27**: 882–890.
- Lipska BS, Iatropoulos P, Maranta R *et al*: Genetic screening in adolescents with steroid-resistant nephrotic syndrome. *Kidney Int* 2013; **84**: 206–213.
- Mele C, Iatropoulos P, Donadelli R *et al*: MYO1E mutations and childhood familial focal segmental glomerulosclerosis. *N Engl J Med* 2011; **365**: 295–306.
- Sanna-Cherchi S, Burgess KE, Nees SN *et al*: Exome sequencing identified MYO1E and NEIL1 as candidate genes for human autosomal recessive steroid-resistant nephrotic syndrome. *Kidney Int* 2011; **80**: 389–396.
- Ovunc B, Otto EA, Vega-Warner V *et al*: Exome sequencing reveals cubilin mutation as a single-gene cause of proteinuria. *J Am Soc Nephrol* 2011; **22**: 1815–1820.
- Gupta IR, Baldwin C, Auguste D *et al*: ARHGDI1: a novel gene implicated in nephrotic syndrome. *J Med Genet* 2013; **50**: 330–338.
- Boyer O, Woerner S, Yang F *et al*: LMX1B mutations cause hereditary FSGS without extrarenal involvement. *J Am Soc Nephrol* 2013; **24**: 1216–1222.
- McCarthy HJ, Bierzynska A, Wherlock M *et al*: Simultaneous sequencing of 24 genes associated with steroid-resistant nephrotic syndrome. *Clin J Am Soc Nephrol* 2013; **8**: 637–648.
- Trujillano D, Ramos MD, Gonzalez J *et al*: Next generation diagnostics of cystic fibrosis and CFTR-related disorders by targeted multiplex high-coverage resequencing of CFTR. *J Med Genet* 2013; **50**: 455–462.
- Ye K, Schulz MH, Long Q, Apweiler R, Ning Z: Pindel: a pattern growth approach to detect break points of large deletions and medium sized insertions from paired-end short reads. *Bioinformatics* 2009; **25**: 2865–2871.
- Krumm N, Sudmant PH, Ko A *et al*: Copy number variation detection and genotyping from exome sequence data. *Genome Res* 2012; **22**: 1525–1532.
- Rossetti S, Consugar MB, Chapman AB *et al*: Comprehensive molecular diagnostics in autosomal dominant polycystic kidney disease. *J Am Soc Nephrol* 2007; **18**: 2143–2160.
- Santín S, Ars E, Rossetti S *et al*: *TRPC6* mutational analysis in a large cohort of patients with focal segmental glomerulosclerosis. *Nephrol Dial Transplant* 2009; **24**: 3089–3096.

- 32 Santín S, García-Maset R, Ruiz P *et al*: Nephtrin mutations cause childhood- and adult-onset focal segmental glomerulosclerosis. *Kidney Int* 2009; **76**: 1268–1276.
- 33 Santín S, Tazón-Vega B, Silva I *et al*: Clinical value of NPHS2 analysis in early- and adult-onset steroid-resistant nephrotic syndrome. *Clin J Am Soc Nephrol* 2011; **6**: 344–354.
- 34 Tavtigian SV, Deffenbaugh AM, Yin L *et al*: Comprehensive statistical study of 452 BRCA1 missense substitutions with classification of eight recurrent substitutions as neutral. *J Med Genet* 2006; **43**: 295–305.
- 35 Houdayer C: In silico prediction of splice-affecting nucleotide variants. *Methods Mol Biol* 2011; **760**: 269–281.
- 36 Heidet L, Arrondel C, Forestier L *et al*: Structure of the human type IV collagen gene COL4A3 and mutations in autosomal Alport syndrome. *J Am Soc Nephrol* 2001; **12**: 97–106.
- 37 Lovric S, Fang H, Vega-Warner V *et al*: Rapid detection of monogenic causes of childhood-onset steroid-resistant nephrotic syndrome. *Clin J Am Soc Nephrol* 2014; **9**: 1109–1116.
- 38 Ars E, Tazon-Vega B, Ruiz P *et al*: Male-to-male transmission of X-linked Alport syndrome in a boy with a 47,XXY karyotype. *Eur J Hum Genet* 2005; **13**: 1040–1046.
- 39 Katsanis N, Ansley SJ, Badano JL *et al*: Triallelic inheritance in Bardet-Biedl syndrome, a Mendelian recessive disorder. *Science* 2001; **293**: 2256–2259.
- 40 Tonna S, Wang YY, Wilson D *et al*: The R229Q mutation in NPHS2 may predispose to proteinuria in thin-basement-membrane nephropathy. *Pediatr Nephrol* 2008; **23**: 2201–2207.
- 41 Voskarides K, Arsali M, Athanasiou Y, Elia A, Pierides A, Deltas C: Evidence that NPHS2-R229Q predisposes to proteinuria and renal failure in familial hematuria. *Pediatr Nephrol* 2012; **4**: 675–679.
- 42 Bergmann C, von Bothmer J, Ortiz Bruchle N *et al*: Mutations in multiple PKD genes may explain early and severe polycystic kidney disease. *J Am Soc Nephrol* 2011; **22**: 2047–2056.
- 43 Savige J, Rana K, Tonna S, Buzza M, Dagher H, Wang YY: Thin basement membrane nephropathy. *Kidney Int* 2003; **64**: 1169–1178.
- 44 Torra R, Tazon-Vega B, Ars E, Ballarin J: Collagen type IV (alpha3-alpha4) nephropathy: from isolated haematuria to renal failure. *Nephrol Dial Transplant* 2004; **19**: 2429–2432.
- 45 Fallerini C, Dosa L, Tita R *et al*: Unbiased next generation sequencing analysis confirms the existence of autosomal dominant Alport syndrome in a relevant fraction of cases. *Clin Genet* 2014; **86**: 252–257.
- 46 Barua M, Brown EJ, Charoonratana VT, Genovese G, Sun H, Pollak MR: Mutations in the INF2 gene account for a significant proportion of familial but not sporadic focal and segmental glomerulosclerosis. *Kidney Int* 2013; **83**: 316–322.

Supplementary Information accompanies this paper on European Journal of Human Genetics website (<http://www.nature.com/ejhg>)