



Published in final edited form as:

Eur J Pain. 2015 October ; 19(9): 1267–1276. doi:10.1002/ejp.656.

Nerve injury-induced calcium channel alpha-2-delta-1 protein dysregulation leads to increased pre-synaptic excitatory input into deep dorsal horn neurons and neuropathic allodynia

Chunyi Zhou¹ and Z. David Luo^{1,2}

¹Department of Pharmacology, School of Medicine, University of California Irvine, Irvine, CA, USA

²Department of Anesthesiology & Perioperative Care, School of Medicine, University of California Irvine, Irvine, CA, USA

Abstract

Background—Upregulation of voltage-gated-calcium-channel $\alpha_2\delta_1$ subunit post spinal nerve ligation injury (SNL) or in $\alpha_2\delta_1$ -overexpressing transgenic (Tg) mice correlates with tactile allodynia, a pain state mediated mainly by A β sensory fibers forming synaptic connections with deep dorsal horn neurons. It is not clear however whether dysregulated $\alpha_2\delta_1$ alters deep dorsal horn synaptic neurotransmission that underlies tactile allodynia development post nerve injury.

Methods—Tactile allodynia was tested in the SNL and $\alpha_2\delta_1$ Tg models. Miniature excitatory/inhibitory postsynaptic currents were recorded in deep dorsal horn (DDH) neurons from these animal models using whole cell patch clamp slice recording techniques..

Results—There was a significant increase in the frequency, but not amplitude, of miniature excitatory postsynaptic currents (mEPSC) in DDH neurons that correlated with tactile allodynia in SNL and $\alpha_2\delta_1$ Tg mice. Gabapentin, an $\alpha_2\delta_1$ ligand that is known to block tactile allodynia in these models, also normalized mEPSC frequency dose-dependently in DDH neurons from SNL and $\alpha_2\delta_1$ Tg mice. In contrast, neither frequency nor amplitude of miniature inhibitory postsynaptic currents (mIPSC) was altered in DDH neurons from SNL and $\alpha_2\delta_1$ Tg mice.

Conclusion—Our data suggest that $\alpha_2\delta_1$ dysregulation is highly likely contributing to tactile allodynia through a pre-synaptic mechanism involving facilitation of excitatory synaptic neurotransmission in deep dorsal horn of spinal cord.

Corresponding author: Z. David Luo, Rm 3113, Gillespie Bldg, 837 Health Sci. Rd., School of Medicine, University of California Irvine, Irvine, CA 92697, USA, Phone: (949) 824-7469, Fax: (949) 824-7447, zluo@uci.edu.

Author Contributions:

In addition to the following contributions from each author, both authors discussed the results, commented on the manuscript and approved for the submission.

C.Z. contributed to conception, design of the study, data acquisition, analysis, and interpretation, drafting, editing the manuscript.

Z.D.L. contributed to conception, design and overall supervision of the study. He also performed data analysis, interpretation, drafting and revising the manuscript.

All authors have no potential conflicts of interest to this study.

- It is known that elevated $\alpha_2\delta_1$ proteins in spinal cord contribute to nerve injury induced neuropathic pain states.
- Findings from this study support that elevated spinal $\alpha_2\delta_1$ proteins increase excitatory pre-synaptic input to deep dorsal horn neurons, underlying the development of neuropathic allodynia post nerve injury.

Introduction

Neuropathic pain, or pain states derived from injuries of the peripheral or central nervous system, usually manifests hypersensitivity to innocuous (allodynia) or noxious (hyperalgesia) stimulation (Woolf & Mannion, 1999; Zimmermann, 2001; Costigan *et al.*, 2009; Baron *et al.*, 2010). It remains as an unmet clinical problem because current treatments are not effective for many patients, and often associated with severe side effects. Development of the next generation of medications for neuropathic pain management relies on our better understanding of neuropathic pain mechanisms.

Dysregulation of the voltage-gated-calcium-channel (VGCC) $\alpha_2\delta_1$ subunit ($\alpha_2\delta_1$) in sensory pathways is believed to play a role in mediating neuropathic pain states. $\alpha_2\delta_1$ is one of auxiliary subunits of VGCC that modulate calcium channel properties *in vivo* and *in vitro* (Tanabe *et al.*, 1987; Klugbauer *et al.*, 2003; Yaksh, 2006; Park & Luo, 2010). Spinal nerve ligation (SNL) injury induces $\alpha_2\delta_1$ upregulation in dorsal root ganglia and associated segments of dorsal spinal cord that contributes to pain modulation (Luo, 2000; Luo *et al.*, 2001; Li *et al.*, 2004; Li *et al.*, 2006; Bauer *et al.*, 2009; Nguyen *et al.*, 2009). Blocking this injury-induced neuroplasticity by intrathecal treatments with $\alpha_2\delta_1$ antisense oligodeoxynucleotide or gabapentin (GBP), a ligand for $\alpha_2\delta_1$, has been shown to block spinal nerve injury induced pain states (Luo *et al.*, 2002; Li *et al.*, 2004; Bauer *et al.*, 2009). Moreover, $\alpha_2\delta_1$ overexpression in transgenic (Tg) mice mimics behavioral hypersensitivities observed in the SNL model (Luo *et al.*, 2002; Li *et al.*, 2006; Nguyen *et al.*, 2009). Together, it is highly likely that increased $\alpha_2\delta_1$ expression contributes to behavioral hypersensitivities post nerve injury.

Importantly, central sensitization is a common mechanism of different chronic pain states (Woolf & Salter, 2000). Factors contributing to central sensitization may include enhanced primary afferent excitability (Gracely *et al.*, 1992; Matzner & Devor, 1994), glutamatergic input to dorsal spinal cord (Kohno *et al.*, 2003; Wang *et al.*, 2007), or impaired spinal inhibitory transmission (Moore *et al.*, 2002b; Baba *et al.*, 2003; Coull *et al.*, 2003). We recently found that $\alpha_2\delta_1$ upregulation post SNL enhances excitatory pre-synaptic input into superficial dorsal horn (SDH, laminae I–II) that contributes to pain states transmitted through nociceptive sensory afferents such as myelinated A δ - and unmyelinated C-type sensory afferents (Zhou & Luo, 2014). However, it is not known if similar changes also occur in deep dorsal horn (DDH, laminae III–V), where neurons receive mainly innocuous tactile input from large A β -sensory afferents. Maladaptive changes of synaptic neurotransmission in DDH post injury may also contribute to specific pain states such as tactile allodynia (Brown, 1982; Basbaum *et al.*, 2009; Todd, 2010).

In this study, we examined the miniature excitatory (mEPSC) and inhibitory (mIPSC) post-synaptic currents in DDH neurons from spinal cord slices of SNL mice with tactile allodynia (Kim & Chung, 1992). In order to determine if injury-induced changes, if any, are modulated by elevated $\alpha_2\delta_1$, but not other injury factors, we compared the data with that from injury-free $\alpha_2\delta_1$ Tg mice with similar tactile allodynia (Li *et al.*, 2006).

Methods and materials

Animals

Adult male mice (8–12 wks) with 129sv background for sham and SNL procedures were purchased from Charles River laboratories, Inc. (Wilmington, MA). The $\alpha_2\delta_1$ Tg mice were generated and characterized as described in our previous studies (Li et al., 2006). Briefly, a neuronal specific thy-1 promoter, derived from deletion of exon 3 and its flanking introns from the murine *thy-1.2* gene (Vidal et al., 1990), was used to drive over expression of the mouse brain $\alpha_2\delta_1$ cDNA (Genbank accession number U73484) in neuronal tissues of the $\alpha_2\delta_1$ Tg mice (Feng et al., 2000). Even though the $\alpha_2\delta_1$ level in neuronal tissues of the Tg mice was compatible to that induced by nerve injury in the SNL model (Li et al., 2006), the $\alpha_2\delta_1$ Tg mice appeared fertile, physically normal with respect to grooming, social interactions, and feeding, and showed no detectable signs of abnormality such as motor defects, ataxia, tremor, or seizure. The Tg mice were backcrossed to the 129sv background for over 10 generations. Only age-matched (8–12 wks) male $\alpha_2\delta_1$ Tg and littermate wild-type (WT) mice were used for the experiments. All animal care and experiments were performed according to protocols approved by the Institutional Animal Care Committees of the University of California Irvine.

Spinal nerve ligation surgery

Unilateral SNL was performed as described by Kim and Chung (Kim & Chung, 1992). Briefly, mice were deeply anesthetized with air-mixed isoflurane (5% for induction, and 2% for maintenance). The left L4 spinal nerve, which is equivalent anatomically to L5 spinal nerve in rat (Rigaud et al., 2008), was exposed and ligated tightly with a 6.0 silk suture. We performed sham ligation by exposing the left L4 spinal nerve and loosely winding a suture around the nerve without ligation.

Von Frey filament stimulation

Paw withdrawal thresholds were determined by the up–down method (Dixon, 1980) using a set of von Frey monofilaments (Stoelting Co., Wood Dale, IL). Briefly, each mouse was habituated in a test compartment with a mesh floor for at least 30 min until exploratory behavior has stopped or decreased to a minimal level. The first von Frey filament (0.41g) was applied to the plantar surface of the hindpaw until it buckled slightly. If a withdrawal response was observed within 5 second, the next lower weight filament was used. Conversely, if the filament failed to elicit a withdrawal response, the next filament with a higher weight was applied. After the first change in response occurred, this paradigm continued until a total of six responses, starting from the one before the change in response, were recorded. In the case of four consecutive positive responses to filaments with decreasing weight, or three consecutive negative responses to filaments with increasing weight, a score of 0.01 g or 3.0 g, respectively, was assigned. These responses were used to calculate the 50% withdrawal threshold as described previously (Luo et al., 2001).

Spinal cord slice preparation

Isoflurane anesthetized mice were decapitated, and the L4 lumber region of the spinal cords was removed, transversely cut with a vibratome (VT-1200, Leica Inc.) into 300 μm slices in ice-cold modified, sucrose-based artificial cerebral spinal fluid (SACSF) saturated with 95% O_2 /5% CO_2 (carbogen). The SACSF contained (mM): 250 sucrose, 2.5 KCl, 1.2 NaH_2PO_4 , 1.2 MgCl_2 , 2.4 CaCl_2 , 26 NaHCO_3 and 11 glucose. Slices were first incubated in SACSF at 31 $^\circ\text{C}$ for at least one hr before being kept at room temperature in carbogenated ACSF containing (mM) 126 NaCl, 2.5 KCl, 1.2 NaH_2PO_4 , 1.2 MgCl_2 , 2.4 CaCl_2 , 26 NaHCO_3 and 11 glucose for at least 30 min, then placed in a recording chamber and continually perfused with carbogen-saturated ACSF at a flow rate of 2.0 ml/min. 1 μM tetrodotoxin (TTX) was applied continually when mEPSC/mIPSC were recorded.

Patch clamp recording

DDH neurons in slices were visualized with an upright microscope (Eclipse FN1, Nikon) with near-infrared illumination. All DDH recordings were made in the grey matter dorsal lateral to the central canal and at least 100 μm away from bottom of lamina II. The neuron locations were confirmed under low magnification after recording. The intracellular solution for recording mEPSC contains (mM): 135 potassium gluconate, 5 KCl, 5 EGTA, 0.5 CaCl_2 , 10 HEPES, 2 Mg-ATP, and 0.1 GTP; that for recording mIPSC contains (mM): 70 potassium gluconate, 65 KCl, 5 EGTA, 0.5 CaCl_2 , 10 HEPES, 2 Mg-ATP, 0.1 GTP. These solutions were adjusted to pH 7.2 with Tris-base, osmolarity 300 mOsm. The junction potential was nulled between the patch pipette and bath solution before gigaseal formation. Series resistance was monitored throughout the experiment without compensation (Multiclamp 700B) and changes in series resistance (15–30 $\text{M}\Omega$) more than 20% during whole-cell recording would lead to data elimination. Electrophysiological signals were recorded with MultiClamp 700B amplifiers (Axon Instruments, Molecular Devices, Union City, CA), Digidata 1440 analog-to-digital converters (Axon Instruments) and pClamp 10.2 software (Axon Instruments). Data were sampled at 10 kHz and filtered at 2 kHz. All recordings were done at a temperature of $32 \pm 0.5^\circ\text{C}$.

The slices were stabilized for 5 min, then mEPSC/mIPSC were recorded, counted and analyzed using clampfit 10.3 (Molecular Devices). For drug treatment, mEPSC frequency baseline mean values were obtained for 5 min during the control period, and the mean values for drug application were obtained over the peak drug response for 2 min. The mEPSC frequencies during and after drug applications were compared to their baseline mean values. Drug effects were expressed as percentage changes (mean \pm SEM) over the baseline values.

Statistics analysis

All data were expressed as mean \pm SEM. Differences between two groups were compared with student's *t*-test. One-way or Two-way ANOVA analysis with Bonferroni post hoc test was used for multi-group comparisons where necessary. $p < 0.05$ was considered statistically significant.

Results

Upregulation of $\alpha_2\delta_1$ induces tactile allodynia

We first examined if tactile allodynia, shown as reduced hindpaw thresholds to light touch was manifested similarly in both SNL and $\alpha_2\delta_1$ Tg mice by measuring hindpaw sensitivity to von Frey filament stimuli in both models. In the SNL model, paw withdrawal thresholds were significantly decreased in the injury side of SNL mice within one week post injury while there was no significant change in the injury side of sham mice during the same period (Fig. 1A). The tactile allodynia state correlates with upregulation of $\alpha_2\delta_1$ proteins in dorsal root ganglia and associated spinal dorsal horn reported previously (Zhou & Luo, 2014). The time course of SNL-induced tactile allodynia was similar to that previously reported (Luo et al., 2001; Luo et al., 2002; Li et al., 2004). Similarly, $\alpha_2\delta_1$ overexpression in the Tg mice also caused tactile allodynia compared with their WT littermates (Fig. 1B) (Li et al., 2006). In combination with previous findings that blocking $\alpha_2\delta_1$ expression or activity with intrathecal $\alpha_2\delta_1$ antisense oligodeoxynucleotides, or the $\alpha_2\delta_1$ ligand gabapentin (GBP), respectively, reverses behavioral hypersensitivity in both SNL and $\alpha_2\delta_1$ Tg animals (Li et al., 2004; Li et al., 2006), these findings suggest that elevated spinal $\alpha_2\delta_1$ mediates tactile allodynia.

SNL or elevated $\alpha_2\delta_1$ enhanced excitatory synaptic transmission in DDH neurons through a pre-synaptic mechanism

To the best of our knowledge, the effects of SNL injury in DDH neuron synaptic transmission are not yet investigated or reported, which however could play a significant role in the processing of tactile allodynia. Accordingly, we examined mEPSC in deep dorsal horn neurons from L4 spinal cord slices of the surgery side (SNL or sham) 5–7 days post operation when SNL mice displayed allodynia. AP-5 (50 μ M), bicuculline (10 μ M), strychnine (1 μ M), and tetrodotoxin (TTX 1 μ M) were added to block NMDA receptors, GABA_A receptors, glycine receptors, and TTX-sensitive sodium channels, respectively. Bath application of 20 μ M CNQX, a non-NMDA receptor antagonist, at the end of the recording abolished mEPSCs (data not shown), confirming that mEPSC from DDH neurons was mediated through AMPA/Kainate receptors.

Our data indicated that mEPSC frequency was increased in deep dorsal horn neurons from the injury side of SNL mice compared with that from sham control mice (Fig. 2A), resulting in a leftward shift in the cumulative probability curve of interevent intervals (Fig. 2B), but not amplitude (Fig. 2C) of mEPSC. These corresponded to a ~100% increase in mEPSC average frequency, likely reflecting increased glutamate release frequency from pre-synaptic sites (Fig. 2D), but no significant change in mEPSC amplitude (Fig. 2E), a measurement of post-synaptic neuron excitability. Importantly, GBP (10–100 μ M) could reversibly normalize SNL-induced mEPSC frequency in DDH neurons dose-dependently without affecting that in sham mice (Fig. 2F and G). These data support that SNL enhanced excitatory synaptic transmission in DDH through a pre-synaptic mechanism.

Since SNL caused dysregulation of a lot of genes in DRG and spinal cord (Wang et al., 2002; Valder et al., 2003), it was not clear if SNL-induced $\alpha_2\delta_1$ was responsible for the

increased pre-synaptic excitatory input that correlated with tactile allodynia. We addressed this question by examining whether elevated $\alpha_2\delta_1$ alone without the influence from other injury factors is sufficient to enhance mEPSC frequency in DDH of the injury-free $\alpha_2\delta_1$ Tg mice. Overexpression of $\alpha_2\delta_1$ in neuronal tissues, including spinal cord and DRG, in the $\alpha_2\delta_1$ Tg mice leads to similar behavioral hypersensitivity as that observed in the SNL model (Li et al., 2006), thus, the injury-free $\alpha_2\delta_1$ Tg mice could be a complementary model to the SNL model in investigating the role of $\alpha_2\delta_1$ upregulation in pain state processing. Similar to that in SNL mice, $\alpha_2\delta_1$ upregulation in the Tg mice resulted in a significant increase in frequency (Fig. 3A) of mEPSC with a leftward shift in cumulative probability curve of interevent intervals (Fig. 3B), but no change in mEPSC amplitude (Fig. 3C) compared with that from the WT mice. These changes corresponded to a ~128% higher average mEPSC frequency (Fig. 3D), but no significant change in average mEPSC amplitude (Fig. 3E) in the Tg mice compared with that in WT mice. GBP (10–100 μ M) dose-dependently attenuated mEPSC frequency in DDH neurons from the Tg mice without affecting that from WT mice with the highest concentration tested (100 μ M) (Fig. 3F and G). In combination with similar findings from the SNL mice, these data support a role of elevated $\alpha_2\delta_1$ in enhancing excitatory synaptic transmission in DDH through a pre-synaptic mechanism.

Unaltered inhibitory synaptic transmission in DDH neurons from SNL and $\alpha_2\delta_1$ Tg mice

Besides glutamatergic excitatory innervations, DDH neurons also receive GABAergic and Glycinergic innervations from inhibitory interneurons in spinal dorsal horn (Braz et al., 2014). To test if $\alpha_2\delta_1$ upregulation in SNL (5–7 days post injury) and $\alpha_2\delta_1$ Tg mice also affected inhibitory synaptic transmission in DDH neurons, we examined mIPSC in the presence of TTX (1 μ M) from L4 spinal slices of these models. During the recording, mEPSC was blocked with DNQX (20 μ M) and APV (50 μ M). The remaining mIPSC could be eliminated by bicuculline (10 μ M) and strychnine (1 μ M) (data not shown) confirming a suitable recording condition for examining mIPSC (Fig. 4A).

Interestingly, there is no significant difference in mIPSC either between the SNL and Sham mice, or between the $\alpha_2\delta_1$ Tg and WT mice. The distribution of interevent intervals and amplitude of mIPSC did not shift in SNL mice compared with that from the sham mice (Fig. 4B, C), nor in $\alpha_2\delta_1$ Tg mice compared with that from the WT mice (Fig. 4F, G). The average mIPSC frequencies and amplitudes in DDH neurons were similar between SNL/sham (Fig. 4D, E) or Tg/WT mice (Fig. 4H, I). These data suggest that increased $\alpha_2\delta_1$ in neither the SNL nor the $\alpha_2\delta_1$ Tg mouse model affects inhibitory synaptic transmission in deep dorsal spinal cord.

Discussion

Peripheral nerve injury induces neuroplasticity changes in deep dorsal horn of spinal cord that may contribute to neuropathic allodynia, a painful state of hypersensitivity to innocuous light touch sensation, which is normally transmitted by A β -sensory afferents projecting to and forming synapse connections with DDH neurons (Braz et al., 2014). These changes may include, but are not limited to, dysregulation of genes (Dobremez et al., 2005), suppressing inhibitory and/or enhancing facilitatory descending modulations (Suzuki et al., 2004;

Rahman et al., 2008), attenuated glutamate uptake (Binns et al., 2005), and increased pain-inducing peptides in DDH neurons (Nitzan-Luques et al., 2011). However, to the best of our knowledge, it is not yet known if peripheral nerve injury alters excitatory and/or inhibitory miniature synaptic transmission in deep dorsal horn, which can play a critical role in mediating neuropathic allodynia. We provide data here to indicate that spinal nerve ligation injury leads to increased frequency of mEPSC in deep dorsal horn neurons, which is gabapentin-reversible, and neuropathic tactile allodynia. Similar maladaptive changes in DDH mEPSC and tactile allodynia also occur in injury-free transgenic mice with $\alpha_2\delta_1$ overexpression in neuronal tissues. In contrast, neither spinal nerve ligation injury nor $\alpha_2\delta_1$ overexpression causes any significant change in DDH mIPSC. Together, these data support that peripheral nerve injury enhances excitatory pre-synaptic input into DDH neurons under non-stimulating conditions, most likely through $\alpha_2\delta_1$ upregulation, that may play a role in neuropathic allodynia.

Several lines of evidence support our conclusion. First, $\alpha_2\delta_1$ proteins are highly upregulated in DRG after SNL injury, then translocated to primary afferent pre-synaptic terminals in dorsal spinal cord during the development of neuropathic allodynia (Li et al., 2004; Bauer et al., 2009). Second, blocking this injury-induced neuroplasticity with intrathecal treatments of $\alpha_2\delta_1$ antisense oligodeoxynucleotides, or GBP prevents (Boroujerdi et al., 2008) and reverses (Luo et al., 2002; Li et al., 2004; Bauer et al., 2009) neuropathic allodynia. Third, $\alpha_2\delta_1$ overexpression in injury-free Tg mice is sufficient to induce behavioral hypersensitivity and dorsal horn neuron hyperexcitability with similar pharmacological profiles as that observed in the SNL model (Li et al., 2006; Chang et al., 2013; Chang et al., 2014; Zhou & Luo, 2014).

To our knowledge, this is the first study revealing maladaptive changes in DDH miniature synaptic transmission underlying nerve injury-induced mechanical allodynia. Even though we have recently reported that peripheral nerve injury induces similar changes in superficial dorsal horn (SDH) and behavioral nociceptions in SNL and $\alpha_2\delta_1$ Tg mice (Zhou & Luo, 2014), SDH and DDH have functional differences in mediating sensory signal processing since neurons in these locations receive distinct sensory input from different primary afferents (Light & Perl, 1979; Brown, 1982; Millan, 1999; Graham et al., 2007; Braz et al., 2014). The SDH neurons receive primary afferent inputs mainly from small myelinated A δ and unmyelinated C-fibers carrying nociceptive, including thermal and cold, information. In contrast, DDH neurons receive inputs mainly from large A β myelinated fibers carrying tactile information and synaptic inputs from SDH neurons. Neuron populations and synaptic connections may also differ in superficial and deep laminae. Compared with a high percentage of nociceptive specific neurons in the superficial dorsal horn (lamina I–II), wide dynamic range (WDR) neurons in deep dorsal horn are responsive to a large range of sensory modalities (thermal, chemical and mechanical) and represent an important component in spinal sensory/pain transmission (Brown, 1982; Basbaum et al., 2009; Todd, 2010; Braz et al., 2014). WDR neurons exhibit a frequency-dependent, progressive increase in neuronal excitability in response to repeated electrical stimulation of afferent C-fibers (windup phenomenon) (Herrero et al., 2000). In addition, excitatory and inhibitory interneurons also form specific circuitries in SDH and DDH with projection, WDR neurons or between each other to modulate sensory information processing at the spinal cord level,

which in turn, is subjected to descending modulation from the higher central nervous system (Braz et al., 2014).

Our previous data from the injury-free $\alpha_2\delta_1$ Tg mice indicate that elevated neuronal $\alpha_2\delta_1$ expression leads to exaggerated and prolonged WDR neuron firing in response to mechanical and thermal stimuli, but normal windup response to electrical stimuli, suggesting that elevated $\alpha_2\delta_1$ mediates DDH neuron aberrant activities and pain states through a mechanism independent of abnormal C-fiber stimulation (Li et al., 2006). Findings from this study indicate that increased mEPSC frequency is found in both SNL and $\alpha_2\delta_1$ Tg mice that have common correlates of elevated $\alpha_2\delta_1$ proteins and behavioral hypersensitivities (Luo et al., 2002; Li et al., 2004; Li et al., 2006). GBP can dose-dependently normalize aberrant mEPSC frequency in both models (Figs. 2F, 2G, 3F, 3G) in a dose-range achievable in plasma when GBP is given orally to patients (Ben-Menachem et al., 1992; Bryans & Wustrow, 1999). While GBP binds to both $\alpha_2\delta_1$ and $\alpha_2\delta_2$ proteins (Klugbauer et al., 2003), it is less likely that $\alpha_2\delta_2$ is involved in this process since SNL only induces DRG $\alpha_2\delta_1$, but not $\alpha_2\delta_2$, upregulation (Bauer et al., 2009) and GBP is not affecting mEPSC frequency in sham and WT control mice (Figs. 2F, 3F). This is consistent with previous studies suggesting that GBP only exhibits therapeutic effect under pathological conditions (Ben-Menachem et al., 1992; Field et al., 1997; Stanfa et al., 1997; Bryans & Wustrow, 1999; Moore et al., 2002a). The fast action of GBP on mEPSC frequency supports its effects on synaptic transmission, rather than trafficking of VGCC (Hendrich et al., 2008). In addition, GBP at this dose-range is able to inhibit evoked Ca^{2+} influx in cortical synaptosomal suspensions (Fink et al., 2000; Meder & Dooley, 2000), and excitatory synaptic transmission in hyperalgesic spinal cord (Patel et al., 2000). Together, these findings support that peripheral nerve injury-induced $\alpha_2\delta_1$ mediates neuropathic allodynia at least at the spinal cord level by enhancing excitatory pre-synaptic input into deep dorsal horn.

The mechanism underlying $\alpha_2\delta_1$ modulation on excitatory synaptic transmission in deep dorsal spinal cord reminds elusive. A large body of emerging evidence indicates that $\alpha_2\delta_1$ subunit is a multifunctional protein. It not only regulates VGCC functions, but is also critical for VGCC-independent functions. Since miniature glutamate release does not require Ca^{2+} influx into pre-synaptic terminals, $\alpha_2\delta_1$ may have other regulatory functions or interactions with pre-synaptic proteins involved in synaptic neurotransmission. Data from recent studies show that $\alpha_2\delta_1$ is the receptor for thrombospondin, an extracellular matrix protein secreted by astrocytes, in promoting excitatory synaptogenesis in the CNS, which is GBP sensitive, but VGCC-independent (Eroglu et al., 2009). In addition, $\alpha_2\delta_1$ interacts with a serotonergic descending facilitation pathway at the spinal level that is critical in mediating central sensitization and neuropathic allodynia/thermal hyperalgesia (Chang et al., 2013), as well as GBP efficacy in reversing pain states (Suzuki et al., 2005). Therefore, it is possible that $\alpha_2\delta_1$ modulation on mEPSC frequency requires integration of multiple pathways that could be GBP-sensitive, but VGCC-independent. This is supported by recent findings that $\alpha_2\delta_1$ modulates EPSC frequency in ventromedial hypothalamus neurons through a VGCC-independent mechanism that underlies GBP-induced weight gain (Cordeira et al., 2014).

The contribution of spinal dorsal horn inhibitory circuitry to pain processing is equally important as that from excitatory circuitry. However, findings related to contribution of

spinal inhibitory circuitry to neuropathic pain state processing remain inconsistent. While suppression of inhibitory synaptic transmission in the dorsal spinal cord has been reported in certain chronic pain states such as tactile allodynia and hyperalgesia (Yaksh, 1989; Kontinen et al., 2001), other studies have shown that neither GABAergic nor glycinergic IPSC is changed in neuropathic rodents (Polgar et al., 2003; Polgar et al., 2005; Wang et al., 2007), which is similar to our findings that behavioral hypersensitivity in neither the SNL nor $\alpha_2\delta_1$ Tg mice correlates with any change in DDH mIPSC (Fig. 4). This discrepancy could be due to the differences in experimental conditions, neuron sampling locations, animal strains and pain models. In addition, our data cannot exclude the possibility that evoked IPSC, but not mIPSC, is altered in these allodynic models. Nevertheless, our findings provide new evidences indicating that $\alpha_2\delta_1$ upregulation in spinal cord differentially modulate excitatory and inhibitory synaptic transmission.

In summary, our data indicate that peripheral nerve injury can enhance excitatory synaptic neurotransmission in DDH by enhancing mEPSC through a pre-synaptic mechanism, most likely mediated by elevated $\alpha_2\delta_1$ expression, which in turn contributes to neuropathic allodynia. Blocking this maladaptive change post nerve injury may provide an alternative strategy in managing modality specific nociception.

Acknowledgments

Funding Sources: This study was supported in part by grants NS40135, NS064341 and DE021847 from the National Institutes of Health (Z.D. Luo).

References

- Baba H, Ji RR, Kohno T, Moore KA, Ataka T, Wakai A, Okamoto M, Woolf CJ. Removal of GABAergic inhibition facilitates polysynaptic A fiber-mediated excitatory transmission to the superficial spinal dorsal horn. *Mol Cell Neurosci*. 2003; 24:818–830. [PubMed: 14664828]
- Baron R, Binder A, Wasner G. Neuropathic pain: diagnosis, pathophysiological mechanisms, and treatment. *Lancet Neurol*. 2010; 9:807–819. [PubMed: 20650402]
- Basbaum AI, Bautista DM, Scherrer G, Julius D. Cellular and molecular mechanisms of pain. *Cell*. 2009; 139:267–284. [PubMed: 19837031]
- Bauer CS, Nieto-Rostro M, Rahman W, Tran-Van-Minh A, Ferron L, Douglas L, Kadurin I, Sri Ranjan Y, Fernandez-Alacid L, Millar NS, Dickenson AH, Lujan R, Dolphin AC. The increased trafficking of the calcium channel subunit $\alpha_2\delta_1$ to presynaptic terminals in neuropathic pain is inhibited by the $\alpha_2\delta_1$ ligand pregabalin. *J Neurosci*. 2009; 29:4076–4088. [PubMed: 19339603]
- Ben-Menachem E, Persson LI, Hedner T. Selected CSF biochemistry and gabapentin concentrations in the CSF and plasma in patients with partial seizures after a single oral dose of gabapentin. *Epilepsy Res*. 1992; 11:45–49. [PubMed: 1373372]
- Binns BC, Huang Y, Goettl VM, Hackshaw KV, Stephens RL Jr. Glutamate uptake is attenuated in spinal deep dorsal and ventral horn in the rat spinal nerve ligation model. *Brain Res*. 2005; 1041:38–47. [PubMed: 15804498]
- Boroujerdi A, Kim HK, Lyu YS, Kim DS, Figueroa KW, Chung JM, Luo ZD. Injury discharges regulate calcium channel $\alpha_2\delta_1$ subunit upregulation in the dorsal horn that contributes to initiation of neuropathic pain. *Pain*. 2008; 139:358–366. [PubMed: 18571852]
- Braz J, Solorzano C, Wang X, Basbaum AI. Transmitting pain and itch messages: a contemporary view of the spinal cord circuits that generate gate control. *Neuron*. 2014; 82:522–536. [PubMed: 24811377]

- Brown AG. The dorsal horn of the spinal cord. *Q J Exp Physiol.* 1982; 67:193–212. [PubMed: 6281848]
- Bryans JS, Wustrow DJ. 3-substituted GABA analogs with central nervous system activity: a review. *Med Res Rev.* 1999; 19:149–177. [PubMed: 10189176]
- Chang E, Chen X, Kim M, Gong N, Bhatia S, Luo ZD. Differential Effects of Voltage-Gated Calcium Channel Blockers on Calcium Channel Alpha-2-Delta-1 Subunit Protein Mediated Nociception. *Eur J Pain.* 2014 In press.
- Chang EY, Chen X, Sandhu A, Li CY, Luo ZD. Spinal 5-HT₃ receptors facilitate behavioural hypersensitivity induced by elevated calcium channel alpha-2-delta-1 protein. *Eur J Pain.* 2013; 17:505–513. [PubMed: 23065867]
- Cordeira JW, Felsted JA, Teillon S, Daftary S, Panessiti M, Wirth J, Sena-Esteves M, Rios M. Hypothalamic dysfunction of the thrombospondin receptor alpha2delta-1 underlies the overeating and obesity triggered by brain-derived neurotrophic factor deficiency. *J Neurosci.* 2014; 34:554–565. [PubMed: 24403154]
- Costigan M, Scholz J, Woolf CJ. Neuropathic pain: a maladaptive response of the nervous system to damage. *Annu Rev Neurosci.* 2009; 32:1–32. [PubMed: 19400724]
- Coull JA, Boudreau D, Bachand K, Prescott SA, Nault F, Sik A, De Koninck P, De Koninck Y. Trans-synaptic shift in anion gradient in spinal lamina I neurons as a mechanism of neuropathic pain. *Nature.* 2003; 424:938–942. [PubMed: 12931188]
- Dixon WJ. Efficient analysis of experimental observations. *Annu Rev Pharmacol Toxicol.* 1980; 20:441–462. [PubMed: 7387124]
- Dobremez E, Bouali-Benazzouz R, Fossat P, Monteils L, Dulluc J, Nagy F, Landry M. Distribution and regulation of L-type calcium channels in deep dorsal horn neurons after sciatic nerve injury in rats. *Eur J Neurosci.* 2005; 21:3321–3333. [PubMed: 16026470]
- Eroglu C, Allen NJ, Susman MW, O'Rourke NA, Park CY, Ozkan E, Chakraborty C, Mulinyawe SB, Annis DS, Huberman AD, Green EM, Lawler J, Dolmetsch R, Garcia KC, Smith SJ, Luo ZD, Rosenthal A, Mosher DF, Barres BA. Gabapentin receptor alpha2delta-1 is a neuronal thrombospondin receptor responsible for excitatory CNS synaptogenesis. *Cell.* 2009; 139:380–392. [PubMed: 19818485]
- Feng G, Mellor RH, Bernstein M, Keller-Peck C, Nguyen QT, Wallace M, Nerbonne JM, Lichtman JW, Sanes JR. Imaging Neuronal Subsets in Transgenic Mice Expressing Multiple Spectral Variants of GFP. *Neuron.* 2000; 28:41–51. [PubMed: 11086982]
- Field MJ, Oles RJ, Lewis AS, McCleary S, Hughes J, Singh L. Gabapentin (neurontin) and S-(+)-3-isobutylgaba represent a novel class of selective antihyperalgesic agents. *Brit J Pharmacol.* 1997; 121:1513–1522. [PubMed: 9283683]
- Fink K, Meder W, Dooley DJ, Gothert M. Inhibition of neuronal Ca(2+) influx by gabapentin and subsequent reduction of neurotransmitter release from rat neocortical slices. *Br J Pharmacol.* 2000; 130:900–906. [PubMed: 10864898]
- Gracely RH, Lynch SA, Bennett GJ. Painful neuropathy: altered central processing maintained dynamically by peripheral input. *Pain.* 1992; 51:175–194. [PubMed: 1484715]
- Graham BA, Brichta AM, Callister RJ. Moving from an averaged to specific view of spinal cord pain processing circuits. *J Neurophysiol.* 2007; 98:1057–1063. [PubMed: 17567772]
- Hendrich J, Van Minh AT, Hebllich F, Nieto-Rostro M, Watschinger K, Striessnig J, Wratten J, Davies A, Dolphin AC. Pharmacological disruption of calcium channel trafficking by the alpha2delta ligand gabapentin. *Proc Natl Acad Sci U S A.* 2008; 105:3628–3633. [PubMed: 18299583]
- Herrero JF, Laird JM, Lopez-Garcia JA. Wind-up of spinal cord neurones and pain sensation: much ado about something? *Prog Neurobiol.* 2000; 61:169–203. [PubMed: 10704997]
- Kim SH, Chung JM. An experimental model for peripheral neuropathy produced by segmental spinal nerve ligation in the rat. *Pain.* 1992; 50:355–363. [PubMed: 1333581]
- Klugbauer N, Marais E, Hofmann F. Calcium channel alpha2delta subunits: differential expression, function, and drug binding. *J Bioenerg Biomembr.* 2003; 35:639–647. [PubMed: 15000524]
- Kohn T, Moore KA, Baba H, Woolf CJ. Peripheral nerve injury alters excitatory synaptic transmission in lamina II of the rat dorsal horn. *J Physiol.* 2003; 548:131–138. [PubMed: 12576493]

- Kontinen VK, Stanfa LC, Basu A, Dickenson AH. Electrophysiologic evidence for increased endogenous gabaergic but not glycinergic inhibitory tone in the rat spinal nerve ligation model of neuropathy. *Anesthesiology*. 2001; 94:333–339. [PubMed: 11176099]
- Li CY, Song YH, Higuera ES, Luo ZD. Spinal dorsal horn calcium channel alpha2delta-1 subunit upregulation contributes to peripheral nerve injury-induced tactile allodynia. *J Neurosci*. 2004; 24:8494–8499. [PubMed: 15456823]
- Li CY, Zhang XL, Matthews EA, Li KW, Kurwa A, Boroujerdi A, Gross J, Gold MS, Dickenson AH, Feng G, Luo ZD. Calcium channel alpha2delta1 subunit mediates spinal hyperexcitability in pain modulation. *Pain*. 2006; 125:20–34. [PubMed: 16764990]
- Light AR, Perl ER. Reexamination of the dorsal root projection to the spinal dorsal horn including observations on the differential termination of coarse and fine fibers. *J Comp Neurol*. 1979; 186:117–131. [PubMed: 447880]
- Luo ZD. Rat dorsal root ganglia express distinctive forms of the alpha2 calcium channel subunit. *Neuroreport*. 2000; 11:3449–3452. [PubMed: 11095497]
- Luo ZD, Calcutt NA, Higuera ES, Valder CR, Song YH, Svensson CI, Myers RR. Injury type-specific calcium channel alpha 2 delta-1 subunit up-regulation in rat neuropathic pain models correlates with antiallodynic effects of gabapentin. *J Pharmacol Exp Ther*. 2002; 303:1199–1205. [PubMed: 12438544]
- Luo ZD, Chaplan SR, Higuera ES, Sorkin LS, Stauderman KA, Williams ME, Yaksh TL. Upregulation of dorsal root ganglion (alpha)2(delta) calcium channel subunit and its correlation with allodynia in spinal nerve-injured rats. *J Neurosci*. 2001; 21:1868–1875. [PubMed: 11245671]
- Matzner O, Devor M. Hyperexcitability at sites of nerve injury depends on voltage-sensitive Na⁺ channels. *J Neurophysiol*. 1994; 72:349–359. [PubMed: 7965019]
- Meder WP, Dooley DJ. Modulation of K⁽⁺⁾-induced synaptosomal calcium influx by gabapentin. *Brain Res*. 2000; 875:157–159. [PubMed: 10967310]
- Millan MJ. The induction of pain: an integrative review. *Prog Neurobiol*. 1999; 57:1–164. [PubMed: 9987804]
- Moore KA, Baba H, Woolf CJ. Gabapentin-- actions on adult superficial dorsal horn neurons. *Neuropharmacology*. 2002a; 43:1077–1081. [PubMed: 12504913]
- Moore KA, Kohno T, Karchewski LA, Scholz J, Baba H, Woolf CJ. Partial peripheral nerve injury promotes a selective loss of GABAergic inhibition in the superficial dorsal horn of the spinal cord. *J Neurosci*. 2002b; 22:6724–6731. [PubMed: 12151551]
- Nguyen D, Deng P, Matthews EA, Kim DS, Feng G, Dickenson AH, Xu ZC, Luo ZD. Enhanced pre-synaptic glutamate release in deep-dorsal horn contributes to calcium channel alpha-2-delta-1 protein-mediated spinal sensitization and behavioral hypersensitivity. *Mol Pain*. 2009; 5:6. [PubMed: 19216737]
- Nitzan-Luques A, Devor M, Tal M. Genotype-selective phenotypic switch in primary afferent neurons contributes to neuropathic pain. *Pain*. 2011; 152:2413–2426. [PubMed: 21872992]
- Park J, Luo ZD. Calcium channel functions in pain processing. *Channels (Austin)*. 2010; 4:510–517. [PubMed: 21150297]
- Patel MK, Gonzalez MI, Bramwell S, Pinnock RD, Lee K. Gabapentin inhibits excitatory synaptic transmission in the hyperalgesic spinal cord. *Br J Pharmacol*. 2000; 130:1731–1734. [PubMed: 10952660]
- Polgar E, Hughes DI, Arham AZ, Todd AJ. Loss of neurons from laminae I–III of the spinal dorsal horn is not required for development of tactile allodynia in the spared nerve injury model of neuropathic pain. *J Neurosci*. 2005; 25:6658–6666. [PubMed: 16014727]
- Polgar E, Hughes DI, Riddell JS, Maxwell DJ, Puskar Z, Todd AJ. Selective loss of spinal GABAergic or glycinergic neurons is not necessary for development of thermal hyperalgesia in the chronic constriction injury model of neuropathic pain. *Pain*. 2003; 104:229–239. [PubMed: 12855333]
- Rahman W, D’Mello R, Dickenson AH. Peripheral nerve injury-induced changes in spinal alpha(2)-adrenoceptor-mediated modulation of mechanically evoked dorsal horn neuronal responses. *J Pain*. 2008; 9:350–359. [PubMed: 18226963]

- Rigaud M, Gemes G, Barabas ME, Chernoff DI, Abram SE, Stucky CL, Hogan QH. Species and strain differences in rodent sciatic nerve anatomy: implications for studies of neuropathic pain. *Pain*. 2008; 136:188–201. [PubMed: 18316160]
- Stanfa LC, Singh L, Williams RG, Dickenson AH. Gabapentin, ineffective in normal rats, markedly reduces C-fibre evoked responses after inflammation. *Neuroreport*. 1997; 8:587–590. [PubMed: 9106728]
- Suzuki R, Rahman W, Hunt SP, Dickenson AH. Descending facilitatory control of mechanically evoked responses is enhanced in deep dorsal horn neurones following peripheral nerve injury. *Brain Res*. 2004; 1019:68–76. [PubMed: 15306240]
- Suzuki R, Rahman W, Rygh LJ, Webber M, Hunt SP, Dickenson AH. Spinal-supraspinal serotonergic circuits regulating neuropathic pain and its treatment with gabapentin. *Pain*. 2005; 117:292–303. [PubMed: 16150546]
- Tanabe T, Takeshima H, Mikami A, Flockerzi V, Takahashi H, Kangawa K, Kojima M, Matsuo H, Hirose T, Numa S. Primary structure of the receptor for calcium channel blockers from skeletal muscle. *Nature*. 1987; 328:313–318. [PubMed: 3037387]
- Todd AJ. Neuronal circuitry for pain processing in the dorsal horn. *Nat Rev Neurosci*. 2010; 11:823–836. [PubMed: 21068766]
- Valder CR, Liu JJ, Song YH, Luo ZD. Coupling gene chip analyses and rat genetic variances in identifying potential target genes that may contribute to neuropathic allodynia development. *J Neurochem*. 2003; 87:560–573. [PubMed: 14535940]
- Vidal M, Morris R, Grosveld F, Spanopoulou E. Tissue-specific control elements of the Thy-1 gene. *Embo Journal*. 1990; 9:833–840. [PubMed: 1968831]
- Wang H, Sun H, Della Penna K, Benz RJ, Xu J, Gerhold DL, Holder DJ, Koblan KS. Chronic neuropathic pain is accompanied by global changes in gene expression and shares pathobiology with neurodegenerative diseases. *Neuroscience*. 2002; 114:529–546. [PubMed: 12220557]
- Wang XL, Zhang HM, Chen SR, Pan HL. Altered synaptic input and GABAB receptor function in spinal superficial dorsal horn neurons in rats with diabetic neuropathy. *J Physiol*. 2007; 579:849–861. [PubMed: 17218355]
- Woolf CJ, Mannion RJ. Neuropathic pain: aetiology, symptoms, mechanisms, and management. *Lancet*. 1999; 353:1959–1964. [PubMed: 10371588]
- Woolf CJ, Salter MW. Neuronal plasticity: increasing the gain in pain. *Science*. 2000; 288:1765–1769. [PubMed: 10846153]
- Yaksh TL. Behavioral and autonomic correlates of the tactile evoked allodynia produced by spinal glycine inhibition: effects of modulatory receptor systems and excitatory amino acid antagonists. *Pain*. 1989; 37:111–123. [PubMed: 2542867]
- Yaksh TL. Calcium channels as therapeutic targets in neuropathic pain. *J Pain*. 2006; 7:S13–30. [PubMed: 16426997]
- Zhou C, Luo ZD. Electrophysiological characterization of spinal neuron sensitization by elevated calcium channel alpha-2-delta-1 subunit protein. *Eur J Pain*. 2014; 18:649–658. [PubMed: 24151064]
- Zimmermann M. Pathobiology of neuropathic pain. *Eur J Pharmacol*. 2001; 429:23–37. [PubMed: 11698024]




Figure 1. Similar tactile allodynia were observed in unilateral L4 SNL and $\alpha_2\delta_1$ Tg mice

Tactile allodynia was measured by applying von Frey filaments to the plantar surface of the hindpaw as described. **A.** Hindpaw withdrawal thresholds (PWT) to light touch in the ipsilateral hindpaws of sham and SNL mice were tested in a blinded fashion at designated time points post surgery. **B.** Similarly tested adult $\alpha_2\delta_1$ Tg mice also showed decreased PWT compared with their WT littermates. Data presented are the means \pm SEM from the number of animals indicated. ** $p < 0.01$ compared with control values by one-way ANOVA test.




Figure 2. L4 spinal nerve ligation in mice led to enhancement of mEPSC frequency in DDH neurons that could be blocked by gabapentin
 mEPSC from injury side of L4 deep dorsal horn neurons of sham and SNL mice was recorded 5 – 7 days post surgery. **A.** Representative traces of mEPSC from DDH neurons of sham and SNL mice, respectively. **B and C.** Cumulative probability curves of mEPSC interevent intervals (**B**) and amplitude (**C**) from Sham and SNL mice, respectively. $p = 0.001$ (k-s test) for a comparison of mEPSC interevent intervals between sham and SNL mice. $p = 0.21$ for a comparison of mEPSC amplitude between sham and SNL mice. **D and E.** Summary of mEPSC frequency (**D**) and amplitude (**E**) in L4 DDH neurons from injury side of SNL and sham mice. $**p < 0.01$ compared with sham L4 neurons by Student's t -test. **F.** Time courses for the effects of $100\mu\text{M}$ GBP on mEPSC frequency from sham and SNL neurons. **G.** Dose-dependent normalization by GBP on mEPSC frequency on SNL neurons. $*p < 0.05$, $**p < 0.01$ compared with pre-GBP treatment; $\#p < 0.05$ compared with $10\mu\text{M}$

treatment by one-way ANOVA test. Dotted line represents percentage of mEPSC frequency in sham mice shown in D. Summarized data are shown as the means \pm SEM from the number of neurons indicated in the parentheses.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 3. Enhanced mEPSC in DDH neurons from $\alpha_2\delta_1$ Tg mice that could be blocked by gabapentin
 mEPSC from L4 DDH neurons of adult WT and $\alpha_2\delta_1$ Tg mice was recorded as described. **A.** Representative traces of mEPSC from WT and $\alpha_2\delta_1$ Tg DDH neurons, respectively. **B** and **C.** Cumulative probability curves of mEPSC interevent intervals (**B**) and amplitude (**C**) from WT and Tg DDH neurons, respectively. $p = 0.001$ (k-s test) for a comparison of mEPSC interevent intervals between WT and Tg mice. $p = 0.53$ for a comparison of mEPSC amplitude between WT and Tg mice. **D** and **E.** Summary data of average mEPSC frequency (**D**) and amplitude (**E**), respectively, from WT and Tg DDH neurons. ** $p < 0.01$ compared with WT group by Student's t -test. **F.** Time courses for the effects of $100\mu\text{M}$ GBP on mEPSC frequency from WT and Tg L4 DDH neurons. **G.** Dose-dependent normalization of mEPSC frequency in L4 Tg neurons by GBP. ** $p < 0.01$ compared with pre-GBP treatment; # $p < 0.01$ compared with $10\mu\text{M}$ GBP treatment by one-way ANOVA test. Dotted

line represents percentage of mEPSC frequency in sham mice shown in D. Summarized data are shown as the means \pm SEM from the number of neurons indicated in the parentheses.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 4. mIPSC was not altered in DDH neurons from SNL and $\alpha_2\delta_1$ Tg mice
 mIPSC from L4 dorsal horn neurons of WT and $\alpha_2\delta_1$ Tg mice or injury side of sham and SNL mice 5–7 days post surgery was recorded as described. **A.** Representative traces of mIPSC from a sham and a SNL DDH neuron, respectively. **B, C.** Cumulative probability curves of mIPSC interevent intervals and amplitude, respectively, from sham and SNL mice. **D, E.** Summary of average mIPSC frequency (**D**) and amplitude (**E**), respectively, from DDH neurons of sham and SNL mice. **F, G.** Cumulative probability curves of mIPSC interevent intervals (**F**) and amplitude (**G**), respectively, from WT and $\alpha_2\delta_1$ Tg DDH neurons. **H, I.** Summary of average mIPSC frequency (**H**) and amplitude (**I**), respectively,

from WT and $\alpha_2\delta_1$ Tg DDH neurons. $p > 0.1$ (k-s test) for all the comparisons between sham and SNL as well as between WT and $\alpha_2\delta_1$ Tg mice.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript