



Published in final edited form as:

J Invest Dermatol. 2010 August ; 130(8): 2141–2144. doi:10.1038/jid.2010.74.

Is the Filaggrin–Histidine–Urocanic Acid Pathway Essential for Stratum Corneum Acidification?

Joachim W. Fluhr^{1,2,3}, Peter M. Elias^{1,4}, Mao-Qiang Man¹, Melanie Hupe^{1,4}, Clare Selden⁵, John P. Sundberg⁶, Erwin Tschachler⁷, Leopold Eckhart⁷, Theodora M. Mauro^{1,4}, and Kenneth R. Feingold^{1,4}

¹Dermatology Service, Veterans Affairs Medical Center, San Francisco, California, USA

²Bioskin, Berlin, Germany

³Department of Dermatology, Charité University Clinic, Berlin, Germany

⁴Department of Dermatology, University of California, San Francisco, San Francisco, California, USA

⁵Department of Medicine, Center for Hepatology, Royal Free and University College Medical School, UCL, London, UK

⁶The Jackson Laboratories, Bar Harbor, Maine, USA

⁷Department of Dermatology, University of Vienna, Austria

TO THE EDITOR

Acidification of the surface of the stratum corneum (SC), the acid mantle, was initially thought to be important in the defense against infection. The growth of pathogenic microorganisms, such as *Staphylococcus aureus* and *Streptococcus pyogenes*, is inhibited by an acidic skin pH whereas the growth of resident (normal) skin flora is stimulated (Puhvel *et al.*, 1975; Korting *et al.*, 1990, 1992). However, recent studies have shown that acidification of the SC has additional functions, including regulating several key SC functions. A major function of the skin is to form a permeability barrier between the dry external environment and the moist interior of the body (Elias, 2007). This permeability barrier resides in the extracellular lipid membranes of the SC, and studies have shown that an acidified SC is required for the formation of a functionally competent permeability barrier (Mauro *et al.*, 1998; Fluhr *et al.*, 2001; Hachem *et al.*, 2003). Specifically, in the SC β -glucocerebrosidase and acid sphingomyelinase metabolize glucosylceramides and sphingomyelin, respectively, to ceramides, which is the major family of lipids in the extracellular membranes that mediate permeability barrier function (Feingold, 2007). Both the enzymes require an acidic milieu for optimal enzymatic activity; hence, when the pH of the SC increases, the metabolism of

joachim.fluhr@gmx.net

CONFLICT OF INTEREST The authors state no conflict of interest in relation to this study except JS (who has a research contract with Procter and Gamble on an unrelated project).

SUPPLEMENTARY MATERIAL

Supplementary material is linked to the online version of the paper at <http://www.nature.com/jid>

glucosylceramides and sphingomyelin to ceramides is impaired, resulting in abnormal permeability barrier homeostasis (Holleran *et al.*, 1992, 1993; Feingold, 2007). In addition, an acidic SC pH inhibits the activity of serine proteases thereby maintaining the cohesiveness and integrity of the SC (Hachem *et al.*, 2005). With an increase in SC pH, the activities of these serine proteases are stimulated resulting in the degradation of corneodesmosomes and a decrease in SC integrity and cohesion (Fluhr *et al.*, 2004b; Hachem *et al.*, 2005). Thus, an acidic SC is important in regulating the metabolism and function of the SC, and alterations in SC pH could have numerous adverse effects.

A variety of different pathways are postulated to contribute to the acid mantle of the skin. Exogenous mechanisms, such as free fatty acids of pilosebaceous origin (Puhvel *et al.*, 1975; Bibel *et al.*, 1989), microbial metabolites (Di Marzio *et al.*, 1999), and eccrine gland-derived products, such as lactic acid (Ament *et al.*, 1997; Thueson *et al.*, 1998), are thought to decrease SC pH. Recent studies have shown that endogenous mechanisms also contribute to SC acidification (Fluhr *et al.*, 2001, 2004a; Behne *et al.*, 2002). Both free fatty acid generation from phospholipid hydrolysis catalyzed by one or more isoforms of sPLA₂ and a sodium/proton pump antiporter, the sodium/hydrogen antiporter-1 (Behne *et al.*, 2002, 2003; Fluhr *et al.*, 2004a), lower SC pH. A third endogenous mechanism, urocanic acid generation from histidine by the deiminating enzyme, histidase, has been shown to acidify SC *in vitro* (Krien and Kermici, 2000), but its impact on SC acidification and function *in vivo* remains uncertain. The histidase pathway of acidification is dependent upon previous proteolysis of filaggrin to histidine, a reaction triggered by a reduction in external humidity (Rawlings and Matts, 2005).

We therefore hypothesized that if the filaggrin–histidine–urocanic acid cascade is crucial in regulating SC pH *in vivo*, then SC pH should increase with either a reduction in substrate (filaggrin) and/or a decrease in histidase activity. Moreover, as SC acidification occurs over the first few days after birth in parallel with activation of filaggrin proteolysis (Fluhr *et al.*, 2004a), we reasoned that histidase activity should increase simultaneously with SC acidification.

Our initial studies examined histidase activity in newborn albino rats immediately after birth and at 4–5 days after birth (details regarding animals and methods are provided in Supplementary Material online). As reported previously, during this period of time the pH of the SC decreases markedly (Fluhr *et al.*, 2004a). If the filaggrin–histidine–urocanic acid cascade makes a major contribution to this reduction in SC pH, then the activity of histidase in the SC should increase in parallel. As shown in Figure 1a, there was no difference in SC histidase activity at day 0–1 after birth compared to 4–5 days after birth.

To determine the role of histidase in SC acidification more definitively, we next studied mice that were deficient in histidase activity, the Peruvian mouse (Selden *et al.*, 1995). As shown in Figure 1b, SC pH was similar in animals deficient in histidase activity, suggesting again that the histidase pathway is not essential for SC acidification.

We next studied animals deficient in filaggrin production due to impaired proteolytic processing of profilaggrin to filaggrin, the flaky tail (ft/ft) mouse (Presland *et al.*, 2000). As

shown in Figure 2a, SC pH was actually slightly decreased, not increased, in flaky tail mice, indicating that filaggrin is not essential for SC acidification. In addition, SC hydration (Figure 2b) is also not altered in the flaky tail mouse, suggesting that this additional function also does not absolutely require filaggrin.

We next asked whether other acidifying mechanisms are upregulated in ft/ft mice to compensate for reduced urocanic acid generation. As shown in Supplementary Figure S1 (Supplementary Material), the immunostainable protein content of both NHE1 and sPLA2A is increased in ft/ft mice whereas two other isoforms of sPLA2, sPLA2F and sPLA2X1, do not change (data not shown). These results suggest that compensatory upregulation of these alternate acidifying pathways could account for the normal to decreased SC pH in ft/ft mice.

Taken together, the above results do not provide support for the hypothesis that the filaggrin–histidine–urocanic acid cascade is essential for SC acidification. Increases in sPLA2 activity and/or NHE1 levels, or other yet to be elucidated mechanisms, appear to acidify the SC in the absence of the filaggrin–histidine–urocanic acid cascade. Yet, when either the sPLA2 or NHE1 pathways of acidification are compromised, the bulk pH of SC rises (Fluhr *et al.*, 2001, 2004a; Behne *et al.*, 2002), indicating that other acidifying mechanisms cannot compensate for them. One can easily envision that because of the importance of an acid mantle in SC function (Fluhr *et al.*, 2001, 2004b; Schmid-Wendtner and Korting, 2006) that numerous pathways contribute to its formation, and that the absence of any particular pathway will result in minor or no changes in SC pH. Thus, although both sPLA2 and NHE1 activities are required for the formation of the acid mantle (Fluhr *et al.*, 2001, 2004a; Behne *et al.*, 2002), the filaggrin–histidine–urocanic acid cascade is not essential for SC acidification.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

This study was supported by National Institutes of Health Grants AR39448, HD029706, and AR049932, and the Research Service, Department of Veterans Administration at San Francisco. The study has received institutional approval for animal experiments.

Abbreviation

SC stratum corneum

REFERENCES

- Ament W, Huizenga JR, Mook GA, et al. Lactate and ammonia concentration in blood and sweat during incremental cycle ergometer exercise. *Int J Sports Med.* 1997; 18:35–9. [PubMed: 9059903]
- Behne MJ, Barry NP, Hanson KM, et al. Neonatal development of the stratum corneum pH gradient: localization and mechanisms leading to emergence of optimal barrier function. *J Invest Dermatol.* 2003; 120:998–1006. [PubMed: 12787127]

- Behne MJ, Meyer JW, Hanson KM, et al. NHE1 regulates the stratum corneum permeability barrier homeostasis. Microenvironment acidification assessed with fluorescence lifetime imaging. *J Biol Chem.* 2002; 277:47399–406. [PubMed: 12221084]
- Bibel DJ, Miller SJ, Brown BE, et al. Antimicrobial activity of stratum corneum lipids from normal and essential fatty acid-deficient mice. *J Invest Dermatol.* 1989; 92:632–8. [PubMed: 2649598]
- Di Marzio L, Cinque B, De Simone C, et al. Effect of the lactic acid bacterium *Streptococcus thermophilus* on ceramide levels in human keratinocytes *in vitro* and stratum corneum *in vivo*. *J Invest Dermatol.* 1999; 113:98–106. [PubMed: 10417626]
- Elias PM. The skin barrier as an innate immune element. *Semin Immunopathol.* 2007; 29:3–14. [PubMed: 17621950]
- Feingold KR. The importance of lipids in cutaneous function. *J Lipid Res.* 2007; 48:2529–30. [PubMed: 18024706]
- Fluhr JW, Behne MJ, Brown BE, et al. Stratum corneum acidification in neonatal skin: secretory phospholipase A2 and the sodium/hydrogen antiporter-1 acidify neonatal rat stratum corneum. *J Invest Dermatol.* 2004a; 122:320–9. [PubMed: 15009712]
- Fluhr JW, Kao J, Jain M, et al. Generation of free fatty acids from phospholipids regulates stratum corneum acidification and integrity. *J Invest Dermatol.* 2001; 117:44–51. [PubMed: 11442748]
- Fluhr JW, Mao-Qiang M, Brown BE, et al. Functional consequences of a neutral pH in neonatal rat stratum corneum. *J Invest Dermatol.* 2004b; 123:140–51. [PubMed: 15191554]
- Hachem JP, Crumrine D, Fluhr J, et al. pH directly regulates epidermal permeability barrier homeostasis, and stratum corneum integrity/cohesion. *J Invest Dermatol.* 2003; 121:345–53. [PubMed: 12880427]
- Hachem JP, Man MQ, Crumrine D, et al. Sustained serine proteases activity by prolonged increase in pH leads to degradation of lipid processing enzymes and profound alterations of barrier function and stratum corneum integrity. *J Invest Dermatol.* 2005; 125:510–20. [PubMed: 16117792]
- Holleran WM, Takagi Y, Imokawa G, et al. beta-Glucocerebrosidase activity in murine epidermis: characterization and localization in relation to differentiation. *J Lipid Res.* 1992; 33:1201–9. [PubMed: 1431599]
- Holleran WM, Takagi Y, Menon GK, et al. Processing of epidermal glucosylceramides is required for optimal mammalian cutaneous permeability barrier function. *J Clin Invest.* 1993; 91:1656–64. [PubMed: 8473508]
- Korting HC, Hubner K, Greiner K, et al. Differences in the skin surface pH and bacterial microflora due to the long-term application of synthetic detergent preparations of pH 5.5 and pH 7.0. Results of a crossover trial in healthy volunteers. *Acta Derm Venereol.* 1990; 70:429–31. [PubMed: 1980979]
- Korting HC, Lukacs A, Vogt N, et al. Influence of the pH-value on the growth of *Staphylococcus epidermidis*, *Staphylococcus aureus* and *Propionibacterium acnes* in continuous culture. *Zentralbl Hyg Umweltmed.* 1992; 193:78–90. [PubMed: 1503605]
- Krien PM, Kermici M. Evidence for the existence of a self-regulated enzymatic process within the human stratum corneum—an unexpected role for urocanic acid. *J Invest Dermatol.* 2000; 115:414–20. [PubMed: 10951277]
- Mauro T, Holleran WM, Grayson S, et al. Barrier recovery is impeded at neutral pH, independent of ionic effects: implications for extracellular lipid processing. *Arch Dermatol Res.* 1998; 290:215–22. [PubMed: 9617442]
- Presland RB, Boggess D, Lewis SP, et al. Loss of normal profilaggrin and filaggrin in flaky tail (ft/ft) mice: an animal model for the filaggrin-deficient skin disease ichthyosis vulgaris. *J Invest Dermatol.* 2000; 115:1072–81. [PubMed: 11121144]
- Puhvel SM, Reisner RM, Sakamoto M. Analysis of lipid composition of isolated human sebaceous gland homogenates after incubation with cutaneous bacteria. Thin-layer chromatography. *J Invest Dermatol.* 1975; 64:406–11. [PubMed: 237966]
- Rawlings AV, Matts PJ. Stratum corneum moisturization at the molecular level: an update in relation to the dry skin cycle. *J Invest Dermatol.* 2005; 124:1099–110. [PubMed: 15955083]
- Schmid-Wendtner MH, Korting HC. The pH of the skin surface and its impact on the barrier function. *Skin Pharmacol Physiol.* 2006; 19:296–302. [PubMed: 16864974]

Selden C, Calnan D, Morgan N, et al. Histidinemia in mice: a metabolic defect treated using a novel approach to hepatocellular transplantation. *Hepatology*. 1995; 21:1405–12. [PubMed: 7537713]
Thueson DO, Chan EK, Oechsli LM, et al. The roles of pH and concentration in lactic acid-induced stimulation of epidermal turnover. *Dermatol Surg*. 1998; 24:641–5. [PubMed: 9648571]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 1. Role of histidase in SC acidification

(a) Histidase activity in the stratum corneum (SC) was measured in newborn rats and rats at age 4–5 days ($n = 6$). (b) The SC surface pH in Peruvian and matched controls was measured with a flat, glass surface electrode (Mettler-Toledo, Giessen, Germany), attached to a pH meter (Skin pH Meter PH 900; Courage & Khazaka, Cologne, Germany). HIS^{-/-}, $n = 8$; HIS^{+/-}, $n = 14$; wild type, $n = 6$.




Figure 2. Effect of filaggrin deficiency on SC function

(a) In flaky tail (ft/ft) mice, surface pH was measured with a flat, glass surface electrode as described in Figure 1 legend ($n = 16$; $P < 0.0001$). (b) Stratum corneum (SC) hydration was measured with corneometer (Corneometer CM 820; Courage & Khazaka); $n = 12$; t -test, NS.