



HHS Public Access

Author manuscript

J Acad Nutr Diet. Author manuscript; available in PMC 2016 September 01.

Published in final edited form as:

J Acad Nutr Diet. 2015 September ; 115(9): 1408–1416. doi:10.1016/j.jand.2015.01.011.

Eating and Weight-related Parenting of Adolescents in the Context of Food Insecurity

Katherine W. Bauer, PhD, MS,

Assistant Professor of Public Health, Center for Obesity Research and Education, Temple University, 3223 N. Broad Street, Suite 175, Philadelphia, PA 19140, Ph: 215-707-8659, Fax: 215-707-6462

Rich MacLehose, PhD,

Associate Professor, Division of Epidemiology and Community Health, School of Public Health, University of Minnesota, 1300 South Second Street Suite 300 Minneapolis, MN 55454, Ph: 612-624-1932, Fax: 612-626-7103

Katie A. Loth, PhD, MPH, RD,

Post-doctoral Fellow, Department of Psychiatry, University of Minnesota, F282/2A West 2450 Riverside Avenue South, Minneapolis, Minnesota 55454

Jennifer O. Fisher, PhD,

Professor of Public Health, Center for Obesity Research and Education, Temple University, 3223 N. Broad Street, Suite 175, Philadelphia, PA 19140, Ph: 215-707-0921, Fax: 215-707-6462

Nicole I. Larson, PhD, MPH, RDN, and

Senior Research Associate, Division of Epidemiology and Community Health, School of Public Health, University of Minnesota, 1300 South Second Street Suite 300 Minneapolis, MN 55454, Ph: 612-625-5881, Fax: 612-626-7103

Dianne Neumark-Sztainer, PhD, MPH, RD

Professor, Division of Epidemiology and Community Health, School of Public Health, University of Minnesota, 1300 South Second Street Suite 300 Minneapolis, MN 55454, Ph: 612-624-0880, Fax: 612-626-7103

Katherine W. Bauer: Katherine.Bauer@Temple.edu; Rich MacLehose: macl0029@umn.edu; Katie A. Loth: fall0075@umn.edu; Jennifer O. Fisher: JOFisher@Temple.edu; Nicole I. Larson: Larsonn@umn.edu; Dianne Neumark-Sztainer: neuma011@umn.edu

Abstract

Background—Food insecurity is hypothesized to influence mothers' use of parenting strategies to regulate children's eating. Little is known about the parenting practices directed toward adolescents in food insecure households.

Correspondence to: Katherine W. Bauer, Katherine.Bauer@Temple.edu.

Conflict of Interest Statement

No authors report a conflict of interest

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Objective—Examine differences in use of eating- and weight-related parenting practices among mothers of adolescents by household food security status.

Design—Cross-sectional

Participants/setting—A socio-demographically diverse sample of mothers and adolescents from the Minneapolis/St. Paul metropolitan area who participated in the EAT 2010 and Project F-EAT studies in 2009–2010 (dyad n=2,087). Seventy percent of mothers identified as non-white.

Main outcome measures—Mother-reported use of parenting practices including pressuring children to eat, restricting high-calorie foods, and encouraging dieting.

Statistical analyses performed—Logistic regression models were used to determine the predicted probabilities of parenting practices among food secure, low food secure, and very low food secure households. Socio-demographic characteristics, mothers' body mass index (BMI), and adolescents' BMI-for-age percentile were examined as confounders.

Results—In unadjusted models, food insecure mothers were more likely than food secure mothers to frequently encourage their children to diet, comment on their child's weight, be concerned about their child's weight, use restrictive feeding practices, and use pressured feeding practices. After adjustment for socio-demographic characteristics and mothers' and children's BMI, compared to food secure mothers, mothers with low food security were more likely to frequently comment on their son's weight (41.5% vs. 32.9%, prevalence difference (PD)=8.6 (0.9, 16.3)) and mothers with very low food security were more likely to be concerned about their son's weight (48.8% vs. 35.1%, PD=13.7 (3.5, 23.9)). Mothers with very low food security were more likely to frequently use restrictive feeding practices with their daughters compared to food secure mothers (33.0% vs. 20.5%, PD=12.4 (4.2, 20.7)).

Conclusions—Interventions to improve food insecure adolescents' eating behaviors may benefit from supporting mothers' use of health-promoting parenting practices.

Keywords

parenting; adolescents; food security; nutrition; obesity

Introduction

Parents play an important role in promoting healthful dietary intake, eating behaviors, and strategies to control weight among children and adolescents. For adolescents specifically, parenting practices such as encouraging and modeling healthful eating have been associated with higher dietary quality and more healthful weight,^{1–4} while use of practices such as excessively restricting food access, encouraging youth to diet for weight control, and engaging in weight talk and teasing have been associated with use of maladaptive weight control and disordered eating behaviors, as well as higher weight, among adolescents.^{5–11} Despite the increasing appreciation of the role of families and parenting practices to youth's eating- and weight-related outcomes, little is known about how the use of these parenting practices may be influenced by families' social, cultural, or economic contexts.^{12,13} Identifying the contextual factors that produce or maintain use of specific parenting practices can provide insight into the mechanisms via which socio-economic inequalities

affect children's behavior and inform family-based interventions that aim to improve youth's dietary intake and reduce risk of disordered eating and obesity.

Food insecurity, or having limited or uncertain access to safe, nutritionally adequate, and culturally- acceptable food,¹⁴ has been associated with numerous poor mental and physical health outcomes.¹⁵ Food insecure youth have less healthful dietary intake and are at greater risk for nutrient deficiencies than food secure youth.¹⁶⁻²⁵ Food insecurity has also been associated with a higher prevalence of obesity among children and adolescents in some,²⁵⁻²⁸ but not all,^{17,29-31} studies. Parenting practices are increasingly being recognized as mediators of the relationship between food insecurity and child-level nutritional and health outcomes.³² Parents experiencing food insecurity use a number of strategies to stretch food resources and ensure that family members, especially young children, do not experience hunger.³³⁻³⁷ For example, mothers who are either currently food insecure or have a history of food insecurity are less likely to adhere to recommended infant feeding practices, breastfeed for shorter duration, and are more likely to over-feed their young children or provide lax monitoring of sweets and snack intake.³³⁻³⁶ While the majority of studies have focused on mothers of children preschool age and younger, Feinberg, et al.³⁷ observed that among mothers of children ages 2 to 13, food insecurity was associated with greater use of high-calorie supplements and appetite stimulants with children. It has been suggested that mothers use these feeding strategies out of concern for children's under nutrition and as a way to treat or appease children.³⁸

Nationwide, adolescents are twice as likely to experience food insecurity and four times as likely to experience very low food security, a more extreme level of food insecurity, compared to young children.³⁹ While young children are protected from food insecurity in many families, adolescents are often are not afforded the same protection, perhaps due to beliefs that adolescents have the ability to obtain food easily outside the home or can better withstand hunger.^{17,40} This suggests that eating- and weight-related parenting of adolescents in food insecure homes are different from what has been observed among parents of young children. For example, parents of adolescents may be more likely to encourage dietary restriction in an effort to conserve household food rather than encouraging over-eating. However, to our knowledge, no studies have examined eating- and weight-related parenting practices directed toward adolescents among food insecure families.

Given this knowledge gap, the current study utilizes data from EAT (Eating and Activity Among Teens) 2010 and Project F-EAT (Families and Eating and Activity among Teens), two complementary studies of middle and high school-aged adolescents and their parents/ caregivers. The primary aim was to examine differences the eating- and weight-related parenting practices that mothers use with their adolescent-aged children among food secure versus insecure families. We hypothesize that mothers from families experiencing food insecurity will be more likely to use parenting practices that encourage dietary restriction and dieting, including commenting on their child's weight, as compared to mothers from food secure families.

Methods

Cross-sectional data were drawn from the EAT 2010 and Project F-EAT studies. EAT 2010 is an observational investigation of socio-ecological correlates of eating, physical activity, and weight-related topics among a diverse sample of 2,793 adolescents from 20 public middle and high schools in the Minneapolis/St. Paul metropolitan area of Minnesota. Students from selected health, physical education, and science classes were recruited to participate. Interested participants completed surveys and had their height and weight measured by trained research staff using standardized procedures during the school day. During participation in EAT 2010, adolescents provided contact information for up to two of their parents/guardians. These parents/guardians were then invited to participate in Project F-EAT, a survey-based study of parents that aimed to learn more about the food, physical activity and home environments of the young people participating in EAT 2010. Data collection for Project F-EAT occurred via mail and phone between October 2009 and October 2010 and was conducted by the Wilder Research Foundation. The parent response rate in Project F-EAT was high; 77.6% of invited parents (n=3,709) completed study measures. The University of Minnesota Institutional Review Board approved the study protocol; all parents provided written informed consent and children provided written assent. Additional information about EAT 2010 and Project F-EAT have been reported elsewhere.^{41–43} Given interest in parenting around eating and weight, only those data from female parents/guardians participating in Project F-EAT who reported living with their child at least 50% of the time were included in the current analytic sample. For families in which two female parents/guardians responded (e.g. two moms, mom and grandma, etc.), an algorithm was utilized that took into account parent/child relationship with preference for biological parent over stepparent and parent over grandparent. The final analytic sample consisted of 2,087 dyads of female parents/guardians, henceforth referred to as mothers, and their adolescent-aged children.

Measures

Eating- and weight-related parenting practices and household food security status were measured among mothers as part of Project F-EAT. Project F-EAT survey items were drawn from several sources including items developed and previously validated for earlier Project EAT surveys⁴⁴ and measures developed and tested by other researchers. After an initial draft of the Project FEAT survey was developed, steps were undertaken to assess face validity of included items, ensure the questions were appropriate for the intended population, and make adjustments to the overall length of the survey. The draft survey was reviewed by content experts, bi-cultural staff from the Wilder Research Foundation, and field tested with 28 socioeconomically and ethnically/racially diverse parents. Finally, test–retest reliability for survey items reported below was assessed using an additional sample of 102 parents who completed the survey twice over two weeks.

Food security

Food security was reported by mothers using the 6-item US Household Food Security Survey Module adapted for self-administration, which assesses household food security over the past 12 months. This scale has been shown to correctly classify 97.7% of families when

compared with the full 18-item scale included in the Current Population Survey.^{45,46} Parents who responded in the affirmative to fewer than 2 items were categorized as experiencing “food security,” those that responded in the affirmative to 2 to 4 items were categorized as experiencing “low food security,” and those that responded in the affirmative to 5 or more items were categorized as experiencing “very low food security.” Two-week percent agreement for household food security (food secure vs. insecure) = 0.91 and kappa = 0.64 for three levels of food security (food secure, low food secure, very low food secure).

Eating- and Weight-Related Parenting Practices

Frequent encouragement to diet—Encouragement to diet was assessed with the following question: “To what extent do you encourage your child to diet to control his/her weight?” Four response options ranged from ‘not at all’ to ‘very much’.⁴⁷ Two week test-retest for this item was $r=0.68$. Mothers were identified as frequently encouraging their child to diet if they responded ‘somewhat’ or ‘very much’ to this item.

Frequent encouragement for healthy eating—Encouragement for healthy eating was assessed with the following question: “How often in the past year have you had a conversation with your child about healthy eating habits?” Response options included, ‘Never/rarely’, ‘A few times a year’, ‘A few times a month’, ‘A few times a week’, and ‘Almost every day’ (two-week test-retest $r=0.48$) This item was adapted from similar items used in the Parental Energy Index.⁴⁸ Frequent encouragement of healthy eating was defined as encouraging healthful eating a few times a month or more often.

Frequent comments about child’s weight—Comments about child’s weight were assessed with the following two questions: “How often in the past year have you had a conversation with your child about his/her weight or size?” (two-week test-retest $r=0.59$) and “Have you mentioned to your child that he/she weighs too much?” (two-week test-retest $r=0.73$) Response options for both included, ‘Never/rarely’, ‘A few times a year’, ‘A few times a month’, ‘A few times a week’, and ‘Almost every day’. This item was adapted from similar items used in the Parental Energy Index.⁴⁸ Parents were identified as frequently commenting on their child’s weight if they responded ‘a few times a month’ or more often to either of these two questions.

Concern about child’s weight—Concern about child’s weight was measured with one question: “How concerned are you about your child’s weight?” Response options included, ‘Not at all concerned’, ‘A little concerned’, ‘Quite concerned’, and ‘Very concerned’. This item was originally developed for the Nepean Kids Growing Up Parent Questionnaire⁴⁹ and two week test-retest was 0.68. Parents were identified as being concerned about their child’s weight if they responded that they were ‘Quite concerned’ or ‘Very concerned’.

Restrictive feeding practices—Restrictive feeding practices were measured using six items from the eight-item Restriction Subscale of the Child Feeding Questionnaire (CFQ).⁵⁰ Two items were dropped from the original subscale based on recommendations from a validation study conducted within a diverse population of parents of adolescents⁵¹ Test-retest among Project F-EAT participants was $r = 0.72$ and Cronbach’s $\alpha = 0.86$. Agreement

with individual items was assessed using a 4-point Likert scale, with anchors to indicate strength of agreement or disagreement. Scores across the 6 items were averaged and mothers who scored in the highest quartile of responses on the Restriction Subscale were defined as using a high level of restrictive feeding practices.

Pressured feeding practices—Pressured feeding practices were measured using the full four-item Pressure-to-Eat Subscale of the CFQ.⁵⁰ Test-retest among Project F-EAT participants was $r = 0.73$ and Cronbach's Alpha = 0.70. The scale is scored from 1 (low pressure) to 4 (high pressure). Individual items were measured using a 4-point Likert scale, with anchors to indicate strength of agreement or disagreement. Scores across the 4 items were averaged and mothers who scored in the highest quartile of responses on the pressure-to-eat subscale were defined as using a high level of pressured feeding practices.

Socio-demographic characteristics—*Household income* was assessed with one item asked of the mothers: "What was the total income of your household before taxes in the past year?" Six response option categories were offered: "less than \$20,000", "\$20,000 to \$34,999", "\$35,000–\$49,000", "\$50,000 to \$74,999", "\$75,000 – \$99,999", and "\$100,000 or more" (two-week test-retest agreement = 74%). Due to the small number of mothers who reported a household income above \$75,000, the highest two response options were collapsed. Mothers were also asked how many children under the age of 18 lived in their household (two-week test-retest $r = 0.99$). *Maternal educational attainment* was assessed with the question: "What is the highest grade or year of school that you have completed?" Response options included "Did not finish high school", "Finished high school or got GED", "Some college or training after high school", "Finished college" and "Advanced degree". (two-week test-retest agreement = 84%). *Maternal employment status* was assessed with one item: "Which of the following best describes your current work situation?" Five response options were available: working full-time, working part-time, stay-at-home caregiver, currently unemployed but actively seeking work, and not working for pay (two-week test-retest agreement = 82%). *Mothers' race/ethnicity* was assessed by the following item: "Do you think of yourself as: 1) White; 2) Black or African American; 3) Hispanic or Latino; 4) Asian American; 5) Hawaiian or Pacific Islander; 6) American Indian or Native American; and 7) Other." If a mother selected "Other" there was a space to fill in the racial/ethnic category with which they identified. Mothers were given the option to choose more than one category, and those with multiple responses were coded as "mixed/other" for analyses. (two-week test-retest agreement = 99%). Adolescents' gender and age were determined by their birthdate as self-reported on the EAT 2010 survey and the date of survey administration.

Maternal body mass index (BMI)—Mothers' height and weight were assessed by self-report. Adult participants were asked to report their height to the nearest feet and inches and their weight to the nearest pound on the Project F-EAT survey. Self-reported height and weight has been shown to be highly correlated with objectively measured values in adults.⁵² BMI was calculated using the formula weight in kilograms divided by height in meters squared (two-week test-retest = 0.97 for height, 0.95 for weight).

Adolescent BMI percentile—Adolescents' height and weight were measured as part of the EAT 2010 study. Height to the nearest 0.1cm and weight to the nearest 0.1kg were assessed in a private area at schools by trained research staff using standardized equipment and procedures. Age- and gender-specific percentiles were based on the 2000 CDC Growth Charts.⁵³

Statistical Analysis

Crude demographic and weight related variables were compared across level of food security (food security, low food security and very low food security) using chi square and F-tests. To assess whether the prevalence of eating- and weight-related parenting practices differed by level of food security, logistic regression models were used. The three-level food security variable was modeled using indicator variables and included as the main predictor. Separate models were fit for each of the 6 dependent variables. For each outcome, we fit an unadjusted, crude, regression (Table 2, Model 1), a model adjusting for socio-demographic characteristics (Tables 3/4, Model 2), and a model additionally adjusted for mothers' BMI, and children's BMI percentile (Tables 3/4, Model 3). Individuals missing data for the outcome or any covariates were dropped from the analysis. Analyses were stratified by adolescent gender to allow estimation of separate associations of food security with eating and weight-related parenting practices for boys and girls. Following each of the logistic models, the adjusted prevalence of the dependent variable (parenting practice) was computed for each level of food security. The associations between food security and parenting practices were calculated as the difference between these prevalences (using food secure as the reference group) with corresponding 95% confidence intervals. All analyses were performed in Stata (version 13, 2013, StataCorp LP).

Results

Among mothers participating in Project F-EAT, 60.2% reported that they were food secure, 26.0% reported low food security, and 13.9% reported very low food security within the past year (Table 1). Differences in level of food security were observed by mothers' race/ethnicity, employment status, household income, educational attainment, and mean number of children in the home (all $p < 0.01$). Mothers' mean BMI was positively associated with food insecurity ($p < 0.01$) and a statistically significant positive association was observed between level of food insecurity and BMI percentile among boys ($p = 0.02$), but not girls ($p = 0.15$).

In unadjusted analyses, several differences in the frequent use of eating- and weight-related parenting practices among mothers with boys in the EAT 2010 study were observed by food security status (Table 2). For example, 30.5% of mothers with low food security and 31.3% of mothers with very low food security reported frequently using restrictive feeding practices with their sons, compared to 21.9% of food secure mothers (prevalence difference (PD) for food secure vs. low food secure = 8.6 (95% CI = 1.9, 15.3), PD for food secure vs. very low food secure = 9.4 (95% CI = 7.0, 18.0). After adjustment for socio-demographic characteristics (Table 3, Model 2), some of these differences became non-significant. Additional adjustment for mothers' BMI and sons' BMI percentile did not greatly alter the

relationships between food security and parenting practices. Compared to food secure mothers, mothers experiencing low food security were more likely to frequently comment on their son's weight (41.5% vs. 32.9%, $PD = 8.6$ (95% $CI = 0.9, 16.3$)) and mothers with very low food security were more likely to be concerned about their sons' weight (48.5% vs. 35.1%, $PD = 13.7$ (95% $CI = 3.5, 23.9$)).

Similar to mothers of boys, food insecure mothers with adolescent girls in EAT 2010 were more likely to report using several eating- and weight-related parenting practices which have been correlated with poor outcomes, as compared to food-secure mothers in unadjusted models (Table 2). After adjustment for socio-demographic characteristics (Table 4, Model 2) some of these associations remained. However, after additional adjustment for maternal BMI and daughters' BMI percentile (Table 4, Model 3) only the association between food security status and restrictive feeding practices remained. Among mothers of daughters, those who were very low food secure were more likely to frequently use restrictive feeding practices compared to food secure mothers (33.0% vs. 20.5%, $PD = 12.4$ (95% $CI = 4.2, 20.7$)). Among both mothers of sons and daughters, no associations were observed between food security status and encouragement of healthful eating in the crude or adjusted models.

Discussion

It has been suggested that food insecurity influences parents' use of specific parenting practices to ensure that food resources are appropriately distributed among families and specifically, often to ensure that young children have sufficient amounts to eat and do not experience hunger.³² In the current study that drew from a socio-demographically diverse, urban population, a large proportion of families reported being food insecure. Several significant differences were observed in mothers' use of eating- and weight-related parenting practices by families' food security status. Mothers from households experiencing low or very low food security were more likely to report that they engage in parenting practices that have been linked to higher child BMI and disordered eating, such as encouraging children to diet, frequently commenting on children's weight, and using restrictive feeding practices.^{8,9,47} Several of these associations became non-significant after adjustment for socio-demographic characteristics and mothers' and adolescents' BMI. However, food insecure mothers remained more likely to be concerned about and comment on their sons' weight and use restrictive feeding practices with their daughters.

Findings from both the crude and adjusted models demonstrate that mothers in food insecure households use different eating- and weight-related parenting practices than mothers from food secure families. However, some of these differences may be explained by other economic, social, or cultural factors that food insecure families are more likely to experience than food secure families. Among mothers in Project F-EAT, the parenting practices used by those experiencing food insecurity reflect a concern of excessive intake and overweight among their children. This runs counter to research conducted among parents of younger children, which has found that food insecure parents are more likely to encourage over-eating or use pressured eating practices with their children.^{34,36,37}

Understanding the broader family food context, in addition to specific parenting practices used, is important to identifying the mechanisms through which food insecurity influences children's dietary and health outcomes. In an earlier analysis of data from Project F-EAT by Bruening, et al.,⁴³ food insecure parents reported several characteristics of the family environment that promote less healthful eating and may increase children's obesity risk. For example, families with low or very low food security reported more frequent fast food for family meals; less frequent inclusion of salad, vegetables, and fruit for family meals; and an increased frequency of serving sugar-sweetened beverages with meals. Additionally, food insecure parents reported poorer dietary quality themselves and a greater likelihood of binge eating. These family environment characteristics, in combination with use of parenting practices that encourage dieting and dietary restriction, may increase food insecure adolescents' risk of dietary inadequacy, disordered eating, and obesity.

The current study has a number of strengths including a large sample size with a sufficient number of participants reporting low and very low food security, which provided the ability to examine differences between these two food insecure groups and food secure families. Data were collected from diverse, urban families in the Midwest, therefore findings are generalizable to similar populations. Additionally, several socio-demographic characteristics were measured in Project F-EAT allowing for covariate adjustment of these variables. A limitation of the study is that both food security and parenting practices were self-reported by mothers, therefore social desirability bias may be introduced by this assessment method. Further, while the 6-item measure of household food security has demonstrated very high validity compared to the gold standard 18-item U.S. Household Food Security Survey Module,⁴⁵ minor misclassification of food insecure households as food secure may have occurred, which likely would bias results toward the null. Additionally, several comparisons were tested in this study, increasing the chance that the statistically significant associations were observed by chance. Finally, this study was descriptive in nature and cross-sectional in design, therefore causality cannot be inferred. Future longitudinal research that follows children through childhood and includes children's perspectives on their families' food security would provide greater insight into how food scarcity alters both parental and child behavior.

In conclusion, several eating- and weight-related parenting practices differ between food secure and insecure mothers and specifically, food insecure mothers were more likely to comment on and be concerned about their son's weight and use restrictive feeding practices with their daughters, even after accounting for socio-economic factors and differences in BMI among food secure and insecure mothers and adolescents. In concert with prior findings from Project F-EAT regarding the home food environment of food insecure families, the current study suggests that the family social environment during the adolescent period may play a role in the development of poor eating behaviors and obesity among food insecure adolescents. Given evidence that mothers from food insecure families are more likely to use maladaptive eating- and weight-related parenting practices with their adolescent-aged children, addressing maternal behavior in the context of interventions that address healthful eating and weight control among food insecure families may increase the likelihood that such programs lead to sustainable dietary intake, eating behavior, and/or weight changes.

Acknowledgments

Funding/Support Disclosure:

EAT 2010 and Project F-EAT were supported by grants R01HL084064 and R01HL093247-02 from the National Heart, Lung, and Blood Institute, National Institutes of Health.

References

1. Berge JM, Wall M, Bauer KW, Neumark-Sztainer D. Parenting characteristics in the home environment and adolescent overweight: a latent class analysis. *Obesity*. 2010; 18(4):818–825. [PubMed: 19816417]
2. Pearson N, Biddle SJ, Gorely T. Family correlates of fruit and vegetable consumption in children and adolescents: a systematic review. *Public Health Nutr*. Feb; 2009 12(2):267–283. [PubMed: 18559129]
3. Savage JS, Fisher JO, Birch LL. Parental influence on eating behavior: conception to adolescence. *J Law Med Ethics*. Spring;2007 35(1):22–34. [PubMed: 17341215]
4. Bauer KW, Neumark-Sztainer D, Fulkerson JA, Hannan PJ, Story M. Familial correlates of adolescent girls' physical activity, television use, dietary intake, weight, and body composition. *Int J Behav Nutr Phys Act*. 2011; 8:25. [PubMed: 21453516]
5. Field AE, Carmego C, Taylor CB, Berkey CS, Roberts SB, Colditz GA. Peer, parent, and media influences on the development of weight concerns and frequent dieting among preadolescent and adolescent girls and boys. *Pediatrics*. 2001; 107(1):54–60. [PubMed: 11134434]
6. Galloway AT, Fiorito LM, Francis LA, Birch LL. 'Finish your soup': counterproductive effects of pressuring children to eat on intake and affect. *Appetite*. May; 2006 46(3):318–323. [PubMed: 16626838]
7. Bauer KW, Berge JM, Neumark-Sztainer D. The importance of families to adolescents' physical activity and dietary intake. *Adolesc Med State Art Rev*. Dec; 2011 22(3):601–613. xiii. [PubMed: 22423466]
8. Loth KA, MacleHose RF, Fulkerson JA, Crow S, Neumark-Sztainer D. Are food restriction and pressure-to-eat parenting practices associated with adolescent disordered eating behaviors? *Int J Eat Disord*. Sep 18.2013
9. Loth KA, MacLeHose RF, Fulkerson JA, Crow S, Neumark-Sztainer D. Food-related parenting practices and adolescent weight status: a population-based study. *Pediatrics*. May; 2013 131(5):e1443–1450. [PubMed: 23610202]
10. Berge JM, MacleHose R, Loth KA, Eisenberg M, Bucchianeri MM, Neumark-Sztainer D. Parent conversations about healthful eating and weight: associations with adolescent disordered eating behaviors. *JAMA Pediatr*. Aug 1; 2013 167(8):746–753. [PubMed: 23797808]
11. Bauer KW, Bucchianeri MM, Neumark-Sztainer D. Mother-reported parental weight talk and adolescent girls' emotional health, weight control attempts, and disordered eating behaviors. *J Eat Disord*. 2013; 1:45. [PubMed: 24999423]
12. Faith MS, Van Horn L, Appel LJ, et al. Evaluating parents and adult caregivers as "Agents of Change" for treating obese children: Evidence for parent behavior change strategies and research Gaps: A scientific statement from the American Heart Association. *Circulation*. 2012; 125(9): 1186–1207. [PubMed: 22271754]
13. Davison KK, Jurkowski JM, Lawson HA. Reframing family-centred obesity prevention using the Family Ecological Model. *Public Health Nutr*. 2012:1–9. [PubMed: 23294865]
14. Anderson SA. Core indicators of nutritional state for difficult-to-sample populations. *J Nutr*. Nov; 1990 120 (Suppl 11):1559–1600. [PubMed: 2243305]
15. Cook JT, Frank DA. Food security, poverty, and human development in the United States. *Ann N Y Acad Sci*. 2008; 1136:193–209. [PubMed: 17954670]
16. Bauer KW, Widome R, Himes JH, et al. High food insecurity and its correlates among families living on a rural American Indian Reservation. *Am J Public Health*. 2012; 102(7):1346–1352. [PubMed: 22594740]

17. Bhattacharya J, Currie J, Haider S. Poverty, food insecurity, and nutritional outcomes in children and adults. *J Health Econ.* Jul; 2004 23(4):839–862. [PubMed: 15587700]
18. Dave JM, Evans AE, Saunders RP, Watkins KW, Pfeiffer KA. Associations among food insecurity, acculturation, demographic factors, and fruit and vegetable intake at home in Hispanic children. *J Am Diet Assoc.* Apr; 2009 109(4):697–701. [PubMed: 19328265]
19. Eicher-Miller HA, Mason AC, Weaver CM, McCabe GP, Boushey CJ. Food insecurity is associated with iron deficiency anemia in US adolescents. *Am J Clin Nutr.* Nov; 2009 90(5):1358–1371. [PubMed: 19776137]
20. Eicher-Miller HA, Mason AC, Weaver CM, McCabe GP, Boushey CJ. Food insecurity is associated with diet and bone mass disparities in early adolescent males but not females in the United States. *J Nutr.* Sep; 2011 141(9):1738–1745. [PubMed: 21795427]
21. Grutzmacher S, Gross S. Household food security and fruit and vegetable intake among low-income fourth-graders. *J Nutr Educ Behav.* Nov-Dec; 2011 43(6):455–463. [PubMed: 21855417]
22. Kaiser LL, Melgar-Quinonez HR, Lamp CL, Johns MC, Sutherlin JM, Harwood JO. Food security and nutritional outcomes of preschool-age Mexican-American children. *J Am Diet Assoc.* Jul; 2002 102(7):924–929. [PubMed: 12146552]
23. Kirkpatrick SI, Tarasuk V. Food insecurity is associated with nutrient inadequacies among Canadian adults and adolescents. *J Nutr.* Mar; 2008 138(3):604–612. [PubMed: 18287374]
24. Sharkey JR, Nalty C, Johnson CM, Dean WR. Children's very low food security is associated with increased dietary intakes in energy, fat, and added sugar among Mexican-origin children (6–11 y) in Texas border Colonias. *BMC Pediatrics.* 2012; 12:16. [PubMed: 22348599]
25. Widome R, Neumark-Sztainer D, Hannan PJ, Haines J, Story M. Eating when there is not enough to eat: eating behaviors and perceptions of food among food-insecure youths. *Am J Public Health.* May; 2009 99(5):822–828. [PubMed: 19299675]
26. Casey PH, Simpson PM, Gossett JM, et al. The association of child and household food insecurity with childhood overweight status. *Pediatrics.* Nov; 2006 118(5):e1406–1413. [PubMed: 17079542]
27. Jyoti DF, Frongillo EA, Jones SJ. Food insecurity affects school children's academic performance, weight gain, and social skills. *J Nutr.* Dec; 2005 135(12):2831–2839. [PubMed: 16317128]
28. Dubois L, Farmer A, Girard M, Porcherie M. Family food insufficiency is related to overweight among preschoolers. *Soc Sci Med.* 2006; 63(6):1503–1516. [PubMed: 16777308]
29. Bhargava A, Jolliffe D, Howard LL. Socio-economic, behavioural and environmental factors predicted body weights and household food insecurity scores in the Early Childhood Longitudinal Study-Kindergarten. *Br J Nutr.* Aug; 2008 100(2):438–444. [PubMed: 18275621]
30. Rose D, Bodor JN. Household food insecurity and overweight status in young school children: results from the Early Childhood Longitudinal Study. *Pediatrics.* Feb; 2006 117(2):464–473. [PubMed: 16452367]
31. Gundersen C, Lohman BJ, Eisenmann JC, Garasky S, Stewart SD. Child-specific food insecurity and overweight are not associated in a sample of 10- to 15-year-old low-income youth. *J Nutr.* Feb; 2008 138(2):371–378. [PubMed: 18203906]
32. Nord M. What have we learned from two decades of research on household food security? *Public Health Nutr.* 2014; 17(01):2–4.
33. Kuyper EM, Smith D, Kaiser LL. Does food insecurity influence child feeding practices? *J Hunger Environ Nutr.* 2009; 4(2):147–157.
34. Kaufman L, Karpati A. Understanding the sociocultural roots of childhood obesity: food practices among Latino families of Bushwick, Brooklyn. *Soc Sci Med.* Jun; 2007 64(11):2177–2188. [PubMed: 17383060]
35. Bronte-Tinkew J, Zaslow M, Capps R, Horowitz A, McNamara M. Food insecurity works through depression, parenting, and infant feeding to influence overweight and health in toddlers. *J Nutr.* 2007; 137(9):2160–2165. [PubMed: 17709458]
36. Gross RS, Mendelsohn AL, Fierman AH, Racine AD, Messito MJ. Food insecurity and obesogenic maternal infant feeding styles and practices in low-income families. *Pediatrics.* Aug; 2012 130(2):254–261. [PubMed: 22826569]

37. Feinberg E, Kavanagh PL, Young RL, Prudent N. Food insecurity and compensatory feeding practices among urban black families. *Pediatrics*. Oct; 2008 122(4):e854–860. [PubMed: 18829783]
38. Rosas LG, Harley K, Fernald LC, et al. Dietary associations of household food insecurity among children of Mexican descent: results of a binational study. *J Am Diet Assoc*. Dec; 2009 109(12): 2001–2009. [PubMed: 19942017]
39. Nord, M. Food Insecurity in Households with Children: Prevalence, Severity, and Household Characteristics. U.S. Dept. of Agriculture, Economic Research Service; Sep. 2009
40. Fram MS, Frongillo EA, Jones SJ, et al. Children are aware of food insecurity and take responsibility for managing food resources. *J Nutr*. Jun; 2011 141(6):1114–1119. [PubMed: 21525257]
41. Berge JM, MacLehose RF, Loth KA, Eisenberg ME, Fulkerson JA, Neumark-Sztainer D. Parent-adolescent conversations about eating, physical activity and weight: prevalence across sociodemographic characteristics and associations with adolescent weight and weight-related behaviors. *J Behav Med*. Jul 6.2014
42. Bauer KW, Hearst MO, Escoto K, Berge JM, Neumark-Sztainer D. Parental employment and work-family stress: associations with family food environments. *Soc Sci Med*. 2012; 75(3):496–504. [PubMed: 22591825]
43. Bruening M, MacLehose R, Loth K, Story M, Neumark-Sztainer D. Feeding a family in a recession: food insecurity among Minnesota parents. *Am J Public Health*. Mar; 2012 102(3):520–526. [PubMed: 22390517]
44. [Accessed December 7, 2014] Project EAT. 2014. <http://www.sphresearch.umn.edu/epi/project-eat/>
45. Blumberg SJ, Bialostosky K, Hamilton WL, Briefel RR. The effectiveness of a short form of the Household Food Security Scale. *Am J Public Health*. Aug; 1999 89(8):1231–1234. [PubMed: 10432912]
46. Bickel, G.; Nord, M.; Hamilton, W. US Household Food Security Survey Module: Three-Stage Design With Screeners. Washington, DC: US Department of Agriculture, Economic Research Service; 2006.
47. Fulkerson JA, McGuire MT, Neumark-Sztainer D, Story M, French SA, Perry CL. Weight-related attitudes and behaviors of adolescent boys and girls who are encouraged to diet by their mothers. *Int J Obes Relat Metab Disord*. 2002; 26(12):1579–1587. [PubMed: 12461674]
48. Lytle LA, Birnbaum A, Boutelle K, Murray DM. Wellness and risk communication from parent to teen: the Parental Energy Index. *Health Educ*. 1999; 99(5):207–214.
49. Campbell KJ, Crawford DA, Salmon J, Carver A, Garnett SP, Baur LA. Associations between the home food environment and obesity-promoting eating behaviors in adolescence. *Obesity*. 2007; 15(3):719–730. [PubMed: 17372323]
50. Birch LL, Fisher JO, Grimm-Thomas K, Markey CN, Sawyer R, Johnson SL. Confirmatory factor analysis of the Child Feeding Questionnaire: a measure of parental attitudes, beliefs and practices about child feeding and obesity proneness. *Appetite*. Jun; 2001 36(3):201–210. [PubMed: 11358344]
51. Kaur H, Li C, Nazir N, et al. Confirmatory factor analysis of the child-feeding questionnaire among parents of adolescents. *Appetite*. Jul; 2006 47(1):36–45. [PubMed: 16624444]
52. Kuczmarski MF, Kuczmarski RJ, Najjar M. Effects of age on validity of self-reported height, weight, and body mass index: findings from the Third National Health and Nutrition Examination Survey, 1988–1994. *J Am Diet Assoc*. Jan; 2001 101(1):28–34. quiz 35–26. [PubMed: 11209581]
53. Kuczmarski RJ, Ogden CL, Guo SS, et al. 2000 CDC Growth Charts for the United States: methods and development. *Vital Health Stat*. 2002; 11(246):1–190.

What is the current knowledge on this topic?

Parenting practices are important to children’s dietary quality, eating behaviors, and obesity risk. Little is known about how food insecurity may influence parents’ use of eating and weight-related parenting practices.

How does this research add to knowledge on this topic?

Food insecure mothers of adolescent-aged children reported more frequent use of several eating and weight-related parenting practices, including encouraging dieting and restricting food, as compared to food secure mothers.

How might this knowledge impact current dietetics practice?

Interventions to improve dietary quality and reduce obesity risk among food insecure families may benefit from addressing parenting practices.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

Food security status by socio-demographic characteristics and body mass index (BMI) of mothers and adolescents participating in Project F-EAT and EAT 2010

	Food Secure	Low Food Secure	Very Low Food Secure	P
TOTAL, % (n)	60.2 (1256)	26.0 (542)	13.9 (289)	
Household income, %				
< \$20,000	26.7	49.7	59.1	<0.01
\$20,000 – \$34,999	21.0	25.5	25.2	
\$35,000 – \$49,999	16.8	15.4	10.8	
\$50,000 – \$74,999	16.0	6.6	4.2	
> \$75,000	19.5	2.7	0.7	
No. of children in home, mean (SD)	2.6 (1.7)	3.2 (1.9)	2.9 (1.5)	<0.01
Mothers' Education, %				
Did not complete high school	25.7	39.7	31.5	<0.01
Completed high school	20.8	21.1	19.9	
Some college	25.0	26.5	35.0	
Completed college	19.7	10.5	12.6	
Post-graduate education	8.9	2.3	1.1	
Mothers' Employment Status, %				
Working full time	51.7	39.4	32.8	<0.01
Working part time	18.6	14.8	16.6	
Stay at home caregiver	11.5	19.9	16.2	
Unemployed but seeking work	7.8	14.2	15.9	
Not working for pay	10.4	11.7	18.7	
Mothers' Race/ethnicity, %				
White	35.8	20.1	26.3	<0.01
African American/black	27.8	28.0	29.8	
Hispanic	17.5	17.9	11.6	
Asian	14.8	27.8	21.8	
Mixed/other	4.1	6.2	10.5	
Mothers' BMI, mean	28.0 (6.1)	28.7 (6.1)	30.4 (7.5)	<0.01
Adolescent Age, mean (SD)	14.5 (2.0)	14.3 (1.9)	14.4 (1.9)	0.20
Adolescent BMI percentile, mean (SD)				
Boys	65.8 (0.3)	72.0 (0.3)	70.2 (0.3)	0.02
Girls	68.6 (0.3)	69.6 (0.3)	73.1 (0.3)	0.15

Table 2 Unadjusted associations between use of eating- and weight-related parenting practices and food security status by gender (Model 1)

	Boys				Girls					
	Prevalence			Prevalence differences (95% CI)		Prevalence		Prevalence differences (95% CI)		
	Food Secure (FS)	Low Food Secure (LFS)	Very Low Food Secure (VLFS)	FS vs. LFS	FS vs. VLFS	Food Secure (FS)	Low Food Secure (LFS)	Very Low Food Secure (VLFS)	FS vs. LFS	FS vs. VLFS
Frequent encouragement to diet	20.9	30.5	32.8	9.6 (3.0, 16.3)	11.9 (3.2, 20.6)	23.6	31.5	35.1	7.9 (1.6, 14.1)	11.4 (3.3, 19.6)
Frequent encouragement for healthy eating	65.3	69.5	71.8	4.2 (-2.8, 11.1)	6.5 (-2.2, 15.1)	61.1	66.7	68.4	5.6 (-0.9, 12.1)	7.3 (-0.8, 15.5)
Frequent comments about child's weight	32.0	47.1	41.5	15.1 (7.8, 22.5)	9.5 (0.2, 18.9)	33.2	42.1	49.0	8.8 (2.1, 15.5)	15.8 (7.1, 24.5)
Concerned about child's weight	35.1	39.1	50.4	4.0 (-3.3, 11.2)	15.2 (5.8, 24.7)	36.0	42.2	48.1	6.2 (-0.5, 13.0)	12.1 (3.4, 20.7)
Frequent use of restrictive feeding practices	21.9	30.5	31.3	8.6 (1.9, 15.3)	9.4 (0.7, 18.0)	20.1	29.1	37.4	8.9 (2.9, 14.9)	17.3 (9.1, 25.5)
Frequent use of pressured feeding practices	18.5	28.1	29.8	9.6 (3.1, 16.0)	11.2 (2.8, 19.7)	17.7	29.0	25.2	11.3 (5.3, 17.2)	7.5 (0.1, 14.9)

Table 3

Predicted probabilities of use of eating and weight-related parenting practices by food security status among mothers of adolescent boys (n=986)

	Model 2				Model 3					
	Estimated Prevalence		Prevalence differences (95% CI)		Estimated Prevalence		Prevalence differences (95% CI)			
	Food Secure (FS)	Low Food Secure (LFS)	Very Low Food Secure (VLFS)	FS vs. LFS	FS vs. VLFS	Food Secure (FS)	Low Food Secure (LFS)	Very Low Food Secure (VLFS)	FS vs. LFS	FS vs. VLFS
Frequent encouragement to diet	24.0	24.1	28.6	0.1 (-6.4, 6.6)	4.6 (-4.1, 13.4)	22.6	23.1	28.5	0.5 (-5.9, 6.8)	5.8 (-2.8, 14.4)
Frequent encouragement for healthy eating	64.2	69.4	72.2	5.3 (-2.3, 12.8)	8.0 (-1.3, 17.3)	63.3	68.5	72.2	5.2 (-2.7, 13.1)	8.9 (-0.5, 18.3)
Frequent comments about child's weight	33.6	42.7	41.4	9.1 (1.5, 16.7)	7.8 (-2.0, 17.6)	32.9	41.5	40.7	8.6 (0.9, 16.3)	7.8 (-2.0, 17.6)
Concerned about child's weight	35.6	35.7	49.2	0.2 (-7.6, 7.9)	13.7 (3.3, 24.0)	35.1	35.8	48.8	0.8 (-7.0, 8.5)	13.7 (3.5, 23.9)
Frequent use of restrictive feeding practices	24.9	25.0	26.3	0.1 (-6.5, 6.8)	1.4 (-7.3, 10.1)	25.3	24.8	25.2	-0.6 (-7.4, 6.3)	-0.2 (-8.7, 8.4)
Frequent use of pressured feeding practices	20.5	23.5	25.8	3.0 (-3.3, 9.2)	5.2 (-3.2, 13.6)	20.0	23.6	26.2	3.5 (-2.9, 9.9)	6.2 (-2.3, 14.7)

Model 2: Adjusted for socio-demographic characteristics (child's age, mothers' race/ethnicity, number of children in home, and educational attainment, and household income)

Model 3: Model 2 with additional adjustment for maternal BMI and adolescent BMI percentile

Table 4

Predicted probabilities of use of eating and weight-related parenting practices by food security status among mothers of adolescent girls (n=1190)

	Model 2						Model 3			
	Estimated Prevalence			Prevalence Differences (95% CI)			Estimated Prevalence		Prevalence Differences (95% CI)	
	Food Secure (FS)	Low Food Secure (LFS)	Very Low Food Secure (VLFS)	FS vs. LFS	FS vs. VLFS	Food Secure (FS)	Low Food Secure (LFS)	Very Low Food Secure (VLFS)	FS vs. LFS	FS vs. VLFS
Frequent encouragement to diet	24.2	28.8	33.1	4.6 (-1.8, 11.1)	8.9 (0.5, 17.4)	25.7	27.3	31.8	1.7 (-4.5, 7.8)	6.1 (-1.8, 14.0)
Frequent encouragement for healthy eating	61.4	65.7	66.2	4.4 (-2.8, 11.5)	4.8 (-4.2, 13.8)	61.3	65.5	64.1	4.2 (-3.2, 11.6)	2.8 (-6.6, 12.2)
Frequent comments about child's weight	35.1	36.8	43.9	1.7 (-5.1, 8.4)	8.8 (0.0, 17.6)	36.1	35.3	42.2	-0.7 (-7.5, 6.0)	6.1 (-2.7, 14.8)
Concerned about child's weight	35.7	40.4	47.4	4.7 (-2.5, 12.0)	11.6 (2.3, 21.0)	36.4	38.5	45.2	2.1 (-4.9, 9.0)	8.8 (-0.1, 17.6)
Frequent use of restrictive feeding practices	20.1	26.9	34.4	6.8 (0.6, 13.0)	14.3 (5.8, 22.8)	20.5	26.3	33.0	5.7 (-0.4, 11.8)	12.4 (4.2, 20.7)
Frequent use of pressured feeding practices	19.1	22.4	22.6	3.4 (-2.2, 8.9)	3.5 (-3.6, 10.6)	18.9	21.6	21.1	2.7 (-2.8, 8.2)	2.3 (-4.7, 9.2)

Model 2: Adjusted for socio-demographic characteristics (child's age, mothers' race/ethnicity and educational attainment, number of children in home, and household income)

Model 3: Model 2 with additional adjustment for maternal BMI and adolescent BMI percentile