



# HHS Public Access

Author manuscript

*Prenat Diagn.* Author manuscript; available in PMC 2016 September 01.

Published in final edited form as:

*Prenat Diagn.* 2015 September ; 35(9): 859–863. doi:10.1002/pd.4622.

## Disparities in the Prenatal Detection of Critical Congenital Heart Disease

GD Hill, JR Block, JB Tanem, and MA Frommelt

Department of Pediatrics, Division of Cardiology, Medical College of Wisconsin, Milwaukee, WI, USA

### Abstract

**Objectives**—Prenatal diagnosis of critical congenital heart disease, that requiring surgical or catheter intervention in the first 30 days of life, allows for delivery at a specialized center which can reduce preoperative morbidity and mortality. We sought to identify risk factors for a missed prenatal diagnosis of critical congenital heart disease.

**Methods**—Patients presenting to the Children’s Hospital of Wisconsin with critical congenital heart disease from 2007-2013 were included. Those with a prenatal diagnosis were compared to those with a postnatal diagnosis.

**Results**—The cohort included 535 patients with prenatal diagnosis made in 326 (61%). The prenatal diagnostic rate improved from 44% in 2007 to 69% in 2013. Independent factors associated with a postnatal diagnosis were a lesion that required a view other than a 4 chamber view to make the diagnosis ( $p<0.0001$ ), absence of another organ system anomaly ( $p<0.0001$ ), and living in a higher poverty ( $p=0.02$ ) or lower population density communities ( $p=0.002$ ).

**Conclusions**—While the prenatal diagnostic rate for critical congenital heart disease is improving, those living in impoverished or rural communities are at highest risk of not having a diagnosis made prenatally. Interventions to improve prenatal detection of congenital heart disease should target these vulnerable areas.

### Introduction

Malformations of the cardiovascular system remain the leading cause of infant mortality from congenital defects<sup>1</sup>. Those with critical congenital heart disease (CHD), defined as disease requiring surgical or catheter based intervention within the first 30 days of life, are at highest risk for early mortality. Prenatal detection allows for delivery at a surgical center capable of caring for these high risk infants and improves preoperative condition with conflicting data on the impact on mortality<sup>2-12</sup>.

While there are risk factors that increase the risk of CHD and therefore are indications for a fetal echocardiogram (i.e. maternal diabetes, family history of CHD), the vast majority of

---

Corresponding Author Garick Hill, Children’s Hospital of Wisconsin 9000 W Wisconsin Ave Milwaukee WI 53226, ghill@chw.org (414) 266-2380.

Disclosures: None declared

CHD occurs in low risk pregnancies<sup>13</sup>. As such, routine obstetric ultrasounds are the primary mechanism to identify CHD before birth. In spite of near universal performance of routine ultrasounds during pregnancy and estimates that over 90% of CHD should be detectable by prenatal ultrasound, actual detection rates remain low, ranging from 36-50% in the United States<sup>13-17</sup>. This study sought to determine the prenatal detection rate for critical CHD and determine risk factors for failure to make a prenatal diagnosis.

## Methods

### Population

After approval from the institutional review board, a retrospective chart review was performed. All patients presenting to the Children's Hospital of Wisconsin requiring surgical or catheter based intervention at 30 days of age between January 1, 2007 and December 31, 2013 were included. Those with isolated patent ductus arteriosus or non-structural heart disease, such as congenital complete heart block, were excluded.

### Definitions

Socioeconomic variables including median household income, percentage below the poverty line and population density were based on the 2010 census data by zip code of the mother's residence at the time of delivery. A population density of <500 people/sq mi was used to dichotomize rural and non-rural areas<sup>18</sup> for graphical presentation; in analysis the continuous variable was used. The types of CHD were divided into three groups based on the echocardiographic views required to accurately identify the disease (Figure 1). Group 1 were those diseases identified by a four chamber view alone, Group 2 were those diseases identified by the addition of an outflow tract view, and Group 3 were those diseases identified by the need for unique views such as an aortic arch view or color and spectral Doppler evaluation of the pulmonary veins. Organ system anomalies were defined as those that could potentially be identified on routine obstetric ultrasound as these may impact referral for further evaluation prior to delivery.

### Statistical analysis

Those with a prenatal diagnosis were compared to those with a postnatal diagnosis using Chi-squared for categorical variables or Wilcoxon rank sum test for continuous variables. Multivariable logistic regression with backwards elimination was performed to determine risk factors for a postnatal diagnosis. Data are presented as number with percent of total or mean  $\pm$  standard deviation. Data analysis was performed using StataIC (StataCorp, College Station, Texas) with a  $p < 0.05$  considered significant.

## Results

The cohort included 535 patients with 326 (61%) having a prenatal diagnosis. The percentage of prenatal diagnosis improved over time with 44% in 2007 and 69% in 2013. Prenatal diagnostic rates were best in tricuspid atresia (14/14, 100%) and worst in total anomalous pulmonary venous connection (1/14, 7%); see Figure 2. Comparison of characteristics between those diagnosed prenatally and those diagnosed postnatally can be

seen in table 1. In multivariable analysis using the factors seen in Table 1, a lesion that required a view other than a 4 chamber view to make the diagnosis ( $p<0.0001$ ), absence of another organ system anomaly ( $p<0.0001$ ), and living in a higher poverty ( $p=0.02$ ) or a lower population density zip code ( $p=0.002$ ) were independently associated with a postnatal diagnosis of critical CHD. In analysis comparing rural and non-rural communities, if the CHD could be defined using the four chamber view alone, there was no significant difference in prenatal diagnostic rate between these communities. When outflow tract or other more complex views were needed to make the diagnosis, prenatal diagnostic rates were significantly lower in rural communities (Figure 3).

## Discussion

This study demonstrates an overall improvement in the frequency of prenatal diagnosis of critical CHD over time, however there continues to be significant disparity in diagnosis between urban and rural communities. When adjusting for the view required for diagnosis and presence of other organ system anomalies, those living in impoverished or rural communities are at greatest risk for a missed prenatal diagnosis of critical CHD. In our study, the disparity between communities was most noticeable in defects that require advanced imaging views for accurate diagnosis. Lesions such as aortic stenosis, double outlet right ventricle, pulmonary atresia, tetralogy of Fallot, transposition of the great arteries, and truncus arteriosus which require an outflow tract view, and coarctation of the aorta, interrupted aortic arch, and total anomalous pulmonary venous return which require an aortic arch view or color and spectral Doppler imaging were less commonly identified in rural communities. It has previously been demonstrated that screening for CHD with only a 4 chamber view is insufficient and results in a sensitivity of only 40% for detection of CHD<sup>19</sup>. Recent changes to the guidelines for performance of obstetric ultrasound screening recognize this limitation, as routine evaluation of both the left and right ventricular outflow tract is recommended<sup>20</sup>.

We speculate that health care personnel working in impoverished, rural areas are less able to acquire the new skills required to adequately evaluate the outflow tracts or other complex views. This is likely a result of the rural location with greater isolation from peers and less ability to pursue further training in both image acquisition and interpretation. Training programs for those performing screening ultrasounds in the United Kingdom and Sweden have demonstrated a significant improvement in the detection of congenital heart disease<sup>21-24</sup>. This improvement has been demonstrated with minimal investment of time, with some programs lasting only 1-3 days<sup>23,24</sup>. Our data suggest that training programs in detection of CHD by screening obstetric ultrasound should be targeted to these impoverished, rural communities for the greatest impact. Supervision of imaging and specialized training on normal findings for more complex views has been demonstrated to be effective by Evans et al and could be a focus for continuing education<sup>25</sup>.

Recently, the United States Department of Health and Human Services recommended universal pulse oximetry screening for all neonates in an effort to improve early detection of CHD<sup>26</sup>. However, prenatal diagnosis offers the distinct advantage of allowing for delivery at a specialized cardiac care center. Morris et al demonstrated that patients with hypoplastic

left heart syndrome born further from a surgical center were at higher risk for neonatal mortality, even independent of prenatal diagnosis<sup>11</sup>. This likely reflects the inability of those physicians and providers in the rural community to appropriately manage a critically ill neonate because of the equipment and expertise required. While pulse oximetry may improve detection, only through prenatal diagnosis can the location of delivery be altered and the risk of mortality and preoperative deterioration minimized.

As a retrospective single center cohort study, these data are subject to some limitations. Most notably, no data on postnatal mortality prior to presentation at our institutional are available. Previous work has shown the number of deaths in Wisconsin from 2002-2006 from undiagnosed CHD to be less than 2 cases per year and this number is likely to be even lower given the improvement in prenatal detection noted over time in our study<sup>27</sup>. However, while postnatal death in undiagnosed patients may have little effect on our study, the rate of prenatal termination is unknown and likely more significant. With the inclusion criteria used, not all patients with each diagnosis were included, for example only severe forms of tetralogy of Fallot, those requiring early intervention, were included. An additional limitation is the use of zip codes to determine population density and poverty as there may be variation within zip codes.

## Conclusions

While the rate of prenatal diagnosis of critical congenital heart disease has improved over time, those residing in impoverished, rural communities remain at greatest risk of being diagnosed postnatally. This difference in diagnostic rate is most notable for diagnoses requiring the outflow tract view and other more complex imaging. Additional training efforts targeted to these vulnerable communities may have the most significant impact.

## Acknowledgments

Funding: This publication was supported by the National Center for Advancing Translational Sciences, National Institutes of Health, through Grant Number 8UL1TR000055. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NIH.

## References

1. Lee K, Khoshnood B, Chen L, Wall SN, Cromie WJ, Mittendorf RL. Infant mortality from congenital malformations in the united states, 1970-1997. *Obstet Gynecol.* 2001; 98(4):620–627. [PubMed: 11576578]
2. Landis BJ, Levey A, Levasseur SM, et al. Prenatal diagnosis of congenital heart disease and birth outcomes. *Pediatr Cardiol.* 2013; 34(3):597–605. [PubMed: 23052660]
3. Kumar RK, Newburger JW, Gauvreau K, Kamenir SA, Hornberger LK. Comparison of outcome when hypoplastic left heart syndrome and transposition of the great arteries are diagnosed prenatally versus when diagnosis of these two conditions is made only postnatally. *Am J Cardiol.* 1999; 83(12):1649–1653. [PubMed: 10392870]
4. Fuchs IB, Muller H, Abdul-Khaliq H, Harder T, Dudenhausen JW, Henrich W. Immediate and long-term outcomes in children with prenatal diagnosis of selected isolated congenital heart defects. *Ultrasound Obstet Gynecol.* 2007; 29(1):38–43. [PubMed: 17200993]
5. Franklin O, Burch M, Manning N, Sleeman K, Gould S, Archer N. Prenatal diagnosis of coarctation of the aorta improves survival and reduces morbidity. *Heart.* 2002; 87(1):67–69. [PubMed: 11751670]

6. Calderon J, Angeard N, Moutier S, Plumet MH, Jambaque I, Bonnet D. Impact of prenatal diagnosis on neurocognitive outcomes in children with transposition of the great arteries. *J Pediatr.* 2012; 161(1):94–8. e1. [PubMed: 22284567]
7. Oster ME, Kim CH, Kusano AS, et al. A population-based study of the association of prenatal diagnosis with survival rate for infants with congenital heart defects. *Am J Cardiol.* 2014; 113(6): 1036–1040. [PubMed: 24472597]
8. Mahle WT, Clancy RR, McGaurn SP, Goin JE, Clark BJ. Impact of prenatal diagnosis on survival and early neurologic morbidity in neonates with the hypoplastic left heart syndrome. *Pediatrics.* 2001; 107(6):1277–1282. [PubMed: 11389243]
9. Bonnet D, Coltri A, Butera G, et al. Detection of transposition of the great arteries in fetuses reduces neonatal morbidity and mortality. *Circulation.* 1999; 99(7):916–918. [PubMed: 10027815]
10. Tworetzky W, McElhinney DB, Reddy VM, Brook MM, Hanley FL, Silverman NH. Improved surgical outcome after fetal diagnosis of hypoplastic left heart syndrome. *Circulation.* 2001; 103(9):1269–1273. [PubMed: 11238272]
11. Morris SA, Ethen MK, Penny DJ, et al. Prenatal diagnosis, birth location, surgical center, and neonatal mortality in infants with hypoplastic left heart syndrome. *Circulation.* 2014; 129(3):285–292. [PubMed: 24135071]
12. Wright LK, Ehrlich A, Stauffer N, Samai C, Kogon B, Oster ME. Relation of prenatal diagnosis with one-year survival rate for infants with congenital heart disease. *Am J Cardiol.* 2014; 113(6): 1041–1044. [PubMed: 24440326]
13. Stumpflen I, Stumpflen A, Wimmer M, Bernaschek G. Effect of detailed fetal echocardiography as part of routine prenatal ultrasonographic screening on detection of congenital heart disease. *Lancet.* 1996; 348(9031):854–857. [PubMed: 8826811]
14. Friedberg MK, Silverman NH, Moon-Grady AJ, et al. Prenatal detection of congenital heart disease. *J Pediatr.* 2009; 155(1):26–31. 31.e1. [PubMed: 19394031]
15. Peiris V, Singh TP, Tworetzky W, Chong EC, Gauvreau K, Brown DW. Association of socioeconomic position and medical insurance with fetal diagnosis of critical congenital heart disease. *Circ Cardiovasc Qual Outcomes.* 2009; 2(4):354–360. [PubMed: 20031861]
16. Pinto NM, Keenan HT, Minich LL, Puchalski MD, Heywood M, Botto LD. Barriers to prenatal detection of congenital heart disease: A population-based study. *Ultrasound Obstet Gynecol.* 2012; 40(4):418–425. [PubMed: 21998002]
17. Israel SW, Roofe LR, Saville BR, Walsh WF. Improvement in antenatal diagnosis of critical congenital heart disease implications for postnatal care and screening. *Fetal Diagn Ther.* 2011; 30(3):180–183. [PubMed: 21525728]
18. U.S. Department of Health and Human Services Rural Health Toolkit. U.S. Department of Health and Human Services Web site. <http://www.hrsa.gov/healthit/toolbox/RuralHealthITtoolbox/Introduction/defined.html>. Updated 2014. Accessed December 12, 2014
19. Gonçaves LF, Bronsteen R, Lee W. Fetal heart: A 4-chamber view is not enough. *Clin Obstet Gynecol.* 2012; 55(1):266–280. [PubMed: 22343243]
20. AIUM Practice Guideline for the Performance of Obstetric Ultrasound Examinations. <http://aium.org/resources/guidelines/obstetric.pdf>. Updated 2013. Accessed May 2, 2013
21. Asplin N, Dellgren A, Conner P. Education in obstetrical ultrasound--an important factor for increasing the prenatal detection of congenital heart disease. *Acta Obstet Gynecol Scand.* 2013; 92(7):804–808. [PubMed: 23521227]
22. Carvalho JS, Mavrides E, Shinebourne EA, Campbell S, Thilaganathan B. Improving the effectiveness of routine prenatal screening for major congenital heart defects. *Heart.* 2002; 88(4): 387–391. [PubMed: 12231598]
23. Hunter S, Heads A, Wyllie J, Robson S. Prenatal diagnosis of congenital heart disease in the northern region of England: Benefits of a training programme for obstetric ultrasonographers. *Heart.* 2000; 84(3):294–298. [PubMed: 10956294]
24. McBrien A, Sands A, Craig B, Dornan J, Casey F. Impact of a regional training program in fetal echocardiography for sonographers on the antenatal detection of major congenital heart disease. *Ultrasound Obstet Gynecol.* 2010; 36(3):279–284. [PubMed: 20205153]

25. Evans W, Castillo W, Rollins R, et al. Moving towards universal prenatal detection of critical congenital heart disease in southern nevada: A community-wide program. *Pediatr Cardiol.* 2015; 36(2):281–288. [PubMed: 25103855]
26. Mahle WT, Martin GR, Beekman RH 3rd, Morrow WR, Section on Cardiology and Cardiac Surgery Executive Committee. Endorsement of health and human services recommendation for pulse oximetry screening for critical congenital heart disease. *Pediatrics.* 2012; 129(1):190–192. [PubMed: 22201143]
27. Ng B, Hokanson J. Missed congenital heart disease in neonates. *Congenit Heart Dis.* 2010; 5(3): 292–296. [PubMed: 20576049]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

What's known:

- Diagnostic rates for congenital heart disease prior to delivery are suboptimal and influenced by socioeconomic factors


What's new:

- Prenatal diagnostic rates for critical congenital heart disease are worse in impoverished rural communities
- The effect is more notable when advanced views are required to make the diagnosis


Group 1 (4 chamber view)	Group 2 (Outflow view)	Group 3 (Other view)
<ul style="list-style-type: none"> <li>• Atrioventricular septal defect</li> <li>• Double inlet left ventricle</li> <li>• Ebstein's anomaly</li> <li>• Hypoplastic left heart syndrome</li> <li>• Pulmonary atresia with intact ventricular septum</li> <li>• Single ventricle</li> <li>• Tricuspid atresia</li> </ul>	<ul style="list-style-type: none"> <li>• Aortic stenosis</li> <li>• Double outlet right ventricle</li> <li>• Pulmonary atresia</li> <li>• Tetralogy of Fallot</li> <li>• Transposition of the great arteries</li> <li>• Truncus arteriosus</li> </ul>	<ul style="list-style-type: none"> <li>• Coarctation of the aorta</li> <li>• Interrupted aortic arch</li> <li>• Total anomalous pulmonary venous return</li> </ul>

**Figure 1.**  
Division of groups based on echocardiographic view required for diagnosis





**Figure 2.** Frequency of prenatal diagnosis by lesion. DILV – Double inlet left ventricle, AVSD – atrioventricular septal defect, HLHS – hypoplastic left heart syndrome, PA/IVS – pulmonary atresia with intact ventricular septum, DORV – double outlet right ventricle, IAA – interrupted aortic arch, TOF – tetralogy of Fallot, TGA – transposition of the great arteries, TAPVR – total anomalous pulmonary venous return



**Figure 3.** Prenatal diagnostic rate by view required to make the lesion and by rural versus urban community

**Table 1**

Comparison of prenatal and postnatal diagnosis groups

	<b>Prenatal n = 326</b>	<b>Postnatal n = 209</b>	<b>p value</b>
Married, n (%)	203 (63%)	111 (54%)	0.04
Gravidity, n (%)			0.12
1	101 (31%)	57 (28%)	
2	80 (25%)	65 (32%)	
3	62 (19%)	30 (15%)	
>3	81 (25%)	49 (24%)	
Male fetus, n (%)	185 (57%)	128 (61%)	0.3
Race/Ethnicity, n (%)			0.16
Black/African American	30 (9%)	17 (8%)	
Hispanic	38 (12%)	21 (10%)	
White/Caucasian	227 (70%)	138 (66%)	
Other	31 (9%)	33 (16%)	
View required for diagnosis, n (%)			<0.001
4 chamber view	156 (48%)	28 (13%)	
Outflow tract view	117 (36%)	91 (44%)	
Other	53 (16%)	90 (43%)	
Other anomaly present, n (%)	70 (21%)	16 (8%)	<0.001
Private insurance, n (%)	219 (68%)	126 (63%)	0.23
Percent below poverty	13.5 ± 9.8	14.1 ± 9.7	0.19
	2518 ±	1935 ±	
Population density (people/sq mi)	3338	3020	<0.001