



Published in final edited form as:

Nurs Res. 2015 ; 64(5): 342–350. doi:10.1097/NNR.000000000000113.

Biological and Sociocultural Differences in Perceived Barriers to Physical Activity among 5th–7th Grade Urban Girls

Amber L. Vermeesch, PhD, MSN, RN, FNP-C [Associate Professor of Nursing]

University of Portland, School of Nursing

Jiying Ling, PhD, MS, RN [Assistant Professor of Nursing]

Michigan State University, College of Nursing

Vicki R. Voskuil, MS, RN, CPNP [Doctoral Candidate]

Michigan State University, College of Nursing

Marion Bakhoya, BS [Former MS Kinesiology Student]

Michigan State University, Department of Kinesiology

Stacey M. Wesolek, MS [Project Manager]

Michigan State University, College of Nursing

Kelly A. Bourne, MS [Measurement Coordinator]

Michigan State University, College of Nursing

Karin A. Pfeiffer, PhD [Associate Professor of Kinesiology]

Michigan State University, Department of Kinesiology

Lorraine B. Robbins, PhD, RN, FAAN, FNP-BC [Associate Professor of Nursing]

Michigan State University, College of Nursing

Abstract

Background—Inadequate physical activity (PA) contributes to the high prevalence of overweight and obesity among U.S. adolescent girls. Barriers preventing adolescent girls from meeting PA guidelines have not been thoroughly examined.

Objectives—The threefold purpose of this study was to: (a) determine pubertal stage, racial/ethnic, and socioeconomic status (SES) differences in ratings of interference of barriers to PA; (b) examine relationships between perceived barriers and age, body mass index (BMI), recreational screen time, sedentary activity, and PA; and (c) identify girls' top-rated perceived barriers to PA.

Methods—Girls ($N = 509$) from eight Midwestern U.S. schools participated. Demographic, pubertal stage, perceived barriers, and recreational screen time data were collected via surveys. Height and weight were measured. Accelerometers measured sedentary activity, moderate-to-vigorous physical activity (MVPA), and light plus MVPA.

Corresponding Author: Lorraine B. Robbins, Michigan State University College of Nursing, Bott Building for Nursing Education and Research, 1355 Bogue Street, C245, East Lansing, MI 48824 (robbin76@msu.edu)..

The authors have no conflicts of interest to report.

Results—Girls of low SES reported greater interference of perceived barriers to PA than those who were not of low SES (1.16 vs. 0.97, $p = .01$). Girls in early/middle puberty had lower perceived barriers than those in late puberty (1.03 vs. 1.24, $p < .001$). Girls' perceived barriers were negatively related to MVPA ($r = -.10$, $p = .03$) and light plus MVPA ($r = -.11$, $p = .02$). Girls' top five perceived barriers included lack of skills, hating to sweat, difficulty finding programs, being tired, and having pain.

Discussion—Innovative interventions, particularly focusing on skill development, are needed to assist girls in overcoming their perceived barriers to PA.

Keywords

adolescent; female; physical exercise; puberty

Inadequate physical activity (PA) contributes to the high prevalence of obesity in 12-19-year-old U.S. girls (20.7%; Ogden, Carroll, Kit, & Flegal, 2014). The pervasive negative effect of inadequate PA participation on body weight status indicates an urgent need to reverse this disconcerting behavioral trend (Kann et al., 2014). Only 17.7% of adolescent girls in the U.S. report attaining recommendations proposed by both the U.S. Department of Health & Human Services (USDHHS; 2008) and the World Health Organization (WHO; 2010) that call for at least 60 minutes of moderate-to-vigorous PA (MVPA) daily (Kann et al., 2014). Consistently, studies support a sharp decline in girls' PA from ages 9 to 12 (Dumith, Gigante, Domingues, & Kohl, 2011). The decrease is particularly notable among urban-dwelling, low socioeconomic status (SES) girls (Wang et al., 2007). Based on 2013 U.S. Youth Risk Behavior Survey data, the prevalence of PA participation for at least 60 minutes daily was higher among White (37.5%), Black (37.2%), and Hispanic (33.9%) adolescent boys than among White (18.7%), Black (16.0%), and Hispanic (17.4%) adolescent girls, respectively (Kann et al., 2014).

Attaining adequate PA during adolescence is critical for reducing risks associated with elevated body mass index (BMI), including the development of type 2 diabetes and cardiovascular disease (Belcher et al., 2010). Over one third (33.8%) of U.S. girls, 12-19 years old, are overweight or obese, and over one fifth (20.7%) are obese (Ogden et al., 2014). Racial and ethnic differences exist in weight status with more Black (22.8%) and Hispanic (19.2%) girls being overweight than White girls (14.3%), and more Black girls (16.7%) than Hispanic (11.2%) and White (9.7%) girls being obese (Kann et al., 2014). Because both PA and obesity can track into adulthood, understanding perceived barriers that prevent girls from establishing a habit of regular PA is vital (Clarke, O'Malley, Johnston, Schulenberg, & Lantz, 2009).

Acquiring a comprehensive understanding of perceived barriers to PA can be challenging due to noted differences among various subgroups of girls. For example, difficulty accessing PA programs or classes is a noted, perceived barrier for low SES youth (Humbert et al., 2006). Cultural considerations and peer opinion must be considered in perceived barriers to PA among various subgroups. Black and Hispanic girls are more accepting of larger body size than White girls, as evidenced by White girls with a BMI closer to obese were found to have less social desirability than White girls with a BMI closer to normal weight by their

peers (Boyington et al., 2008; Lanza, Echols, & Graham, 2013). Peer social environment, including low peer PA levels—especially among female friends—is another perceived barrier to PA for adolescent girls (Larson, Wall, Story, & Neumark-Sztainer, 2013). Kelly et al. (2010) reported that perceived barriers to PA were negatively associated with accelerometer-measured PA for White, but not Black, girls. Dishman, Dunn, Sallis, Vandenberg, and Pratt (2010) found that perceived barriers to PA were correlated inversely with accelerometer-measured PA for girls in 8th grade, but not 6th grade. One quantitative study with girls identified barriers to PA based on academic grade (6th, 7th, & 8th) and race (Black and White; Robbins, Pender, & Kazanis, 2003). Although some differences were evident, the sample size was too small to draw any definitive conclusions ($N = 77$; Robbins et al., 2003). These findings indicate that understanding the relationship between perceived barriers to PA and the behavior itself, and identifying specific barriers to PA, require subgroup analysis.

Despite a strong likelihood of increasing PA when relevant barriers are effectively targeted, gaps in the literature remain regarding the influence of biological and sociocultural factors on girls' perceived barriers to PA and the relationship between barriers and PA, as well as sedentary activity (Camacho-Miñano, LaVoi, & Barr-Anderson, 2011). Interventions designed to overcome perceived barriers to PA by providing access to PA programs or including strategies for promoting positive perceptions regarding PA have shown limited success, indicating that continued investigation of biological and sociocultural differences may offer important insights for tailoring interventions to address the unique needs of varied population subgroups (Camacho-Miñano et al., 2011; Van der Horst, Paw, Twisk, & Van Mechelen, 2007). Because physical transformation of the body during adolescence can modify adolescents' self-perceptions, attitudes, and behavior, examination of pubertal stage differences in perceived barriers to PA is warranted (Waylen & Wolke, 2004). For example, White adolescent girls identify embarrassment or self-consciousness as a barrier more than other races (Robbins et al., 2003). A thorough investigation of the barriers that may underlie the decline in girls' PA that begins around the onset of adolescence, particularly among adolescent girls of low SES and minority backgrounds, is also necessary to understand existing disparities. Therefore, the threefold purpose of this study was to:

1. determine pubertal stage, racial/ethnic, and SES differences in ratings of interference of barriers to PA;
2. examine relationships between perceived barriers and age, body mass index (BMI), recreational screen time, sedentary activity, and PA; and
3. identify girls' top-rated perceived barriers for the total sample and by pubertal stage, race, and SES.

Methods

Study Design, Participants, and Setting

In the first intervention year (2012-2013) of a group randomized controlled trial (2011-2016), eight Midwestern U.S. schools were randomly assigned to either receive a multicomponent PA intervention called "Girls on the Move" or serve as a control (Robbins

et al., 2013). The trial was based on the health promotion model and self-determination theory (Pender, Murdaugh, & Parsons, 2011; Ryan & Deci, 2000); the complete trial protocol has been published (Robbins et al., 2013). Girls meeting the following inclusion criteria were selected for participation on a first-come, first-served basis: (a) 5th-7th grade girls; (b) available and willing to participate in the PA club three days/week for 17 weeks; (c) available for nine-month follow up after the intervention ends; and (d) able to read, understand, and speak English. Exclusion criteria included: (a) involved in or planning to be involved in school or community sports or other organized PAs that require participation three or more days/week after school; and (b) having a health condition precluding safe MVPA (Robbins et al., 2013).

Baseline data collected during fall 2012 from 5th-7th grade girls ($N = 509$) in the eight schools were used for this comparative and correlational study. The schools were located in four cities with 39.7%, 20.2%, 36.8%, and 27.1% of the population being below poverty level compared to 16.3% in the state, and the annual income per capita ranged from \$14,454 to \$20,891 compared to the state level of \$25,547 (U.S. Census Bureau, 2014). Approximately 3.0-14.3% of the population was Hispanic and 23.7-56.6% was Black (U.S. Census Bureau, 2014). The proportion of students involved in free or reduced-price lunch programs (financially disadvantaged) ranged from 59.7% to 95.0% ($M = 74.0%$; State of Michigan, 2014). On average, 30.3% of the students in the schools were White (min-max: 1.2-49.2%), 58.4% were Black (min-max: 23.2-100%), and 9.2% were Hispanic (min-max: 2.9-18.7%; State of Michigan, 2014).

Measures

Demographic survey—The demographic survey had questions that addressed age, academic grade, race, ethnicity, and SES. Enrollment in the free or reduced-price lunch program served as a proxy for SES. Parents/guardians completed the demographic survey in collaboration with their daughters.

Pubertal stage—Pubertal stage was categorized as early, middle, and late puberty, and was assessed by the Pubertal Development Scale (Peterson, Crockett, Richards, & Boxer, 1988). Validity and reliability have been established with girls as young as those in the 5th and 6th grades (Carskadon & Acebo, 1993; Peterson et al., 1988). Pubertal stage was computed by summing self-reported scores for underarm hair growth, breast development, and menarche. Response choices to measure hair and breast development were: 1 = *no*; 2 = *yes, barely*; 3 = *yes, definitely*; and 4 = *development complete*. For menarche, girls reported either: no menstruation (indicating early or middle puberty) or yes, menstruation started (indicating late puberty; Carskadon & Acebo, 1993). Girls reporting no menstruation with summed scores ≥ 3 for underarm hair growth and breast development were considered to be in early puberty, whereas those having summed scores > 3 were categorized as being in middle puberty (Carskadon & Acebo, 1993).

Body mass index (BMI)—BMI in kg/m^2 was calculated from girls' height and weight without shoes using standardized procedures for the study protocol (Robbins et al., 2013). BMI z-scores were determined via use of the SAS Program for the Centers for Disease

Control and Prevention Growth Charts. The program is available online at www.cdc.gov/nccdphp/dnpao/growthcharts/resources/sas.htm

Recreational screen time, sedentary activity, and physical activity—

Recreational screen time was calculated as the amount of time spent viewing television, talking on the phone or sending messages, and playing video games or using the computer or Internet for non-school-related work (Costigan, Barnett, Plotnikoff, & Lubans, 2013). Girls responded to six items when reporting number of hours that they spent engaging in each activity on a typical school day and weekend day. Response choices ranged from 0 = *I do not* (the specific behavior) to 5 = *five or more hours per day*. For this study, Cronbach's alpha was .82 with item-total correlation coefficients ranging from .52 to .63.

To assess minutes per hour of sedentary activity, MVPA, and light plus MVPA (LMVPA), the ActiGraph GT3X-plus accelerometer was used. Girls were asked to wear accelerometers for seven consecutive days except when showering, swimming, and sleeping at night. Data generated for at least eight hours daily on at least three weekdays and one weekend day were considered adequate for analysis (Patnode et al., 2011). Investigators downloaded and processed the data using the ActiLife software program. Evenson, Catellier, Gill, Ondrak, and McMurray (2008) cut points for children were used to analyze data. Trost, Loprinzi, Moore, and Pfeiffer (2011) found Evenson et al. (2008) cut points to be acceptable among 5- to 15-year-old children and adolescents. Average minutes per hour were determined to account for variable wear time to minimize any impact of some girls accruing fewer or greater counts per day by wearing the monitor for a shorter or longer period of time than others, respectively.

Perceived barriers to physical activity—A 9-item Perceived Barriers Scale was previously developed to measure perceptions of obstacles interfering with PA (Robbins, Wu, Sikorskii, & Marley, 2008). Response choices included: 0 = *not at all true*; 1 = *not very true*; 2 = *somewhat true*; and 3 = *very true*. Higher scores indicated a more negative perception. Acceptable face, content, and construct validity, and reliability estimated by Cronbach's alpha of .78 have been reported (Robbins, Sikorskii, Hamel, Wu, & Wilbur, 2009; Robbins et al., 2008). Prior to this study, seven new items were added to the scale based on recommendations from 6th-8th grade girls ($N = 25$) participating in focus groups conducted by one of the authors (Robbins et al., 2013). Examples of items include: "I have some pain from activity," "The weather is bad," and "I am too busy." Additional items enhanced the comprehensiveness of the scale in this study, resulting in an increased Cronbach's alpha of .85 with item-total correlation coefficients ranging from .35 to .57.

Procedures

The University Institutional Review Board (IRB) and school district administrators approved the study. Recruitment procedures and response rates have been reported (Ling, Robbins, Resnicow, & Bakhoya, 2014; Robbins et al., 2013). During data collection, girls completed an iPad-delivered survey that included the perceived barriers to PA and recreational screen time measures. Data collectors measured height and weight to calculate BMI. Each girl completed the Pubertal Development Scale behind a privacy screen.

Afterward, girls watched an instructional video on wearing the accelerometer and received an accelerometer, along with written instructions to share with parents/guardians.

Data Analysis

Data were analyzed using the Statistical Package for the Social Sciences (SPSS 21.0). Independent samples *t*-tests and one-way ANOVA examined biological (i.e., pubertal stage) and sociocultural (i.e., race/ethnicity, SES) differences in BMI, recreational screen time, sedentary activity, PA, and perceived barriers. Pearson product-moment bivariate correlations were calculated to examine the associations among age, BMI, recreational screen time, sedentary activity, PA, and perceived barriers. Identification of the top five perceived barriers was based on percentages of girls selecting “somewhat true” or “very true.” A mixed effects model was applied to examine the effects of school (random factor) and age, ethnicity, race, age, SES, puberty stage, BMI *z*-score, and perceived barriers (fixed effects) on MVPA. Restricted maximum likelihood (REML) estimation was used to deal with missing data with the assumption that the missing response were missing at random.

Results

Demographics

Almost four fifths of the girls were 11-12 years old ($n = 404$, 79.4%), and slightly over half were in the 6th grade. Greater than half were Black, and the majority participated in the free or reduced-price lunch program. Due to the small number of girls in early puberty ($n = 15$, 2.9%), early and middle puberty were collapsed into a new category called early/middle puberty ($n = 300$, 58.9%). The remaining girls were in late puberty. All girls in late puberty, but none in early/middle puberty, had started menstruation. Table 1 presents additional sample characteristics.

BMI, Recreational Screen Time, Sedentary Activity, and Physical Activity

Four hundred sixty-two (90.8%) girls provided acceptable accelerometer data for analysis. On average, the girls wore the accelerometer for about 14 hours per day ($SD = 1.81$, minimum = 10.25, maximum = 23.29).

Over half of the girls were overweight (21.5%) or obese (32.3%). Significant racial differences in BMI ($p = .002$) and BMI *z*-score ($p = .02$) occurred. Specifically, Black girls had higher BMI ($M = 23.55$, $SD = 5.64$ vs. $M = 21.39$, $SD = 4.45$) and BMI *z*-score ($M = 1.08$, $SD = 1.02$ vs. $M = 0.79$, $SD = 0.85$) than White girls. Girls in late puberty had a higher BMI *z*-score than those in early/middle puberty (see Table 2). No SES differences in BMI occurred. BMI was positively correlated with age ($r = .16$, $p < .001$) and self-reported recreational screen time ($r = .11$, $p = .01$), but negatively correlated with MVPA ($r = -.12$, $p = .01$).

Girls reported an average of 6.21 hours of total daily recreational screen time, with a mean of 5.70 hours ($SD = 3.44$) on school days and 6.72 hours ($SD = 3.64$) on weekend days. Average time in sedentary activity was 38.24 minutes/hour. Girls participated in 3.07 minutes/hour of MVPA with a range of 0.73 to 8.56 minutes/hour. Significant racial

differences in recreational screen time emerged ($p = .001$), with Black girls reporting more recreational screen time than White girls ($M = 6.64, SD = 3.40$ vs. $M = 5.35, SD = 3.06, p = .002$). Early/middle puberty girls participated in more LMVPA and MVPA and had less recreational screen time and sedentary activity than those in late puberty (see Table 2). Age was positively correlated with sedentary activity ($r = .25, p < .001$) and recreational screen time ($r = .25, p < .001$), but negatively correlated with MVPA ($r = -.21, p < .001$) and LMVPA ($r = -.25, p < .001$).

Table 3 presents the random and fixed effects of school, age, ethnicity, race, age, SES, pubertal stage, BMI z -score, and perceived barriers on MVPA. The two significant predictors for MVPA were pubertal stage and age. Specifically, girls in early/middle puberty participated in an average of 0.47 minutes/hour more MVPA than their peers in late puberty; as age increased by one year, MVPA decreased by 0.26 minutes/hour. The nested effects of school only accounted for about 2.9% of the variance in random effects, thus, the nested effect of school is not a concern in this study. The fixed effects model explained about 8% variance in MVPA.

Perceived Barriers to Physical Activity

Girls of low SES reported significantly greater perceived barriers to PA ($M = 1.16, SD = 0.58$) than those who were not of low SES ($M = 0.97, SD = 0.57; p = .01$). Perceived barriers to PA were lower among early/middle pubertal girls than late pubertal girls (see Table 2). Age was positively correlated with perceived barriers to PA ($r = .12, p = .008$). No significant racial or ethnic differences emerged in perceived barriers. Perceived barriers were negatively related to LMVPA ($r = -.11, p = .02$) and MVPA ($r = -.10, p = .03$), but positively correlated with sedentary activity ($r = .11, p = .02$) and recreational screen time ($r = .22, p < .001$).

Table 4 presents the top five perceived barriers for the total sample and for each pubertal stage, race, and SES. The only barrier identified by over half of all girls (51.5%) was lack of skills. The majority of girls in late puberty (56.9%) also indicated that lack of skills was a major barrier, followed by hating to sweat during the school day (56.5%) and difficulty finding PA programs or classes they like (53.6%). The majority of Black girls (54.6%) considered hating to sweat during the school day as a barrier followed by lack of skills (51.0%). For low SES girls, the majority indicated that hating to sweat during the school day (54.2%) followed by lack of skills (52.5%) were major barriers. In contrast, percentages related to each barrier were all under 50% for girls in early/middle puberty, girls who were White or mixed race, and girls who were not of low SES.

Discussion

This comparative and correlational study including 5th-7th grade urban Midwestern U.S. girls examined pubertal stage, racial/ethnic, and SES differences in perceived barriers to PA and the relationships among the following variables: age, BMI, recreational screen time, sedentary activity, PA, and perceived barriers to PA. Pubertal stage, racial/ethnic, and SES differences in girls' top perceived barriers to PA were also identified. A broad understanding of biological and sociocultural differences in perceived barriers to PA can be instrumental in

designing interventions using systematic and meaningful personalized strategies to assist diverse groups of urban girls to overcome their perceived barriers to PA.

BMI, Recreational Screen Time, Sedentary Activity, and Physical Activity

The prevalence of girls who were overweight or obese in this study was 53.8%—higher than the U.S. percentage of approximately 33.8% reported for 12- to 19-year-old girls (Ogden et al., 2014). In addition, the study's urban sample mainly comprised girls of low SES. Consistent with previous research, Black girls had a higher BMI and reported more recreational screen time than White girls (Kann et al., 2014). Thus, effective interventions are urgently needed to control the obesity crisis involving urban adolescent girls, especially among Black girls of low SES.

Findings that girls in late puberty had higher BMI and more recreational screen time and sedentary activities than those in early/middle puberty are comparable to other results noted in the literature. Brodersen, Steptoe, Boniface, and Wardle (2007) reported that as girls age, their sedentary behavior increases. Additionally, strong evidence indicates a positive association between recreational screen time and weight status, and a negative association between recreational screen time and participation in PA (Costigan et al., 2013).

Although this study demonstrated that girls in late puberty participated in less MVPA than those in early/middle puberty, conflicting findings concerning the relationship between pubertal stage and PA exist (Finne, Bucksch, Lampert, & Kolip, 2011; Smart et al., 2012). In a study involving 6,813 adolescents, aged 11-17 years, pubertal stage correlated with boys', but not girls', self-reported PA (Finne et al., 2011). Another study with 222 adolescent girls demonstrated pubertal stage had a negative and indirect effect on self-reported PA through perceptions of sports competence, body attractiveness, and physical condition (Smart et al., 2012). Even though positive correlations between adolescent self-report of PA and accelerometer data have been noted, self-report of PA may be one explanation for the inconsistent findings concerning girls' pubertal stage and their PA (Chinapaw, Mokkink, van Poppel, van Mechelen, & Terwee, 2010). Reliable and valid measurements of PA are important for examining the relationship between pubertal stage and PA.

To determine clinical significance of the accelerometer findings in terms of minutes/day of MVPA, mean minutes/hour were multiplied by the girls' average wear time of approximately 14 hours/day in this study. Troiano et al. (2007) also found that approximately 14 hours was the average accelerometer wear time per day for this age group. Based on the calculations, the mean of 3.3 minutes/hour of MVPA for girls in early/middle puberty, as noted in Table 2, translates to 46.2 minutes/day, and the mean of 2.7 minutes/hour for girls in late puberty translates to 37.8 minutes/day. These findings indicate that neither group met the USDHHS and WHO PA recommendations calling for 60 minutes of MVPA per day to acquire health benefits. In addition, the findings that girls in early/middle puberty participated in an average of 0.47 minutes/hour more MVPA than their peers in late puberty; and, as age increased by one year, MVPA decreased by 0.26 minutes/hour, translated to 6.6 minutes/day and 3.6 minutes/day, respectively. To put these numbers into perspective, the multi-center Trial of Activity for Adolescent Girls significantly increased

MVPA by 1.6 minutes/day among girls in the intervention schools, as compared to those in the control schools. The researchers indicated that, although small, an increase of this magnitude could prevent a weight gain of 0.82 kg per year, which could be substantial at the population level (Webber et al., 2008). Also, in terms of clinical significance, 6.6 and 3.6 minutes/day represent 11 and 6% of the recommended 60 minutes of MVPA per day, respectively.

Perceived Barriers to Physical Activity

Findings that perceived barriers were negatively and weakly correlated with both LMVPA and MVPA are both similar and contradictory to results of other studies. Similar to this study's findings, Young et al. (2014) found a negative relationship between barriers to PA and accelerometer-measured MVPA among girls in 6th, 8th, and 11th grades. Although significant, the low correlation of $r = -.10$ between perceived barriers and MVPA in this study may be due to the bias resulting from a common method artifact of self-report. Dishman et al. (2010) explains that low correlations are not surprising when a subjective and objective measure are employed because use of self-report to measure both girls' beliefs, and their PA may lead to inflated relations between the two. Taking a different approach, Kelly et al. (2010) reported racial differences in the correlation between barriers to PA and accelerometer-measured MVPA, whereas Dishman et al. (2010) noted perceived barriers did not differ among girls of varied racial and ethnic backgrounds. Racial and ethnic influences on the relationship between perceived barriers and PA may need further investigation.

Consistent with previous studies, lack of skills followed by intrapersonal factors, such as hating to sweat, emerged as the most frequently reported perceived barriers among girls (Kelly et al., 2010; Rees et al., 2006). Similarly, among Canadian and Spanish adolescent girls, perceived incompetence in sports was found to be one of the top two barriers to PA (Bélanger et al., 2011; Zaragoza, Generelo, Julián, & Abarca-Sos, 2011). Interventions focused on skill development may be essential for girls to enhance their PA self-efficacy and other related factors to promote continued PA engagement (Humbert et al., 2006). Expressions of not being good enough to participate in PA figured prominently in previously conducted focus group discussions among Australian adolescent girls and face-to-face interviews among British adolescent girls (Stanley, Boshoff, & Dollman, 2013; Wetton, Radley, Jones, & Pearce, 2013). In the current study, because over 50% of the girls who were Black, in late puberty, and of low SES, reported lack of skills as a barrier, identifying ways to help girls—particularly those in these three subgroups—increase their PA skills is important for assisting them to achieve PA recommendations.

Several studies support that embarrassment, body image concerns, and physical discomfort, including sweating or fatigue, are barriers frequently reported by girls (Kelly et al., 2010; Wetton et al., 2013). Rees et al. (2006) found that barriers, such as body insecurity, were more concerning for girls than boys. Despite conflicting prior reports, racial or ethnic differences were evident in this study with the majority of Black girls, but not White girls, reporting sweating, as well as lack of skills, as barriers (Dishman et al., 2010; Kelly et al., 2010). Similar to findings reported by Robbins et al. (2003), the highest percentage of White girls in this study (46.9%) identified embarrassment or self-consciousness as a barrier.

Consistent with these results, hating to sweat during the school day emerged as the second most prevalent barrier in our study. This information underscores the importance of assisting girls to avoid negative body image and self-esteem issues, while simultaneously helping them to attain adequate PA, particularly as they progress through puberty.

Although finding PA programs or classes did not emerge as a major problem for girls who were not of low SES, 49.0% of low SES girls reported difficulty in this area. Though research examining SES and perceived barriers to PA among girls is limited, access issues, such as lack of PA facilities and cost of programs, have previously been reported as barriers to PA participation for youth of low SES (Humbert et al., 2006). Lack of access is also a major barrier to PA for Canadian adolescent girls (Bélanger et al., 2011). McCarron et al. (2010) found that across demographics, community members identified the need for external sources, such as schools, to contribute to a built environment that is supportive of PA for adolescents—especially girls—through collaborative partnerships with the community. Therefore, assisting girls of low SES to access PA programs or classes that they like should be a primary focus of future interventions.

Although lack of time for PA, often resulting from school-related academic responsibilities, emerged as a major barrier to PA among British girls participating in a prior study, the majority of girls in the total sample or any subgroup in this study (i.e., pubertal stage, race, or SES) did not identify lack of time as a barrier to PA (Wetton et al., 2013). A high percentage of girls in late puberty (46.4%), however, did report that others want them to do tasks other than PA with their time, perhaps resulting from having greater family obligations than those of early/middle pubertal girls. This barrier, identified in low SES adolescents, presumably increases with advancing maturity, and increased ability to participate in family obligations (Humbert et al., 2006). Researchers may need to consider unique family responsibilities and ways to improve time management skills when aiming to improve the PA of late pubertal girls.

Similar to lack of time, lack of motivation is consistently identified in the literature as a perceived barrier to PA among girls, indicating that interventions focused on enhancing motivation may lead to increased PA (Robbins et al., 2003; Robbins et al., 2009). Contrary to this information, lack of motivation was not reported by the majority of girls in this study. Consistent with a prior study showing that being tired interferes with PA among both male and female adolescents, ages 11 to 14, all subgroups in this study, except for those in late puberty, identified this barrier as a major problem (Robbins et al., 2009). Qualitative research may be needed with urban adolescent girls, particularly those in early or middle puberty, to explore reasons underlying this barrier, as well as social determinants of PA, so that they can be adequately addressed in interventions.

Implications

Future research is needed to design and test interventions that are specifically tailored to meet the needs of various subgroups of girls to target their perceived barriers to PA (Camacho-Miñano et al., 2011; Robbins et al., 2003; Robbins et al., 2009). Tailoring interventions based on biological and sociocultural characteristics, particularly pubertal stage and SES, may be a promising approach. Interventions that focus on skill development,

offer some measure of privacy for PA, provide access to showers and personal hygiene products, and include activities not likely to induce profuse sweating (i.e. brisk walking), may be more appealing to girls and efficacious in increasing their PA.

Adolescents in general and adolescent girls in particular, need to be presented with opportunities to provide input regarding the design of programs to increase their PA (Stanley et al., 2013). This approach may help uncover unique barriers and assist girls in being more invested in PA programs (Humbert et al., 2006; Stanley et al., 2013). Allowing girls to provide input regarding their PA programs may also lead to an increase in activities being offered that they find enjoyable and a decrease in those in which girls feel unskilled. This effort may result in a reduction of several major perceived barriers, including lack of skills, sweating during the school day, and an inability to find enjoyable PA programs. Humbert et al. (2006) suggested that, in interventions for low-SES youth in particular, emphasis needs to be placed on offering enjoyable PA and increasing students' skill development and confidence, while considering environmental factors. Girls of low SES may not have the resources, financial or otherwise, associated with PA participation away from the school venue; thus, physical education, after-school PA programs with transportation home afterward, and other school-based approaches (e.g., before-school PA programs, recess, class breaks for PA) are critical for assisting them to attain adequate PA. Perhaps, increasing parental support for PA by assisting parents to encourage and monitor their daughter's PA throughout the adolescent period may be one solution to overcoming some perceived barriers.

Conclusion

Increasing girls' PA continues to represent a major challenge. Barriers, including lack of skills, hating to sweat, difficulty finding programs, being tired, and having pain interfere with PA in urban girls prior to and during puberty. Designing interventions that assist girls to overcome the barriers they identified as interfering with PA may aid in overcoming the challenge.

Acknowledgments

The authors would like to acknowledge that the research was supported by Grant Number R01HL109101 from the National Heart, Lung, and Blood Institute (NHLBI) at the National Institutes of Health (NIH); PI: L. B. Robbins, Michigan State University (MSU) College of Nursing. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NHLBI or NIH. The "Girls on the Move Intervention" study was also funded by MSU College of Nursing. The funding bodies did not have a role in or influence the various phases of the project, the writing of the manuscript, or the decision to submit it for publication.

References

- Bélangier M, Casey M, Cormier M, Filion AL, Martin G, Aubut S, Beauchamp J. Maintenance and decline of physical activity during adolescence: Insights from a qualitative study. *International Journal of Behavioral Nutrition and Physical Activity*. 2011; 8:117. doi:10.1186/1479-5868-8-117. [PubMed: 22017754]
- Belcher BR, Berrigan D, Dodd KW, Emken BA, Chou C-P, Spruijt-Metz D. Physical activity in US youth: Effect of race/ethnicity, age, gender, and weight status. *Medicine and Science in Sports and Exercise*. 2010; 42:2211–2221. doi:10.1249/MSS.0b013e3181e1fba9. [PubMed: 21084930]

- Boyington JEA, Carter-Edwards L, Piehl M, Hutson J, Langdon D, McManus S. Cultural attitudes toward weight, diet, and physical activity among overweight African American girls. *Preventing Chronic Disease*. 2008; 5:A36. [PubMed: 18341772]
- Brodersen NH, Steptoe A, Boniface DR, Wardle J. Trends in physical activity and sedentary behaviour in adolescence: Ethnic and socioeconomic differences. *British Journal of Sports Medicine*. 2007; 41:140–144. doi:10.1136/bjsm.2006.031138. [PubMed: 17178773]
- Camacho-Miñano MJ, LaVoi NM, Barr-Anderson DJ. Interventions to promote physical activity among young and adolescent girls: A systematic review. *Health Education Research*. 2011; 26:1025–1049. doi:10.1093/her/cyr040. [PubMed: 21680763]
- Carskadon MA, Acebo C. A self-administered rating scale for pubertal development. *Journal of Adolescent Health*. 1993; 14:190–195. doi:10.1016/1054-139X(93)90004-9. [PubMed: 8323929]
- Chinapaw MJM, Mokkink LB, van Poppel MNM, van Mechelen W, Terwee CB. Physical activity questionnaires for youth: A systematic review of measurement properties. *Sports Medicine*. 2010; 40:539–563. doi:10.2165/11530770-000000000-00000. [PubMed: 20545380]
- Clarke PJ, O'Malley PM, Johnston LD, Schulenberg JE, Lantz P. Differential trends in weight-related health behaviors among American young adults by gender, race/ethnicity, and socioeconomic status: 1984–2006. *American Journal of Public Health*. 2009; 99:1893–1901. doi:10.2105/ajph.2008.141317. [PubMed: 19696395]
- Costigan SA, Barnett L, Plotnikoff RC, Lubans DR. The health indicators associated with screen-based sedentary behavior among adolescent girls: A systematic review. *Journal of Adolescent Health*. 2013; 52:382–392. doi:10.1016/j.jadohealth.2012.07.018. [PubMed: 23299000]
- Dishman RK, Dunn AL, Sallis JF, Vandenberg RJ, Pratt CA. Social-cognitive correlates of physical activity in a multi-ethnic cohort of middle-school girls: Two-year prospective study. *Journal of Pediatric Psychology*. 2010; 35:188–198. doi:10.1093/jpepsy/jsp042. [PubMed: 19468040]
- Dumith SC, Gigante DP, Domingues MR, Kohl HW III. Physical activity change during adolescence: a systematic review and a pooled analysis. *International Journal of Epidemiology*. 2011; 40:685–698. doi:10.1093/ije/dyq272. [PubMed: 21245072]
- Evenson KR, Catellier DJ, Gill K, Ondrak KS, McMurray RG. Calibration of two objective measures of physical activity for children. *Journal of Sports Sciences*. 2008; 26:1557–1565. doi:10.1080/02640410802334196. [PubMed: 18949660]
- Finne E, Bucksch J, Lampert T, Kolip P. Age, puberty, body dissatisfaction, and physical activity decline in adolescents. Results of the German Health Interview and Examination Survey (KiGGS). *International Journal of Behavioral Nutrition and Physical Activity*. 2011; 8:119. doi:10.1186/1479-5868-8-119. [PubMed: 22032266]
- Humbert ML, Chad KE, Spink KS, Muhajarine N, Anderson KD, Bruner MW, Gryba CR. Factors that influence physical activity participation among high- and low-SES youth. *Qualitative Health Research*. 2006; 16:467–483. doi:10.1177/1049732305286051. [PubMed: 16513991]
- Kann L, Kinchen S, Shanklin SL, Flint KH, Hawkins J, Harris WA, Zaza S. Youth risk behavior surveillance—United States, 2013. *MMWR Surveillance Summaries*. 2014; 63:1–168.
- Kelly EB, Parra-Medina D, Pfeiffer KA, Dowda M, Conway TL, Webber LS, Pate RR. Correlates of physical activity in Black, Hispanic, and White middle school girls. *Journal of Physical Activity & Health*. 2010; 7:184–193. [PubMed: 20484757]
- Lanza HI, Echols L, Graham S. Deviating from the norm: Body mass index (BMI) differences and psychosocial adjustment among early adolescent girls. *Journal of Pediatric Psychology*. 2013; 38:376–386. doi:10.1093/jpepsy/jss130. [PubMed: 23248348]
- Larson NI, Wall MM, Story MT, Neumark-Sztainer DR. Home/family, peer, school, and neighborhood correlates of obesity in adolescents. *Obesity*. 2013; 21:1858–1869. doi:10.1002/oby.20360. [PubMed: 23512596]
- Ling J, Robbins LB, Resnicow K, Bakhoya M. Social support and peer norms scales for physical activity in adolescents. *American Journal of Health Behavior*. 2014; 38:881–889. doi:10.5993/AJHB.38.6.10. [PubMed: 25207514]
- McCarron DA, Richartz N, Brigham S, White MK, Klein SP, Kessel SS. Community-based priorities for improving nutrition and physical activity in childhood. *Pediatrics*. 2010; 126:S73–S89. doi:10.1542/peds.2010-0482C. [PubMed: 21041288]

- Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of childhood and adult obesity in the United States, 2011–2012. *JAMA*. 2014; 311:806–814. doi:10.1001/jama.2014.732. [PubMed: 24570244]
- Patnode CD, Lytle LA, Erickson DJ, Sirard JR, Barr-Anderson DJ, Story M. Physical activity and sedentary activity patterns among children and adolescents: A latent class analysis approach. *Journal of Physical Activity and Health*. 2011; 8:457–467. [PubMed: 21597117]
- Pender, NJ.; Murdaugh, CL.; Parsons, MA. *Health promotion in nursing practice*. 6th. Prentice Hall; Upper Saddle River, NJ: 2011.
- Peterson AC, Crockett L, Richards M, Boxer A. A self-report measure of pubertal status: Reliability, validity, and initial norms. *Journal of Youth and Adolescence*. 1988; 17:117–133. doi:10.1007/BF01537962. [PubMed: 24277579]
- Rees R, Kavanagh J, Harden A, Shepherd J, Brunton G, Oliver S, Oakley A. Young people and physical activity: A systematic review matching their views to effective interventions. *Health Education Research*. 2006; 21:806–825. doi:10.1093/her/cyl120. [PubMed: 17041020]
- Robbins LB, Pender NJ, Kazanis AS. Barriers to physical activity perceived by adolescent girls. *Journal of Midwifery & Women's Health*. 2003; 48:206–212. doi:10.1016/S1526-9523(03)00054-0.
- Robbins LB, Pfeiffer KA, Vermeesch A, Resnicow K, You Z, An L, Wesolek SM. “Girls on the Move” intervention protocol for increasing physical activity among low-active underserved urban girls: A group randomized trial. *BMC Public Health*. 2013; 13:474. doi: 10.1186/1471-2458-13-474. [PubMed: 23672272]
- Robbins LB, Sikorskii A, Hamel LM, Wu T-Y, Wilbur J. Gender comparisons of perceived benefits of and barriers to physical activity in middle school youth. *Research in Nursing & Health*. 2009; 32:163–176. doi:10.1002/nur.20311. [PubMed: 19086055]
- Robbins LB, Wu T-Y, Sikorskii A, Marley B. Psychometric assessment of the adolescent physical activity perceived benefits and barriers scales. *Journal of Nursing Measurement*. 2008; 16:98–112. doi:10.1891/1061-3749.16.2.98. [PubMed: 18939715]
- Ryan RM, Deci EL. Self-determination theory and the facilitation of intrinsic motivation, social development, and well-being. *American Psychologist*. 2000; 55:68–78. doi:10.1037/0003-066X.55.1.68. [PubMed: 11392867]
- Smart JE, Cumming SP, Sherar LB, Standage M, Neville H, Malina RM. Maturity associated variance in physical activity and health-related quality of life in adolescent females: A mediated effects model. *Journal of Physical Activity & Health*. 2012; 9:86–95. [PubMed: 22232510]
- Stanley RM, Boshoff K, Dollman J. A qualitative exploration of the “critical window”: Factors affecting Australian children’s after-school physical activity. *Journal of Physical Activity & Health*. 2013; 10:33–41. http://journals.humankinetics.com/AcuCustom/SiteName/Documents/DocumentItem/04_stanley_JPAH_20100231-eja.pdf [PubMed: 22397802]
- State of Michigan. MI school data. 2014. Retrieved from <http://www.mischooldata.org>
- Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M. Physical activity in the United States measured by accelerometer. *Medicine and Science in Sports and Exercise*. 2008; 40:181–188. doi: 10.1249/mss.0b013e31815a51b3. [PubMed: 18091006]
- Trost SG, Loprinzi PD, Moore R, Pfeiffer KA. Comparison of accelerometer cut points for predicting activity intensity in youth. *Medicine and Science in Sports and Exercise*. 2011; 43:1360–1368. doi: 10.1249/Mss.0b013e318206476e. [PubMed: 21131873]
- U.S. Census Bureau. State & County QuickFacts. 2014. Retrieved from <http://quickfacts.census.gov/qfd/states/26000.html>
- U.S. Department of Health & Human Services. 2008 Physical Activity Guidelines for Americans. 2008. Retrieved from <http://www.health.gov/paguidelines/guidelines/summary.aspx>
- Van der Horst K, Paw MJCA, Twisk JW, Van Mechelen W. A brief review on correlates of physical activity and sedentariness in youth. *Medicine & Science in Sports & Exercise*. 2007; 39:1241–1250. doi:10.1249/mss.0b013e318059bf35. [PubMed: 17762356]
- Wang Y, Liang H, Tussing L, Braunschweig C, Caballero B, Flay B. Obesity and related risk factors among low socio-economic status minority students in Chicago. *Public Health Nutrition*. 2007; 10:927–938. doi:10.1017/S1368980007658005. [PubMed: 17381954]

- Waylen A, Wolke D. Sex 'n' drugs 'n' rock 'n' roll: The meaning and social consequences of pubertal timing. *European Journal of Endocrinology*. 2004; 151:U151–U159. doi:10.1530/eje.0.151U151. [PubMed: 15554900]
- Webber LS, Catellier DJ, Lytle LA, Murray DM, Pratt CA, Young DR, Pate RR. Promoting physical activity in middle school girls: Trial of Activity for Adolescent Girls. *American Journal of Preventive Medicine*. 2008; 34:173–184. doi:10.1016/j.amepre.2007.11.018. [PubMed: 18312804]
- Wetton AR, Radley R, Jones AR, Pearce MS. What are the barriers which discourage 15-16 year-old girls from participating in team sports and how can we overcome them? *BioMed Research International*. 2013;2013 doi:10.1155/2013/738705.
- World Health Organization. *Global recommendations on physical activity for health*. WHO Press; Geneva, Switzerland: 2010.
- Young D, Saksvig BI, Wu TT, Zook K, Li X, Champaloux S, Treuth MS. Multilevel correlates of physical activity for early, mid, and late adolescent girls. *Journal of Physical Activity & Health*. 2014; 11:950–960. doi:10.1123/jpah.2012-0192. [PubMed: 23676305]
- Zaragoza J, Generelo E, Julián JA, Abarca-Sos A. Barriers to adolescent girls' participation in physical activity defined by physical activity levels. *Journal of Sports Medicine & Physical Fitness*. 2011; 51:128–135. [PubMed: 21297572]

TABLE 1

Sample Characteristics

Characteristic	<i>n</i>	(%)
Age		
10	79	(15.5)
11	252	(49.5)
12	152	(29.9)
13	23	(4.5)
14	3	(0.6)
Grade		
5th	69	(13.6)
6th	284	(55.8)
7th	156	(30.6)
Pubertal stage		
Early/middle puberty	300	(58.9)
Late puberty	209	(41.1)
Hispanic ethnicity (yes) ^a	62	(12.8)
Race		
Black	302	(59.3)
White	113	(22.2)
Mixed/other	94	(18.5)
Free or reduced-price lunch (yes) ^{b,c}	402	(84.8)
Weight status ^d		
Underweight	5	(1.0)
Normal weight	229	(45.2)
Overweight	109	(21.5)
Obese	164	(32.3)

Note. *N* = 509.

^a missing = 25.

^b missing = 35.

^c Participation in free or reduced-price lunch program as an indicator for low SES.

^d missing = 2.

TABLE 2

BMI, Screen Time, Sedentary Activity, Physical Activity, and Perceived Barriers by Pubertal Stage

Variable	All		Early-middle		Late		<i>p</i>
	<i>M</i>	(<i>SD</i>)	<i>M</i>	(<i>SD</i>)	<i>M</i>	(<i>SD</i>)	
BMI ^a	23.0	(5.48)	22.2	(5.34)	24.2	(5.49)	< .001
BMI <i>z</i> -score ^a	1.0	(1.00)	0.87	(1.07)	1.2	(0.85)	< .001
Screen time ^b	6.2	(3.35)	5.7	(3.27)	7.0	(3.32)	< .001
Sedentary activity ^c	38.2	(4.39)	37.2	(4.04)	39.7	(4.46)	< .001
MVPA ^c	3.1	(1.32)	3.3	(1.32)	2.7	(1.24)	< .001
LMVPA ^c	21.8	(4.39)	22.8	(4.04)	20.3	(4.46)	< .001
Perceived barriers	1.1	(0.58)	1.0	(0.56)	1.2	(0.59)	< .001

Note. *N* = 509. BMI = body mass index; LMVPA = light plus moderate-to-vigorous physical activity; MVPA = moderate-to-vigorous physical activity.

^a missing = 2.

^b missing = 3.

^c missing = 47.

TABLE 3

Moderate-to-Vigorous Physical Activity: Mixed Effects Model

Predictor	Estimate	(SE)	<i>p</i>
Intercept	5.91	(1.29)	< .001
Non-Hispanic	.04	(.20)	.86
SES (high) ^a	.04	(.19)	.83
Early/middle puberty	.47	(.15)	.002
Race (Black)	.14	(.17)	.41
Race (mixed/other)	.08	(.21)	.70
Age	-.26	(.11)	.01
BMI z-score	-.02	(.07)	.78
<u>Perceived barriers</u>	-.14	(.11)	.22
<u>Variance component</u>			
School	.05	(.04)	.28
Residual	1.68	(.12)	< .001

Note. *N* = 509. BMI = body mass index; SES = economic status.

^aNo participation in free/reduced-price lunch program was used as an indicator for high SES.

TABLE 4

Top Five Perceived Barriers to Physical Activity for the Total Sample and by Pubertal Stage, Race, and SES

Barrier	Total (N = 509)	Pubertal stage		Race			Low SES ^a	
		Early/mid (n = 300)	Late (n = 209)	Black (n = 302)	White (n = 113)	Mixed (n = 94)	No (n = 72)	Yes (n = 402)
Need better skill	51.5	47.7	56.9	51.0	45.1	--	48.6	52.5
Hate to sweat	49.7	45.0	56.5	54.6	38.9	46.8	36.1	54.2
Hard to find PA programs	45.6	40.0	53.6	48.0	--	47.9	--	49.0
Am tired	42.6	41.7	--	41.7	44.2	43.6	37.5	44.3
Pain from activity	41.7	41.0	--	40.0	44.2	43.6	37.5	--
Embarrassed: looks during exercise	--	--	49.3	--	46.9	44.7	36.1	42.8
Other things to do	--	--	46.4	--	--	--	--	--

Note. Entries are percent responding “somewhat true” or “very true.” The top five most frequently rated items are shown for each group.
SES = socioeconomic status.

^aParticipation in free/reduced-price lunch program was used as an indicator for low SES.