



Published in final edited form as:

Biochem Biophys Res Commun. 2015 September 25; 465(3): 464–470. doi:10.1016/j.bbrc.2015.08.040.

Isoform-specific dynamic translocation of PKC by α_1 -adrenoceptor stimulation in live cells

Jin O-Uchi^{1,2,*},§, Jaime Sorenson²,§, Bong Sook Jhun¹, Jyotsna Mishra¹, Stephen Hurst¹, Kaleef Williams², Shey-Shing Sheu¹, and Coeli M.B. Lopes^{2,*}

¹Center for Translational Medicine, Department of Medicine, Sidney Kimmel Medical College, Thomas Jefferson University, Philadelphia, PA, 19107 USA

²Aab Cardiovascular Research Institute, Department of Medicine, University of Rochester School of Medicine and Dentistry, Rochester, NY, 14642 USA

Abstract

Protein kinase C (PKC) plays key roles in the regulation of signal transduction and cellular function in various cell types. At least ten PKC isoforms have been identified and intracellular localization and trafficking of these individual isoforms are important for regulation of enzyme activity and substrate specificity. PKC can be activated at downstream of G_q-protein coupled receptor (G_qPCR) signaling and translocated to the various cellular compartments including plasma membrane (PM). Recent reports suggested that a different type of G_qPCRs would activate different PKC isoforms (classic, novel and atypical PKCs) with different trafficking patterns. However, the knowledge of isoform-specific activation of PKC by each G_qPCR is limited. α_1 -Adrenoceptor (α_1 -AR) is the one of the G_qPCR highly expressed in the cardiovascular system. In this study, we examined the isoform-specific dynamic translocation of PKC in living HEK293T cells by α_1 -AR stimulation (α_1 -ARS). Rat PKC α , β I, β II, δ , ϵ and ζ fused with GFP at C-term were co-transfected with human α_{1A} -AR into HEK293T cells. The isoform-specific dynamic translocation of PKC in living HEK293T cells by α_1 -ARS using phenylephrine was measured by confocal microscopy. Before stimulation, GFP-PKCs were localized at cytosolic region. α_1 -ARS strongly and rapidly translocated a classical PKC (cPKC), PKC α , (< 30s) to PM, with PKC α returning diffusively into the cytosol within 5 min. α_1 -ARS rapidly translocated other cPKCs, PKC β I and PKC β II, to the PM (<30s), with sustained membrane localization. One of novel PKCs (nPKCs), PKC ϵ , but not another nPKC, PKC δ , was translocated by α_1 -AR stimulation to the PM (<30s) and its membrane localization was also sustained. Finally, α_1 -AR stimulation did not cause a diacylglycerol-insensitive atypical PKC, PKC ζ translocation. Our data suggest that PKC α , β and ϵ activation may underlie physiological and pathophysiological responses of α_1 -AR signaling for

*Correspondence to: Coeli M.B. Lopes, Ph.D., Cardiovascular Research Institute, Department of Medicine, University of Rochester, 601 Elmwood Ave, Box CVRI, Rochester, NY, 14642, USA. Phone: 585-276-9784, Fax: 585-275-9830.

coeli_lopes@urmc.rochester.edu Or Jin O-Uchi, M.D., Ph.D., Center for Translational Medicine, Department of Medicine, Sidney Kimmel Medical College, Thomas Jefferson University, 1020 Locust Street, Suite543G, Philadelphia, PA 19107, Phone: 215-503-1567, Fax: 215-503-5731, jin.ouchi@jefferson.edu.

§Both authors contributed equally.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

the phosphorylation of membrane-associated substrates including ion-channel and transporter proteins in the cardiovascular system.

INTRODUCTION

Protein kinase C (PKC) is a multi-gene family of serine/threonine kinase that plays key roles in the regulation of signal transduction and cellular function in various cell-types/tissues [1-4]. PKC forms a multi-gene family and at least 10 PKC isoforms have been identified that differ in primary structure, tissue distribution, subcellular localization and substrate specificity [1-4]. These isoforms can be sub-grouped into three subfamilies. Members of the first family (include PKC α , β I, β II and γ) are regulated by Ca²⁺, diacylglycerol (DAG) and phospholipids, and are known as Ca²⁺-dependent PKCs [or classical PKCs (cPKC)]; members of the second family (include PKC δ , ϵ , η and θ) have phospholipid-dependent but Ca²⁺-independent activation mechanism, and are known as Ca²⁺-independent PKCs or novel PKCs (nPKC); members of the third family (include PKC ζ and ι/λ) are not regulated by either intracellular Ca²⁺ concentration or phospholipid and their activity are maintained by other mechanisms (e.g. by protein-protein interactions), and are known as atypical PKCs (aPKC). Intracellular localizations of these individual isoforms are important for regulation of isoform-specific enzyme activity and substrate specificity [2,3]. Since the stimulation of G_qPCRs such as α_1 -adrenoceptor (α_1 -AR), endothelin-1 receptor or angiotensin II receptor can generate DAG and also mobilize cytosolic Ca²⁺ elevation by releasing Ca²⁺ from the intracellular stores by stimulating inositol trisphosphate (IP₃) receptors [5], cPKC and nPKC isoforms, but not aPKC are located downstream of G_qPCR signaling pathways. Growing evidence suggests that stimulation of different G_qPCRs exhibits receptor-specific pattern of activation and translocation of PKC isoforms [3]. However, since prior research investigating PKC translocation were mostly conducted using general PKC activators, such as phorbol esters [3], the knowledge of receptor-specific regulation of PKC-isoform activation/translocation is still incomplete.

One of G_qPCRs, α_1 -AR, is stimulated by catecholamines (norepinephrine and epinephrine) [6-9]. α_1 -AR is expressed in variety of human tissue [10] and α_1 -AR stimulation has been shown to play important roles in cellular physiological functions, such as 1) regulation of smooth muscle contraction and tone in vascular system [9], prostate, urethra, bladder [11], uterine [12] and iris [13], 2) myocardial inotropy and chronotropy [14], 3) hepatic glucose metabolism [15], 4) water secretion at salivary gland [16] and 5) neurotransmission in central nerve system [9]. In addition, chronic α_1 -AR stimulation leads to pathophysiological responses in the various cells/tissues via both cPKC and nPKC isoform signaling pathways, including cardiac hypertrophy [17,18], hypertension and atherosclerosis [19,20] in cardiovascular system and portal hypertension and fibrosis in liver [21]. Despite strong interest in the mechanism underlying α_1 -AR signaling-mediated pathology, especially in cardiovascular system, little is known about the molecular mechanisms for the PKC isoform-specific kinetics of activation and translocation.

Here we examined the isoform-specific dynamic translocation of PKC in live HEK293T cells by α_1 -AR stimulation using GFP-tagged PKC isoforms, especially monitoring their

translocation to the plasma membrane (PM). We tested six PKC isoforms (α , β I, β II, δ , ϵ and ζ) in this study which are endogenously expressed in this cell line [22]. Our results show that PKC α , β I, β II and ϵ are able to be translocated to the PM upon α_1 -AR stimulation, while PKC δ did not show any significant PM translocation upon α_1 -ARS, although all isoforms are translocated to PM by a common PKC activator, phorbol 12-myristate 13-acetate (PMA) with a relatively slow time course. Our data suggest that PKC α , β I, β II and ϵ activation may underlie physiological responses of α_1 -AR signaling. In addition, due to a prolonged retention of PKC β I, PKC β II and PKC ϵ at PM by α_1 -AR stimulation, these isoforms may be particularly important for the cellular pathophysiology in various cell types including cardiovascular system during chronic α_1 -AR stimulation.

MATERIAL AND METHODS

An expanded Material and methods section is available in the online supplementary file.

Live cell imaging

HEK293T cells and stable HEK293T cell line [23] carrying HA-tagged α_{1A} -AR (Supplementary Fig.1) were transfected with GFP-tagged and/or DsRed2-tagged PKC isoforms and used for experiments 48 hours after transfection [24,25]. Time-dependent changes in localization of PKC isoforms by either phenylephrine (Phe) or PMA stimulation in HEK293T cells was measured using laser scanning confocal microscopy (Olympus, Tokyo Japan) without fixation at room temperature [25,26]. PM localization of PKC isoform was quantified by line scan intensity measurements through each cell beginning in the cytosol region (avoiding the nucleus area) and ending at the cell periphery. Translocation of each PKC isoform was evaluated by fluorescence intensity ratio between membrane and cytosolic region (membrane/cytosol ratio: M/C ratio) [25,26].

Statistical Analysis

All results are shown as mean \pm standard error (SE). One-way ANOVA followed by Dunnet's test (for multiple comparison) and paired-T-test (for two-group comparison) were done with the significance level set at $P < 0.05$.

RESULTS

PKC α transiently translocates to the PM in response to α_1 -ARS

We co-expressed a GFP-tagged Ca^{2+} -dependent PKC isoform PKC α with α_{1A} -AR in HEK293T cells, stimulated the cells with a specific α_1 -AR agonist 100 μM Phe, and observed time-dependent changes in the subcellular localization of PKC α upon Phe stimulation (Fig.1A). Prior to stimulation (0 min), PKC α -GFP mainly localized in cytosol, with no nuclear localization (Fig.1A and B). After Phe stimulation PKC α -GFP rapidly translocated to the cell membrane (less than 30s) and gradually returned to cytosol. PKC α localization was restored within 5 min despite continuous Phe stimulation. This PKC translocation was completely blocked by the α_1 -AR-selective antagonist, 1 μM prazosin (Fig.1B and Supplementary Fig.2). We also tested the effect of PKC activator PMA on

PKC α -GFP translocation. PKC α -GFP showed relatively slower translocation to PM by PMA treatment compared to Phe (Fig.1B and Supplementary Fig.2).

As a control experiment, we co-expressed GFP with α_{1A} -AR in HEK293 cells and confirmed that GFP localization did not alter by Phe stimulation (Supplementary Fig.3).

PKC β I and PKC β II translocate to the PM in response to α_1 -ARS

We next co-expressed other Ca²⁺-dependent PKC isoforms (GFP-tagged PKC β I or PKC β II) with α_{1A} -AR in HEK293T cells and observed time-dependent changes in the subcellular localization of each PKC isoform by Phe (Fig.1C to F). Prior to stimulation, both PKC β I and PKC β II were localized in the cytosol, with no nuclear localization. Phe stimulation induced translocation of both PKC β I and PKC β II to the PM rapidly (after 30 sec) and PM localization was maintained throughout the course of the Phe application. Both PKC β I-GFP and PKC β II-GFP showed slower translocation to the PM by a PKC activator PMA treatment compared to Phe stimulation (Fig.1D and F), which is a similar time course as the one observed in PKC α (see Fig.1B). Interestingly, magnitude of PKC β II translocation by Phe stimulation was larger than that of PKC β I (Fig.1D and F).

To confirm the different translocation dynamics of PKC β I and β II by either Phe or PMA, we stimulated cells expressing both GFP- and DsRed2-tagged PKC β s. To normalize the magnitude of signal inputs (namely receptor signals via the G_q-protein) between the individual cells, HEK293T cells stably overexpressing α_{1A} -AR (α_{1A} -AR-HEK293T cells) were used for these experiments (Supplementary Fig.1). First, we confirmed that GFP- and DsRed2-tagged PKC β II showed similar translocation profiles under Phe or PMA stimulations in α_{1A} -AR-HEK293T (Fig.1G and H). Using α_{1A} -AR-HEK293T cells co-expressing PKC β I-GFP and DsRed2-PKC β II, we next confirmed that magnitude of PKC β II translocation by Phe stimulation was significantly larger than that of PKC β I (Fig.1I and J). In contrast, PMA stimulation induces larger translocation of PKC β I than that of PKC β II (Fig.1J).

PKC ϵ but not PKC δ translocate to the PM in response to α_1 -ARS

We next tested whether Ca²⁺-independent PKC isoforms translocates to the PM by α_1 -ARS. We co-expressed GFP-tagged PKC δ or PKC ϵ with α_{1A} -AR in HEK293T cells and observed time-dependent changes in the subcellular localization of these PKC isoforms upon Phe application. Prior to stimulation, PKC δ was localized in cytosol (Fig.2A). Some cells have strong punctuated fluorescence areas around the nucleus (Supplementary Fig.4). No significant time-dependent changes were observed in the subcellular localization of PKC δ by Phe treatment (Fig.2A and B). As a positive control, we observed significant translocation of PKC δ in response to treatment with PMA in the same cells (Fig2A and B). After PMA treatment, PKC δ was translocated from the cytosol to the PM and also to the nuclear membrane (Fig.2A, 2B and supplementary Fig.4). We also confirmed the lack of GFP-tagged PKC δ translocation by Phe using α_{1A} -AR-HEK293T cells co-transfected with DsRed2-PKC β II (Fig.2C). In contrast, PKC ϵ was significantly translocated from the cytosol to the PM 3 min after Phe stimulation and its PM localization was maintained during the course of the Phe application (Fig.2D and E). Magnitude of GFP-tagged PKC ϵ translocation

by 10-min Phe stimulation was larger than that in PKC β I and also reached to the similar level of that in PKC β II (see Fig.1D and F). To quantitatively compare the translocation levels of PKC ϵ and PKC β I by Phe, we stimulated α_1 A-AR-HEK293T cells expressing both GFP-PKC ϵ and DsRed2-tagged PKC β I. The magnitude of PKC ϵ translocation was consistently higher after Phe stimulation than that of PKC β I when measured in the same cell (Fig.2F, left). In contrast, PMA stimulation induces similar translocation profiles in these PKC isoforms (Fig.2F, right).

PKC ζ does not show any translocation in response to α_1 -ARS

Finally, we tested whether DAG- and PMA-insensitive isoform (aPKC) change its subcellular localization in response to α_1 -ARS. We co-expressed one of the aPKC isoform PKC ζ in HEK293 cells and monitored time-dependent changes in the subcellular localization of PKC ζ upon Phe application. Prior to stimulation, PKC ζ was localized in cytosol and no nuclear localization was observed (Fig. 3). There was no significant time-dependent change in the subcellular localization of PKC ζ by α_1 -ARS.

DISCUSSION

In this study, we examined the translocation profiles of PKC isoforms to the PM in live cells expressing GFP- and DsRed2-tagged PKC isoforms upon α_1 -AR stimulation. One advantage of using fluorescence protein-tagged PKC expression system is the ability to obtain more precise and quantitative spatiotemporal kinetics information on PKC activation and translocation profiles compared to classical methods, including Western blotting of fractionated proteins and the immunocytochemistry of fixed cells using isoform-specific antibodies [2,3]. Our results clearly indicate that each PKC isoform shows a different spatiotemporal pattern of translocation under α_1 -ARS (Fig. 4). We found the significant translocation of Ca^{2+} -dependent PKCs (α , β I and β II) and one Ca^{2+} -dependent PKC (PKC ϵ) to the PM in response to α_1 -AR stimulation, which may underlie acute responses of α_1 -AR signaling at the PM. However, subcellular localization of another Ca^{2+} -dependent PKC, PKC δ was not significantly modified by α_1 -ARS in the current experiments, despite showing a strong response to PMA (Fig.2).

Physiological and pathophysiological functions of each PKC isoform are diverse even though their substrate specificity is low and multiple isoforms are expressed in each cell-type [2,3]. For instance, in the human myocardium, PKC α was the most abundant isoform present [19; 20] and PKC α activation has been linked to heart failure, hypertrophy and diabetes [21-27]. However, subsequent studies in genetic models of PKC α overexpression or knockout in mice failed to yield significant role of PKC α in generating hypertrophy to PKC α , identifying a more predominant role for PKC α in the regulation of cardiac contraction [28]. On the other hand, expression level of Ca^{2+} -dependent PKC β increases in the cardiovascular system during disease states. PKC β activation has been linked to increased vascular inflammation and atherosclerosis [29]. In addition, genetic models showed that PKC β expression is not necessary for the development of cardiac hypertrophy nor does its absence attenuate the hypertrophic response [30]. PKC ϵ has been suggested to have a protective role in myocardial ischemia-reperfusion injury [31] and arrhythmias [32].

Moreover, the use of specific PKC isoenzyme blockers may lead to novel treatments for heart failure [35; 36] and also for cancers [33] and Alzheimer's disease [34]. Targeting PKC to the PM is crucial for the phosphorylation of membrane-associated substrates, including ion-channel and transporter proteins [27]. PMA is one of the well recognized activators of DAG-sensitive PKC subfamilies and is frequently used to mimic G_q PCR signaling for investigating the effect of PKC translocation [2,3] to the PM, including in the studies for the regulation of ion channel function. However, we showed in live cells that the translocation kinetics of PKCs are largely different in response to either PMA treatment or receptor stimulation. First, α_1 -AR stimulation by Phe initiates PKC translocation much faster than PMA started (≈ 30 sec after stimulation). Second, while G_q PCR-mediated PKC α translocation to the PM is transient, PMA treatment induces this isoform to be retained at the PM (Figs.1 and 4). Finally, isoform-specificity is also different between G_q PCR and PMA responses; PMA caused sustained translocation of all DAG-sensitive isoforms tested to the PM (Figs.1 and 2), while α_1 -AR-mediated PKC δ translocation to the PM was not observed (Fig.2). This observation is consistent with previous data showing that connexin43, a gap junction channel protein, can be phosphorylated by PMA treatment through both PKC δ and PKC ϵ [28,29], but PKC δ does not likely phosphorylate Connexin43 under G_q PCR stimulation [30]. Thus, these results indicate that great caution is required when extending the results obtained with general PKC activators to PKC isoform activation by more physiological signaling pathways.

While we did not observe significant PKC δ translocation by α_1 -AR stimulation from cytosol to the PM using transfected PKC δ -GFP, our data does not exclude the possibility that PKC δ can be activated by α_1 -AR stimulation and translocated to the other cellular compartments including mitochondria and nucleus. For instance, Newton and colleagues detected a relatively small but significant PKC δ translocation to the outer mitochondrial membrane (OMM) after the application of a phorbol ester using fluorescence resonance energy (FRET) by generating an OMM-targeted [31] CFP (mt-CFP) and PKC δ -GFP [32]. They also confirmed that PKC δ activity was increased at the OMM upon phorbol-ester stimulation, using the OMM-targeted FRET-based PKC δ kinase activity reporter. Taken together, the data suggest that G_q PCR stimulation may activate and translocate PKC δ to other intracellular compartments, but not to PM.

In summary, we showed that α_1 -AR induces PKC α , PKC β and PKC ϵ translocation to the PM, but not PKC δ and PKC ζ by tracking the subcellular localization of transfected GFP-tagged PKC after Phe stimulation. In addition, we found that kinetics of α_1 -AR-mediated activation of PKCs is isoform-specific. Our data suggests that activation of PKC α , PKC β and PKC ϵ may underlie physiological and pathophysiological responses of α_1 -AR signaling for the phosphorylation of PM-localized proteins such as ion channels and transporters.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGEMENTS

The authors thank Drs. Naoaki Sato and John Hepler for providing us the plasmids of PKC-isoforms and α_{1A} -adrenoceptor, respectively, and Dr. Elena Fujiwara, Ms. Nobiru Suzuki, Ms. Mehreen Butt, Ms. Sarah Monaco and Mr. Michael Cypress for their technical assistance. This work was partly supported by AHA grants (09POST2310079 and 14BGIA18830032 to J.O.-U.), Medical Research Grant from W.W. Smith Charitable Trust (to J.O.-U.), Foreign Study Grant Award of Kanae Foundation (to J.O.-U.), Irisawa Memorial Promotion Award for Young Physiologists from the Physiological Society of Japan (to J.O.-U.) and NIH grants (5T32AA007463-26 to S.H, R01HL093671 to S.-S.S. and R01HL114792 to C.M.B.L.).

Abbreviations

α_1 -AR	α_1 -adrenoceptor
α_{1A} -AR-HEK293T cells	HEK293T cells stably overexpressing α_{1A} -AR
aPKC	atypical protein kinase C
cPKC	classical protein kinase
DAG	diacylglycerol
nPKC	novel protein kinase
Phe	phenylephrine
G _q PCR	G _q -protein coupled receptor
M/C ratio	membrane/cytosol ratio
PKC	protein kinase C
PM	plasma membrane

Reference List

- [1]. Spitaler M, Cantrell DA. Protein kinase C and beyond. *Nat. Immunol.* 2004; 5:785–790. [PubMed: 15282562]
- [2]. Steinberg SF. Structural basis of protein kinase C isoform function. *Physiol. Rev.* 2008; 88:1341–1378. [PubMed: 18923184]
- [3]. Shirai Y, Saito N. Activation mechanisms of protein kinase C: maturation, catalytic activation, and targeting. *J. Biochem.* 2002; 132:663–668. [PubMed: 12417013]
- [4]. Wu-Zhang AX, Newton AC. Protein kinase C pharmacology: refining the toolbox. *Biochem. J.* 2013; 452:195–209. [PubMed: 23662807]
- [5]. Kang M, Chung KY, Walker JW. G-protein coupled receptor signaling in myocardium: not for the faint of heart. *Physiology (Bethesda).* 2007; 22:174–184. [PubMed: 17557938]
- [6]. O-Uchi J, Sasaki H, Morimoto S, Kusakari Y, Shinji H, Obata T, Hongo K, Komukai K, Kurihara S. Interaction of alpha1-adrenoceptor subtypes with different G proteins induces opposite effects on cardiac L-type Ca²⁺ channel. *Circ. Res.* 2008; 102:1378–1388. [PubMed: 18467629]
- [7]. Brodde OE, Michel MC. Adrenergic and muscarinic receptors in the human heart. *Pharmacol. Rev.* 1999; 51:651–690. [PubMed: 10581327]
- [8]. Zhong H, Minneman KP. Alpha1-adrenoceptor subtypes. *Eur. J. Pharmacol.* 1999; 375:261–276. [PubMed: 10443582]
- [9]. Piascik MT, Perez DM. Alpha1-adrenergic receptors: new insights and directions. *J. Pharmacol. Exp. Ther.* 2001; 298:403–410. [PubMed: 11454900]
- [10]. Price DT, Lefkowitz RJ, Caron MG, Berkowitz D, Schwinn DA. Localization of mRNA for three distinct alpha 1-adrenergic receptor subtypes in human tissues: implications for human alpha-adrenergic physiology. *Mol. Pharmacol.* 1994; 45:171–175. [PubMed: 8114668]

- [11]. Roehrborn CG. Efficacy of alpha-Adrenergic Receptor Blockers in the Treatment of Male Lower Urinary Tract Symptoms. *Rev. Urol.* 2009; 11:S1–8. [PubMed: 20126606]
- [12]. Dupuis M, Houdeau E, Mhaouty-Kodja S. Increased potency of alpha 1-adrenergic receptors to induce inositol phosphates production correlates with the up-regulation of alpha 1d/Gh alpha/phospholipase C delta 1 signaling pathway in term rat myometrium. *Reproduction.* 2008; 135:55–62. [PubMed: 18159083]
- [13]. Michel MC, Okutsu H, Noguchi Y, Suzuki M, Ohtake A, Yuyama H, Yanai-Inamura H, Ukai M, Watanabe M, Someya A, Sasamata M. In vivo studies on the effects of alpha1-adrenoceptor antagonists on pupil diameter and urethral tone in rabbits. *Naunyn Schmiedebergs Arch. Pharmacol.* 2006; 372:346–353. [PubMed: 16489448]
- [14]. Varma DR, Deng XF. Cardiovascular alpha1-adrenoceptor subtypes: functions and signaling. *Can. J. Physiol. Pharmacol.* 2000; 78:267–292. [PubMed: 10772055]
- [15]. Fagan TE, Romani A. Activation of Na(+)- and Ca(2+)-dependent Mg(2+) extrusion by alpha(1)- and beta-adrenergic agonists in rat liver cells. *Am. J. Physiol. Gastrointest. Liver Physiol.* 2000; 279:G943–50. [PubMed: 11052991]
- [16]. Ishikawa Y, Ishida H. Aquaporin water channel in salivary glands. *Jpn. J. Pharmacol.* 2000; 83:95–101. [PubMed: 10928320]
- [17]. Ramirez MT, Zhao XL, Schulman H, Brown JH. The nuclear deltaB isoform of Ca2+/calmodulin-dependent protein kinase II regulates atrial natriuretic factor gene expression in ventricular myocytes. *J. Biol. Chem.* 1997; 272:31203–31208. [PubMed: 9388275]
- [18]. Jensen BC, O'Connell TD, Simpson PC. Alpha-1-adrenergic receptors: targets for agonist drugs to treat heart failure. *J. Mol. Cell. Cardiol.* 2011; 51:518–528. [PubMed: 21118696]
- [19]. deBlois D, Schwartz SM, van Kleef EM, Su JE, Griffin KA, Bidani AK, Daemen MJ, Lombardi DM. Chronic alpha 1-adrenoreceptor stimulation increases DNA synthesis in rat arterial wall. Modulation of responsiveness after vascular injury. *Arterioscler. Thromb. Vasc. Biol.* 1996; 16:1122–1129. [PubMed: 8792765]
- [20]. Polichnowski AJ, Griffin KA, Long J, Williamson GA, Bidani AK. Blood pressure-renal blood flow relationships in conscious angiotensin II- and phenylephrine-infused rats. *Am. J. Physiol. Renal Physiol.* 2013; 305:F1074–84. [PubMed: 23825067]
- [21]. Sancho-Bru P, Bataller R, Gasull X, Colmenero J, Khurdayan V, Gual A, Nicolas JM, Arroyo V, Gines P. Genomic and functional characterization of stellate cells isolated from human cirrhotic livers. *J. Hepatol.* 2005; 43:272–282. [PubMed: 15964095]
- [22]. Kuriyama M, Taniguchi T, Shirai Y, Sasaki A, Yoshimura A, Saito N. Activation and translocation of PKCdelta is necessary for VEGF-induced ERK activation through KDR in HEK293T cells. *Biochem. Biophys. Res. Commun.* 2004; 325:843–851. [PubMed: 15541367]
- [23]. O-Uchi J, Jhun BS, Xu S, Hurst S, Raffaello A, Liu X, Yi B, Zhang H, Gross P, Mishra J, Ainbinder A, Kettlewell S, Smith GL, Dirksen RT, Wang W, Rizzuto R, Sheu SS. Adrenergic signaling regulates mitochondrial Ca2+ uptake through Pyk2-dependent tyrosine phosphorylation of the mitochondrial Ca2+ uniporter. *Antioxid. Redox Signal.* 2014; 21:863–879. [PubMed: 24800979]
- [24]. O-Uchi J, Komukai K, Kusakari Y, Morimoto S, Kawai M, Jhun BS, Hurst S, Hongo K, Sheu SS, Kurihara S. Alpha-adrenoreceptor stimulation inhibits cardiac excitation-contraction coupling through tyrosine phosphorylation of beta-adrenoreceptor. *Biochem. Biophys. Res. Commun.* 2013
- [25]. Jhun BS, O-Uchi J, Wang W, Ha CH, Zhao J, Kim JY, Wong C, Dirksen RT, Lopes CM, Jin ZG. Adrenergic signaling controls RGK-dependent trafficking of cardiac voltage-gated L-type Ca2+ channels through PKD1. *Circ. Res.* 2012; 110:59–70. [PubMed: 22076634]
- [26]. Williams DM, Lopes CM, Rosenhouse-Dantsker A, Connelly HL, Matavel A, O-Uchi J, McBeath E, Gray DA. Molecular basis of decreased Kir4.1 function in SeSAME/EAST syndrome. *J. Am. Soc. Nephrol.* 2010; 21:2117–2129. [PubMed: 21088294]
- [27]. Gentile S. Ion channel phosphorylopathy: a link between genomic variation and human disease. *ChemMedChem.* 2012; 7:1757–1761. [PubMed: 22927196]
- [28]. Cone AC, Cavin G, Ambrosi C, Hakozaiki H, Wu-Zhang AX, Kunkel MT, Newton AC, Sosinsky GE. Protein Kinase Cdelta-mediated Phosphorylation of Connexin43 Gap Junction Channels

- Causes Movement within Gap Junctions followed by Vesicle Internalization and Protein Degradation. *J. Biol. Chem.* 2014; 289:8781–8798. [PubMed: 24500718]
- [29]. Doble BW, Ping P, Fandrich RR, Cattini PA, Kardami E. Protein kinase C-epsilon mediates phorbol ester-induced phosphorylation of connexin-43. *Cell. Commun. Adhes.* 2001; 8:253–256. [PubMed: 12064598]
- [30]. Miura T, Yano T, Naitoh K, Nishihara M, Miki T, Tanno M, Shimamoto K. Delta-opioid receptor activation before ischemia reduces gap junction permeability in ischemic myocardium by PKC-epsilon-mediated phosphorylation of connexin 43. *Am. J. Physiol. Heart Circ. Physiol.* 2007; 293:H1425–31. [PubMed: 17513490]
- [31]. Gallegos LL, Kunkel MT, Newton AC. Targeting protein kinase C activity reporter to discrete intracellular regions reveals spatiotemporal differences in agonist-dependent signaling. *J. Biol. Chem.* 2006; 281:30947–30956. [PubMed: 16901905]
- [32]. Wu-Zhang AX, Murphy AN, Bachman M, Newton AC. Isozyme-specific interaction of protein kinase Cdelta with mitochondria dissected using live cell fluorescence imaging. *J. Biol. Chem.* 2012; 287:37891–37906. [PubMed: 22988234]

Highlights

- Isoform-specific translocation pattern of PKC was observed in live cells.
- PKC α , β I, β II, and ϵ translocates to plasma membrane by α_1 -adrenergic stimulation.
- PKC δ did not show any translocation to plasma membrane by α_1 -adrenergic stimulation.
- PKC translocation kinetics were different between PMA and α_1 -adrenergic stimulation.
- PKC α , β I, β II and ϵ may underlie physiological responses of α_1 -adrenergic signaling.




Fig.1. Ca²⁺-dependent PKC isoforms translocate to the PM in response to α_1 -ARs

A. Representative cell images (top) and fluorescence intensity profiles at white dot lines (bottom) in cells expressing α_{1A} -AR and PKC α -GFP during Phe stimulation. A.U., fluorescence arbitrary units. **B.** Time-dependent changes in the subcellular localization of PKC α -GFP by Phe in the presence or in the absence of α_1 -AR antagonist prazosin. The effect of PMA is also shown. *p < 0.05, compared to control (0 min, before Phe stimulation). #p < 0.05, compared to control (0 min, before PMA stimulation). **C.** Representative cell images (top) and fluorescence intensity profiles (bottom) in cells expressing α_{1A} -AR and PKC β I-GFP during Phe stimulation. **D.** Time-dependent changes in the subcellular localization of PKC β I-GFP by Phe in the presence or in the absence of α_1 -AR antagonist prazosin. The effect of PMA is also shown. *p < 0.05, compared to control (0 min, before Phe stimulation). #p < 0.05, compared to control (0 min, before PMA stimulation). **E.** Representative cell images (top) and fluorescence intensity profiles (bottom) in cells expressing α_{1A} -AR and PKC β II-GFP during Phe stimulation. **F.** Time-dependent changes in the subcellular localization of PKC β II-GFP by Phe in the presence or in the absence of α_1 -AR antagonist prazosin. The effect of PMA is also shown. *p < 0.05, compared to control (0 min, before Phe stimulation). #p < 0.05, compared to control (0 min, before PMA stimulation). **G.** Representative cell images and fluorescence intensity profiles in α_{1A} -AR-HEK293T cells co-expressing PKC β II-GFP and DsRed2-PKC β II during Phe stimulation. **H.** Time-dependent changes in the subcellular localization of PKC β II-GFP and DsRed2-PKC β II by Phe (top) or PMA (bottom) in α_{1A} -AR-HEK293T cells. **I.** Representative cell images and fluorescence intensity profiles in α_{1A} -AR-HEK293T cells co-expressing PKC β I-GFP and DsRed2-PKC β II during Phe stimulation. **J.** Time-dependent changes in the subcellular localization of PKC β I-GFP and DsRed2-PKC β II by Phe (top) or

PMA (bottom) in α_{1A} -AR-HEK293T cells. * $p < 0.05$, compared to PKC β I-GFP at each time point.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Fig.2. Ca^{2+} -independent PKC, PKC ϵ does (but PKC δ does not) translocate to the plasma membrane in response to α_1 -ARS

A. Representative cell images (top) and fluorescence intensity profiles at white dot lines (bottom) in cells expressing α_{1A} -AR and PKC δ -GFP during Phe stimulation. PKC δ translocation by PMA treatment was also observed in same cell as a positive control (right) (see also Supplementary Fig.4). A.U., fluorescence arbitrary units. **B.** Time-dependent changes in the subcellular localization of PKC δ -GFP by the treatment of Phe or PMA. # $p < 0.05$, compared to control (0 min, before PMA stimulation). **C. Left**, representative cell images (top) and fluorescence intensity profiles at white dot lines (bottom) in α_{1A} -AR-HEK293T cells co-transfected with PKC δ -GFP and DsRed2-PKC β II before and after Phe stimulation. The results shown are representative of five cells. **Right**, representative cell images (top) and fluorescence intensity profiles at white dot lines (bottom) in α_{1A} -AR-HEK293T cells co-transfected with PKC δ -GFP and DsRed2-PKC β II before and after PMA stimulation. The results shown are representative of six cells. A.U., fluorescence arbitrary units. **D.** Representative cell images (top) and fluorescence intensity profiles at white dot lines (bottom) in cells expressing α_{1A} -AR and PKC ϵ -GFP during Phe stimulation. **E.** Time-dependent changes in the subcellular localization of PKC ϵ -GFP by 100 μM Phe in the presence or in the absence 1 μM prazosin. The effect of PMA is also shown as a positive control. * $p < 0.05$, compared to control (0 min, before Phe stimulation). # $p < 0.05$, compared to control (0 min, before PMA stimulation). **F.** Time-dependent changes in the subcellular localization of PKC ϵ -GFP and DsRed2-PKC β I by Phe (left) or PMA (right) in α_{1A} -AR-HEK293T cells. * $p < 0.05$, compared to DsRed2-PKC β I at each time point.





Fig.3. Atypical PKC, PKC ζ does not show any translocation by α_1 -ARS

Representative cell images (top) and fluorescence intensity profiles at white dot lines (bottom) in cells expressing α_{1A} -AR and PKC ζ -GFP during α_1 -ARS (100 μ M Phe). The results shown are representative of four cells. A.U., fluorescence arbitrary units.

A.

	localization at rest	translocation by Phe	translocation by PMA
α	cytosol	Yes (Plasma membrane)	Yes (Plasma membrane)
β I	cytosol	Yes (Plasma membrane)	Yes (Plasma membrane)
β II	cytosol	Yes (Plasma membrane)	Yes (Plasma membrane)
δ	cytosol (Golgi?)	No	Yes (Plasma membrane and nuclear membrane)
ϵ	cytosol	Yes (Plasma membrane)	Yes (Plasma membrane)
ζ	cytosol	No	No
GFP	cytosol and nuclear	No	No

B.**Fig.4. Isoform-specific translocation of PKC by α_1 -ARS in HEK293T cells**

A. Summary table of the Isoform-specific translocation of PKC by α_1 -ARS (Phe treatment) or PMA treatment in HEK293T cells. **B.** Summary of the time-dependent translocation of PKC isoforms to the PM by α_1 -ARS in HEK293T cells. M/C ratio in each isoform was normalized by that obtained after 10-min application of PMA (set as a 100% translocation).