

Complete Genome Sequence and Methylome of *Staphylococcus schleiferi*, an Important Cause of Skin and Ear Infections in Veterinary Medicine

Ana M. Mistic,^a Christine L. Cain,^b Daniel O. Morris,^b Shelley C. Rankin,^a Daniel P. Beiting^a

Department of Pathobiology^a and Department of Clinical Studies,^b School of Veterinary Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, USA

***Staphylococcus schleiferi*, a Gram-positive and coagulase-variable organism, is an opportunistic human pathogen and a major cause of skin and soft tissue infections in dogs. Here, we report the first *S. schleiferi* genome sequence and methylome from four canine clinical isolates.**

Received 24 July 2015 Accepted 31 July 2015 Published 10 September 2015

Citation Mistic AM, Cain CL, Morris DO, Rankin SC, Beiting DP. 2015. Complete genome sequence and methylome of *Staphylococcus schleiferi*, an important cause of skin and ear infections in veterinary medicine. *Genome Announc* 3(5):e01011-15. doi:10.1128/genomeA.01011-15.

Copyright © 2015 Mistic et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Daniel P. Beiting, beiting@vet.upenn.edu.

Staphylococcus is an important genus of Gram-positive bacteria that are both commensal and pathogenic for humans and animals. The propensity of *Staphylococcus* species to develop antibiotic resistance, combined with the potential for zoonotic transmission (1–6), highlights the importance of understanding the genetic blueprint of *Staphylococcus* species that are commonly associated with infections in humans and animals. *Staphylococcus schleiferi* is a leading cause of drug-resistant pyoderma and otitis in dogs (7–10) and a source of rare but serious infections in humans (11–14).

Pulsed-field gel electrophoresis of *S. schleiferi* isolated from clinical samples at the Matthew J. Ryan Veterinary Hospital of the University of Pennsylvania showed that the population structure is dominated by four clonal clusters (15). Clonal cluster 1 accounts for the majority of clinical isolates, and 90% of the strains from this cluster have been shown to be methicillin resistant. Clonal cluster 2 is more heterogeneous and contains both methicillin-resistant (70%) and methicillin-susceptible isolates. Representatives from clonal clusters 1, 2, and 3 were selected for whole-genome sequencing from a banked collection of *S. schleiferi* isolates obtained from canine skin and ear canal infections between 2002 and 2013 from patients of the Matthew J. Ryan Veterinary Hospital of the University of Pennsylvania.

Isolates were grown in LB broth at 37°C overnight to an optical density at 600 nm (OD₆₀₀) of approximately 1.2. Unamplified *S. schleiferi* genomic DNA was isolated using a Qiagen Genomic-tip kit (Valencia, CA, USA) and was used to construct SMRTbell sequencing libraries (Pacific Biosciences, Menlo Park, CA, USA). Sequencing was carried out using P4-C2 chemistry on a PacBio RSII sequencing platform with one 120-min. movie per cell. Sequencing each strain on a single PacBio single-molecule real-time (SMRT) cell yielded between 45,000 and 87,000 reads, with an N₅₀ read length of approximately 6.5 to 10 kb. The SMRT Analysis software suite (version 2.0.1) (Pacific Biosciences) was used for *de novo* genome assembly with the HGAP.3 module (16). The sequences for each strain assembled to form a single contig representing the bacterial chromosome, with a coverage depth ranging

from 59 to 140×, with >99.99% consensus accuracy. The genomic size range was 2,419,762 to 2,528,396 bp, with a G+C content of 35.8 to 36.0%. Isolate 5909-02 was additionally found to carry one 24-kb conjugative plasmid (p5909-02). Manual curation was used to resolve gaps and achieve a closed genome. Analysis of the *S. schleiferi* methylomes using the RS_Motif_and_Modification_Detection.1 module revealed that isolates 2142-05, 5909-02, and 2317-03 possessed two type I restriction-modification systems, while isolate 1360-13 possessed a type III restriction-modification system. The National Center for Biotechnology Information (NCBI) Prokaryotic Genome Annotation Pipeline (PGAP) was used to annotate the genomes and plasmid deposited into the NCBI Genomes database (17). A detailed analysis of the *S. schleiferi* genomes and comparisons to other *Staphylococcus* species will be included in a future publication.

Nucleotide sequence accession numbers. Genome sequences for *S. schleiferi* strains 1360-13, 2142-05, 5909-02, a plasmid from 5909-02 (p5909-02), and strain 2317-03 have been deposited in GenBank under accession numbers CP009470, CP009762, CP009676, KP670395, and CP010309. PacBio sequence data for strains 1360-13, 2142-05, 5909-02, and 2317-03 are publicly available in the Sequence Read Archive (SRA) under accession numbers SRX701735, SRX847734, SRX738682, and SRX971499. Methylome data are available in the SRA under accession numbers SRP047175, SRP052584, SRP049178, and SRP056633. The genomes are also available in the Pathogen Resource Integration Center (PATRIC) at <https://www.patricbrc.org>.

ACKNOWLEDGMENTS

We thank Kathleen O'Shea for assistance in growing strains and Roberto Lleras (Pacific Biosciences) for helpful discussion regarding genome assembly and refinement.

This work was supported by a Morris Animal Foundation postdoctoral fellowship to A.M.M. (D14CA-404).

REFERENCES

1. Bramble M, Morris D, Tolomeo P, Lautenbach E. 2011. Potential role of pet animals in household transmission of methicillin-resistant *Staphylo-*

- coccus aureus*: a narrative review. Vector Borne Zoonotic Dis 11:617–620. <http://dx.doi.org/10.1089/vbz.2010.0025>.
2. Rutland BE, Weese JS, Bolin C, Au J, Malani AN. 2009. Human-to-dog transmission of methicillin-resistant *Staphylococcus aureus*. Emerg Infect Dis 15:1328–1330. <http://dx.doi.org/10.3201/eid1508.081635>.
 3. Van Duijkeren E, Wolfhagen MJ, Box AT, Heck ME, Wannet WJ, Fluit AC. 2004. Human-to-dog transmission of methicillin-resistant *Staphylococcus aureus*. Emerg Infect Dis 10:2235–2237. <http://dx.doi.org/10.3201/eid1012.040387>.
 4. Loeffler A, Boag AK, Sung J, Lindsay JA, Guardabassi L, Dalsgaard A, Smith H, Stevens KB, Lloyd DH. 2005. Prevalence of methicillin-resistant *Staphylococcus aureus* among staff and pets in a small animal referral hospital in the UK. J Antimicrob Chemother 56:692–697. <http://dx.doi.org/10.1093/jac/dki312>.
 5. Verkade E, Kluytmans-van den Bergh M, van Benthem B, van Cleef B, van Rijen M, Bosch T, Schouls L, Kluytmans J. 2014. Transmission of methicillin-resistant *Staphylococcus aureus* CC398 from livestock veterinarians to their household members. PLoS One 9:e100823. <http://dx.doi.org/10.1371/journal.pone.0100823>.
 6. Morris DO, Rook KA, Shofer FS, Rankin SC. 2006. Screening of *Staphylococcus aureus*, *Staphylococcus intermedius*, and *Staphylococcus schleiferi* isolates obtained from small companion animals for antimicrobial resistance: a retrospective review of 749 isolates (2003–04). Vet Dermatol 17:332–337. <http://dx.doi.org/10.1111/j.1365-3164.2006.00536.x>.
 7. Griffeth GC, Morris DO, Abraham JL, Shofer FS, Rankin SC. 2008. Screening for skin carriage of methicillin-resistant coagulase-positive staphylococci and *Staphylococcus schleiferi* in dogs with healthy and inflamed skin. Vet Dermatol 19:142–149. <http://dx.doi.org/10.1111/j.1365-3164.2008.00663.x>.
 8. May ER, Kinyon JM, Noxon JO. 2012. Nasal carriage of *Staphylococcus schleiferi* from healthy dogs and dogs with otitis, pyoderma or both. Vet Microbiol 160:443–448. <http://dx.doi.org/10.1016/j.vetmic.2012.06.020>.
 9. Morris DO, Boston RC, O'Shea K, Rankin SC. 2010. The prevalence of carriage of methicillin-resistant staphylococci by veterinary dermatology practice staff and their respective pets. Vet Dermatol 21:400–407. <http://dx.doi.org/10.1111/j.1365-3164.2010.00866.x>.
 10. May ER, Hnilica KA, Frank LA, Jones RD, Bemis DA. 2005. Isolation of *Staphylococcus schleiferi* from healthy dogs and dogs with otitis, pyoderma, or both. J Am Vet Med Assoc 227:928–931. <http://dx.doi.org/10.2460/javma.2005.227.928>.
 11. Hernández JL, Calvo J, Sota R, Agüero J, García-Palomo JD, Fariñas MC. 2001. Clinical and microbiological characteristics of 28 patients with *Staphylococcus schleiferi* infection. Eur J Clin Microbiol Infect Dis 20: 153–158. <http://dx.doi.org/10.1007/PL00011251>.
 12. Kumar D, Cawley JJ, Irizarry-Alvarado JM, Alvarez A, Alvarez S. 2007. Case of *Staphylococcus schleiferi* subspecies *coagulans* endocarditis and metastatic infection in an immune compromised host. Transpl Infect Dis 9:336–338. <http://dx.doi.org/10.1111/j.1399-3062.2007.00222.x>.
 13. Jindal A, Shivpuri D, Sood S. 2015. *Staphylococcus schleiferi* meningitis in a child. Pediatr Infect Dis J 34:329. <http://dx.doi.org/10.1097/INF.0000000000000561>.
 14. Thawabi M, Jerome M, Slim J, Shamooun F, Boghossian J. 2015. Pericardial effusion and sepsis caused by *Staphylococcus schleiferi*. J Infect Public Health 8:392–393. <http://dx.doi.org/10.1016/j.jiph.2015.01.006>.
 15. Cain CL, Morris DO, O'Shea K, Rankin SC. 2011. Genotypic relatedness and phenotypic characterization of *Staphylococcus schleiferi* subspecies in clinical samples from dogs. Am J Vet Res 72:96–102. <http://dx.doi.org/10.2460/ajvr.72.1.96>.
 16. Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. Nat Methods 10:563–569. <http://dx.doi.org/10.1038/nmeth.2474>.
 17. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Ciufu S, Li W. 2013. Prokaryotic genome annotation pipeline. The NCBI handbook, 2nd ed. National Center for Biotechnology Information, Bethesda, MD.