

Genetics of Rapid and Extreme Size Evolution in Island Mice

Melissa M. Gray,* Michelle D. Parmenter,* Caley A. Hogan,* Irene Ford,* Richard J. Cuthbert,[†]
Peter G. Ryan,[‡] Karl W. Broman,[§] and Bret A. Payseur*¹

*Laboratory of Genetics and [§]Department of Biostatistics and Medical Informatics, University of Wisconsin, Madison, Wisconsin 53706, [†]Royal Society for the Protection of Birds, The Lodge, Sandy, Bedfordshire, SG19 2DL, United Kingdom, and [‡]Percy FitzPatrick Institute of African Ornithology, Department of Science and Technology/National Research Foundation Centre of Excellence, University of Cape Town, Rondebosch 7701, South Africa

ABSTRACT Organisms on islands provide a revealing window into the process of adaptation. Populations that colonize islands often evolve substantial differences in body size from their mainland relatives. Although the ecological drivers of this phenomenon have received considerable attention, its genetic basis remains poorly understood. We use house mice (subspecies: *Mus musculus domesticus*) from remote Gough Island to provide a genetic portrait of rapid and extreme size evolution. In just a few hundred generations, Gough Island mice evolved the largest body size among wild house mice from around the world. Through comparisons with a smaller-bodied wild-derived strain from the same subspecies (WSB/EIJ), we demonstrate that Gough Island mice achieve their exceptional body weight primarily by growing faster during the 6 weeks after birth. We use genetic mapping in large F_2 intercrosses between Gough Island mice and WSB/EIJ to identify 19 quantitative trait loci (QTL) responsible for the evolution of 16-week weight trajectories: 8 QTL for body weight and 11 QTL for growth rate. QTL exhibit modest effects that are mostly additive. We conclude that body size evolution on islands can be genetically complex, even when substantial size changes occur rapidly. In comparisons to published studies of laboratory strains of mice that were artificially selected for divergent body sizes, we discover that the overall genetic profile of size evolution in nature and in the laboratory is similar, but many contributing loci are distinct. Our results underscore the power of genetically characterizing the entire growth trajectory in wild populations and lay the foundation necessary for identifying the mutations responsible for extreme body size evolution in nature.

KEYWORDS island syndrome; island evolution; phenotypic extreme; body size; complex trait

THE question of how organisms adapt to new environments continues to captivate biologists. Because adaptation requires genetic change, discovering the mutations responsible for adaptive phenotypes is a key step toward understanding the mechanisms of this process. There is a growing list of traits for which adaptive differences in nature have been directly traced to specific genes. Examples include adaptive coloration in pocket mice (Nachman *et al.* 2003), deer mice (Hoekstra *et al.* 2006), *Drosophila melanogaster* (Rebeiz *et al.* 2009), and peppered moths (van't Hof *et al.* 2011); armor plate patterning and pelvic spine reduction in stickleback fish

(Colosimo *et al.* 2005; Chan *et al.* 2010; Jones *et al.* 2012); and defense chemistry in *Boechea stricta* (Prasad *et al.* 2012). Despite these advances, the genetic architecture of adaptation in nature remains poorly understood. Most progress has focused on traits with simple genetic bases, where one or a few loci explain observed phenotypic variation (Rockman 2012). But the majority of trait differences between populations inhabiting contrasting environments are quantitative, suggesting that adaptation often involves more complex inheritance.

While it is notoriously difficult to demonstrate adaptation (Endler 1986), populations that rapidly evolve to phenotypic extremes following major environmental shifts provide a revealing window into the process. Evidence for adaptive change is often observed in island populations, where the limited geographic scope, sharp boundaries, and simplified biotas of islands facilitate the interpretation of evolutionary patterns (Losos and Ricklefs 2009). Insular mammals show elevated rates of morphological evolution (Pergams and Ashley 2001;

Copyright © 2015 by the Genetics Society of America

doi: 10.1534/genetics.115.177790

Manuscript received May 1, 2015; accepted for publication July 18, 2015; published Early Online July 20, 2015.

Supporting information is available online at www.genetics.org/lookup/suppl/doi:10.1534/genetics.115.177790/-/DC1.

¹Corresponding author: Laboratory of Genetics, 425-G Henry Mall, Madison, WI 53706. E-mail: payseur@wisc.edu

Millien 2006) and include several examples of gigantism and dwarfism (Stock 1935; Freudenthal 1972; Roth 1992; Moncunill-Solé *et al.* 2014). Populations that colonize islands often experience substantial changes in predation risk, competition, and resource availability that together generate strong selection for shifts in body size (Sondaar 1977; Case 1978; Heaney 1978; Lawlor 1982; Lomolino 1985; Lomolino *et al.* 2012). Although the question of whether island mammals in general follow directional patterns in the evolution of body size (Foster 1964; Van Valen 1973; Lomolino 1985) has inspired debate (Lawlor 1982; Lomolino 1985, 2005; Meiri *et al.* 2004, 2008, 2011; Lomolino *et al.* 2005, 2012; Raia and Meiri 2006; Bromham and Cardillo 2007), murid rodents usually evolve larger sizes on islands (Adler and Levins 1994; Meiri *et al.* 2008). This pattern, combined with the remarkable success of house mice (*Mus musculus*) in colonizing islands from around the world, positions these rodents as an especially promising system for understanding adaptive size change. Considerable morphological diversity within and among island populations of house mice has been documented, especially in the research of R. J. Berry (Berry 1964, 1965, 1968, 1996; Berry and Jakobson 1975; Berry *et al.* 1987, 1978a,b, 1979, 1982; Berry and Scriven 2005).

The largest wild house mice in the world reside on Gough Island (GI), a remote volcanic island in the middle of the South Atlantic Ocean (Rowe-Rowe and Crafford 1992). Gough Island mice weigh approximately twice as much as their mainland relatives (Jones *et al.* 2003). Mice were introduced to Gough Island by teams of seal hunters (Wace 1961; Jones *et al.* 2003) between 130 and 200 years ago (Verrill 1895; Gray *et al.* 2014), suggesting that body size evolution has been very rapid.

Gough Island mice belong to the same subspecies as the laboratory mouse (*Mus musculus domesticus*) (Gray *et al.* 2014), providing access to an expansive genetic toolkit for investigating their phenotypic evolution. In this article, we focus on the rapid evolution of body size, for two reasons. First, body size is highly correlated with aspects of physiology, life history, morphology, behavior, and ecology (Peters 1983; Calder 1984; Schmidt-Nielsen 1984). As a result, size is perhaps the best singular phenotypic measurement of an organism's overall adaptive profile (Damuth and MacFadden 1990). Second, body size is a canonical complex trait from a genetic perspective. Induced mutations in a large fraction of genes noticeably alter size (Reed *et al.* 2008), many quantitative trait loci (QTL) contribute to existing size differences among laboratory strains (Corva and Medrano 2001), and a variety of environmental factors influence growth. By genetically dissecting body size in Gough Island mice, we can learn how an island population in a new environment rapidly and recently achieved a marked evolutionary shift in a complex trait.

The major characteristics and determinants of body size growth in mice are understood. The growth trajectory is sigmoidal, with an increasing rate during the first few weeks after birth, peak growth between 3 and 5 weeks, and a decreasing rate thereafter (Cheverud 2005). Growth *in utero* and soon after birth is primarily promoted by insulin-like growth factor 2

(IGF2) (Fowden 2003), whereas the growth hormone (GH)-IGF1 axis drives growth after weaning (Lupu *et al.* 2001). Early growth mostly reflects increases in cell number, while later growth is mostly due to expansions in cell size (Riska and Atchley 1985; Atchley *et al.* 2000; Lui and Baron 2011). Genetic correlations between age-specific body weights decline with increasing time between ages (Cheverud *et al.* 1983a; Riska *et al.* 1984). Many QTL that contribute to body weight differences between laboratory mouse strains have been identified (Dragani *et al.* 1995; Cheverud *et al.* 1996; Keightley *et al.* 1996; Rance *et al.* 1997; Brockmann *et al.* 1998, 2004; Kirkpatrick *et al.* 1998; Morris *et al.* 1999; Vaughn *et al.* 1999; Corva and Medrano 2001; Rocha *et al.* 2004; Bennett *et al.* 2005; Kenney-Hunt *et al.* 2006; Shao *et al.* 2007; Casellas *et al.* 2009; Mollah and Ishikawa 2011; Ishikawa and Okuno 2014). A common pattern is that strain differences in weight and growth rate are due to a large number of loci with modest phenotypic effects. Consistent with the physiology of mouse growth, QTL effects vary among ages, an observation that motivates the genetic dissection of size evolution throughout the entire ontogenetic trajectory.

Whether similar genetic patterns characterize exceptional size evolution in nature remains an open empirical question. Two key contrasts prevent extrapolation of findings in laboratory mice to evolution in the wild. First, the dynamics of selection differ. Artificial selection for divergent body sizes in laboratory mice has generated impressive responses (Goodale 1938; MacArthur 1944; Falconer 1953; Eisen 1989; Bünger and Hill 1999), providing useful systems for mapping QTL. But wild mice encounter selective agents that are completely absent from the laboratory environment, and it is unclear whether artificial and natural selection are similar in intensity. Different selective pressures and intensities could lead to different genetic architectures. Second, laboratory mice have a complex history, with uncertain connections to wild mice (Silver 1995). Many genetic variants observed in natural populations are missing from the panel of commonly used inbred strains (Salcedo *et al.* 2007), suggesting that evolution in nature and in the laboratory could involve different sets of mutations. Here, we use mice from Gough Island to provide a genetic portrait of rapid and extreme size evolution in a wild, island population.

Materials and Methods

Gough Island mice

Gough Island belongs to the United Kingdom Overseas Territory of Tristan da Cunha, has an area of 65 km², and is situated in the South Atlantic Ocean almost halfway between South Africa and South America (40° 19'S and 9° 55'W). Fifty mice were live trapped on Gough Island near the research station in September of 2009. Four mice died and two litters were born (five pups) during transport from Gough Island to the Charman Instructional Facility at the University of Wisconsin-Madison. Twenty-five mature female mice and 21 mature male mice were used to start a breeding colony.

All mice were housed in microisolator cages and separated by sex. Males were grouped only if they were from the same litter and had not previously mated. The room was temperature controlled (68–72° F) and set on a 12 hr light/dark cycle. Corn cobs ground to 1/8th inch were used for bedding (Waldschmidt and Sons, Madison, WI). Water and Teklad 6% fat mouse/rat diet was provided *ad libitum* (Harlan Laboratories, Madison, WI). Irradiated sunflower seeds (Harlan Laboratories) and nesting material were provided as enrichment and were replaced weekly during cage changes.

Selected individuals were mated after 8 weeks of age. Breeders were supplemented with extra bedding and red huts and were fed breeder chow (Teklan Global 19% protein/9% fat; Harlan Laboratories) *ad libitum*. Mated pairs were only disturbed during weekly cage changes. Weekly weights to the nearest milligram were recorded, beginning 1 week after birth and ending at 16 weeks. To individually identify pups starting at 1 week, toe tattooing was performed using sterile lancets and tattoo paste. Ear punches were given at weaning (3–4 weeks) to individually identify adult mice.

Once the mice arrived in Madison, we first performed a random breeding common garden experiment with the wild founders. The purposes of this experiment were to establish a colony and to verify the large weight of Gough Island (GI) mice was inherited and not solely the result of environmental factors.

Intercross experiments

Beginning with the laboratory-born offspring of the wild mice, we created several partially inbred lines by brother–sister mating for four generations. While inbreeding was far from complete, four generations of brother–sister mating is expected to increase homozygosity to 60% within lines. We chose one pair of male and female littermates (full siblings) from each of two partially inbred lines (denoted line/cross A and B). These four mice were each crossed with WSB/EiJ (The Jackson Laboratory, Bar Harbor, ME; subsequently abbreviated as WSB) to generate four independent F₂ intercrosses (Figure 1). Although we focus on QTL that are fixed between the panel of four GI mice and WSB in this study, our design will allow us to detect loci that are polymorphic within the GI population in the future. WSB was chosen because it is a wild-derived strain, has a body size typical of wild house mice, belongs to the same subspecies as GI mice, is fully inbred, has a sequenced genome (Keane *et al.* 2011), and is featured in large-scale efforts to genetically dissect complex traits such as the Collaborative Cross (Churchill *et al.* 2004). We generated a total of 1374 F₂ mice: 497 from cross A (WSB × GI = 279 and GI × WSB = 218; maternal strain denoted first) and 877 from cross B (WSB × GI = 494 and GI × WSB = 383).

Phenotyping

Weekly weights were collected for all mice from 1 week through 16 weeks of age, unless they were paired for breeding. Once mice reached 16 weeks (±1 day), they were killed by CO₂ asphyxiation or by decapitation. Liver samples were collected and stored at –80° C until genetic analysis. This study was




Figure 1 Design of crosses for genetic mapping. Wild, first generation of Gough Island mice born in the laboratory; FS, filial self-generation; W, WSB.

approved by the Institutional Animal Care and Use Committee at the University of Wisconsin (protocol no. V01447).

Genotyping

All mice were genotyped on the Mega Mouse Universal Genotyping Array (MegaMUGA; <http://www.neogen.com/Agrigenomics/pdf/Slicks/MegaMUGAFlyer.pdf>). The MegaMUGA is an Illumina Infinium array that contains ~77,800 markers. Markers are spaced at ~33 kb across the genome and are mostly single nucleotide polymorphisms (SNPs) with a few structural variants and transgenic markers. The markers cover all autosomes, sex chromosomes, and the mitochondria. Marker ascertainment was carried out to be maximally informative for studies involving the Collaborative Cross (Aylor *et al.* 2011; Threadgill and Churchill 2012), Diversity Outbred Cross (Svenson *et al.* 2012), wild populations of house mice, and other *Mus* species. Liver tissue was sent to GeneSeek (NeoGene, Lincoln, NE) for DNA extraction and genotyping with the MegaMUGA arrays. A total of 1536 samples were sent, including controls and mice that died before 16 weeks of age (which are still informative for genetic map estimation).

Multiple controls were used during extraction and genotyping to increase our ability to identify technical and biological





Figure 2 Comparison of body weight in Gough Island mice to other mice from the wild, from wild-derived inbred strains, and from classic inbred strains. Weights were compiled from sources listed in Table S1.



errors. Liver tissue was organized into 16 (96 deep well) plates in such a way as to minimize array batch effects on related sets of samples. Tissue from WSB was placed in identical wells on every plate to account for plate extraction effects. The four GI parental samples were replicated four times each across the 16 plates. Lastly, replicate samples of the first well of each plate were placed in a more distant well and run on different arrays.

Genotype diagnostics

We examined the genotypes to identify biological, technical, and data entry errors. We omitted markers with high rates of

missing data and those that were not informative in our crosses. We removed a small number of individuals that appeared to be unresolved sample mix-ups based on large numbers of Mendelian inconsistencies or mismatched sex (inferred from array intensities of markers on the X and Y chromosomes). We also removed a few mice with high rates of missing genotype data. Following these initial screens, the cleaned data included the four GI parents of the crosses, 70 F₁ individuals, 1346 F₂ individuals, and 33,191 markers.

In all subsequent analyses, we focused on a subset of 11,833 markers that were fixed in the four GI parents and




Figure 3 Body weight (top panels) and growth rate (bottom panels) for males (left) and females (right), as a function of age in weeks, for a sample of Gough Island mice (GI, green) and WSB mice (purple) raised in the lab. Individual body weight curves were lightly smoothed using cubic splines; the growth rate curves were estimated as the first derivative of the fitted splines. Thicker curves follow the group averages.

therefore segregated as in a standard F_2 intercross between inbred lines. We estimated the intermarker genetic distances assuming a genotyping error rate of 0.2% and converted estimated recombination fractions to map distances with the Carter–Falconer map function (Carter and Falconer 1951).

Smoothed phenotypes

We omitted F_2 mice missing body weights at ≥ 4 of the 16 time points, resulting in a final set of 1212 F_2 mice with nearly complete data. To reduce measurement error in the body weights and to interpolate some remaining missing values, we used the R package *pspline* to fit a 7 degrees-of-freedom cubic spline to the data for each mouse. The degrees of freedom were chosen by visual inspection, considering the trade-off between smoothness and the fidelity of the underlying curves. We estimated growth rate as the first derivative of the fitted cubic splines.

QTL analysis

QTL analysis was performed by Haley–Knott regression (Haley and Knott 1992) on a 0.5-cM grid across the genome. We used conditional genotype probabilities (given the available marker data) calculated assuming a genotyping error rate of 0.2% and the Carter–Falconer map function. While we considered data on 11,833 markers, QTL calculations were ultimately based on a fixed grid of 2665 pseudo-markers. Smoothed body weights and estimated growth rates were analyzed separately.

Single-QTL analysis was performed at each of the 16 time points, individually, including indicators for each F_1 mother (to account for sibship differences) and sex as fixed additive covariates. The results across time points were combined by averaging the LOD scores across time points at each genomic

position to give SLOD scores (Kwak *et al.* 2014). Thresholds for statistical significance were calculated using a permutation test (Churchill and Doerge 1994), permuting the rows in the phenotype data relative to the rows in the genotype data (that is, maintaining the correlation structure within the phenotype data and within the genotype data, but breaking the association between the genotypes and phenotypes). Autosome- and X-chromosome-specific significance thresholds were derived using the approach of Broman *et al.* (2006), with 1500 permutation replicates for the autosomes and 28,200 replicates for the X chromosome.

Multiple-QTL analysis was performed using strictly additive QTL models, using the penalized-SLOD criterion of Kwak *et al.* (2014), an extension of the penalized-LOD score criterion of Broman and Speed (2002). Consider a multiple-QTL model γ (a vector specifying the locations of some set of QTL). We considered each time point individually and calculated a LOD score measuring the quality of fit of the model at time point i , $LOD_i(\gamma)$. We then obtained an overall measure of the fit of the model by taking the average, $SLOD(\gamma) = \sum_i LOD_i(\gamma)/16$, where the sum is over the 16 time points. We subsequently considered a penalized version of this statistic to balance quality of fit with model size. The penalized statistic was $pSLOD(\gamma) = SLOD(\gamma) - T|\gamma|$, where $|\gamma|$ is the number of QTL in the model γ , and T is a penalty, taken to be a quantile from the null distribution of SLOD in a single-QTL genome scan, estimated by a permutation test (here using a common threshold for the autosomes and the X chromosome). Model searching was performed by a stepwise algorithm: forward selection up to a model with 15 QTL, followed by backward elimination to the null model. The chosen model was that which maximized pSLOD, among all models visited. As with the single-QTL analysis, the multiple-QTL




Figure 4 Body weight (top panels) and growth rate (bottom panels) for males (left) and females (right), as a function of age in weeks, for the F₂ mice. In each panel there are three shaded regions; the darkest region covers the middle third of the individuals; the next-darkest region all mice. The blue curve is for the average of the F₂ mice. The green and purple curves are for the averages of Gough Island and WSB mice, respectively, as in Figure 3.

analysis included indicators for each F₁ mother and sex as fixed additive covariates.

To indicate the strength of evidence for the QTL in the context of the selected multiple-QTL model, we calculated profile SLOD curves (Kwak *et al.* 2014) based on an idea from Zeng *et al.* (2000). We varied the position of each QTL, keeping all other QTL fixed at their estimated locations, and compared the full model with that QTL in varying position to the reduced model with that QTL omitted. To define approximate confidence intervals for the QTL, we fit the inferred multiple-QTL model at each time point and calculated 1.5-LOD support intervals (using the profile LOD scores) for the location of each QTL at each time point. For each QTL, we cite the interval corresponding to the time point for which the evidence for that QTL is maximum (that is, the time point at which the profile LOD score is highest).

To assess the relationship between sample size and the number of inferred QTL, we took random subsets of the F₂ mice and applied the stepwise QTL analysis, using SLOD with the 5% significance threshold calculated for the full data. (The significance threshold is not much influenced by sample size.) We performed 37–38 replicates at sample sizes of 300, 600, 800, and 1000 F₂ mice.

Data availability

Phenotypes and genotypes from this study are available from the QTL Archive at the Jackson Laboratory, at <http://phenome.jax.org/db/q?rtn=projects/projdet&reqprojid=539>. Supplemental information includes: p-values from strain comparisons (Figure S1), histograms of F₂ weights (Figure S2), average F₂ weights and growth rates (Figure S3), p-values from comparisons among crosses and mothers (Figure S4), genetic and physical maps of

markers used in QTL analyses (Figure S5), F₂ frequency of WSB allele (Figure S6), and candidate genes in QTL regions (Table S1).

Results

Patterns of phenotypic variation

Wild-caught GI mice are exceptionally heavy compared to wild mice of the same subspecies (*M. m. domesticus*) from the mainland and other islands (Figure 2). GI mice raised in the laboratory are heavier than wild-derived inbred strains of *M. m. domesticus* (Figure 2). Lab-reared GI mice are similar in weight to classical inbred strains (Figure 2), which are bigger than most wild mice (presumably because of artificial selection during domestication). The weight difference between GI mice sampled from nature and those raised in the lab likely reflects the contrasting environments these mice experienced and the older ages of the wild-caught individuals.

GI and WSB mice raised in the same environment exhibit contrasting growth trajectories (Figure 3), demonstrating that trait divergence has a heritable component. Weights are significantly different at all ages in both sexes (Supporting Information, Figure S1; *t*-tests, maximum *P*-values across time points; total: $P < 10^{-8}$; females: $P < 10^{-5}$; males: $P < 10^{-3}$). The rate of growth differs over a narrower time window, pinpointing early growth as the primary driver of phenotypic divergence. Growth rates estimated from the first derivatives of fitted growth curves in individual mice are significantly faster in GI mice from 1–6 weeks (Figure S1; total: $P < 10^{-4}$; females: $P < 0.05$; males: $P < 0.01$). At ~4–6 weeks, the growth rate difference between GI and WSB mice begins to decline. GI and WSB mice also show divergent growth rates at 11–16 weeks (total: $P < 0.05$;




Figure 5 Pairwise correlations across ages for body weight (left) and growth rate (right). Black pixels indicate a correlation near 1; white pixels indicate a correlation near 0.

males: $P < 0.05$), though females are not significantly different at 14–16 weeks. Male GI mice are heavier than female GI mice beginning at 3 weeks (Figure S1; $P < 10^{-3}$), and grow significantly faster at 1–5 weeks ($P < 10^{-3}$). Phenotypic patterns are similar in the partially inbred GI mice (and their offspring) we used for genetic mapping.

F_1 mean weights from crosses between GI and WSB mice are closer to GI means for the first 3 weeks (Figure S2). At subsequent ages, F_1 weights are near the mid-parent value, with a bias toward WSB values (especially in males).

F_2 's from crosses between GI and WSB mice vary widely in weight at all ages (Figure S2). F_2 weight distributions are continuous and weight ranges exceed parental averages. Weight variances increase with age, whereas coefficients of variation are relatively stable across ages. Growth rate also shows substantial F_2 variances at all ages, with coefficients of variation tending to increase with age. Collectively, these observations suggest that multiple mutations contributed to the evolution of weight and growth rate differences between GI and WSB mice.

The collection of 16-week growth trajectories for individual F_2 's is shown in Figure 4. Mirroring patterns in the parental lines, F_2 males and females show large differences in body weight (Figure S3; $P < 0.015$; $< 10^{-20}$ from week 3 onward) and growth rate at all ages ($P < 10^{-4}$). Faster growth of males during weeks 1–5 (Figure 3) is primarily responsible for weight differences among the sexes. F_2 's from different F_1 mothers vary significantly in weight (Figure S4; $P < 10^{-18}$) and growth rate ($P < 0.05$, with the exception of females at 14–15 weeks) at all ages, providing evidence for maternal effects (which could be genetic or environmental). F_2 weights at all ages depend on GI parent (Figure S4; $P < 10^{-8}$), as do growth rates at most ages ($P < 10^{-5}$ in weeks 1–4; $P < 0.037$, aside from weeks 6–7 and 13–14). These results motivated us to account for effects of sex, mother, and line in all QTL analyses.

F_2 weights show strong pairwise correlations across ages, with a steady decline in autocorrelation as the difference in ages increases (Figure 5). In contrast, correlations between growth rates are only high for neighboring ages (Figure 5). For example, the growth rate at one time point is nearly

uncorrelated with the growth rate at an age 4 weeks later, whereas weights are highly correlated at these ages.

Genetic map characteristics

The genetic map constructed from F_2 genotypes (Figure S5) is 1327 cM in total length. Characteristics of the map resemble known recombination patterns in house mice (Cox *et al.* 2009), including an average of 26 crossovers per meiosis, elevated rates in subtelomeric regions, and decreased rates in subcentromeric regions. A region on chromosome 2 shows remarkable transmission ratio distortion in favor of the WSB allele (allele frequency = 0.65; peak marker located at 79.6 Mb) in F_2 's descended from one GI line but not the other (Figure S6). Biased transmission of the WSB allele in this interval is also observed in the Collaborative Cross (Aylor *et al.* 2011) and seems to be generated by maternal meiotic drive (Didion *et al.* 2015).

QTL number and location

By combining information across ages, our multiple QTL mapping approach allowed us to identify sets of QTL that affect divergence in overall growth trajectories. We focus on results from this method. Major patterns are similar to those obtained by single QTL mapping at individual ages (results not shown).

Eight QTL contribute to weight differences between GI and WSB mice when a 5% genome-wide significance threshold is used (Figure 6; Table 1). For growth rate, 11 QTL are detected (Figure 6; Table 1). Five weight QTL overlap with growth rate QTL on chromosomes 1, 6, 7, and 10 (two QTL) (Table 1).

To gauge our ability to accurately estimate the number of QTL, we randomly subsampled F_2 's, reran QTL analyses, and counted the number of QTL in the best fitting model (at the 5% significance threshold). Two key patterns emerge (Figure 7). First, the number of detected QTL steadily increases with sample size for both weight and growth rate, suggesting that a large number of loci with small effects contribute to trait divergence between GI and WSB. Second, there is considerable variance in the number of QTL detected with different F_2 samples of the same size. Collectively, these results underscore

the difficulty of estimating QTL number, even with large sample sizes such as ours.

QTL effects

We measured QTL effects using two parameters: the additive effect of substituting one allele (calculated as half the difference in genotypic means between alternative homozygotes), and the dominance effect (calculated as the difference between the genotypic mean of the heterozygote and the average genotypic mean of alternative homozygotes).

Additive and dominance effects for all QTL are presented in Table 2 and Table 3. Individual QTL effects are small-to-moderate in size relative to the difference in mean phenotypes between GI and WSB mice. The largest additive effect across all weight QTL and ages is 0.66 g (chromosome 8 QTL at 16 weeks); homozygotes carrying the GI vs. WSB allele at this QTL differ by an average of 1.32 g. Additive effects of weight QTL tend to increase with age, whereas age-related trends in additive effects of growth rate QTL are less apparent. Looking across development, chromosomes 10, 7, and 8 harbor QTL with the largest effects for both weight and growth rate.

QTL vary in the timing of their largest effects. For example, QTL on chromosomes 10 and 11 affect weight beginning at week 1, whereas the chromosome 9 weight QTL has little effect until week 6. Once QTL begin to affect weight, they continue to do so at subsequent ages. In contrast, the biggest effects of growth rate QTL tend to be concentrated early in ontogeny. The growth rate QTL found distally on chromosome 10 exhibits its strongest effects during the first 4 weeks, and the QTL on chromosome 7 principally affects growth at 3–5 weeks of age. These patterns underscore the power of dissecting the genetics of size throughout the entire growth trajectory.

Most dominance effects of detected QTL are small. The two exceptions are weight QTL on chromosomes 8 and 9. The weight QTL on chromosome 9 shows the largest dominance deviations, which exceed its additive effects at all ages. Heterozygotes at this locus are shifted toward the GI mean. QTL with the biggest phenotypic effects primarily act in an additive fashion.

Collectively, the eight weight QTL explain between 6% and 24% of the F_2 variance in weight across ages (Figure 8). Overall heritability due to weight QTL starts low (at 1 week), increases to its maximum at 7 weeks, and decreases slightly through 16 weeks. The 11 growth rate QTL jointly explain between 3% and 20% of the F_2 variance (Figure 8). In contrast to weight QTL, the overall heritability due to growth rate QTL peaks at 3 weeks, quickly drops during weeks 4 and 5, and remains low through 16 weeks. This pattern mirrors the temporal divergence in growth rate between GI and WSB mice (Figure 3).

We used an additional approach to measure the ability of detected QTL to explain body size evolution. Phenotypic means of GI and WSB mice represent their genetic values. In a purely additive model, the difference between phenotypic

means of these parents at a particular age should equal twice the sum of the additive effects of all mutations that contribute to weight or growth rate differences. Figure 9 compares the summed additive effects of detected QTL to phenotypic differences between GI and WSB. These results raise the possibility that detected QTL collectively explain a substantial proportion of the genetic difference in size between GI and WSB mice.

Further characterization of QTL

To better understand the biological context of QTL, we conducted targeted interaction tests involving sex, mother, line, and other loci. Though sex and mother are important contributors to F_2 weight and growth rate, there is no evidence that QTL action depends on these factors. These results indicate that the QTL we detected are likely shared by males and females, and across F_1 mothers. Most QTL also show no sign of being affected by line. One QTL—on chromosome 7—displays an interaction with line for body weight (but not growth rate). Because the two lines of GI mice that contributed to our intercross were founded by different wild parents, this result suggests that the chromosome 7 QTL may be polymorphic in the island population. Pairwise interaction tests revealed no evidence for epistasis between




Figure 6 Profile SLOD curves for the selected multiple-QTL models for body weight and growth rate. The location of each QTL was varied, one at a time, with all other QTL fixed at their estimated locations, and the multiple-QTL model was compared to the model with the given QTL omitted.

Table 1 Genomic positions of QTL

Phenotype	Chromosome	cM	cM lower CI	cM upper CI	Mb	Mb lower CI	Mb upper CI
Weight	1	37.5	22.0	53.5	78.9	42.4	136.9
Weight	6	4.0	0.0	6.0	17.9	3.4	23.5
Weight	7	44.0	35.5	45.0	118.9	86.1	121.9
Weight	8	29.0	25.5	39.0	81.5	69.1	106.1
Weight	9	17.0	15.0	40.5	47.3	42.7	97.5
Weight	10	27.0	23.0	31.0	68.1	58.6	77.4
Weight	10	66.5	55.0	67.0	128.1	119.0	129.6
Weight	11	27.0	13.5	42.5	51.5	31.5	81.5
Growth rate	1	39.5	33.5	41.5	84.3	73.0	92.9
Growth rate	4	42.5	40.5	65.5	104.6	103.1	141.6
Growth rate	6	4.0	3.5	8.5	17.9	17.0	27.3
Growth rate	7	32.0	13.5	36.0	80.1	38.6	86.9
Growth rate	7	44.0	47.5	55.5	118.9	125.4	137.0
Growth rate	8	17.5	11.0	34.0	45.9	32.4	92.7
Growth rate	10	26.0	4.0	34.5	66.3	16.9	88.7
Growth rate	10	62.0	59.5	67.0	125.3	123.3	129.6
Growth rate	13	52.0	47.0	54.5	116.5	111.2	119.3
Growth rate	15	9.0	3.5	12.0	36.4	19.6	42.2
Growth rate	16	10.0	0.0	13.0	26.4	3.5	29.8

Confidence interval is the 1.5-LOD interval for the age with the highest LOD score.

detected QTL. Comprehensive searches for epistasis involving other loci are difficult to conduct and interpret because we are considering the entire growth trajectory. Consequently, we focus on the additive and dominance effects of individual QTL in this article. To identify candidate genes, we looked for genes in the centers of QTL intervals (including 2 Mb on either side of the peak) that are known to affect body size or related processes. We used Co-Citer software (Qiao *et al.* 2013) to count references that mention each gene and “body weight,” “body mass,” or “growth.” Candidate genes are listed in Table S1.

QTL evolution

Additive effects indicate that GI alleles increase weight at most QTL and at most ages. To evaluate whether this pattern is consistent with neutral evolution of body size, we asked whether the number of QTL at which the GI allele increases size is consistent with chance expectations, conditioning on the observed weight difference between GI and WSB mice (Orr 1998). Considering the maximum effect for each weight QTL across ages reveals suggestive evidence for a contribution of natural selection (Orr’s test; $P = 0.07$), with GI alleles increasing weight at all eight QTL. Again using the maximum effect as a summary, GI alleles increase growth rate at 10 of 11 QTL ($P = 0.15$; 11 of 11 QTL would have yielded an estimated $P = 0.02$).

Discussion

The ability of house mice to invade islands from around the world and their status as a genetic model organism combine to make them an unusually powerful system for addressing the question of how natural populations evolve extreme phenotypes. Our results provide a rare genetic portrait of this process in a wild population that has achieved a dramatic increase in body size during just a few hundred generations.

The suite of QTL we identified and the large magnitude of our cross allow us to confidently conclude that GI mice did not evolve their unusual body size via a few genetic changes of major effect, perhaps contrary to intuition. Instead, multiple mutations with a range of modest, mostly additive effects were responsible. The contrast between this result and the simple genetic architecture underlying fitness-related phenotypic changes in other rodents and in stickleback fish (Nachman *et al.* 2003; Colosimo *et al.* 2005; Hoekstra *et al.* 2006) may primarily reflect differences in the focal traits rather than differences between species. One potential explanation for the genetic complexity of body size evolution is that weight and growth rate are both composite phenotypes affected by a diverse array of biological processes. Decomposing these traits into a series of intermediate phenotypes could reveal simpler genetic mechanisms.

From an evolutionary perspective, the distribution of QTL effects deserves further attention. The modest additive effects of individual QTL and the low percentage of F_2 phenotypic variance jointly explained by all QTL suggest limited ability to predict body size from detected loci. Our subsampling experiment indicates that QTL with smaller effects were missed, and undetected epistasis also may have contributed to F_2 variance. At the same time, there are reasons to be optimistic that detected QTL capture much of the genetic basis of body size evolution in this case. In addition to being affected by genetic variation, F_2 phenotypes reflect uncontrolled environmental and developmental variance. For example, mice that ingested different amounts of food would be expected to grow to different sizes. The heritability of body size is the maximum fraction of phenotypic variance that QTL could theoretically explain. It is therefore interesting to note that the combined heritability for the group of body weight QTL we detected roughly matches overall heritability estimates for weight in laboratory mouse populations (range of point




Figure 7 Number of inferred QTL as a function of sample size, for random subsets of the data, and for the full dataset, for body weight (top) and growth rate (bottom).

estimates = 0.22–0.43; Lynch 1992), although heritability is a population-specific quantity. In contrast to individual F_2 phenotypes, average values for the partially inbred GI mice and the fully inbred WSB reflect *genetic* values, with random environmental deviations contributing to the variance about these averages. In this context, it is revealing that the summed additive effects of detected QTL account for most of the phenotypic differences between GI mice and WSB. An important caveat to this point is that QTL mapping generates upward biases in additive effect estimates (Beavis 1994, 1998). Nevertheless, the suite of QTL we discovered may explain much of the genetic component of body size evolution, despite its complexity.

When viewed in combination with previous findings, our results provide an unusual opportunity to compare the genetics of size evolution in nature to that in the laboratory. Here, we focus on a study that is especially comparable in terms of experimental design. Cheverud *et al.* (1996) identified QTL for age-specific (weekly) weights and growth rates up to 10 weeks in an F_2 intercross between the large (LG) and small (SM) strains. These strains were developed

by (i) crossing a variety of inbred lines, (ii) artificially selecting for increased (Goodale 1938) or decreased (MacArthur 1944) body weight at 60 days of age, and (iii) inbreeding for >125 generations (Chai 1956a,b; Festing 1996). Selection was very effective, producing a 20- to 24-g weight difference between LG and SM at 10 weeks of age (Chai 1956b; Kramer *et al.* 1998).

The divergence in growth trajectories between GI mice and WSB resembles that between LG and SM. In GI mice and LG, accelerated growth during the first 6 weeks is the primary driver of increased weight gain. The magnitude of the acceleration is greater in males. In F_2 's, correlations between weekly weights decline as the time between ages increases. Early and late growth rates are weakly correlated.

The overall genetic architectures of LG–SM and GI–WSB size differences are also similar. The number of QTL affecting LG–SM weight differences ranges from 7 (1 week) to 17 (10 weeks), whereas 8 QTL control GI–WSB weight differences at some time during ontogeny. Between 11 (early) and 12 (late) QTL contribute to growth rate differences between LG and SM; 11 QTL are responsible for growth rate variation among GI and WSB. In both cases, summed genotypic effects of QTL reveal mostly additive action, with some dominance toward the larger strain at early ages. The average additive effect of a weight QTL ranges from 0.05 g (1 week) to 0.60 g (10 weeks) for LG–SM and from 0.03 g (1 week) to 0.51 g (16 weeks) for GI–WSB. QTL effects vary throughout ontogeny, with different QTL shaping early and later growth rates. The allele from the larger strain increases body weight at most QTL in both crosses. The overall genetic response to selection for increased body size—whether selection is natural or artificial—has a common profile.

Why might rapid size evolution show similar properties in nature and in the laboratory? The agents of selection are fundamentally distinct. It also seems likely that artificial selection targeting body size is usually much stronger than natural selection. Perhaps the common profile instead reflects shared genetic, developmental, and evolutionary constraints on growth. Laboratory studies of random-bred mice revealed higher heritabilities for growth rate between 2 and 5 weeks of age than during other intervals, as well as reduced genetic correlations between early and late growth (Cheverud *et al.* 1983a,b; Riska *et al.* 1984). These observations predict that responses to selection for increased size will often accelerate early growth. Furthermore, the pervasive correlations between body size and other traits suggest that mutations affecting size will routinely have deleterious pleiotropic consequences. This logic could explain why mutations with small effects contribute most of the response to selection in nature and in the laboratory.

Although these comparisons reiterate the value of artificial selection experiments, understanding the genetics of size evolution in nature will continue to require the examination of wild populations. An important demonstration of this fact comes from the comparison of QTL location. Only 7 of the 19 QTL intervals we discovered contain estimated QTL

Table 2 Additive and dominance effects (in grams) of weight QTL

Week		1@37.5	6@4.0	7@44.0	8@29.0	9@17.0	10@27.0	10@66.5	11@27.0
1	a	-0.008	-0.045	-0.057	0.024	-0.033	0.177	0.043	0.127
	d	0.054	0.054	0.009	0.009	0.106	0.043	0.029	0.028
2	a	0.045	-0.004	0.047	0.090	-0.057	0.200	0.171	0.159
	d	0.054	0.076	0.061	-0.035	0.157	0.050	0.043	0.007
3	a	0.117	0.050	0.197	0.148	-0.044	0.226	0.337	0.182
	d	0.074	0.078	0.118	-0.109	0.210	0.061	0.030	-0.030
4	a	0.214	0.146	0.381	0.206	-0.016	0.281	0.453	0.221
	d	0.074	0.040	0.148	-0.137	0.239	0.056	0.014	-0.037
5	a	0.291	0.237	0.519	0.241	0.052	0.351	0.503	0.266
	d	0.063	0.001	0.143	-0.149	0.234	0.046	0.001	-0.046
6	a	0.321	0.298	0.561	0.284	0.120	0.406	0.507	0.307
	d	0.047	-0.013	0.095	-0.173	0.226	0.049	-0.001	-0.068
7	a	0.334	0.334	0.554	0.337	0.169	0.445	0.510	0.342
	d	0.035	-0.008	0.027	-0.206	0.239	0.057	0.001	-0.083
8	a	0.344	0.359	0.538	0.390	0.206	0.471	0.521	0.365
	d	0.039	-0.004	-0.022	-0.235	0.269	0.048	0.001	-0.081
9	a	0.347	0.389	0.533	0.443	0.235	0.488	0.535	0.377
	d	0.055	-0.023	-0.033	-0.250	0.309	0.035	-0.005	-0.072
10	a	0.333	0.431	0.538	0.501	0.250	0.509	0.535	0.380
	d	0.071	-0.066	-0.020	-0.251	0.343	0.035	-0.010	-0.069
11	a	0.316	0.473	0.551	0.549	0.245	0.534	0.519	0.382
	d	0.082	-0.106	-0.007	-0.249	0.365	0.039	-0.011	-0.083
12	a	0.315	0.509	0.564	0.586	0.233	0.570	0.505	0.385
	d	0.102	-0.128	0.003	-0.244	0.365	0.037	-0.012	-0.102
13	a	0.327	0.536	0.575	0.615	0.235	0.603	0.509	0.389
	d	0.134	-0.133	0.014	-0.229	0.348	0.044	-0.015	-0.114
14	a	0.332	0.537	0.601	0.639	0.251	0.624	0.526	0.383
	d	0.154	-0.145	0.005	-0.206	0.345	0.058	-0.008	-0.135
15	a	0.341	0.546	0.626	0.660	0.267	0.617	0.556	0.396
	d	0.187	-0.162	-0.004	-0.182	0.342	0.076	-0.008	-0.147
16	a	0.346	0.546	0.644	0.660	0.257	0.592	0.619	0.419
	d	0.224	-0.168	-0.006	-0.159	0.347	0.109	-0.045	-0.157

Effects are adjusted for variation due to sex, line, and mother. QTL are designated by chromosome and cM position (chromosome@cM).

positions from Cheverud *et al.* (1996). Although our judgment of QTL overlap is rough, this result suggests that the evolution of body size involved different genetic changes in GI mice and laboratory strains. This conclusion is bolstered by comparisons to other QTL known to affect body weight in mice (Dragani *et al.* 1995; Cheverud *et al.* 1996; Keightley *et al.* 1996; Rance *et al.* 1997; Brockmann *et al.* 1998, 2004; Kirkpatrick *et al.* 1998; Morris *et al.* 1999; Vaughn *et al.* 1999; Corva and Medrano 2001; Rocha *et al.* 2004; Bennett *et al.* 2005; Kenney-Hunt *et al.* 2006; Shao *et al.* 2007; Casellas *et al.* 2009; Mollah and Ishikawa 2011; Ishikawa and Okuno 2014). Although a few of these QTL overlap with those we identified, most do not. For example, a major X-linked QTL for body weight discovered from another artificial selection experiment (Liu *et al.* 2001; Oliver *et al.* 2005) is not found in our study.

Nevertheless, those QTL that are shared across strains and populations may be repeatedly targeted by selection and represent promising loci for identifying the causative mutations (Chan *et al.* 2012). A series of genetic mapping studies identified a large number of loci that contribute to weight differences between *M. m. castaneus* from the Philippines (a smaller form) and the C57BL/6 laboratory strain (primarily *M. m. domesticus* in origin) (Ishikawa *et al.* 2000;

Ishikawa and Namikawa 2004; Mollah and Ishikawa 2010), including one in the region of the QTL we identified in the distal part of chromosome 10. Although the considerable divergence time between the two subspecies *M. m. castaneus* and *M. m. domesticus* makes it difficult to associate QTL with island evolution in this case, the resemblance between these results and ours is encouraging. Furthermore, a population genetic study that compared the genomes of laboratory mice artificially selected for divergent body sizes discovered genomic regions showing consistent signatures of adaptive evolution and unusual patterns of variation in large mice from the Faroe Islands (Chan *et al.* 2012). One of these genomic regions is located within the distal QTL on chromosome 10. Genetic characterization of mice from other islands will reveal whether shared patterns of size evolution across islands reflect a common genetic basis.

QTL mapped in crosses between a single pair of strains could represent genetic changes along either lineage. As a result, some QTL we identified might have contributed to size evolution in WSB or its ancestors rather than conferring phenotypic changes in GI mice. If the time to the most recent common ancestor of GI mice and WSB was earlier than the colonization time of Gough Island, using WSB as a reference

Table 3 Additive and dominance effects (in grams per week) of growth rate QTL

Week		1@39.5	4@42.5	6@4.0	7@32.0	7@44.0	8@17.5	10@26.0	10@62.0	13@52.0	15@9.0	16@10.0
1	a	0.047	0.031	0.021	0.072	0.025	0.047	0.001	0.124	-0.034	0.065	0.069
	d	0.019	0.033	0.039	-0.027	0.052	0.002	0.007	0.040	0.006	0.036	0.047
2	a	0.071	0.018	0.045	0.093	0.058	0.049	-0.001	0.165	-0.050	0.075	0.077
	d	0.036	0.046	0.028	-0.058	0.093	0.000	0.004	0.006	-0.022	0.077	0.040
3	a	0.099	-0.003	0.084	0.108	0.097	0.052	0.025	0.152	-0.091	0.053	0.070
	d	0.024	0.057	-0.015	-0.061	0.081	-0.001	-0.012	-0.006	-0.043	0.071	0.025
4	a	0.097	-0.018	0.102	0.061	0.132	0.050	0.063	0.091	-0.108	0.022	0.030
	d	0.001	0.043	-0.041	-0.038	0.042	-0.012	-0.021	-0.013	-0.035	0.043	0.008
5	a	0.052	-0.016	0.077	-0.033	0.118	0.045	0.071	0.014	-0.073	0.005	-0.013
	d	-0.016	-0.003	-0.028	0.007	-0.029	-0.025	-0.009	-0.012	-0.006	0.014	-0.015
6	a	0.013	0.001	0.044	-0.078	0.066	0.047	0.054	-0.018	-0.023	-0.009	-0.032
	d	-0.014	-0.054	-0.007	0.024	-0.078	-0.025	0.003	-0.014	0.013	0.004	-0.026
7	a	0.004	0.019	0.024	-0.056	0.021	0.043	0.040	-0.014	0.000	-0.031	-0.038
	d	-0.001	-0.073	0.002	0.012	-0.072	-0.014	-0.007	-0.018	-0.002	-0.005	-0.023
8	a	0.000	0.029	0.022	-0.008	-0.011	0.041	0.026	-0.005	0.001	-0.042	-0.030
	d	0.011	-0.061	-0.010	0.010	-0.035	-0.005	-0.022	-0.020	-0.017	-0.006	-0.024
9	a	-0.011	0.037	0.033	0.025	-0.023	0.047	0.022	-0.004	-0.001	-0.031	-0.021
	d	0.013	-0.036	-0.037	0.008	0.002	0.014	-0.013	-0.017	0.001	-0.002	-0.022
10	a	-0.022	0.042	0.040	0.023	-0.013	0.052	0.025	-0.013	0.001	-0.023	-0.017
	d	0.008	-0.013	-0.046	-0.016	0.027	0.034	0.001	-0.011	0.025	0.005	-0.016
11	a	-0.013	0.042	0.035	-0.003	0.013	0.053	0.032	-0.016	-0.004	-0.034	-0.020
	d	0.010	0.005	-0.028	-0.039	0.038	0.030	0.000	-0.004	0.024	0.018	-0.006
12	a	0.004	0.043	0.027	-0.042	0.042	0.052	0.038	-0.004	-0.012	-0.037	-0.033
	d	0.021	0.014	-0.007	-0.039	0.039	0.019	0.006	0.000	0.014	0.013	0.006
13	a	0.011	0.044	0.017	-0.067	0.060	0.042	0.028	0.014	-0.008	-0.031	-0.034
	d	0.039	0.004	0.001	-0.022	0.030	0.024	0.019	-0.003	0.004	-0.010	0.015
14	a	0.009	0.026	0.007	-0.056	0.064	0.027	0.008	0.025	0.004	-0.022	-0.024
	d	0.042	-0.016	-0.013	-0.024	0.014	0.021	0.021	-0.001	-0.018	-0.016	0.019
15	a	0.006	0.019	0.006	-0.033	0.051	0.016	-0.020	0.038	0.010	-0.007	-0.002
	d	0.042	-0.021	-0.016	-0.038	0.013	0.033	0.026	-0.007	-0.029	-0.023	0.029
16	a	0.000	0.025	0.000	-0.026	0.033	0.010	-0.043	0.059	0.018	0.003	0.011
	d	0.042	-0.031	-0.005	-0.047	0.025	0.072	0.034	-0.041	-0.012	-0.031	0.026

Effects are adjusted for variation due to sex, line, and mother. QTL are designated by chromosome and cM position (chromosome@cM).

strain could have exacerbated this problem. Although the mice used to found the WSB strain (which were caught in Maryland) were likely descended from Western European mice, determining the precise geographic origin of the GI mice will help to elucidate the dynamics of size evolution on the island.

Our results usefully focus the search for the genetic, developmental, and molecular mechanisms that ultimately underlie the evolution of body size in nature. GI mice evolved extreme size by accelerating growth during the first 6 weeks of life. The observation that GI mice are heavier at 1 week of age suggests that accelerated growth begins *in utero* or soon after birth. In addition to eliminating alternative evolutionary scenarios—including the notion that size was increased by extending the growth trajectory—our results suggest that natural selection on size or size-correlated traits targeted early development. This research is a first, necessary, and foundational step toward pinpointing the genetic variants responsible for increased size in GI mice. Identifying the causative genes and mutations will allow several intriguing evolutionary questions to be answered. Did selection on Gough Island target standing variants or new mutations? Do the causative loci show

signatures of adaptive evolution predicted by strong selection on individual loci or by selection spread across many genes?

How did the genetic changes that increased body size in GI mice evolve? Rapid phenotypic evolution in a novel environment suggests natural selection as the primary evolutionary force. The observed bias toward GI mouse alleles increasing body size provides modest support for this conclusion. Differences in the environments experienced by mice on Gough Island and on the mainland are potential selective agents. Mice on the island may experience significantly colder temperatures because they live outside year round (in contrast to mice on the mainland that often live indoors). It is possible that the availability of an energetically rich food resource in the winter period (albatross and petrel chicks) and the predatory behavior of mice to exploit this resource (Jones *et al.* 2003; Cuthbert and Hilton 2004; Wanless *et al.* 2007) were important selective factors in the rapid evolution of body size. The size increase of GI mice also could have been catalyzed by the removal of selective constraints. GI mice have few natural predators or interspecific competitors (Verrill 1895; Heaney and Holdgate 1957; Jones *et al.* 2003; Wanless *et al.* 2009). Both reduced predation and reduced competition have been invoked to




Figure 8 Estimated heritability due to the set of QTL in the multiple-QTL models for body weight (blue) and growth rate (pink).

explain the tendency of small mammals to grow larger on islands (Sondaar 1977; Heaney 1978; Lawlor 1982), and recent comparative work supports this conclusion (Lomolino *et al.* 2012). In principle, GI mice could have achieved their unusual body size without natural selection. Perhaps the evolution of body size was enabled by stronger genetic drift resulting from the reduction in population size that accompanied island colonization (Gray *et al.* 2014). Understanding the relative contributions of different evolutionary processes to extreme body size evolution in mice on Gough Island will ultimately require additional field studies.

Acknowledgments

We thank Trevor Glass and the administrator of Tristan da Cunha for permission to live capture and remove mice from Gough Island and Henk Louw and Paul Visser for capturing mice. We thank Lauren Brooks for sharing her unpublished compilation of body weights for mice from different islands. We thank Hannah Buchanan, Valeri Lapacek, Spencer Compton, Arielle Henderson, Elizabeth Linder, Lauren Hennelly, and Lauren Agnew for their devotion to mouse husbandry and phenotyping. We acknowledge staff at the Charmany Instructional Facility for assistance with all aspects of mouse care. We thank Mark Nolte for useful comments on the manuscript. Mark Nolte, Alan Attie, and Mark Keller provided helpful guidance about genetic mechanisms of mouse growth. We acknowledge Chris Vinyard for sharing ideas about the connection between fitness and body size. We thank Peicheng Jing for computational support. We appreciate enthusiasm and encouragement from Payseur lab members, including Mark Nolte, Beth Dumont, Michael White, Ryan Haasl, Jenny Andrie, Leslie Turner, John Hvala, Richard Wang, Lauren Brooks, April Peterson, and Peicheng Jing, throughout the course of this project. We thank Leonie Moyle and two anonymous reviewers for their constructive comments on the manuscript. The research was supported by National Institutes of Health (NIH) R01 GM100426A to B.A.




Figure 9 Comparison of differences in average GI and WSB values with summed additive effects of QTL.

P., NIH National Research Service Award 1F32GM090685 to M.M.G., and a National Science Foundation graduate research fellowship to M.D.P.

Literature Cited

- Adler, G. H., and R. Levins, 1994 The island syndrome in rodent populations. *Q. Rev. Biol.* 69: 473–490.
- Atchley, W. R., R. Wei, and P. Crenshaw, 2000 Cellular consequences in the brain and liver of age-specific selection for rate of development in mice. *Genetics* 155: 1347–1357.
- Aylor, D. L., W. Valdar, W. Foulds-Mathes, R. J. Buus, R. A. Verdugo *et al.*, 2011 Genetic analysis of complex traits in the emerging Collaborative Cross. *Genome Res.* 21: 1213–1222.
- Beavis, W. D., 1994 The power and deceit of QTL experiments: lessons from comparative QTL studies. In: *Proceedings of the Forty-Ninth Annual Corn and Sorghum Industry Research Conference*, Washington, D.C., pp. 250–266.
- Beavis, W. D., 1998 QTL analyses: power, precision, and accuracy, pp. 145–162 in *Molecular Dissection of Complex Traits*, edited by A. H. Paterson. CRC Press, Boca Raton, FL.

- Bennett, B., P. J. Carosone-Link, L. Lu, E. J. Chesler, and T. E. Johnson, 2005 Genetics of body weight in the LXS recombinant inbred mouse strains. *Mamm. Genome* 16: 764–774.
- Berry, R. J., 1964 The evolution of an island population of the house mouse. *Evolution* 18: 468–483.
- Berry, R. J., 1965 Genetical change in an island mouse population. *Heredity* 20: 651.
- Berry, R. J., 1968 The ecology of an island population of house mouse. *J. Anim. Ecol.* 37: 445–470.
- Berry, R. J., 1996 Small mammal differentiation on islands. *Philos. Trans. R. Soc. B* 351: 753–764.
- Berry, R. J., and M. E. Jakobson, 1975 Ecological genetics of an island population of house mouse (*Mus-musculus*). *J. Zool.* 175: 523–540.
- Berry, R. J., and P. N. Scriven, 2005 The house mouse: a model and motor for evolutionary understanding. *Biol. J. Linn. Soc. Lond.* 84: 335–347.
- Berry, R. J., M. E. Jakobson, and J. Peters, 1978a The house mice of the Faroe Islands: a study in microdifferentiation. *J. Zool.* 185: 573–592.
- Berry, R. J., J. Peters, and R. J. Van Aarde, 1978b Sub-antarctic house mice: colonization, survival and selection. *J. Zool.* 184: 127–141.
- Berry, R. J., W. N. Bonner, and J. Peters, 1979 Natural-selection in house mice (*Mus-musculus*) from South Georgia (South-Atlantic Ocean). *J. Zool.* 189: 385–398.
- Berry, R. J., A. Cuthbert, and J. Peters, 1982 Colonization by house mice: an experiment. *J. Zool.* 198: 329–336.
- Berry, R. J., M. E. Jakobson, and J. Peters, 1987 Inherited differences within an island population of the house mouse (*Mus-domesticus*). *J. Zool. (Lond.)* 211: 605–618.
- Brockmann, G. A., C. S. Haley, U. Renne, S. A. Knott, and M. Schwerin, 1998 Quantitative trait loci affecting body weight and fatness from a mouse line selected for extreme high growth. *Genetics* 150: 369–381.
- Brockmann, G. A., E. Karatayli, C. S. Haley, U. Renne, O. J. Rottmann *et al.*, 2004 QTLs for pre- and postweaning body weight and body composition in selected mice. *Mamm. Genome* 15: 593–609.
- Broman, K. W., and T. P. Speed, 2002 A model selection approach for the identification of quantitative trait loci in experimental crosses. *J. R. Stat. Soc., B* 64: 641–656.
- Broman, K. W., S. Sen, S. E. Owens, A. Manichaikul, E. M. Southard-Smith *et al.*, 2006 The X chromosome in quantitative trait locus mapping. *Genetics* 174: 2151–2158.
- Bromham, L., and M. Cardillo, 2007 Primates follow the “island rule”: implications for interpreting *Homo floresiensis*. *Biol. Lett.* 3: 398–400.
- Bünger, L., and W. G. Hill, 1999 Inbred lines of mice derived from long-term divergent selection on fat content and body weight. *Mamm. Genome* 10: 645–648.
- Calder, W. A., 1984 *Size, Function, and Life History*. Harvard University Press, Cambridge, MA.
- Carter, T. C., and D. S. Falconer, 1951 Stocks for detecting linkage in the mouse, and the theory of their design. *J. Genet.* 50: 307–323.
- Case, T. J., 1978 General explanation for insular body size trends in terrestrial vertebrates. *Ecology* 59: 1–18.
- Casellas, J., C. R. Farber, R. J. Gualarte, K. A. Haus, C. H. Warden *et al.*, 2009 Evidence of maternal QTL affecting growth and obesity in adult mice. *Mamm. Genome* 20: 269–280.
- Chai, C., 1956a Analysis of quantitative inheritance of body size in mice. I. Hybridization and maternal influence. *Genetics* 41: 157–164.
- Chai, C., 1956b Analysis of quantitative inheritance of body size in mice. II. Gene action and segregation. *Genetics* 41: 167–178.
- Chan, Y. F., M. E. Marks, F. C. Jones, G. Villarreal, and M. D. Shapiro *et al.*, 2010 Adaptive evolution of pelvic reduction in sticklebacks by recurrent deletion of a *Pitx1* enhancer. *Science* 327: 302–305.
- Chan, Y. F., F. C. Jones, E. McConnell, J. Bryk, L. Bünger *et al.*, 2012 Parallel selection mapping using artificially selected mice reveals body weight control loci. *Curr. Biol.* 22: 794–800.
- Cheverud, J. M., J. J. Rutledge, and W. R. Atchley, 1983a Quantitative genetics of development: genetic correlations among age-specific trait values and the evolution of ontogeny. *Evolution* 37: 895–905.
- Cheverud, J. M., L. Leamy, J. Rutledge, and W. Atchley, 1983b Quantitative genetics and the evolution of ontogeny. I. Ontogenetic changes in quantitative genetic variance components in randombred mice. *Genet. Res.* 42: 65–75.
- Cheverud, J. M., E. J. Routman, F. A. M. Duarte, B. Van Swinderen, K. Cothran *et al.*, 1996 Quantitative trait loci for murine growth. *Genetics* 142: 1305–1319.
- Cheverud, J. M., 2005 Genetics of growth in the mouse, pp. 113–130 in *The Mouse in Animal Genetics and Breeding Research*, edited by E. J. Eisen. Imperial College Press, Singapore.
- Churchill, G. A., and R. W. Doerge, 1994 Empirical threshold values for quantitative trait mapping. *Genetics* 138: 963–971.
- Churchill, G. A., D. C. Airey, H. Allayee, J. M. Angel, A. D. Attie *et al.*, 2004 The Collaborative Cross, a community resource for the genetic analysis of complex traits. *Nat. Genet.* 36: 1133–1137.
- Colosimo, P. F., K. E. Hosemann, S. Balabhadra, G. Villarreal, M. Dickson *et al.*, 2005 Widespread parallel evolution in sticklebacks by repeated fixation of *Ectodysplasin* alleles. *Science* 307: 1928–1933.
- Corva, P. M., and J. F. Medrano, 2001 Quantitative trait loci (QTLs) mapping for growth traits in the mouse: a review. *Genet. Sel. Evol.* 33: 105–132.
- Cox, A., C. Ackert-Bicknell, B. L. Dumont, Y. Ding, J. Tzenova Bell *et al.*, 2009 A new standard genetic map for the mouse. *Genetics* 182: 1335–1344.
- Cuthbert, R., and G. Hilton, 2004 Introduced house mice *Mus musculus*: A significant predator of threatened and endemic birds on Gough Island, South Atlantic Ocean? *Biol. Conserv.* 117: 483–489.
- Damuth, J., and B. J. MacFadden, 1990 *Body Size in Mammalian Paleobiology*, Cambridge University Press, New York.
- Didion, J. P., A. P. Morgan, A. M.-F. Clayshulte, R. C. McMullan, L. Yadgary *et al.*, 2015 A multi-megabase copy number gain causes maternal transmission ratio distortion on mouse chromosome 2. *PLoS Genet.* 11: e1004850.
- Dragani, T. A., Z. B. Zeng, F. Canzian, M. Gariboldi, M. T. Ghilarducci *et al.*, 1995 Mapping of body weight loci on mouse Chromosome X. *Mamm. Genome* 6: 778–781.
- Eisen, E. J., 1989 Selection experiments for body composition in mice and rats: a review. *Livest. Prod. Sci.* 23: 17–32.
- Endler, J. A., 1986 *Natural Selection in the Wild*, Princeton University Press, Princeton, NJ.
- Falconer, D. S., 1953 Selection for large and small size in mice. *Genetics* 51: 470–501.
- Festing, M. F. W., 1996 Origins and characteristics of inbred strains of mice, *Genetic Variants and Strains of the Laboratory Mouse*, edited by M. F. Lyon, S. Raston, and S. D. M. Brown. Oxford University Press, New York.
- Foster, J. B., 1964 The evolution of mammals on islands. *Nature* 202: 234–235.
- Fowden, A. L., 2003 The insulin-like growth factors and fetoplacental growth. *Placenta* 24: 803–812.
- Freudenthal, M., 1972 *Deinogalerix koenigswaldi* nov. gen., nov. spec., a giant insectivore from the Neogene of Italy. *Scr. Geol.* 14: 1–19.

- Goodale, H., 1938 A study of the inheritance of body weight in the albino mouse by selection. *J. Hered.* 29: 101–112.
- Gray, M. M., D. Wegmann, R. J. Haas, M. A. White, S. I. Gabriel *et al.*, 2014 Demographic history of a recent invasion of house mice on the isolated Island of Gough. *Mol. Ecol.* 23: 1923–1939.
- Haley, C. S., and S. A. Knott, 1992 A simple regression method for mapping quantitative trait loci in line crosses using flanking markers. *Heredity* 69: 315–324.
- Heaney, J. T., and M. W. Holdgate, 1957 The Gough Island scientific survey. *Geogr. J.* 123: 20–31.
- Heaney, L. R., 1978 Island area and body size of insular mammals: evidence from the tri-colored squirrel (*Callosciurus prevosti*) of Southeast Asia. *Evolution* 32: 29–44.
- Hoekstra, H. E., R. J. Hirschmann, R. A. Bunday, P. A. Insel, and J. P. Crossland, 2006 A single amino acid mutation contributes to adaptive beach mouse color pattern. *Science* 313: 101–104.
- Ishikawa, A., Y. Matsuda, and T. Namikawa, 2000 Detection of quantitative trait loci for body weight at 10 weeks from Philippine wild mice. *Mamm. Genome* 11: 824–830.
- Ishikawa, A., and T. Namikawa, 2004 Mapping major quantitative trait loci for postnatal growth in an intersubspecific backcross between C57BL/6J and Philippine wild mice by using principal component analysis. *Genes Genet. Syst.* 79: 27–39.
- Ishikawa, A., and S. Okuno, 2014 Fine mapping and candidate gene search of quantitative trait loci for growth and obesity using mouse intersubspecific subcongenic intercrosses and exome sequencing. *PLoS One* 9: e113233.
- Jones, A. G., S. L. Chown, and K. J. Gaston, 2003 Introduced house mice as a conservation concern on Gough Island. *Biodivers. Conserv.* 12: 2107–2119.
- Jones, F. C., M. G. Grabherr, Y. F. Chan, P. Russell, and E. Mauceli *et al.*, 2012 The genomic basis of adaptive evolution in threespine sticklebacks. *Nature* 484: 55–61.
- Keane, T. M., L. Goodstadt, P. Danecek, M. A. White, K. Wong *et al.*, 2011 Mouse genomic variation and its effect on phenotypes and gene regulation. *Nature* 477: 289–294.
- Keightley, P. D., T. Hardge, L. May, and G. Bulfield, 1996 A genetic map of quantitative trait loci for body weight in the mouse. *Genetics* 142: 227–235.
- Kenney-Hunt, J. P., T. T. Vaughn, L. S. Pletscher, A. Peripato, E. Routman *et al.*, 2006 Quantitative trait loci for body size components in mice. *Mamm. Genome* 17: 526–537.
- Kirkpatrick, B. W., A. Mengelt, N. Schulman, and I. C. Martin, 1998 Identification of quantitative trait loci for prolificacy and growth in mice. *Mamm. Genome* 9: 97–102.
- Kramer, M. G., T. T. Vaughan, L. S. Pletscher, K. King-Ellison, E. Adams *et al.*, 1998 Genetic variation in body weight gain and composition in the intercross of Large (LG/J) and Small (SM/J) inbred strains of mice. *Genet. Mol. Biol.* 21: 211–218.
- Kwak, I.-Y., C. R. Moore, E. P. Spalding, and K. W. Broman, 2014 A simple regression-based method to map quantitative trait loci underlying function-valued phenotypes. *Genetics* 197: 1409–1416.
- Lawlor, T. E., 1982 The evolution of body size in mammals: evidence from insular populations in Mexico. *Am. Nat.* 119: 54–72.
- Liu, X. J., F. Oliver, S. D. M. Brown, P. Denny, and P. D. Keightley, 2001 High-resolution quantitative trait locus mapping for body weight in mice by recombinant progeny testing. *Genet. Res.* 77: 191–197.
- Lomolino, M. V., 1985 Body size of mammals on islands: the island rule reexamined. *Am. Nat.* 125: 310–316.
- Lomolino, M. V., 2005 Body size evolution in insular vertebrates: generality of the island rule. *J. Biogeogr.* 32: 1683–1699.
- Lomolino, M. V., B. R. Riddle, and J. H. Brown, 2005 *Biogeography*, Sinauer Associates, Sunderland, MA.
- Lomolino, M. V., D. F. Sax, M. R. Palombo, and A. A. van der Geer, 2012 Of mice and mammoths: evaluations of causal explanations for body size evolution in insular mammals. *J. Biogeogr.* 39: 842–854.
- Losos, J. B., and R. E. Ricklefs, 2009 Adaptation and diversification on islands. *Nature* 457: 830–836.
- Lui, J. C., and J. Baron, 2011 Mechanisms limiting body growth in mammals. *Endocr. Rev.* 32: 422–440.
- Lupu, F., J. D. Terwilliger, K. Lee, G. V. Segre, and A. Efstatiadis, 2001 Roles of growth hormone and insulin-like growth factor 1 in mouse postnatal growth. *Dev. Biol.* 229: 141–162.
- Lynch, C., 1992 Clinal variation in cold adaptation in *Mus domesticus*: verification of predictions from laboratory populations. *Am. Nat.* 139: 1219–1236.
- MacArthur, J., 1944 Genetics of body size and related characters. I. Selection of small and large races of the laboratory mouse. *Am. Nat.* 78: 142–157.
- Meiri, S., T. Dayan, and D. Simberloff, 2004 Body size of insular carnivores: little support for the island rule. *Am. Nat.* 163: E469–E479.
- Meiri, S., N. Cooper, and A. Purvis, 2008 The island rule: Made to be broken? *Proc. Biol. Sci.* 275: 141–148.
- Meiri, S., P. Raia, and A. B. Phillimore, 2011 Slaying dragons: limited evidence for unusual body size evolution on islands. *J. Biogeogr.* 38: 89–100.
- Millien, V., 2006 Morphological evolution is accelerated among island mammals. *PLoS Biol.* 4: 1863–1868.
- Mollah, B. R., and A. Ishikawa, 2010 A wild derived quantitative trait locus on mouse chromosome 2 prevents obesity. *BMC Genet.* 11: 1–10.
- Mollah, M. B. R., and A. Ishikawa, 2011 Intersubspecific subcongenic mouse strain analysis reveals closely linked QTLs with opposite effects on body weight. *Mamm. Genome* 22: 282–289.
- Moncunill-Solà, B., X. Jordana, N. Marín-Moratalla, S. Moyà-Solà, and M. Köhler, 2014 How large are the extinct giant insular rodents? New body mass estimations from teeth and bones. *Integr. Zool.* 9: 197–212.
- Morris, K. H., A. Ishikawa, and P. D. Keightley, 1999 Quantitative trait loci for growth traits in C57BL/6J x DBA/2J mice. *Mamm. Genome* 10: 225–228.
- Nachman, M. W., H. E. Hoekstra, and S. L. D'Agostino, 2003 The genetic basis of adaptive melanism in pocket mice. *Proc. Natl. Acad. Sci. USA* 100: 5268–5273.
- Oliver, F., J. K. Christians, X. Liu, S. Rhind, V. Verma *et al.*, 2005 Regulatory variation at glypican-3 underlies a major growth QTL in mice. *PLoS Biol.* 3: e135.
- Orr, H. A., 1998 Testing natural selection vs. genetic drift in phenotypic evolution using quantitative trait locus data. *Genetics* 149: 2099–2104.
- Pergams, O. R. W., and M. V. Ashley, 2001 Microevolution in island rodents. *Genetica* 112: 245–256.
- Peters, R. H., 1983 *The Ecological Implications of Body Size*, Cambridge University Press, Cambridge.
- Prasad, K. V. S. K., B.-H. Song, C. Olson-Manning, J. T. Anderson, C.-R. Lee *et al.*, 2012 A gain-of-function polymorphism controlling complex traits and fitness in nature. *Science* 337: 1081–1084.
- Qiao, N., Y. Huang, H. Naveed, C. D. Green, and J.-D. J. Han, 2013 CoCiter: an efficient tool to infer gene function by assessing the significance of literature co-citation. *PLoS One* 8: e74074.
- Raia, P., and S. Meiri, 2006 The island rule in large mammals: paleontology meets ecology. *Evolution* 60: 1731–1742.
- Rance, K. A., W. G. Hill, and P. D. Keightley, 1997 Mapping quantitative trait loci for body weight on the X chromosome in mice. I. Analysis of a reciprocal F2 population. *Genet. Res.* 70: 117–124.

- Rebeiz, M., J. E. Pool, V. A. Kassner, C. F. Aquadro, and S. B. Carroll, 2009 Stepwise modification of a modular enhancer underlies adaptation in a *Drosophila* population. *Science* 326: 1663–1667.
- Reed, D. R., M. P. Lawler, and M. G. Tordoff, 2008 Reduced body weight is a common effect of gene knockout in mice. *BMC Genet.* 9: 4.
- Riska, B., W. R. Atchley, and J. J. Rutledge, 1984 A genetic analysis of targeted growth in mice. *Genetics* 107: 79–101.
- Riska, B., and W. R. Atchley, 1985 Genetics of growth predict patterns of brain-size evolution. *Science* 229: 668–671.
- Rocha, J. L., E. J. Eisen, L. D. Van Vleck, and D. Pomp, 2004 A large-sample QTL study in mice: I. Growth. *Mamm. Genome* 15: 83–99.
- Rockman, M. V., 2012 The QTN program and the alleles that matter for evolution: all that's gold does not glitter. *Evolution* 66: 1–17.
- Roth, V. L., 1992 Inferences from allometry and fossils: dwarfing of elephants on islands. *Oxford Surv. Evol. Biol.* 8: 259–288.
- Rowe-Rowe, D. T., and J. E. Crafford, 1992 Density, body size, and reproduction of feral house mice on Gough Island. *S. Afr. J. Zool.* 27: 1–5.
- Salcedo, T., A. Geraldes, and M. W. Nachman, 2007 Nucleotide variation in wild and inbred mice. *Genetics* 177: 2277–2291.
- Schmidt-Nielsen, K., 1984 *Scaling: Why Is Animal Size so Important?* University of Cambridge Press, Cambridge.
- Shao, H., D. R. Reed, and M. G. Tordoff, 2007 Genetic loci affecting body weight and fatness in a C57BL/6J x PWK/PhJ mouse intercross. *Mamm. Genome* 18: 839–851.
- Silver, L. M., 1995 *Mouse Genetics: Concepts and Applications*. Oxford University Press, New York.
- Sondaar, P. Y., 1977 Insularity and its effects on mammal evolution, pp. 671–707 in *Major Patterns of Vertebrate Evolution*, edited by M. K. Hecht, P. C. Goody, and B. M. Hecht. Plenum Press, New York.
- Stock, C., 1935 Exiled elephants of the Channel Islands. *Sci. Mon.* 41: 205–214.
- Svenson, K. L., D. M. Gatti, W. Valdar, C. E. Welsh, R. Cheng *et al.*, 2012 High-resolution genetic mapping using the Mouse Diversity outbred population. *Genetics* 190: 437–447.
- Threadgill, D. W., and G. A. Churchill, 2012 Ten years of the Collaborative Cross. *Genetics* 190: 291–294.
- Van Valen, L., 1973 Pattern and the balance of nature. *Evol. Theory* 1: 31–49.
- van't Hof, A. E., N. Edmonds, M. Dalíková, F. Marec, and I. J. Saccheri, 2011 Industrial melanism in British peppered moths has a singular and recent mutational origin. *Science* 332: 958–960.
- Vaughn, T. T., L. S. Pletscher, A. Peripato, K. King-Ellison, E. Adams *et al.*, 1999 Mapping quantitative trait loci for murine growth: a closer look at genetic architecture. *Genet. Res.* 74: 313–322.
- Verrill, G. E., 1895 Notes on birds and eggs from the islands of Gough, Kerguelen, and South Georgia. *Trans. Connect. Acad.* 11: 429–480.
- Wace, N. M., 1961 Vegetation of Gough Island. *Ecol. Monogr.* 31: 337–367.
- Wanless, R. M., A. Angel, R. J. Cuthbert, G. M. Hilton, and P. G. Ryan, 2007 Can predation by invasive mice drive seabird extinctions? *Biol. Lett.* 3: 241–244.
- Wanless, R. M., P. G. Ryan, R. Altwegg, A. Angel, J. Cooper *et al.*, 2009 From both sides: dire demographic consequences of carnivorous mice and longlining for the critically endangered Tristan albatrosses on Gough Island. *Biol. Conserv.* 142: 1710–1718.
- Zeng, Z. B., J. Liu, L. F. Stam, C. H. Kao, J. M. Mercer *et al.*, 2000 Genetic architecture of a morphological shape difference between two *Drosophila* species. *Genetics* 154: 299–310.

Communicating editor: L. C. Moyle

GENETICS

Supporting Information

www.genetics.org/lookup/suppl/doi:10.1534/genetics.115.177790/-/DC1

Genetics of Rapid and Extreme Size Evolution in Island Mice

**Melissa M. Gray, Michelle D. Parmenter, Caley A. Hogan, Irene Ford, Richard J. Cuthbert,
Peter G. Ryan, Karl W. Broman, and Bret A. Payseur**




Figure S1: $-\log_{10}$ P-values for the comparison of Gough Island (GI) and WSB mice for body weight (top panel) and growth rate (center panel), and for the comparison of the two sexes within GI (bottom panel).




Figure S2: Histograms of the body weight of F2 mice, by week. Vertical lines are plotted at the averages for the WSB (purple), F1 (blue), and Gough (green) parents.




Figure S3: Average \pm 2 SE of body weight (top panel) and growth rate (center panel) within the F2, and $-\log_{10}$ P-values for comparison of the two sexes.




Figure S4: $-\log_{10}$ P-values for the comparison among the four crosses (top two panels) and among all mothers (bottom two panels).





Figure S5: Genetic map (top panel) and physical map (bottom panel) of the 11,833 genetic markers considered in the QTL analyses.





Figure S6: Estimated frequency of the WSB allele in the F2 mice, separated by line of origin, along the autosomes (top panel), and along chromosome 2 (bottom panel).

Table S1. Candidate genes in QTL regions.

Gene Abbreviation	Gene Full Name	Co-Citer #	Chr	Start (Mb)	Genome Browser ID	Function and connection to body size	References
Cav1	Caveolin 1	261	6	17.3	uc009azo.1	Functions in lipid metabolism and in adipocytes	(RAZANI <i>et al.</i> 2002; COHEN <i>et al.</i> 2004; WERNSTEDT ASTERHOLM <i>et al.</i> 2012)
Met	Hepatocyte growth factor receptor precursor	762	6	17.4	uc009azr.1	Regulated by growth factors; promotes differentiation and proliferation of hematopoietic stem cells	(ORGAN and TSAO 2011; ISHIKAWA <i>et al.</i> 2012)
Cftr	Cystic fibrosis transmembrane conductance	195	6	18.1	uc009bai.1	Involved in secretory cell proliferation and differentiation	(GRUBB and BOUCHER 1999)
Pth	Parathyroid hormone	135	7	120.5	uc009jhm.1	Regulates serum calcium levels (important for energy metabolism, signaling, and skeletal growth)	(ROVERI <i>et al.</i> 1978; IWANIEC <i>et al.</i> 2007; YUAN <i>et al.</i> 2012)
Casp3	Caspase 3	428	8	47.7	uc009lql.1	Functions in apoptosis; involved in embryonic and	(MCILWAIN <i>et al.</i> 2013)

						hematopoietic stem cell differentiation	
Smad1	Mothers against decapentaplegic homolog 1	409	8	81.9	uc012ggl.1	Mediates signaling in association with bone morphogenetic proteins (BMPs); involved in cell growth, apoptosis, and morphogenesis	(RETTING <i>et al.</i> 2009; CHEN <i>et al.</i> 2012)
Apoa1	Apolipoprotein A1	158	9	46.0	uc009phb.2	Facilitates removal of cholesterol and other fats from tissues by transporting them to the liver for excretion; associated with obesity	(WAN <i>et al.</i> 2003; SLAGTER <i>et al.</i> 2013)
Cdk1	Cell division protein kinase 1	227	10	68.8	uc007fmr.1	Regulates cell cycle and cell proliferation	(KIYOKAWA <i>et al.</i> 1996; SANTAMARÍA <i>et al.</i> 2007; LUI and BARON 2011)
Cdk4	Cell division protein kinase 4	197	10	126.5	uc007hhv.2	Regulates cell cycle and cell proliferation	(KIYOKAWA <i>et al.</i> 1996; SANTAMARÍA <i>et al.</i> 2007; LUI and BARON 2011)
Gli1	Zinc finger	181	10	126.8	uc007hjf.1	Controls	(PARK <i>et al.</i>

	protein GLI1					transcription in Hedgehog pathway (important in skeletal formation, neuronal development, and overall body size)	2000; KIMURA <i>et al.</i> 2005)
ErbB3	Receptor tyrosine protein kinase erbB 3	220	10	128.0	uc007hnm.1	Affects cell growth, apoptosis and differentiation	(ZHANG <i>et al.</i> 2008)
Cdk2	Cell division protein kinase 2 isoform 1	247	10	128.1	uc007hny.2	Regulates cell cycle and cell proliferation	(KIYOKAWA <i>et al.</i> 1996; SANTAMARÍA <i>et al.</i> 2007; LUI and BARON 2011)
Prop1	Homeobox protein prophet of Pit1	104	11	50.8	uc007itg.1	Regulates production of circulating growth hormones; associated with ames dwarfism	(BARTKE 2000; NASONKIN <i>et al.</i> 2011)
Il4	Interleukin 4	224	11	53.4	uc007iwq.2	Involved in Hedgehog pathway (important in skeletal formation, neuronal development, and overall body size)	(KÜHN <i>et al.</i> 1991; SADLACK <i>et al.</i> 1994)
Isl1	Insulin gene enhancer protein ISL 1	145	13	117.1	uc007ryf.2	Transcription factor that mediates	(GUO <i>et al.</i> 2011; HO <i>et al.</i> 2014)

						proliferation of pancreatic islets maintenance of endocrine cells	
--	--	--	--	--	--	--	--

References Cited

- BARTKE A., 2000 Delayed aging in Ames dwarf mice. Relationships to endocrine function and body size. *Results Probl. Cell Differ.* **29**: 181–202.
- CHEN G., DENG C., LI Y.-P., 2012 TGF- β and BMP signaling in osteoblast differentiation and bone formation. *Int. J. Biol. Sci.* **8**: 272–288.
- COHEN A. W., RAZANI B., SCHUBERT W., WILLIAMS T. M., WANG X. B., IYENGAR P., BRASAEMLE D. L., SCHERER P. E., LISANTI M. P., 2004 Role of Caveolin-1 in the Modulation of Lipolysis and Lipid Droplet Formation. *Diabetes* **53**: 1261–1270.
- GRUBB B. R., BOUCHER R. C., 1999 Pathophysiology of gene-targeted mouse models for cystic fibrosis. *Physiol. Rev.* **79**: S193–S214.
- GUO T., WANG W., ZHANG H., LIU Y., CHEN P., MA K., ZHOU C., 2011 ISL1 promotes Pancreatic islet cell proliferation. *PLoS One* **6**: e22387.
- HO M. M., HU X., KARUNAKARAN S., JOHNSON J. D., CLEE S. M., 2014 Altered pancreatic growth and insulin secretion in WSB/EiJ mice. *PLoS One* **9**: e88352.
- ISHIKAWA T., FACTOR V. M., MARQUARDT J. U., RAGGI C., SEO D., KITADE M., CONNER E. A., THORGEIRSSON S. S., 2012 Hepatocyte growth factor/c-met signaling is required for stem-cell-mediated liver regeneration in mice. *Hepatology* **55**: 1215–1226.
- IWANIEC U. T., WRONSKI T. J., LIU J., RIVERA M. F., ARZAGA R. R., HANSEN G., BROMMAGE R., 2007 PTH stimulates bone formation in mice deficient in Lrp5. *J. bone Miner. Res.* **22**: 394–402.

- KIMURA H., STEPHEN D., JOYNER A., CURRAN T., 2005 Gli1 is important for medulloblastoma formation in Ptc1^{+/-} mice. *Oncogene* **24**: 4026–4036.
- KIYOKAWA H., KINEMAN R. D., MANOVA-TODOROVA K. O., SOARES V. C., HOFFMAN E. S., ONO M., KHANAM D., HAYDAY A. C., FROHMAN L. A., KOFF A., 1996 Enhanced growth of mice lacking the cyclin-dependent kinase inhibitor function of p27(Kip1). *Cell* **85**: 721–732.
- KÜHN R., RAJEWSKY K., MÜLLER W., 1991 Generation and analysis of interleukin-4 deficient mice. *Science*. **254**: 707–710.
- LUI J. C., BARON J., 2011 Mechanisms limiting body growth in mammals. *Endocr. Rev.* **32**: 422–440.
- MCILWAIN D. R., BERGER T., MAK T. W., 2013 Caspase functions in cell death and disease. *Cold Spring Harb. Perspect. Biol.* **5**: a008656.
- NASONKIN I. O., WARD R. D., BAVERS D. L., BEUSCHLEIN F., MORTENSEN A. H., KEEGAN C. E., HAMMER G. D., CAMPER S. A., 2011 Aged PROP1 deficient dwarf mice maintain ACTH production. *PLoS One* **6**: e28355.
- ORGAN S. L., TSAO M.-S., 2011 An overview of the c-MET signaling pathway. *Ther. Adv. Med. Oncol.* **3**: S7–S19.
- PARK H. L., BAI C., PLATT K. A., MATISE M. P., BEEGLY A., HUI C. C., NAKASHIMA M., JOYNER A. L., 2000 Mouse Gli1 mutants are viable but have defects in SHH signaling in combination with a Gli2 mutation. *Development* **127**: 1593–1605.
- RAZANI B., COMBS T. P., WANG X. B., FRANK P. G., PARK D. S., RUSSELL R. G., LI M., TANG B., JELICKS L. A., SCHERER P. E., LISANTI M. P., 2002 Caveolin-1-deficient mice are lean, resistant to diet-induced obesity, and show hypertriglyceridemia with adipocyte abnormalities. *J. Biol. Chem.* **277**: 8635–8647.
- RETTING K. N., SONG B., YOON B. S., LYONS K. M., 2009 BMP canonical Smad signaling through Smad1 and Smad5 is required for endochondral bone formation. *Development* **136**: 1093–1104.
- ROVERI N., RIPAMONTI A., GARBISA S., VOLPIN D., 1978 Structural organization of collagen fibrils in media aortic wall. *Connect. Tissue Res.* **5**: 249–253.

- SADLACK B., KÜHN R., SCHORLE H., RAJEWSKY K., MÜLLER W., HORAK I., 1994 Development and proliferation of lymphocytes in mice deficient for both interleukins-2 and -4. *Eur. J. Immunol.* **24**: 281–284.
- SANTAMARÍA D., BARRIÈRE C., CERQUEIRA A., HUNT S., TARDY C., NEWTON K., CÁCERES J. F., DUBUS P., MALUMBRES M., BARBACID M., 2007 Cdk1 is sufficient to drive the mammalian cell cycle. *Nature* **448**: 811–815.
- SLAGTER S. N., VLIET-OSTAPTCHOUK J. V VAN, VONK J. M., BOEZEN H. M., DULLAART R. P. F., KOBOLD A. C. M., FESKENS E. J., BEEK A. P. VAN, KLAUW M. M. VAN DER, WOLFFENBUTTEL B. H. R., 2013 Associations between smoking, components of metabolic syndrome and lipoprotein particle size. *BMC Med.* **11**: 195.
- WAN Y. J. Y., HAN G., CAI Y., DAI T., KONISHI T., LENG A. S., 2003 Hepatocyte retinoid X receptor- α -deficient mice have reduced food intake, increased body weight, and improved glucose tolerance. *Endocrinology* **144**: 605–611.
- WERNSTEDT ASTERHOLM I., MUNDY D. I., WENG J., ANDERSON R. G. W., SCHERER P. E., 2012 Altered mitochondrial function and metabolic inflexibility associated with loss of caveolin-1. *Cell Metab.* **15**: 171–185.
- YUAN Q., SATO T., DENSMORE M., SAITO H., SCHÜLER C., ERBEN R. G., LANSKE B., 2012 Deletion of PTH rescues skeletal abnormalities and high osteopontin levels in *Klotho*^{-/-} mice. *PLoS Genet.* **8**: e1002726.
- ZHANG Y., LU Y., ZHOU H., LEE M., LIU Z., HASSEL B. A., HAMBURGER A. W., 2008 Alterations in cell growth and signaling in ErbB3 binding protein-1 (Ebp1) deficient mice. *BMC Cell Biol.* **9**: 69.