

Cyclin-dependent kinase 5 activates guanine nucleotide exchange factor GIV/Girdin to orchestrate migration–proliferation dichotomy

Deepali Bhandari^{a,b,1}, Inmaculada Lopez-Sanchez^c, Andrew To^a, I-Chung Lo^a, Nicolas Aznar^c, Anthony Leyme^d, Vijay Gupta^a, Ingrid Niesman^a, Adam L. Maddox^b, Mikel Garcia-Marcos^d, Marilyn G. Farquhar^{a,1}, and Pradipta Ghosh^{a,c,1}

^aDepartment of Cellular and Molecular Medicine, University of California at San Diego, La Jolla, CA 92093-0651; ^bDepartment of Chemistry and Biochemistry, California State University Long Beach, Long Beach, CA 90840-9507; ^cDepartment of Medicine, University of California at San Diego, La Jolla, CA 92093-0651; and ^dDepartment of Biochemistry, Boston University School of Medicine, Boston, MA 02118

Contributed by Marilyn G. Farquhar, July 20, 2015 (sent for review May 16, 2015; reviewed by Danny Dhanasekaran, Henrik Gunnar Dohlman, and Kavita Shah)

Signals propagated by receptor tyrosine kinases (RTKs) can drive cell migration and proliferation, two cellular processes that do not occur simultaneously—a phenomenon called “migration–proliferation dichotomy.” We previously showed that epidermal growth factor (EGF) signaling is skewed to favor migration over proliferation via noncanonical transactivation of G α i proteins by the guanine exchange factor (GEF) GIV. However, what turns on GIV-GEF downstream of growth factor RTKs remained unknown. Here we reveal the molecular mechanism by which phosphorylation of GIV by cyclin-dependent kinase 5 (CDK5) triggers GIV’s ability to bind and activate G α i in response to growth factors and modulate downstream signals to establish a dichotomy between migration and proliferation. We show that CDK5 binds and phosphorylates GIV at Ser1674 near its GEF motif. When Ser1674 is phosphorylated, GIV activates G α i and enhances promigratory Akt signals. Phosphorylated GIV also binds G α s and enhances endosomal maturation, which shortens the transit time of EGFR through early endosomes, thereby limiting mitogenic MAPK signals. Consequently, this phosphoevent triggers cells to preferentially migrate during wound healing and transmigration of cancer cells. When Ser1674 cannot be phosphorylated, GIV cannot bind either G α i or G α s, Akt signaling is suppressed, mitogenic signals are enhanced due to delayed transit time of EGFR through early endosomes, and cells preferentially proliferate. These results illuminate how GIV-GEF is turned on upon receptor activation, adds GIV to the repertoire of CDK5 substrates, and defines a mechanism by which this unusual CDK orchestrates migration–proliferation dichotomy during cancer invasion, wound healing, and development.

migration–proliferation dichotomy | heterotrimeric G protein | growth factor receptor tyrosine kinase | GIV/Girdin | guanine nucleotide exchange factor

Upon growth factor stimulation, cells initiate signaling cascades favoring either migration or proliferation (migration–proliferation dichotomy) depending on the extracellular environmental cues and/or cellular needs. This dichotomy (also known as “go-or-grow mechanism”) plays a crucial role during a variety of normal and pathophysiologic processes, including development, wound healing, and cancer progression (1–5).

Of the multiple molecular mechanisms implicated in the orchestration of migration–proliferation dichotomy, G α -interacting vesicle associated protein (GIV) (also known as Girdin) is one such player that helps tilt the signaling network to favor migration over proliferation downstream of stimulated growth factor receptors as well as G protein-coupled receptors (GPCRs) (6–12). In the case of EGF stimulation, GIV is recruited to the plasma membrane (PM) where it directly binds ligand-activated EGFR via its SH2-like domain (13) and serves as a platform for the assembly of receptor tyrosine kinase (RTK)-GIV-G α i complexes and transactivation of G α i via its guanine exchange factor (GEF)

motif (8, 14). Such transactivation of Gi in the vicinity of RTKs at the PM enhances RTK autophosphorylation, prolongs RTK signaling from the PM, enhances PI3K/Akt signals and actin reorganization, and triggers cell migration (7–9, 13). Besides its role in the assembly of RTK-GIV-G α i complexes at the PM, GIV also assembles EGFR-GIV-G α s complexes on early endosomes where it facilitates down-regulation of EGFR via endosomal maturation, ensures finiteness of mitogenic signaling from that compartment, and limits cell proliferation (15).

Much of the experimental evidence supporting GIV’s ability to skew the phenotypic response toward migration has been generated using a GEF-deficient mutant (F1685A, FA) of GIV, which cannot bind either G α i at the PM or G α s on endosomes (8, 15). Without the formation of RTK-GIV-G α i/s complexes, ligand-activated EGFR spends a shorter time at the cell surface, but takes longer to transit through endosomes due to delayed endosomal maturation. Consequently, cells expressing GIV-FA suppress promigratory PI3K-Akt signals at the PM and enhance mitogenic signals from endosomes to preferentially trigger proliferation (8, 15). Despite the insights gained, the GIV-FA mutant did not illuminate how GIV-GEF may be reversibly switched “on/off” in physiology until recently, when that question was partially

Significance

The guanine nucleotide exchange factor (GEF) GIV/Girdin has previously been shown to trigger noncanonical activation of trimeric G proteins in response to multiple chemical stimuli at the plasma membrane and at various locations within cells. In this work we identified a single phosphoevent and pinpointed cyclin-dependent kinase 5 (CDK5) as the responsible kinase that activates GIV’s GEF function and initiates non-canonical G protein signaling via GIV. These studies provide evidence that CDK5 is essential for maximal activation of GIV-GEF in cells and insights into a hitherto unrecognized crosstalk between CDK5 and G proteins via GIV. Such crosstalk may shape migration–proliferation dichotomy during several physiological and pathological scenarios such as wound healing and cancer metastasis.

Author contributions: D.B., I.L.-S., A.T., I.-C.L., N.A., A.L., V.G., I.N., A.L.M., M.G.-M., and P.G. designed research; D.B., I.L.-S., A.T., I.-C.L., N.A., A.L., V.G., I.N., A.L.M., and M.G.-M. performed research; M.G.F. and P.G. contributed new reagents/analytic tools; D.B., I.L.-S., A.T., I.-C.L., N.A., A.L., V.G., I.N., M.G.-M., M.G.F., and P.G. analyzed data; and D.B., M.G.F., and P.G. wrote the paper.

Reviewers: D.D., University of Oklahoma Health Sciences Center; H.G.D., University of North Carolina, Chapel Hill; and K.S., Purdue University.

The authors declare no conflict of interest.

¹To whom correspondence may be addressed. Email: prghosh@ucsd.edu, deepali.bhandari@csulb.edu, or mfarquhar@ucsd.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1514157112/-DCSupplemental.

answered by the discovery of a key phosphoevent triggered by PKC θ at S1689 on GIV, which turns GIV-GEF “off” (16). However, what turns it “on” upon receptor activation still remained unknown.

Here, we identify a key phosphoevent triggered by cyclin-dependent kinase 5 (CDK5) which turns on GIV’s GEF activity by phosphorylating it at Ser1674 and thereby increases GIV’s ability to bind *Gai*s and enhances its ability to activate *Gai*. We provide




Fig. 1. Phosphomimetic S1674D (74D) mutation enhances GIV’s ability to bind and activate *Gai*3, whereas the nonphosphorylatable S1674A (74A) mutation abolishes binding. (A) Schematic of the domain architecture of GIV and sequence alignment of its C-terminal GEF motif. (Top) Various domains of GIV are shown including the N-terminal microtubule-binding hook domain, coiled-coil domain, $G\alpha$ -binding domain (GBD), and the C-terminal domain containing the GEF motif. The residue numbers marking the boundaries of each domain are shown. (Bottom) The sequence encompassing the GEF motif (in red rectangle) and surrounding residues was aligned among various species (accession numbers are shown in brackets) using ClustalW. Conserved residues are shaded in black and similar residues in gray. The most heavily phosphorylated residue in the C terminus as determined by mass spectrometry (S1674 in human GIV) is boxed in red. (B) Equimolar amounts of purified His-*Gai*3 were incubated with purified GST or GST-GIV-CT proteins immobilized on glutathione-Sepharose beads. Bound His-*Gai*3 was analyzed by immunoblotting with anti-His antibody. (C) Lysates of Cos7 cells expressing WT or mutant (74D or 74A) GST-GIV-CT were incubated with glutathione-Sepharose beads. Bound proteins (Upper) were analyzed by immunoblotting for endogenous *Gai*3. Equal loading was confirmed by immunoblotting the pull-downs for GST and GST-GIV CT and the lysates for *Gai*3 and actin. (D) Immunoprecipitation (IP) was carried out with anti-FLAG antibody on equal aliquots of lysates of Cos7 cells coexpressing FLAG-*Gai*3 and full-length Myc-tagged GIV-WT, 74D, or 74A followed by incubation with protein-G Sepharose beads. Bound immune complexes (IP) were analyzed for FLAG (*Gai*3) and Myc (GIV) by immunoblotting, and equal loading of lysates was confirmed by immunoblotting for tubulin. (E) The steady-state GTPase activity of His-*Gai*3 (50 nM) was determined in the presence of increasing amounts (0–2 μ M) of purified His-GIV-CT WT (solid circles) and 74D (open circles). *Gai*3 activation is expressed as percent of the steady-state GTPase activity of *Gai*3 alone. Results are shown as mean \pm SEM of three independent experiments. Activation of *Gai*3 by GIV-CT is enhanced in the case of the 74D mutant compared with its WT counterpart.

mechanistic insight into how GIV-GEF is activated and also define a previously unidentified substrate by which CDK5 triggers cell migration. Thus, this study illustrates a physiologic event to activate GIV's GEF function via a promigratory kinase, CDK5, which in turn dictates the orchestration of migration–proliferation dichotomy.

Results

Identification of a Key Phosphosite That Enhances GIV's GEF Activity.

We have previously shown that GIV binds G α i/s and activates G α i via an evolutionarily conserved short stretch of residues (amino acids 1678–1694), i.e., the GEF motif within its C terminus (Fig. 1A) (9, 15). We noted that there are 10 Ser/Thr within or flanking the GEF sequence, of which 9 are conserved, and several of them are phosphorylated as shown by multiple high-throughput phosphoproteomic studies (www.phosphosite.org). Because phosphorylation at one such site, S1689, is known to turn off GIV-GEF (16), we hypothesized that similar phospho-events may serve to turn on GIV-GEF. To explore what those phosphosites may be, we performed a mass spectrometric analysis on GIV's C terminus (amino acids 1660–1870, expressed and purified from Cos7 cells as GST-tagged protein; GST-GIV-CT). We used this construct because prior studies have demonstrated that this fragment of GIV (tagged with an equally bulky CFP-tag) is sufficient for assembly of RTK-GIV-G α i complexes at the PM and transactivation of Gi (14). We found that S1674, an evolutionarily conserved residue that lies adjacent to the GEF motif, is the most abundantly phosphorylated residue (~30–35% of all peptides), both at steady state in 10% (vol/vol) FBS and after EGF stimulation (Fig. 1A).

Given its proximity to the GEF motif, next we asked if phosphorylation of S1674 affects GIV's ability to bind G α i/s in vitro. We generated a phosphomimetic mutation at S1674 (74D) in GST-GIV-CT and carried out in vitro pulldown assays with recombinant 6xHis-tagged G α i3 and G α s. Compared with GIV-CT-WT, the phosphomimetic 74D mutant bound G α i3 (Fig. 1B) and G α s (Fig. S1A) with approximately three- to fivefold increased affinity. To determine how this phosphosite affects GIV-G protein interactions in cells, we generated both phosphomimetic (S1674D) and nonphosphorylatable (S1674A) GST-GIV-CT mutants, expressed them in mammalian cells, and tested their ability to bind endogenous G α i3 and G α s. Compared with GIV-WT, binding of GIV-74D to both G α i3 (Fig. 1C) and G α s (Fig. S1B) was enhanced, but the increase was not as robust as observed in in vitro binding assays carried out with recombinant proteins (Fig. 1B and Fig. S1A). A likely explanation for such a discrepancy came from mass spectrometry data, which showed that the S1674 residue in GIV-CT-WT is abundantly phosphorylated (~35%) in cells at steady state. The nonphosphorylatable GIV-74A mutant, on the other hand, showed little to no binding to either G α i3 (Fig. 1C) or G α s (Fig. S1B). These results indicate that S1674 is a key determinant of GIV's ability to bind G α i/s subunits and that phosphorylation at that site significantly improves GIV-G α interactions. Similar results were obtained when we immunoprecipitated FLAG-tagged G α i3 or G α s and looked for binding to Myc-tagged full-length GIV-74D, or 74A mutants compared to GIV-WT. Binding of the phosphomimetic 74D mutant was increased, and binding of the nonphosphorylatable 74A mutant to both G α i3 (Fig. 1D) and G α s (Fig. S1C) was decreased.

To determine if enhanced binding of the phosphomimetic mutant to G α i3 translates into higher GEF activity, we measured the steady-state GTPase activity of G α i3 in the presence of increasing amounts of recombinant His-GIV-CT-WT and 74D proteins as described previously (9). The 74D mutant was significantly more potent than GIV-CT-WT as a GEF at each concentration tested (Fig. 1E), suggesting that phosphorylation at S1674 augments GIV's ability to bind and activate G α i. As for G α s, no significant difference was observed between WT and

74D mutant His-GIV-CT proteins; both showed a weak inhibitory effect on the steady-state GTPase activity of G α s in vitro compared with the G protein binding-deficient FA mutant (Fig. S1D).

Phosphorylation of S1674 Regulates EGFR Trafficking and Signaling.

Because GIV-G α i/s interactions have been shown to shape RTK trafficking, which is a key determinant of downstream signals (8, 13, 15), next we asked if phosphorylation of GIV at S1674 affects EGFR trafficking. To answer this question, we generated GIV-depleted HeLa cell lines stably expressing full-length GIV-FLAG (WT, 74D, and 74A) at levels ~1.5- to 2-fold higher than endogenous GIV (Fig. S2A). Previous studies using this approach have characterized the phenotypic effects of various mutants of GIV in a clean background devoid of endogenous GIV (8, 13). Upon serum starvation, EGFR localized at the PM in all three cell lines (Fig. 2A, 0 min), but after EGF stimulation they showed differential responses. At 5 min after ligand stimulation, the amount of EGFR on the cell surface was higher in GIV-74D cells compared with GIV-WT, but little or no EGFR was on the cell surface in GIV-74A cells (Fig. 2A), indicating that the duration EGFR spends at the PM is increased when GIV is phosphorylated at S1674 and can effectively enhance Gi activity. At 15 min, EGFR colocalized with early endosome antigen 1 (EEA1) endosomes in GIV-WT and in both the mutants. By 30 min, the receptor was no longer seen in EEA1 endosomes in GIV-WT and GIV-74D cells, but in GIV-74A cells, EGFR still persisted on EEA1-positive endosomes. We further confirmed that such differences in transit time through EEA1-positive endosomes are associated with differential rates of down-regulation of receptor autophosphorylation. Autophosphorylated receptor as determined by anti-EGFR pY1068 (Fig. S2B) was rapidly cleared from endosomes in 74D cells (fewer pixels by 15 min, compared with GIV-WT and GIV-74A cells), but lingered on until 30 min in GIV-74A cells. Together, these findings indicate that the phosphomimicking GIV-74D mutant, which enhances G α i activation and binds more strongly to G α s, prolongs receptor stay at the PM, but once endocytosed, ensures rapid clearance of the ligand-activated receptor through endosomes and the finiteness of signaling from that site. By contrast, the nonphosphorylatable GIV-74A mutant, which impairs binding to G α s and G α i, shortens receptor stay at the PM, delays endosomal maturation, and prolongs the transit time and signaling from ligand-activated EGFR on EEA1 endosomes.

Phosphorylation of GIV at S1674 Orchestrates Migration–Proliferation Dichotomy During Epithelial Wound Healing and Morphogenesis.

The distribution of ligand-activated EGFR between the PM and endosomes is known to affect EGFR signaling (8, 17) in that promigratory PI3K-Akt signals are initiated by ligand-activated receptor largely or exclusively at the PM, whereas mitogenic MAPK-ERK1/2 signals can be propagated from endosomes (7, 18). Because trafficking of EGFR is closely intertwined with signals downstream of the receptor, next we asked if phosphomodulation of GIV's GEF motif at S1674 also affects signals that originate from the PM versus endosomes. Compared with the GIV-WT cells, those expressing the GIV-74A mutant, in which EGFR spends less time at the PM but stays longer on endosomes, suppress PM-based promigratory PI3K/Akt signals (as determined by phosphorylation of Akt at S473) and enhance endosome-based mitogenic MAPK/ERK signals [as determined by phosphorylation of ERK1/2 and cAMP response element-binding protein (CREB)] (Fig. 2B). By contrast, a mirror image response (high PI3K/Akt, low ERK1/2/CREB) was seen in cells expressing GIV-74D, in which EGFR spends more time at the PM but less on endosomes.

Next we hypothesized that the contrasting promigratory and prometotic signaling profiles in cells expressing GIV-74D and GIV-74A may translate into a differential preference for cell migration versus proliferation. We found that indeed such was

the case, because cells expressing GIV-74D, which enhance promigratory PI3K/Akt signals, displayed a higher migration index (~50% faster than WT), as determined by scratch-wound assays (Fig. 2C and Fig. S2C). Conversely, cells expressing GIV-74A,

which suppress promigratory signals, displayed a lower migration index (~25–30% slower compared with WT, and ~55% slower than GIV-74D). These cells also showed contrasting chemotaxis toward EGF, as determined by Transwell migration assay using a




Fig. 2. Phosphorylation of GIV at S1674 influences EGFR trafficking, signaling, and preference for migration versus proliferation of HeLa cells. (A) Serum starved (0.2% FBS, overnight) GIV-depleted HeLa cells stably expressing GIV-WT, 74D, and 74A (Fig. S2A) were stimulated with 50 nM EGF before fixation. Fixed cells were stained for total EGFR (green), the early endosome marker EEA1 (red), and DAPI (nucleus; blue) and examined by confocal microscopy. (Insets) Enlargement (3 \times) of boxed regions. (Scale bars, 10 μ m.) Phosphomimetic 74D mutant GIV enhances the duration EGFR spends at the PM (arrowheads, green pixels at 5 min), whereas the nonphosphorylatable 74A mutant GIV prolongs the duration EGFR spends on early endosomes (arrow, yellow pixels at 30 min). Bar graphs on the *Right* show quantification of pixels in each set of cells at the indicated time points. (B) Serum starved, GIV-depleted (by shRNA) HeLa cell lines stably expressing shRNA-resistant GIV-WT, 74D, and 74A were stimulated with 50 nM EGF before lysis. Equal aliquots of whole-cell lysates were analyzed for phospho(pY1068)EGFR, total(t) and phospho(p)-Akt, -ERK, and -CREB and actin by immunoblotting. (C) HeLa cells in B were subjected to scratch wounding and examined immediately (0 h) and after 24 h by light microscopy (Fig. S2C). The migration index (y axis) was calculated as described in *Experimental Procedures*, normalized to cells expressing GIV-WT (set at 100%) and plotted as a bar graph. (D) HeLa cells in B were analyzed for chemotaxis toward EGF using a Transwell migration assay (Fig. S2D). Cells were allowed to migrate for 6 h, fixed, and stained with Toluidine Blue. The number of migrating cells was averaged from 20 field-of-view images per experiment. Data are presented as mean \pm SEM; $n = 3$. HPF, high power field. (E) HeLa cells in B were grown in low serum [2% (vol/vol) FBS] overnight, incubated in BrdU for 30 min, fixed [3% (wt/vol) paraformaldehyde], stained for BrdU, and analyzed by confocal microscopy (Fig. S2E). Bar graph shows percentage of cells with BrdU uptake (y axis). Data are presented as mean \pm SEM; $n = 3$.

modified Boyden chamber apparatus (Fig. S2D) (19). GIV-74D cells showed a ~65% increase, whereas GIV-74A cells showed a ~30% decrease compared with GIV-WT cells (Fig. 2D and Fig. S2D). Noteworthy, the GIV-74A cells were ~50–60% less motile than GIV-74D cells. Consistent with the fact that GIV-74A cells enhanced mitogenic MAPK/ERK/CREB signals, the rate of proliferation of these cells, as determined by BrDU uptake assays (Fig. 2E and Fig. S2E) was also increased (~50% compared with GIV-WT cells). Conversely, cells expressing GIV-74D, which suppressed mitogenic signals, proliferated less (~40% less than WT, and ~60% less than GIV-74A). Together, these observations demonstrate that phosphorylation of GIV at S1674 not only alters receptor localization, but also alters the profile of downstream signals propagated and the resultant phenotype. Phosphorylation at S1674 and activation of GIV-GEF led to increased migratory and suppressed mitogenic signals, and cells preferentially migrated. By contrast, dephosphorylation at S1674 and inactivation of GIV-GEF led to increased mitogenic and suppressed migratory signals, and cells preferentially proliferated.

Next we asked how phosphomodification of GIV at S1674 affects epithelial morphogenesis, which relies on many cellular processes, including a delicate balance between migration and proliferation. To study GIV's effect on epithelial morphogenesis, we generated Madin–Darby canine kidney (MDCK) cell lines stably expressing GIV-WT or 74A/74D mutants (Fig. S3A) and used them in 3D cultures of MDCK epithelial cysts, which mimic epithelial organization in vivo. Consistent with the need for directional cell migration during epithelial morphogenesis (20, 21), GIV-WT cells and the promigratory GIV-74D cells, but not the proproliferative GIV-74A cells formed mature cysts with a central lumen (Fig. S3B). In the case of GIV-74A cells, cells grew as solid masses, without a central lumen. Together, these results demonstrate that phosphorylation of GIV-GEF at S1674, which is a key determinant of whether cells preferentially migrate or divide, is also required for the formation of MDCK epithelial cysts and suggest a role for this phosphoevent during epithelial morphogenesis.

Phosphorylation of GIV at S1674 Orchestrates Migration–Proliferation Dichotomy in Invasive Cancer Cells. Migration–proliferation dichotomy was first described in invasive tumors that exhibit high and spatially heterogeneous cell proliferation and motility rates (1), and the ability to switch between the two phenotypes is a hallmark of invasive cancer cells. We asked if phosphorylation of GIV at S1674 can affect differential activation of signaling pathways in highly metastatic breast cancer cell line MDA-MB231. MDA-MB231 cells stably expressing GIV-74D (Fig. S4) enhanced promigratory PI3K/Akt signals as determined by Akt phosphorylation at 5 and 15 min after EGF stimulation, whereas 74A cells enhanced and sustained promotive ERK and CREB signals (as determined by phosphorylation of ERK1/2 and CREB at 15 and 30 min) (Fig. 3A). Consistent with their opposing signaling programs, GIV-74D cells preferentially migrated, as determined by Transwell chemotaxis assays (Fig. 3B), but failed to proliferate into colonies as determined by anchorage-dependent cell growth assays (Fig. 3C). By contrast, GIV-74A cells were less motile (Fig. 3B) and instead, preferentially grew into colonies (Fig. 3C). These results demonstrate that reversible phosphorylation of GIV at S1674, which directly impacts GIV's ability to bind $\text{G}\alpha\text{i/s}$, is sufficient to impart migration–proliferation dichotomy in invasive cancer cells.

CDK5 Binds and Phosphorylates GIV at S1674. To identify the kinase responsible for phosphorylating GIV at S1674, we analyzed the sequence flanking it for known consensus sites of kinases. Because the sequence resembled the consensus for multiple kinases, i.e., PKA/B/C (K/RXXS/T consensus), MAPK/ERK and CDKs (SP/TP consensus) (22), we screened multiple recombinant kinases (PKA, PKB, all PKC isoforms, CDK1-6, ERK/MAPK, Braf




Fig. 3. Phosphorylation of GIV at S1674 influences EGF signaling and influences preference for migration versus proliferation of MDA-MB231 cancer cells. (A) Serum-starved MDA-MB231 cells stably expressing GIV-WT or 74D/A mutants were stimulated with EGF before lysis. Whole-cell lysates were analyzed for total(t) and phospho(p) Akt, ERK, CREB, and actin by immunoblotting. (B) MDA-MB231 cells used in A were subjected to chemotaxis toward EGF, and migrating cells were visualized and analyzed as in Fig. 2D. Results are expressed as mean \pm SEM; $n = 3$. HPF, high power field. (Magnification: 10 \times .) (C) MDA-MB231 cells used in A were analyzed for their ability to form adherent colonies on plastic plates for 2–3 wk before fixation and staining with crystal violet. (Left) Photograph of the crystal violet-stained wells. (Right) Bar graphs showing the number of colonies/cell line counted by ImageJ (colony counter).

kinase) in in vitro kinase assays for their ability to phosphorylate His-GIV-CT at S1674. We found that among all of the kinases tested, CDK5 emerged as the only kinase that could specifically phosphorylate GIV-CT-WT, but not 74A, indicating that S1674 is a substrate site for CDK5 in vitro (Fig. 4A).

Because CDK5 is known to be activated early (within seconds) after EGF stimulation (23) and confirmed by us to be the case (Fig. S5A), we hypothesized that this kinase should be able to phosphorylate S1674 on GIV and activate its GEF function for subsequent coupling of G α i proteins to RTKs and trigger G-protein activation at the PM by 5 min (14). To determine if CDK5 indeed phosphorylates GIV-GEF in cells and modulates its ability to bind G α i/s, we generated a truncated form of GST-GIV-CT (1660–1736), which contains only one “SP” motif (S¹⁶⁷⁴P¹⁶⁷⁵) as the potential site for CDK5 and retains its ability to interact with G α i3 (Fig. S5B). Moreover, when the S1674A mutation was introduced into this construct, it reduced the binding between GIV and G α i3 (Fig. S5B). When expressed in Cos7 cells, immunoblots revealed that GIV-CT_{1660–1736} appears as two differentially migrating bands—one faster major band and one slower minor band (Fig. 4B, lane 1). Because phosphorylation of

proteins tends to retard their electrophoretic mobility, we hypothesized that the slower migrating band may represent phosphorylated GIV-CT. We found such is indeed the case because the band disappeared when the samples were treated with Calf intestinal alkaline phosphatase (CIP) (Fig. 4B, lane 2) but persisted when a similar CIP treatment was carried out in the presence of its inhibitor, EDTA (Fig. 4B, lane 3). The phosphorylated form also decreased when cells expressing GST-GIV-CT_{1660–1736} were either treated with Roscovitine, a CDK5-specific inhibitor (Fig. 4B, lanes 4 and 5) or depleted of CDK5 by siRNA (Fig. 4B, lanes 6 and 7). Finally, the relative abundance of the phosphorylated band was enhanced when GST-GIV-CT_{1660–1736} was coexpressed with WT CDK5, but not a kinase inactive/dominant negative mutant of CDK5(D144N) (24) (Fig. 4B, lanes 8 and 9). Although these data confirmed that CDK5 phosphorylates GIV at S1674 *in vivo*, we noted that the extent of phosphorylation




Fig. 4. CDK5 phosphorylates GIV at S1674. (A) *In vitro* kinase assays were carried out on bacterially expressed and purified His-GIV-CT (WT) or non-phosphorylatable 74A mutant with recombinant CDK5 kinase in the presence of [³²P]ATP followed by autoradiography (Upper). Equal loading of active kinase and His-GIV-CT substrate was analyzed by immunoblotting (Middle) and Ponceau-S staining (Lower), respectively. (B) GST-GIV-CT 1660–1736 expressed in Cos7 cells was subjected to different treatments as indicated and analyzed by immunoblotting using anti-GST antibody. The faster migrating form of the doublet represents the unmodified form, and the slower migrating form represents the phosphorylated form of GIV-CT 1660–1736 as determined by treatment of bead-bound GIV-CT with calf intestinal phosphatase (CIP) (lanes 1–3). CDK5 inactivation using Roscovitine (lanes 4 and 5), siRNA mediated depletion (lanes 6 and 7), and expression of a dominant negative (DN) form of CDK5 (lanes 8 and 9) reduced phosphorylation of GIV-CT 1660–1736. (C) Control or CDK5-depleted HeLa cells were fixed and analyzed for interactions between GIV and G α i3 (Upper Left) or GIV and G α s (Lower Left) by *in situ* PLA (red). Nuclei, DAPI (blue). Whole cell lysates of HeLa cells used for PLA assays were analyzed for efficiency of CDK5 depletion by immunoblotting (Upper Right). PLA dots were quantified per cell from a total of 25–30 cells per experiment (Lower Right). Results are expressed as mean \pm SEM; $n = 3$. (D and E) Serum-starved control (si Scr) or CDK5-depleted (si CDK5) HeLa cells were stimulated with 50 nM EGF, fixed, and stained for total EGFR (green; D) or pY1068-EGFR (green; E), EEA1 (red), and DAPI (nucleus; blue), and analyzed by confocal microscopy as in Fig. 2A. Yellow pixels, indicative of total (D) and active EGFR (E) in endosomes were increased \sim 2.8-fold in D and \sim 8.3-fold in E, respectively, as determined using ImageJ. (Scale bar, 10 μ m).

was modest. The substoichiometric phosphorylation we observe in cells likely represents an underestimation due to overexpression related artifact (saturated system) or due to the synthetic nature of a suboptimal substrate that does not resemble endogenous protein. It is also possible that kinases other than CDK5 may phosphorylate GIV at that site. Regardless, these results strongly suggest that CDK5 is able to phosphorylate GIV at S1674 in cells. CDK5 also coimmunoprecipitated with full-length GIV (Fig. S5C), indicating that CDK5 binds its substrate GIV.

To determine if CDK5-dependent phosphorylation of GIV-GEF translates to increased binding of GIV to G α i/s, we analyzed the abundance of GIV-G α i and GIV-G α s complexes in control vs. CDK5-depleted cells by the *in situ* proximity ligation assay (PLA). Depletion of CDK5 (~95% efficacy; Fig. 4C, Right) significantly reduced the interaction of endogenous GIV with G α i3

(Fig. 4C, Top Left) and G α s (Lower Left). These data confirm that phosphorylation of GIV-GEF by CDK5 at S1674 is required for enhanced interactions between GIV and G α i/s subunits.

Because phosphorylation of GIV at S1674 plays a role in trafficking of EGFR through early endosome compartments, we hypothesized that CDK5 may be required for efficient trafficking of EGFR. Depletion of CDK5 by siRNA resulted in a phenotype identical to that observed in cells expressing the non-phosphorylatable GIV-74A mutant, i.e., ligand-activated EGFR stayed longer in EEA1-positive endosomes (at 30 min), indicating a delay in transit (Fig. 4D and E). Taken together, these data suggest that phosphorylation of GIV at S1674 by CDK5, which enhanced GIV-G α s interaction, is required for efficient EGFR trafficking through the endosomal system.




Fig. 5. Cell migration after scratch-wounding and cancer cell chemotaxis were inhibited by Roscovitine in cells that express GIV-WT, but not in those expressing the phosphomimic 74D GIV mutant. (A) GIV-depleted HeLa cell lines stably expressing vector control, GIV-WT, and GIV-74D were treated or not with EGF in the presence or absence of Roscovitine and analyzed for cell migration after scratch wounding as in Fig. 2C. Bar graph shows the migration index (wound closure) of each cell line under each condition. Results are expressed as mean \pm SEM; $n = 3$. (B) MDA-MB231 cells were evaluated for chemotaxis toward EGF as in Fig. 3B in the presence (+) or absence (-) of Roscovitine. The number of migrating cells was averaged from 20 field-of-view images per experiment. Results are expressed as mean \pm SEM; $n = 3$; n.s., not significant. (C) Schematic of a working model. (Left) EGF stimulation activates CDK5, which turns GIV's GEF activity "on" by phosphorylating it at S1674. This phosphoevent enhances GIV's ability to bind and activate G α i at the PM, which in turn prolongs mitogenic EGF signaling at the PM, likely via a delay in endocytosis. Such phosphorylation also enhances GIV's ability to bind G α s (and perhaps modulate its activation; "?") on endosomes, which speeds up endosomal maturation and suppresses mitogenic EGF signaling from endosomes. Overall, phosphorylation of GIV by CDK5 at S1674 triggers preferential cell migration over proliferation. (Right) Inhibition or depletion of CDK5 maintains GIV's GEF function in the "off" state, reducing GIV's ability to bind and modulate G α i/s at both locations. Consequently, EGF triggers an opposite response, such that inhibition of CDK5 favors proliferation over cell migration.

CDK5 Triggers Cell Migration via Phosphoactivation of GIV-GEF. Prior studies have implicated CDK5 in directional cell migration via phosphoregulation of multiple substrates (25–28). We asked if phosphorylation and activation of its newly identified substrate GIV-GEF contributes to the promigratory role of CDK5, and if so, to what extent. To answer this question, we analyzed the effect of the CDK5 inhibitor Roscovitine on cells expressing GIV-WT or GIV-74D. If GIV is a major substrate and a promigratory effector of CDK5, we predicted that inhibition of CDK5 with Roscovitine will inhibit phosphorylation of GIV at S1674 and thereby inhibit migration of GIV-WT cells. However, similar treatment with Roscovitine should spare the promigratory, constitutively activated GIV-74D cells, which essentially “bypass” the kinase inhibition (schematically depicted in Fig. S6A). If GIV is not a major substrate/effector of CDK5, Roscovitine should inhibit cell migration in both cell lines. We found that Roscovitine inhibited wound healing in control and GIV-WT HeLa cells (Fig. 5A), but did not have any effect on GIV-74D cells. Similarly, Roscovitine impaired the migration of MDA-MB231 cells expressing GIV-WT in Transwell chemotaxis assays, but had no significant effect on GIV-74D cells (Fig. 5B and Fig. S6B). These results indicate that phosphorylation of GIV is a key mechanism by which CDK5 triggers cell migration.

Discussion

Summary and Working Model. The major finding in this work is the identification of CDK5 as a kinase that phosphoactivates GIV-GEF and initiates noncanonical G protein signaling downstream of growth factor RTKs. Whether or not such activation occurs determines if cells migrate or proliferate. Based on our findings here and the literature, we propose the following model (Fig. 5C, *Left*): EGF stimulation activates CDK5 within seconds (23), which in turn phosphorylates GIV-GEF at S1674 and turns on its ability to bind G α i/s and activate G α i. Binding and activating G α i prolongs the duration that EGFR spends on the cell surface and enhances PM-based promigratory PI3K-Akt signals (8). Binding G α s, on the other hand, shortens the duration EGFR spends in endosomes and diminishes endosome-based promitotic MAPK/ERK signals (15). Consequently, when CDK5 phosphoactivates GIV-GEF and enhances its ability to bind both G α -subunits, cells preferentially migrate while suppressing proliferation. Inhibition or depletion of CDK5 or targeted inhibition of this phosphoevent (Fig. 5C, *Right*) renders GIV-GEF unresponsive to growth factors and maintains it in an off state: each of the above responses are reversed, i.e., GIV does not bind either G α i/s, EGFR stays shorter at the PM and longer in endosomes, PI3K-Akt signals are suppressed, MAPK/ERK signals are enhanced, and cells preferentially divide.

Dynamic Phosphorylation of GIV at Ser1674 Provides a Physiologic Mechanism for Modulating GIV's GEF Activity. Previously, using a battery of synthetic mutants generated on the basis of a structure-guided approach we showed that the potency of GIV's GEF activity has an *in vivo* threshold effect on cellular response (29), that its presence triggers cell migration, and its absence triggers mitosis (8). Together these results provided a potential explanation of how growth factor signals undergo bifurcation at the immediate postreceptor level to choose between two distinct fates (go or grow) in response to a single stimulus and how their activation levels may be modulated. However, both studies used an approach involving artificial mutations to irreversibly modulate the GEF function of GIV. Here we show that phosphorylation of GIV at S1674 is a physiologic mechanism by which GIV's GEF function overcomes the previously described *in vivo* threshold effect because phosphorylation was essential to trigger cell migration. We also show that dynamic phospho-dephosphorylation events at a single site can enable or disable, respectively, GIV's GEF motif and dictate whether cells migrate or divide.

Although the kinase-substrate relationship between CDK5 and GIV is specifically responsible for triggering the phosphoevent, the identity of the antagonistic phosphatase remains unknown.

Phosphorylation of GIV at Ser1674 Modulates EGFR Signaling and Trafficking via Two G α Subunits at Two Distinct Locations. We previously showed that GIV's ability to bind and activate G α i at the PM and bind G α s on endosomes are key determinants of when and where the activated receptor is compartmentalized, for how long, and which downstream signals are propagated and which are attenuated (8, 15, 30). Here we show that the phosphorylation status of S1674 enhances GIV's ability to bind both G α i and G α s. Based on our prior work and the findings of this study, the consequences of enhanced GIV-G α i interaction triggered by CDK5 are expected to increase the assembly of EGFR-GIV-G protein complexes and transactivation of G α i in the vicinity of ligand-activated EGFR (as shown recently by live cell FRET studies) (14), increase PM-association of EGFR, enhance PM-based PI3K-Akt signals, and enhance cell migration. Therefore, CDK5-dependent phosphorylation of GIV-GEF may influence both G protein and growth factor signaling.

In the case of G α s, we previously showed that the GIV-G α s interaction is critical for endosomal maturation and clearance of ligand-activated receptors (15). In order for the GIV-G α s complex to promote rapid clearance of EGFR out of early endosomes and impose finiteness to promitotic signaling from that compartment, G α s must remain inactive (15). Here we show that GIV's GEF motif has a weak inhibitory effect on nucleotide exchange by G α s *in vitro*. Whether this weak effect is responsible for keeping G α s inactive on early endosomes and whether additional post-translational modifications on either GIV or G α s further enhance the inhibitory effect remain unclear. What is clear are the consequences of enhanced GIV-G α s interaction triggered by CDK5 on EGFR signaling and trafficking: the duration of EGFR on early endosomes is shortened and the endosome-based promitotic MAPK/ERK1/2/CREB signals are suppressed. When CDK5 cannot phosphorylate GIV, and GIV cannot bind G α s, EGFR stays longer in early endosomes and endosome-based promitotic signals are enhanced. Together, the GIV-G α i and GIV-G α s interactions promoted by CDK5 coordinately enhance promigratory signals from the PM while dampening the mitogenic signals from endosomes with the net outcome being persistent cell migration. Despite these insights, how GIV may interact with two G α subunits (that have opposing effects on adenylyl cyclase) at two subcellular locations and whether such interactions are simultaneous or sequential remain unknown.

The Functional Interplay Between GIV and CDK5 Unravel the Molecular Mechanisms for Their Previously Known Functions. Since its discovery two decades ago, CDK5, the atypical member of the CDK family, has emerged as a trigger for cell migration in diverse cell types during angiogenesis, neurite outgrowth, epithelial wound healing, cancer cell invasion, etc., and is known to simultaneously arrest cell cycle progression via unclear mechanisms (31). Consistent with our finding that phosphorylation of GIV by CDK5 enhances Akt phosphorylation and cell migration, others have shown that *Cdk5-null* mice die prenatally due to massive failure of neuronal cell migration, and the embryonic extracts show reduced levels of phospho-Akt (32–34). Like GIV, which is a bona fide metastasis-related protein (35), CDK5 has also been found to be overexpressed in malignant human cancers (prostate, pancreatic, and breast) where it has been implicated in promoting metastasis (36–40). The enhanced migration of MDA-MB231 cells containing a constitutively phosphorylated GIV mutant (S1674D) suggests that the degree of phosphorylation of GIV at S1674 may provide an additive value as a biomarker (besides the use of GIV-positivity score) (41) to accurately prognosticate outcome in patients who have a GIV-positive tumor.

Furthermore, inhibition of the CDK5-GIV axis may be a viable/promising strategy to curb the aggressive behavior of highly metastatic cancers.

Although CDK5 is known to trigger cell migration via phosphorylation of several substrates (31), how this kinase suppresses the cell cycle is poorly understood. Some reports have indicated that CDK5 inhibits cell cycle entry via suppression of MAPK/ERK signaling (42, 43); however, little or nothing is known about how this might occur. We showed that CDK5 suppresses mitogenic MAPK/ERK1/2/CREB signals in part by phosphorylating GIV's GEF motif and enhancing GIV-G α s interaction, which speeds up endosomal maturation and thereby limits the duration of EGFR signals from endosomes.

This work also illuminates a few spatiotemporal aspects of G protein activation by GIV. Using FRET studies in living cells, we have recently shown (14) that within 3–5 min after EGF stimulation, GIV gets recruited to ligand-activated EGFR. Within 5 min EGFR–GIV–G α i ternary complexes are assembled at the PM, which is followed by activation of Gi by \sim 5 min and suppression of cAMP by \sim 6 min. Because CDK5 is activated within seconds after EGF stimulation (23), it is likely that once activated, CDK5 can promptly phosphorylate GIV at S1674 before or during the latter's recruitment to the activated receptor at the PM, ensuring maximal coupling to and activation of G α i within 5 min after ligand stimulation. Because GIV-GEF has also been found to be functional on two types of intracellular membranes, e.g., triggers secretion from the Golgi (44) and inhibits the formation/maturation of autophagosomes (45), and because activation of CDK5 exerts similar effects on both processes (46, 47), it is tempting to speculate that activation of GIV-GEF on internal membranes is also via CDK5. Further studies are required to determine if such is the case.

In conclusion, we provide insights into a hitherto unrecognized crosstalk between CDK5 and G α i/s proteins via GIV and how such crosstalk may shape migration–proliferation dichotomy during several physiological and pathological scenarios such as development, wound healing, and cancer metastasis.

Experimental Procedures

Detailed methods are described in *SI Experimental Procedures*.

Cell Culture, Transfection, Immunoblotting, Immunofluorescence, and Protein-Protein Interaction Assays (GST Pulldowns and Immunoprecipitations). These assays were carried out exactly as described before (7, 8, 15). All transfections

were performed using *TransIT-LT1* reagent (Mirus). All Western blotting (Odyssey-LICOR) images were processed and assembled for presentation using Image Studio Lite, Photoshop, and Illustrator software (Adobe).

Scratch Wounding, BrdU Incorporation, and Steady-State GTPase Activity Assays. These assays were performed as described previously (8, 9, 15).

PLA. In situ interactions of endogenous GIV with G α i3 or G α s were detected using a proximity ligation assay kit Duolink (Olink Biosciences) as per the manufacturer's instructions.

Anchorage-Dependent Colony Formation Assay. Anchorage-dependent growth was monitored on solid (plastic) surface as described previously (48). Briefly, 1,000 MDA-MB231 cells stably expressing WT, 74D, and 74A GIV-FLAG constructs were grown in six-well tissue culture plates at 37 °C for 2 wk in medium supplemented with 0.2% FBS before staining with 0.005% crystal violet for 1 h. Images were acquired by light microscopy.

MDCK Cyst Formation Assay. This assay was performed as previously described (49). Briefly, 3×10^4 MDCK type II cells stably expressing GIV constructs were added to a collagen solution at pH 7 and placed in a well of a four-well chamber slide. After the collagen was polymerized, culture media [DMEM 1 \times , FBS 2% (vol/vol)] was added, and the plate was incubated at 37 °C in a CO₂ incubator. The medium was changed every 2 d for 2 wk. Cyst formation was monitored by phase-contrast microscopy.

Statistical Analysis. Each experiment presented in the figures is representative of at least three independent experiments. Statistical significance between the differences of means was calculated by unpaired Student's *t* test. A two-tailed *P* value of <0.05 at 95% confidence interval is considered as statistically significant. **P* < 0.05; ***P* < 0.01; ****P* < 0.001; *****P* < 0.0001. All graphical data presented were prepared using GraphPad or Matlab software.

ACKNOWLEDGMENTS. We thank Karen Sykes for technical assistance. This work was supported by NIH grants CA100768 (to M.G.F.), and CA160911 and DK099226 (to P.G.). D.B. was supported by the California State University Program for Education and Research in Biotechnology (CSUPERB) New Investigator Grant GF00631143. P.G. was also supported by the Burroughs Wellcome Fund [Career Awards for Medical Scientists (CAMS) award], the American Cancer Society (ACS-IRG 70-002), and by the University of California at San Diego Moores Cancer Center. M.G.-M. was supported by the American Cancer Society (RGS-13-362-01-TBE) and NIH (R01GM108733). I.L.-S. was supported by a fellowship from the American Heart Association (AHA 14POST20050025) and I.-C.L. by a fellowship (NSC 100-2917-1-564-032) from the National Science Council of Taiwan.

- Fedotov S, Iomin A (2007) Migration and proliferation dichotomy in tumor-cell invasion. *Phys Rev Lett* 98(11):118101–118104.
- Gerhardt H, et al. (2003) VEGF guides angiogenic sprouting utilizing endothelial tip cell filopodia. *J Cell Biol* 161(6):1163–1177.
- Gaylarde PM, Sarkany I (1975) Cell migration and DNA synthesis in organ culture of human skin. *Br J Dermatol* 92(4):375–380.
- Bonneton C, Sibarita JB, Thiery JP (1999) Relationship between cell migration and cell cycle during the initiation of epithelial to fibroblastoid transition. *Cell Motil Cytoskeleton* 43(4):288–295.
- Chung EH, Hutcheon AE, Joyce NC, Zieske JD (1999) Synchronization of the G1/S transition in response to corneal debridement. *Invest Ophthalmol Vis Sci* 40(9):1952–1958.
- Enomoto A, et al. (2005) Akt/PKB regulates actin organization and cell motility via Girdin/APE. *Dev Cell* 9(3):389–402.
- Ghosh P, Garcia-Marcos M, Bornheimer SJ, Farquhar MG (2008) Activation of Galphai3 triggers cell migration via regulation of GIV. *J Cell Biol* 182(2):381–393.
- Ghosh P, et al. (2010) A G α i-GIV molecular complex binds epidermal growth factor receptor and determines whether cells migrate or proliferate. *Mol Biol Cell* 21(13):2338–2354.
- Garcia-Marcos M, Ghosh P, Farquhar MG (2009) GIV is a nonreceptor GEF for G alpha i with a unique motif that regulates Akt signaling. *Proc Natl Acad Sci USA* 106(9):3178–3183.
- Jiang P, et al. (2008) An actin-binding protein Girdin regulates the motility of breast cancer cells. *Cancer Res* 68(5):1310–1318.
- Kitamura T, et al. (2008) Regulation of VEGF-mediated angiogenesis by the Akt/PKB substrate Girdin. *Nat Cell Biol* 10(3):329–337.
- Garcia-Marcos M, Ghosh P, Ear J, Farquhar MG (2010) A structural determinant that renders G α (i) sensitive to activation by GIV/girdin is required to promote cell migration. *J Biol Chem* 285(17):12765–12777.
- Lin C, et al. (2014) Structural basis for activation of trimeric Gi proteins by multiple growth factor receptors via GIV/Girdin. *Mol Biol Cell* 25(22):3654–3671.
- Midde KK, et al. (2015) Multimodular biosensors reveal a novel platform for activation of G proteins by growth factor receptors. *Proc Natl Acad Sci USA* 112(9):E937–E946.
- Beas AO, et al. (2012) Gus promotes EEA1 endosome maturation and shuts down proliferative signaling through interaction with GIV (Girdin). *Mol Biol Cell* 23(23):4623–4634.
- López-Sánchez I, et al. (2013) Protein kinase C- θ (PKC θ) phosphorylates and inhibits the guanine exchange factor, GIV/Girdin. *Proc Natl Acad Sci USA* 110(14):5510–5515.
- Haugh JM (2002) Localization of receptor-mediated signal transduction pathways: The inside story. *Mol Interv* 2(5):292–307.
- Murphy JE, Padilla BE, Hasdemir B, Cottrell GS, Bunnett NW (2009) Endosomes: A legitimate platform for the signaling train. *Proc Natl Acad Sci USA* 106(42):17615–17622.
- Richards KL, McCullough J (1984) A modified microchamber method for chemotaxis and chemokinesis. *Immunol Commun* 13(1):49–62.
- Wang Y, et al. (2011) Girdin is an intrinsic regulator of neuroblast chain migration in the rostral migratory stream of the postnatal brain. *J Neurosci* 31(22):8109–8122.
- Ohara K, et al. (2012) Involvement of Girdin in the determination of cell polarity during cell migration. *PLoS One* 7(5):e36681.
- Amanchy R, et al. (2007) A curated compendium of phosphorylation motifs. *Nat Biotechnol* 25(3):285–286.
- Lee HY, Jung H, Jang IH, Suh PG, Ryu SH (2008) Cdk5 phosphorylates PLD2 to mediate EGF-dependent insulin secretion. *Cell Signal* 20(10):1787–1794.
- Nikolic M, Dudek H, Kwon YT, Ramos YFM, Tsai L-H (1996) The cdk5/p35 kinase is essential for neurite outgrowth during neuronal differentiation. *Genes Dev* 10(7):816–825.

25. Hosoi T, et al. (1995) Evidence for cdk5 as a major activity phosphorylating tau protein in porcine brain extract. *J Biochem* 117(4):741–749.
26. Veeranna SKT, et al. (1995) Neuronal cyclin-dependent kinase-5 phosphorylation sites in neurofilament protein (NF-H) are dephosphorylated by protein phosphatase 2A. *J Neurochem* 64(6):2681–2690.
27. Beffert U, et al. (2004) Reelin and cyclin-dependent kinase 5-dependent signals cooperate in regulating neuronal migration and synaptic transmission. *J Neurosci* 24(8):1897–1906.
28. Huang C, et al. (2009) Talin phosphorylation by Cdk5 regulates Smurf1-mediated talin head ubiquitylation and cell migration. *Nat Cell Biol* 11(5):624–630.
29. Garcia-Marcos M, et al. (2012) Functional characterization of the guanine nucleotide exchange factor (GEF) motif of GIV protein reveals a threshold effect in signaling. *Proc Natl Acad Sci USA* 109(6):1961–1966.
30. Ghosh P, Garcia-Marcos M, Farquhar MG (2011) GIV/Girdin is a rheostat that fine-tunes growth factor signals during tumor progression. *Cell Adhes Migr* 5(3):237–248.
31. Dhavan R, Tsai L-H (2001) A decade of CDK5. *Nat Rev Mol Cell Biol* 2(10):749–759.
32. Gilmore EC, Ohshima T, Goffinet AM, Kulkarni AB, Herrup K (1998) Cyclin-dependent kinase 5-deficient mice demonstrate novel developmental arrest in cerebral cortex. *J Neurosci* 18(16):6370–6377.
33. Ohshima T, et al. (1996) Targeted disruption of the cyclin-dependent kinase 5 gene results in abnormal corticogenesis, neuronal pathology and perinatal death. *Proc Natl Acad Sci USA* 93(20):11173–11178.
34. Li BS, et al. (2003) Cyclin-dependent kinase-5 is involved in neuregulin-dependent activation of phosphatidylinositol 3-kinase and Akt activity mediating neuronal survival. *J Biol Chem* 278(37):35702–35709.
35. Garcia-Marcos M, Ghosh P, Farquhar MG (2015) GIV/Girdin transmits signals from multiple receptors by triggering trimeric G protein activation. *J Biol Chem* 290(11):6697–6704.
36. Strock CJ, et al. (2006) Cyclin-dependent kinase 5 activity controls cell motility and metastatic potential of prostate cancer cells. *Cancer Res* 66(15):7509–7515.
37. Hsu FN, et al. (2011) Regulation of androgen receptor and prostate cancer growth by cyclin-dependent kinase 5. *J Biol Chem* 286(38):33141–33149.
38. Feldmann G, et al. (2010) Inhibiting the cyclin-dependent kinase CDK5 blocks pancreatic cancer formation and progression through the suppression of Ras-Ral signaling. *Cancer Res* 70(11):4460–4469.
39. Eggers JP, et al. (2011) Cyclin-dependent kinase 5 is amplified and overexpressed in pancreatic cancer and activated by mutant K-Ras. *Clin Cancer Res* 17(19):6140–6150.
40. Liang Q, et al. (2013) CDK5 is essential for TGF- β 1-induced epithelial-mesenchymal transition and breast cancer progression. *Sci Rep* 3:2932.
41. Garcia-Marcos M, et al. (2011) Expression of GIV/Girdin, a metastasis-related protein, predicts patient survival in colon cancer. *FASEB J* 25(2):590–599.
42. Modi PK, Komaravelli N, Singh N, Sharma P (2012) Interplay between MEK-ERK signaling, cyclin D1, and cyclin-dependent kinase 5 regulates cell cycle reentry and apoptosis of neurons. *Mol Biol Cell* 23(18):3722–3730.
43. Zheng YL, et al. (2007) Cdk5 Modulation of mitogen-activated protein kinase signaling regulates neuronal survival. *Mol Biol Cell* 18(2):404–413.
44. Lo IC, et al. (2015) Activation of Gxi at the Golgi by GIV/Girdin imposes finiteness in Arf1 signaling. *Dev Cell* 33(2):189–203.
45. Garcia-Marcos M, Ear J, Farquhar MG, Ghosh P (2011) A GDI (AGS3) and a GEF (GIV) regulate autophagy by balancing G protein activity and growth factor signals. *Mol Biol Cell* 22(5):673–686.
46. Paglini G, Peris L, Diez-Guerra J, Quiroga S, Cáceres A (2001) The Cdk5-p35 kinase associates with the Golgi apparatus and regulates membrane traffic. *EMBO Rep* 2(12):1139–1144.
47. Furuya T, et al. (2010) Negative regulation of Vps34 by Cdk mediated phosphorylation. *Mol Cell* 38(4):500–511.
48. Franken NA, Rodermond HM, Stap J, Haveman J, van Bree C (2006) Clonogenic assay of cells in vitro. *Nat Protoc* 1(5):2315–2319.
49. Elia N, Lippincott-Schwartz J (2009) Culturing MDCK cells in three dimensions for analyzing intracellular dynamics. *Curr Protoc Cell Biol*. Chapter 4:Unit 4.22.