

# IL-15 suppresses colitis-associated colon carcinogenesis by inducing antitumor immunity

Rajia Bahri<sup>1,2</sup>, Ioannis S Pateras<sup>3</sup>, Orietta D'Orlando<sup>2</sup>, Diego A Goyeneche-Patino<sup>2</sup>, Michelle Campbell<sup>1</sup>, Julia K Polansky<sup>2</sup>, Hilary Sandig<sup>1</sup>, Marilena Papaioannou<sup>3</sup>, Kostas Evangelou<sup>3</sup>, Periklis G Foukas<sup>3,4</sup>, Vassilis G Gorgoulis<sup>3,5,6</sup>, and Silvia Bulfone-Paus<sup>1,\*</sup>

<sup>1</sup>Institute of Inflammation and Repair and MCCIR; University of Manchester; Manchester, UK; <sup>2</sup>Priority Area Asthma and Allergies; Research Center Borstel; Borstel, Germany;

<sup>3</sup>Molecular Carcinogenesis Group; Department of Histology & Embryology; School of Medicine; University of Athens; Athens, Greece; <sup>4</sup>2nd Department of Pathology; University of Athens Medical School; "Attikon" University Hospital; Athens, Greece; <sup>5</sup>Biomedical Research Foundation; Academy of Athens; Athens, Greece; <sup>6</sup>Faculty Institute for Cancer Sciences; University of Manchester; Manchester Academic Health Science Centre; Manchester, UK

**Keywords:** AOM/DSS, colitis associated cancer, cytokines, IL-15, immunity, tumor

**Abbreviations:** AOM, Azoxymethane; CAC, Colitis associated carcinoma; CRC, Colorectal cancer; DSS, Dextran sulfate sodium; EAE, Experimental autoimmune encephalomyelitis; IBD, Irritable bowel disease; IEL, Intestinal epithelial cell; IFN, Interferon; IL, Interleukin; LN, Lymph node; NK, Natural killer cell; SCRC, Spontaneous colorectal cancer; TNM, TNM classification of malignant tumors; WT, Wild type

IL-15 regulates the development, survival, and proliferation of multiple innate and adaptive immune cells and plays a dual role, inducing both tumor cell growth and antitumor immunity. However, the role of IL-15 in inflammation-induced cancer remains unclear. To explore this, we have compared the colon carcinoma burden of *Il15*<sup>-/-</sup> and *Il15ra*<sup>-/-</sup> mice with wild type (WT) mice after induction of colitis-associated colon carcinogenesis utilizing the AOM/DSS model. Compared to WT mice, *Il15*<sup>-/-</sup> but not *Il15ra*<sup>-/-</sup> mice showed reduced survival, along with higher tumor incidence, colon weight, and tumor size. This suggests that low affinity IL-15 signaling via the shared IL-2Rβ/γc decreases the risk for developing colitis-associated cancer. *CD11c-Il15* mice, in which IL-15 expression is reconstituted in *Il15*<sup>-/-</sup> mice under the control of the CD11c-promoter, showed that selective reconstitution of IL-15 in antigen-presenting cells restored the CD8<sup>+</sup> T and NK cell compartments, serum levels of IFNγ, G-CSF, IL-10, and CXCL1 and reduced tumor burden. After demonstrating IL-15 expression in human colorectal cancer (CRC) cells *in situ*, we investigated the role of this cytokine in the modulation of key colonic oncogenic pathways in the tumor. While these pathways were found to be unaltered in the absence of IL-15, tumor transcriptome analysis showed that the loss of IL-15 upregulates key inflammatory mediators associated with colon cancer progression, such as *IL-1β*, *IL-22*, *IL-23*, *Cxcl5*, and *Spp1*. These findings provide evidence that IL-15 suppresses colitis-associated colon carcinogenesis through regulation of antitumor cytotoxicity, and modulation of the inflammatory tumor microenvironment.

## Introduction

IL-15 has garnered attention as a potential therapeutic agent in cancer immunotherapy, with multiple studies attributing antitumor effects to IL-15, along with enhanced NK and CD8<sup>+</sup> T cell cytotoxicity.<sup>1</sup> IL-15 is a pleiotropic cytokine constitutively expressed by dendritic cells (DC), macrophages, fibroblasts, and epithelial cells,<sup>2</sup> which following inflammatory stimuli is upregulated in macrophages,<sup>3</sup> enterocytes and CD11c<sup>+</sup>CD11b<sup>+</sup> DCs.<sup>4</sup> IL-15 signals via a hetero-trimer composed of the IL-2 receptor β (IL-2Rβ), the common gamma chain (γc) and the unique high affinity receptor α chain (IL-15Rα),<sup>5-7</sup> with the latter presenting IL-15 in *trans* to IL-2Rβ/γc on neighboring cells.<sup>8,9</sup> IL-15- and IL-15Rα-deficient mice exhibit a common phenotype that has demonstrated an essential role of IL-15 in

both the development and homeostasis of memory CD8<sup>+</sup> T cells, natural killer cells (NK), invariant NKT (iNKT) cells and intestinal intraepithelial lymphocytes (IEL).<sup>10,11</sup>

Conversely, IL-15 has been shown to be pro-tumorigenic, through promotion of tumor growth, invasion and metastasis, while also protecting tumor cells from apoptosis.<sup>12-16</sup> Furthermore, in humans, overexpression of IL-15 promotes development of large granular lymphocytic leukemia.<sup>12,17</sup> Thus, the clinical value of targeting IL-15 in cancer immunotherapy remains uncertain and may be highly context-dependent. Of interest to this study, it is ill understood whether IL-15 is required for the development or suppression of inflammation-induced cancer.

Cytokines produced by tumor-infiltrating immune cells modify tumor growth and survival, promote angiogenesis and inhibit

\*Correspondence to: Silvia Bulfone-Paus; Email: silvia.bulfone-paus@manchester.ac.uk

Submitted: 10/14/2014; Revised: 12/18/2014; Accepted: 12/20/2014

<http://dx.doi.org/10.1080/2162402X.2014.1002721>

antitumor immune responses to promote the development of pre-malignant cells. The balance between pro- and anti-inflammatory activities is critical for an efficient antitumor immune response,<sup>18</sup> with dysregulation of IL-6, IL-17, IL-23, and IL-22 cytokine signaling pathways playing an important inflammatory role in the development and progression of intestinal tumors.<sup>19</sup> Furthermore, genetic variations within pro- and anti-inflammatory interleukins may confer risk of developing or impact survival rates in colon cancer.<sup>20,21</sup> Among these, single nucleotide polymorphisms (SNPs) from IL-3, IL-6R, IL-8 and, interestingly, IL-15 were found to be associated with an increased colon cancer risk,<sup>22</sup> while IL-1 or IL-1 receptor antagonist polymorphisms negatively impact on CRC survival and recurrence.<sup>23,24</sup>

The role of the inflammatory milieu is of particular importance in CRC as the risk of developing dysplasia and CRC positively correlates with the duration and degree of inflammation in patients with irritable bowel disease (IBD).<sup>25</sup> While there is abundant evidence indicating the causal relationship of chronic mucosal inflammation with CRC,<sup>26</sup> the mechanism(s) underlying this correlation, namely the role of IL-15-mediated signaling, remain largely unknown.

In light of the complex and dichotomous role of IL-15 in an oncology setting, summarized above, this current study seeks to clarify the contribution of IL-15 in a well-defined experimental model of inflammation-induced CRC. For this, a single injection of the carcinogen azoxymethane (AOM) was combined with the dextran sulfate sodium (DSS) mouse model of chronic colon inflammation.<sup>27,28</sup> This generates a relatively non-invasive and reproducible model of colitis-associated carcinoma (CAC). Utilizing this model, we have investigated the role that IL-15 plays on tumor development and growth *in vivo* using IL-15- or IL-15R $\alpha$ -deficient mice as well as a newly generated strain of *Il15*<sup>-/-</sup> mice, in which IL-15 is selectively expressed in predominantly antigen presenting cells under the control of the CD11c-promoter. In this model, we demonstrate that IL-15 deficiency increases tumor burden as a result of deficient NK and CD8<sup>+</sup> T cell immunity and a tumor-supporting inflammatory milieu, indicating that intestinal homeostasis and suppression of inflammation induced-tumorigenesis are dependent on IL-15.

## Results

### IL-15 deficiency promotes inflammation-induced colorectal tumorigenesis in mice

To determine the role of IL-15 in inflammation-induced colorectal tumorigenesis we utilized the AOM/DSS model. Briefly, mice were injected with AOM followed by three cycles of 1% DSS in drinking water.<sup>27,28</sup> Firstly, we compared the survival and colon carcinoma burden between *Il15*<sup>-/-</sup> and *Il15r $\alpha$* <sup>-/-</sup> to WT mice after induction of colitis-associated colon carcinogenesis.

IL-15- but not IL-15R $\alpha$ -deficient mice, had significantly reduced survival rates compared to WT mice (Fig. 1A), with 25% of *Il15*<sup>-/-</sup> mice dying after completion of the second DSS administration cycle. Tumor incidence was 100% in *Il15*<sup>-/-</sup> mice, and

colon weight and tumor size was significantly higher in *Il15*<sup>-/-</sup> compared to *Il15r $\alpha$* <sup>-/-</sup> and WT mice (Fig. 1B–F). Adenomas greater than 4mm in diameter were exclusively found in mice with disrupted IL-15 signaling (*Il15*<sup>-/-</sup> 34% and *Il15r $\alpha$* <sup>-/-</sup> 22%) (Fig. 1D).

Interestingly the majority of tumors in both *Il15*<sup>-/-</sup> and *Il15r $\alpha$* <sup>-/-</sup> mice were localized within the mid and distal region of the colon (Fig. 1E and F), mirroring the predominant occurrence of colorectal tumors in humans, raising the question whether tumor development in some intestinal compartments is more IL-15-dependent than others. All *Il15*<sup>-/-</sup> and a significant number of *Il15r $\alpha$* <sup>-/-</sup> mice developed tumors with low- and high-grade dysplasia (Fig. 1G). Although invasive intramucosal carcinomas were observed in all genotypes, pT1 stage adenocarcinomas were found exclusively in *Il15*<sup>-/-</sup> mice (Fig. 1G). IL-15 can reduce mucosal damage through a reduction in intestinal epithelial cell (IEC) apoptosis,<sup>29</sup> and in our AOM/DSS treated *Il15*<sup>-/-</sup> and *Il15r $\alpha$* <sup>-/-</sup> mice we found significantly increased apoptosis by cleaved caspase-3 immunostaining in both tumor free epithelium (Fig. S1A) and adenomas (Fig. S1B) compared with WT mice. Furthermore *Il-15tg* mice showed levels of apoptosis comparable with that of WT mice (Fig. S1B and S1C). We observed no significant differences in the proliferative index in adenomas between the different strains (data not shown).

Naïve *Il15*<sup>-/-</sup> and *Il15r $\alpha$* <sup>-/-</sup> mice exhibit consistently reduced numbers of NK and CD8<sup>+</sup> T cells in both spleen and mesenteric LN,<sup>10,11</sup> and this phenotype is maintained after AOM/DSS induction (Fig. 2A and 2B). The number of CD4<sup>+</sup> T cells and B220<sup>+</sup> B cells are slightly but significantly increased both in percentage (CD4<sup>+</sup> T cells 0.37%  $\pm$  0.02 vs. 0.30%  $\pm$  0.02,  $p = 0.0002$ , B220<sup>+</sup> B cells 0.438%  $\pm$  0.04 vs. 0.385%  $\pm$  0.046,  $p = 0.0188$ ) and absolute number (CD4<sup>+</sup> T cells 4.38  $\pm$  1.64 vs. 2.92  $\pm$  1,  $p = 0.0434$ , B220<sup>+</sup> B cells 5.49  $\pm$  2.22 vs. 3.41  $\pm$  0.96,  $p = 0.0314$ ) was observed in *Il15*<sup>-/-</sup> mLN compared with WT. Furthermore, compared with WT mice, the sera of *Il15*<sup>-/-</sup> mice showed higher concentrations of IFN $\gamma$ , IL-10, IL-17, CXCL1, and G-CSF with decreased levels of CCL5 and IL-12p40 (Fig. 2C). Serum levels of eotaxin, IL-1 $\alpha$ , IL-1 $\beta$ , IL-2, GM-CSF, IL-3, IL-4, IL-6, IL-12p70, IL-13, MCP1, MIP1 $\alpha$ , and MIP1 $\beta$ , were not found to be significantly different between the groups (Fig. S2).

These data demonstrate that IL-15 deficiency profoundly affects survival and accelerates tumor growth and development. In contrast, loss of IL-15R $\alpha$  showed a less pronounced effect on tumor burden, indicating that intact IL-15 signaling via IL-2R $\beta$ / $\gamma$  may partially compensate for absent IL-15R $\alpha$ -mediated signaling with respect to controlling tumor growth.

### CD11c-restricted IL-15 expression shows reduced tumor formation


IL-15 is constitutively expressed and highly inducible in DC and trans-presented to CD8<sup>+</sup> T and NK cells which induces proliferation and activation of both cell populations.<sup>8,9</sup> We therefore sought to determine, whether recovery of NK and CD8<sup>+</sup> T cell populations in *Il15*<sup>-/-</sup> mice would be sufficient to inhibit tumor growth in the AOM/DSS model. To address this, we generated a

novel *CD11c-Il15* mouse strain, in which IL-15 expression is reconstituted in antigen presenting cells of *Il15*<sup>-/-</sup> mice under the control of the CD11c-promoter (strain 65). (Polansky et al., in preparation).


As expected these mice have a recapitulated NK and CD8<sup>+</sup> T cell compartment with percentages of NK and CD8<sup>+</sup> T cells comparable with WT mice, along with essentially unchanged relative percentages of total immune cells (Fig. 2A and 2B). Moreover, the serum levels of IFN $\gamma$ , IL-10, CXCL1, and G-CSF in these mice approached that of WT levels (Fig. 2C). In particular, IL-10 and CXCL1 levels are significantly different in the sera of *Il15*<sup>-/-</sup> and *CD11c-Il15* mice.

Interestingly, *CD11c-Il15* mice showed a re-established control of tumor growth after AOM/DSS induction in comparison with the global IL-15 deficient mouse strain. Selective IL-15 expression in CD11c<sup>+</sup> cells resulted in a tumor burden comparable to that of WT mice and while tumor size was similar the macroscopically assessed tumor number was significantly decreased in comparison to *Il15*<sup>-/-</sup> mice (Fig. 3A–C). Histological analysis confirmed a decrease in the number of tumor lesions, reduced dysplasia severity and an absence of invasive carcinoma in *CD11c-Il15* mice compared with *Il15*<sup>-/-</sup> mice following AOM/DSS (Fig. 3D and 3E).


These data demonstrate that the restoration of IL-15-dependent CD8<sup>+</sup> T and NK cell immune responses reduce tumor formation, congruent with similar effects of IL-15-mediated signaling in other tumor models.<sup>1,30-32</sup> To determine if the antitumor immunity-restoring effect of IL-15 can be further enhanced if IL-15 is ubiquitously



**Figure 1.** Increased tumorigenesis in *Il15*<sup>-/-</sup> mice in a colitis-associated cancer model. Survival rates (A) of WT, *Il15*<sup>-/-</sup> and *Il15ra*<sup>-/-</sup> mice subjected to AOM/DSS treatment, were monitored. Colon weight (B), numbers of tumors per mouse (C), size (D) and location (E) of tumors were determined. Data represented as mean  $\pm$  SEM of two independent experiments (WT n = 14, *Il15*<sup>-/-</sup> n = 11, *Il15ra*<sup>-/-</sup> n = 13). (F) Representative images of longitudinally cut colons from WT (n = 2), *Il15*<sup>-/-</sup> (n = 2) and *Il15ra*<sup>-/-</sup> (n = 2). (G) Representative H&E images of formalin-fixed tissues from proximal, mid and distal colonic regions. Dotted lines indicate adenomas with high-grade dysplasia. Insert enlarges a cancerous gland invading muscularis mucosa, with the overlying epithelium which is dysplastic. Scale bar: 600  $\mu$ m, within the insert 200  $\mu$ m. Mann-Whitney tests were used for statistical analysis.



**Figure 2.** *CD11c-Il15* mice showed restored CD8<sup>+</sup> T and NK cells and serum levels of IFN $\gamma$ , G-CSF, IL-10 and CXCL1. Mesenteric lymph nodes (mLN) and spleens were isolated from mice after completion of the AOM/DSS protocol. Total cell numbers and percentage of CD8<sup>+</sup> T cells and NK cells were analyzed by flow cytometry in mLNs (A) and spleens (B). Lines show mean values with each point representing an individual mouse from three independent experiments (left panels) (WT  $n = 14$ , *Il15*<sup>-/-</sup>  $n = 14$ , *Il15ra*<sup>-/-</sup>  $n = 14$ , *CD11c-Il15*  $n = 15$ ) and two experiments (all other panels) (WT  $n = 9$ , *Il15*<sup>-/-</sup>  $n = 9$ , *Il15ra*<sup>-/-</sup>  $n = 7$ , *CD11c-Il15*  $n = 5-8$ ). Mann-Whitney tests were used for statistical analysis. (C) Sera were collected from mice after completion of the AOM/DSS protocol and cytokine/chemokines were quantified using the Bio-Plex array system (Bio-Rad). Lines indicate mean values with each point representing an individual mouse from two independent experiments (WT  $n = 22$ , *Il15*<sup>-/-</sup>  $n = 24$ , *Il15ra*<sup>-/-</sup>  $n = 20$ , *CD11c-Il15*  $n = 18$ ). Mann-Whitney tests were used for statistical analysis.



**Figure 3.** Restored IL-15 expression reduces tumor formation. Colons were isolated from mice following completion of the AOM/DSS protocol. Numbers of tumors per mouse (**A**) and size (**B**) were determined. In **A**, each point represents an individual mouse and in **B** data represent mean  $\pm$  SEM of three independent experiments (WT  $n = 31$ , *Il15*<sup>-/-</sup>  $n = 32$ , *CD11c-Il15*  $n = 28$ , *Il15tg*  $n = 11$ ). Mann-Whitney test was used for statistical analysis. (**C**) Representative images of colons from WT ( $n = 2$ ), *Il15*<sup>-/-</sup> ( $n = 2$ ) and *CD11c-Il15* ( $n = 2$ ). (**D**) Representative H&E images of formalin-fixed tissues from proximal, mid and distal colon. Dotted line indicates the muscularis mucosa with the overlying colon adenoma showing high-grade dysplasia. Dashed lines enclose colon adenomas with high-grade dysplasia. Scale bar: 600  $\mu$ m. (**E**) Percentages of tumor-types (low-grade dysplasia, high-grade dysplasia, intramucosal carcinoma, adenocarcinoma) in *Il15*<sup>-/-</sup> vs. *CD11c-Il15* mice (*Il15*<sup>-/-</sup>  $n = 26$ , *CD11c-Il15*  $n = 12$ ).

overexpressed, AOM/DSS was induced in *Il15tg* mice.<sup>33</sup> These mice had a significantly reduced tumor load compared to both *CD11c-Il15*, and WT mice (Fig. 3A). Thus, as expected, antitumor immunity is maximal if IL-15 is expressed, secreted, and/or trans-presented by both immune and non-immune cells. However, IL-15 overexpression restricted to antigen presenting cells, e.g., in DC therapy strategies, in a physiological situation where other cells are present that constitutively express IL-15, may well suffice to induce therapeutic antitumor immunity.

#### IL-15 is expressed in human colorectal cancer

In addition to its antitumor effect, IL-15 has also been reported to show tumor growth-enhancing properties<sup>16</sup> and is also expressed by epithelial cell populations.<sup>14,34</sup> We therefore sought to determine if IL-15 expression was altered in a pathophysiological setting. IL-15 expression was profiled in a large cohort of patients with CRC, including 10 cases of CAC and tissue arrays encompassing 74 cases of spontaneous colorectal carcinoma (SCRC).

In almost all of the examined CRC patients, CAC and SCRC cells exhibited IL-15 immunoreactivity with varying extent and intensity (Fig. 4 and Table S1 and S2). IL-15 localization was mainly membranous/cytoplasmic (Fig. 4B) with little IL-15 nuclear staining (Fig. 4C and 4D) the latter having been reported previously for macrophages.<sup>35</sup> However, no significant correlation was found between the semi-quantitative IL-15 protein expression score and the TNM stage (Table S1). This shows that while IL-15 protein is prominently expressed in human CRC, its relationship with cytokine expression and tumor characteristics remains unclear.

#### Key colonic oncogenic pathways do not appear to be altered by IL-15

As a first step toward dissecting the potential dual role of IL-15 in tumor immunology, we asked whether IL-15 impacts on the expression of key colonic oncogenic pathways. To this purpose, we analyzed the protein expression of key molecules known to associate with murine colon tumorigenesis,<sup>36</sup> namely  $\beta$ -catenin, p53 and p21WAF1 in colon tumors from WT, *Il15*<sup>-/-</sup>, *Il15 $\alpha$* <sup>-/-</sup>, and *CD11c-Il15* mice following AOM/DSS induction.

All colon adenomas of AOM/DSS-treated mice exhibited cytoplasmic/nuclear  $\beta$ -catenin staining, irrespective of their genotype and IL-15 expression status (Fig. S3A). In all genotypes, p53 immunostaining was limited and faint, accompanied by strong p21WAF1 staining within colon adenomas, again without appreciable differences between the various mice strains examined (Fig. S3B). Collectively, these *ex vivo*-results from established colon tumors in IL-15 deletion mouse strains suggest that IL-15 does not induce classical pathways implicated in murine colon tumorigenesis.

#### Tumor transcriptional profile analysis correlate IL-15 loss with an increase in pro-inflammatory cytokines associated with colon cancer progression

To further determine the role of IL-15 in colon cancer, we looked at its ability to modulate the inflammatory tumor microenvironment. We utilized microarray analysis of tumor,

perilesional colon, and non-tumor-affected colon tissue from IL-15 disrupted and WT AOM/DSS mice. Principal component analysis (PCA) based on 2D scaling divided samples into three different groups of comparable genes expression data space: group 1 included tumor tissue from *Il15*<sup>-/-</sup> mice, group 2 included near-tumor and tumor tissue from WT and group 3 included tumor-free tissue from WT and *Il15*<sup>-/-</sup> mice and the *Il15*<sup>-/-</sup> near-tumor tissue (Fig. S4A and S4B).


Interestingly, while tumor-free and perilesional colon showed no major gene expression differences between WT and *Il15*<sup>-/-</sup> mice, 1856 genes were dysregulated in *Il15*<sup>-/-</sup> compared with WT tumors: 201 genes with fold change of  $\geq 2$ , of which 106 were upregulated and 95 downregulated (Fig. 5A and Fig. S4C and Table S3). These results suggest that IL-15 loss exerts a major transcriptional impact on the tumor, on tumor-infiltrating cells, or both.

The upregulated genes in *Il15*<sup>-/-</sup> tumors were grouped into functionally different clusters: hydrolases/proteases (e.g., mast cell proteases and matrix metalloproteases), cytokines and cytokine/cytokine receptor interactions (e.g., IL-1 $\alpha$ , IL-1 $\beta$ , chemokine CXCL14, CXCL5, and SPP1), signaling pathways implicated in cell proliferation/growth/Wnt signaling, apoptosis, oxidation/reduction reactions, and transcription factors and membrane components (Fig. S4C and Table S4). Within the top-50 upregulated genes, the majority were found in the hydrolases/proteases cluster, followed by the cytokine/cytokine receptors interaction gene cluster. Since the upregulation of hydrolases/proteases could be a secondary effect associated with tissue remodeling induced by the increased tumor growth in *Il15*<sup>-/-</sup> mice, we focused on the upregulation of cytokine/chemokine expression as a potentially direct IL-15-dependent phenomenon.


To confirm the microarray results, the expression levels of selected cytokine and chemokine genes that were exclusively upregulated in tumor tissue were quantified by qRT-PCR. This confirmed that not only *Spp1* but also *Il1b* and *Cxcl5* expression were strongly and significantly upregulated in *Il15*<sup>-/-</sup> tumors compared to WT tumors and non-tumor bearing, AOM/DSS-treated colon (Fig. 5C and 5D). In addition, mRNA expression of the TH17 cytokines, *Il22* and *Il23* in lesional tissue, albeit transcribed at low levels, was significantly upregulated in *Il15*<sup>-/-</sup> compared to WT tumors (Fig. 5C and 5D). These gene expression profiling data from IL-15-deficient mice suggest that IL-15 impacts on the expression of key inflammatory mediators associated with colon cancer progression and that IL-15 deficiency facilitates the creation of a tumor growth-enhancing cytokine/chemokine tumor microenvironment.

## Discussion

Collectively, the current study contributes to dissecting the as yet unclear, dual role of IL-15 as a potential regulator of both pro- and antitumor immunity. We have shown that the deletion of IL-15 has a dual effect in colitis-associated murine colon carcinoma: while IL-15 deletion impairs NK/CD8<sup>+</sup> T cell-mediated



**Figure 4.** IL-15 is expressed in human colorectal cancer. (A and B) IL-15 immunostaining in CAC cases. In A, faint cytoplasmic IL-15 immunostaining in CAC4 (IL-15 Score:1). In B, arrows denote cancerous glands with intense membranous/cytoplasmic immunostaining. (CAC  $n = 10$ ). (C and D) Representative images of CAC expressing intense cytoplasmic/membranous IL-15 immunostaining (C, CAC6, Score:9), SCRC carcinoma exhibiting faint IL-15 immunostaining (D, SColon48, Score:2), Case SColon6 (E), exhibiting intense IL-15 immunostaining (low IL-15 score; score:9) and case SColon16 (F) displaying faint IL-15 immunostaining (high IL-15 score; score:2). (CAC  $n = 10$  and SColon  $n = 74$ ). Arrows indicate cancerous glands with intense membranous/cytoplasmic and double arrows depict nuclear IL-15 immunostaining. Scale bar: 50  $\mu$ m.



**Figure 5.** Differentially-expressed genes profiles in WT and *Il15*<sup>-/-</sup> colon tissues. RNA from no tumor, near tumor and tumor colon tissue from WT and *Il15*<sup>-/-</sup> mice were used for gene expression analysis by microarrays. **(A)** LIMMA analysis of differentially expressed genes showing a  $q$  value  $< 0.05$  and a  $\log_2$  fold change (FC)  $\geq 2$ . Pie charts show the number of significant genes (yellow) and the number significant genes that exhibit a fold change  $\geq 2$  (green). The green gene pool was used to determine the proportion of upregulated (red) and downregulated (blue) genes. **(B)** Fold change of top 50 upregulated and downregulated genes. Comparative heat map shows the fold change of the gene expression between WT and *Il15*<sup>-/-</sup> mice from no tumor, near tumor and tumor tissue from colon sections. Red and blue color intensity indicates the degree of upregulated and downregulated genes, respectively (WT  $n = 3$ , *Il15*<sup>-/-</sup>  $n = 3$ ). The lists of genes are shown in **Table S3**. **(C)** qRT-PCR analysis showing the level of mRNA expression of indicated genes in normal and tumor tissue from WT and *Il15*<sup>-/-</sup> mice relative to *Hprt*. Data are shown as mean  $\pm$  SEM of WT  $n = 14-17$ , *Il15*<sup>-/-</sup>  $n = 14-17$ . Results are from three independent experiments. **(D)** Mean fold change increase of expression levels of genes in *Il15*<sup>-/-</sup> vs. WT tumor tissues normalized to their respective non-lesional tissue. Mann-Whitney tests were used for statistical analysis.



anti-colon carcinoma immunity, this may also create a cytokine/chemokine microenvironment that facilitates tumor growth.

IL-15 deficient mice have been used extensively to delineate the multifaceted role that IL-15 plays in inflammatory conditions. Most recently, deletion of IL-15 has been associated with increased disease severity in the experimental autoimmune encephalomyelitis (EAE) model of multiple sclerosis with increased CD8<sup>+</sup> T cell influx and cytokines levels such as IL-6,<sup>37</sup> and in the TAX-LUC model of human lymphoma characterized instead by a reduction of CD8<sup>+</sup> infiltrating T cells and IL-6.<sup>38</sup> Therefore, these findings highlight the duality of IL-15 function with respect to the CD8<sup>+</sup> T cell compartment and the influence on IL-6 levels. Furthermore, the contribution of IL-15 toward the final disease outcome is dependent on the inflammatory context, with deletion of the cytokine in a *Toxoplasma gondii* infection model resulting in reduced disease severity and mortality, with lower levels of IL-6, TNF $\alpha$  and IL-1 $\beta$  produced compared to WT mice.<sup>4</sup> Of interest to this present study, IL-15 deletion has been investigated in the DSS colitis model, with IL-15 deletion resulting in a protective effect with lower numbers of lamina propria CD8<sup>+</sup> T cells, NK cells, and pro-inflammatory cytokines.<sup>39</sup>

Our data provide novel insight into the role of IL-15 in colorectal carcinogenesis, by showing that in mice, disruption of the IL-15 signaling pathway correlates with enhanced tumorigenesis which is marked by increased serum levels of IFN $\gamma$ , IL-10, IL-17, CXCL1, and G-CSF. In a different model using metastatic murine CT26 or MC38 colon carcinoma, IL-15 alone or in combination with other molecules exert a protective role in cancer.<sup>32,40</sup> Additionally, in human tissue we observed markedly increased IL-15 expression in colon neoplastic cells suggesting a regulatory effect of IL-15 in CRC.

Under physiologic conditions IL-15 exerts a tumor suppressive function; however in transformed tumor epithelial cells IL-15 promotes their survival ultimately exerting a tumor-promoting role. This provides an explanation for the initially confounding findings of Kuniyasu and colleagues (2001) supporting an oncogenic role for IL-15 by promoting the proliferation of IECs and the production of angiogenic factors.<sup>14,16</sup> Interestingly, the antitumor effect of IL-15 appears to display only partial dependence on IL-15R $\alpha$  signaling. Our data show that *Il15r $\alpha$* <sup>-/-</sup> mice are significantly less susceptible to tumor development than *Il15*<sup>-/-</sup>, suggestive of alternative and redundant IL-15 signaling via IL-2R $\beta$  and IL-2R $\gamma$  pathways.

IL-15 is expressed by epithelial and non-epithelial cells and the increased incidence and accelerated onset of tumorigenesis in *Il15*<sup>-/-</sup> mice potentially results from a combination of epithelial damage and non-epithelial contributions and deregulated colon repair. IL-15 in chronic colitis, besides acting as a pro-inflammatory cytokine reduces mucosal damage by preventing IEC apoptosis.<sup>29</sup> Neutralization of IL-15 in chronic DSS-induced colitis increased the score of disease.<sup>29</sup> Furthermore, it has been shown that IL-15 is required for intestinal epithelial barrier integrity by regulating tight junction formation in an IL-2R $\beta$ -dependent manner.<sup>41</sup> In this present study we find increased apoptosis in IL-15 deficient mice, with no change in

proliferation, supporting the previously described role for IL-15 in colon repair.<sup>29</sup>

We show that selective recovery of IL-15 expression in CD11c<sup>+</sup> cells of *Il15*<sup>-/-</sup> mice significantly lowers tumor burden and decreases the severity of dysplasia compared with *Il15*<sup>-/-</sup> mice. CD11c is a marker of DC, which are known to trans-present IL-15 to NK and CD8<sup>+</sup> cytotoxic lymphocytes, augmenting the “tumor-suppressor” activity of the latter.<sup>1,42</sup> Our data suggest that during AOM/DSS (a model of carcinogenesis) mice deficient in IL-15 are more susceptible to tumor development as a consequence of an increased inflammatory promoting effect on tumor growth. In our hands the restoration of CD8<sup>+</sup> T cells and NK antitumor immunity is sufficient to overcome the protumorigenic effects of the inflammatory milieu seen in the IL-15 deficient mice. However, we do recognize that reconstitution of IL-15 in antigen presenting cells could have an additional effect on other immune cells such as macrophages. In tumor transplantation model Gillgrass and colleagues recently demonstrated that *Il15*<sup>-/-</sup> CD4<sup>+</sup> T cells display a Th2 phenotype, which may be capable of polarizing macrophages to a pro-metastatic M2 phenotype while lung macrophages of *Il15tg* mice produce high levels of nitric oxide and IL-12, indicative of an anti-metastatic M1 polarized phenotype.<sup>43,44</sup>

The increased tumorigenesis in *Il15*<sup>-/-</sup> mice was found to be accompanied by increased levels of specific cytokines in serum, among them, IL-17, IFN $\gamma$  and IL-10. TH17 cytokines are expressed by TH17 cells and by innate lymphoid cells (ILCs) in response to IL-23 in the inflamed colon,<sup>45</sup> furthermore IL-17 has been described to promote angiogenesis and tumor growth<sup>46</sup> while IL-22 plays a dual protective and pro-inflammatory role in intestinal inflammation.<sup>47,48</sup> Interestingly, IL-22 was recently found to play a significant role for CAC development<sup>49</sup> and in the absence of IL-15, we have detected an induction of IL-23 and IL-23-induced cytokines. We observed an increase in IL-22 transcripts in colonic tumor together with IL-17A and IFN $\gamma$  levels in sera of *Il15*<sup>-/-</sup> mice. In comparison with WT, we also observed a significant decrease in IL-12(p40), a shared subunit between IL-23 and IL-12 possibly indicating a more prominent role for the cytokine in this setting. Our results compliment previous studies demonstrating IL-17 correlation with worse prognosis in patients affected by colon carcinomas,<sup>50</sup> furthermore, increased IL-17 levels have been described in the mouse colonic cancer model.<sup>51</sup> Taken together, decreased IL-15 correlates with an increase in IL-22 as well as IL-17 and IFN $\gamma$ , cytokines that promote active chronic colitis and progression to colon cancer. Our results therefore, support an inhibitory effect of IL-15 on type 17-responses.

The IL-15 signaling in the presence of its receptor, IL-15R $\alpha$  regulates CCL5 production by T cells, DC and myeloid cells.<sup>52-54</sup> In accordance, in our model, the T and NK cell recruiting chemokine, CCL5, is strongly reduced in the sera of *Il15*<sup>-/-</sup> mice compared to the WT level, therefore inhibiting antitumor immunity.<sup>55,56</sup>

The role of IL-15 on regulatory T cells (Tregs) is complex and controversial. The increased levels of IL-10, the main suppressive cytokine produced by Tregs, in the sera of IL-15 deficient AOM/

DSS mice agrees with the inhibitory role of IL-15 on intestinal regulatory activities shown by Depaolo and colleagues.<sup>57</sup> Our data suggest that the loss of IL-15 leads to increased levels of the regulatory IL-10 and the pro-tumoral chemokine, CXCL1<sup>58</sup> which in turn results in suppression of antitumor immune responses and promotes tumor progression. Moreover, this is significantly inhibited when CD11c<sup>+</sup> cells produce IL-15 selectively.

Microarray analysis revealed that IL-15 deficiency in tumor tissue correlates with an increased expression of inflammatory promoting genes, such as proteases, cytokines/chemokines/cytokines receptors. Collectively the results obtained from our analysis indicate that since IL-15 deficiency promotes colon-associated carcinogenesis, classical tumor markers are amplified, but that specific inflammatory mediators are induced in an IL-15-dependent manner. To confirm the latter hypothesis, it will be interesting to examine in subsequent studies whether, for example, a signaling milieu dominated by *Spp1*, *Il1b*, *Cxcl5*, *Il22*, and/or *Il23*, does indeed promote colon tumor development/growth. On this matter, currently we can only speculate that the differences observed in cytokine expression in the tumor microenvironment are the final result of multiple events and indirectly controlled by IL-15. IL-22, IL-23, and IL-1 $\beta$  have been shown to cross-regulate their expression, be pro-inflammatory and exert both tumor inhibitory and promoting effects in colon cancer and are most likely the product of infiltrating immune cells, while *Spp1* and *CXCL5* are likely the result of increased epithelial activity and/or colonic tumor cells. We therefore suggest that the dysregulation of inflammatory cells accompanying tumor growth seen in IL-15 deficient mice results in the skewing of the cytokine milieu.

In summary, based on the data presented here we conclude that IL-15 plays a crucial and multifaceted role in gut homeostasis. This study sheds light on the role of IL-15 in colorectal carcinogenesis, providing supporting data for the employment of IL-15 for immune boosting in cancer immunotherapy, although the anti-apoptotic function of IL-15 on epithelial tumor cells should be carefully considered.

## Materials and Methods

### Mice

6–10 week old aged-matched female mice were used for all experiments. C57BL/6j (WT), *Il15*<sup>-/-10</sup>, *Il15tg* (strain B1)<sup>33</sup>, *Il15 $\alpha$* <sup>-/-11</sup>, and *CD11c-Il15* were used for the study, with all mice on the C57BL/6 background. *CD11c-Il15* mice were generated by breeding transgenic mice, which express IL-15 under the CD11c promoter, with *Il15*<sup>-/-</sup> mice (Polansky et al., in preparation). Mice were maintained in specific pathogen-free conditions at the Research Center Borstel animal facilities with all *in vivo* procedures performed in compliance with national animal care and institutional guidelines. Protocols were approved by the “Ministerium für Landwirtschaft, Umwelt und ländliche Räume des Landes Schleswig-Holstein” (# AZ V312-72241.123-3 (12-2/10)).

### Human tissue samples

Formalin-fixed paraffin-embedded specimens from 10 patients with CAC along with tissue arrays encompassing 74 cases with SCRC were examined. Patients had not received any immuno-, chemo- or radiotherapy interventions prior to surgical resection of the lesions. All samples were obtained from the Laboratory of Histology and Embryology, Athens Medical School according to local ethical committee approvals.

### AOM/DSS protocol

Colitis-induced tumorigenesis was induced by a single intraperitoneal injection of AOM (Sigma–Aldrich) (10 mg/kg) 7 d prior to three 5-d cycles of 1% (w/v) DSS (MP Biomedicals, molecular weight of 36,000–50,000) in the drinking water. DSS was alternated with 14-d cycles of drinking water. At the endpoint (day 100–120) colons were harvested, incised longitudinally, and washed with PBS. Colon weight was measured, photographs were taken (Sony Cyber-shot DSC-W55S Compact 7.2 Mpix), tumors were counted and their diameter was measured.

### FACS analysis

Cells were isolated from spleen and mesenteric lymph nodes, resuspended in FACS-buffer (2% newborn calf serum, 0.1% NaN<sub>3</sub>, 2 mM EDTA in PBS), stained with mAb against CD8 $\alpha$  (clone: 53-6.7, 100701), NKp46 (clone: 29A1.4, 137601) (all Biologend), for 20 min at 4°C and subsequently washed with FACS buffer containing 1  $\mu$ g/mL propidium iodide (Sigma–Aldrich). Cells were analyzed using a FACSCalibur flow cytometer (BD Biosciences) and Flowjo software (TreeStar Inc.).

### Cytokines and chemokines measurement

Cytokines (IL-1 $\alpha$ , IL-1 $\beta$ , IL-2, IL-3, IL-4, IL-5, IL-6, IL-9, IL-10, IL-12 (p40), IL-12 (p70), IL-13, IL-17A, Eotaxin, G-CSF, GM-CSF, IFN $\gamma$ , KC, MCP-1 (MCAF), MIP-1 $\alpha$ , MIP-1 $\beta$ , CCL5, and TNF- $\alpha$ ) were quantified from mouse sera using the Bio-Plex Mouse Cytokine Group 1 23-plex Assay (Bio-Rad) according to manufacturer's instructions.

### Histopathological analysis of mouse tissue

Colon tissue was fixed in 3.7% buffered formalin and embedded in paraffin and sections examined following H&E staining. Colitis severity was evaluated (designated as histological activity index) according to Ullman and Itzkowitz grading: grade 1: inactive colitis; grade 2: mild colitis; grade 3: moderate colitis; grade 4: severe colitis. Assessment of dysplasia was based on the two-tiered (low- and high-grade) system.<sup>59</sup> Histopathological changes were evaluated by three molecular pathologists blinded to the identity of the treatment group.

### Immunohistochemical analysis

Immunohistochemical staining was performed on paraffin-embedded tissues as previously described,<sup>60</sup> using the following antibodies against: IL-15 (500-p15, Peprotech; 1:500),  $\beta$ -catenin (E-5, sc7963, Santa Cruz; 1:100), p53 (NCL p53-CM5p, Novocastra; 1:400), p21 (F-5, sc-6246, Santa Cruz; 1:200), cleaved

caspase-3 (9661L, Cell Signaling; 1:100). Heat-mediated antigen retrieval was performed in a microwave using 10 mM citric acid (pH 6.0) for 25 min. For the immunohistochemical analysis the UltraVision LP Detection System (#TL-060-HD, Thermo Scientific) was employed according to the manufacturer's instructions. A mixed score (MS) combining a staining index (SI) and a labeling index (LI) was used for the evaluation of the IL-15 expression. The SI is subdivided into four ordinal values: 0: negative, 1: weak, 2: moderate, 3: strong. LI is defined as follows: 0: negative, 1: 1–20%, 2: 21–50%, 3: 51–100% positive cells. We classified IL-15 score as follows: 0–2: low, 3–4: medium, 6–9: high. The MS is obtained by multiplying SI with LI (MS = SI × LI). Evaluation of p53, p21,  $\beta$ -catenin, and cleaved caspase-3 was performed as previously described.<sup>60–64</sup> Previously characterized cases served as positive controls for p53 and p21.<sup>60–62</sup> Three independent observers carried out slide examination, with minimal inter-observer variability.

### RNA extraction and microarray analysis

Colon tissue (30 mg) from WT and *Il15*<sup>-/-</sup> mice was homogenized in 600  $\mu$ L of RLT buffer using the Omni TH2 homogenizer (Omni International). RNA extraction was performed using the RNeasy mini Kit (Qiagen) as per manufacturer's instructions. First strand cDNA was synthesized using 2  $\mu$ g of RNA, oligo-dT primer (5'-TTTTGTACAAGC(TTT)10-3') and the Superscript III reverse transcriptase (Life Technologies) according to manufacturer's instructions.

Gene expression from non-tumor, near tumor and tumor sections from 3 WT and 3 *Il15*<sup>-/-</sup> mice were examined by microarray analysis using the Affymetrix Genechip<sup>®</sup> Mouse Gene 1.0 ST array (28,853 gene transcripts). RNA quality control, gene chip hybridization and data acquisition were performed at the Expression Core Facility, TU Munich. Arrays data were normalized using the Exon Robust Multichip Average (RMA) algorithm from the software Partek Genomic Suite (Partek Inc.) and analyzed with the Linear Model for Microarray Data (LIMMA) to identify differentially expressed genes with a q value (FDR) < 0.05 using the open-source R software, v.3.0.1. and the LIMMA and QVALUE Bioconductor Packages. Genes with a log<sub>2</sub> fold change  $\geq 2$  were taken as input to perform gene functional clustering analysis using the Database for Annotation, Visualization and Integrated Discovery (DAVID).

### qRT-PCR

Gene expression from Tumor tissues was measured by qRT-PCR using the LightCycler 480 (Roche Diagnostics), the ready-

to-use hot start reaction mix (LightCycler<sup>®</sup> 480 probe master), the specific hydrolysis probes (Universal Probe Library, UPL) and primers, designed through the Universal Probe Library Assay Designed Center ([www.universalprobelibrary.com](http://www.universalprobelibrary.com)) (Table S5). Gene expression values were normalized to the housekeeping gene *Hprt*.

### Statistical analysis

Statistically significant differences were calculated by using the nonparametric Mann–Whitney test (Prism software; GraphPad Software Inc.) with \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$  considered significant.

### Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

### Acknowledgments

We thank Hanno Ewers, Katrin Westphal, Katrin Streek, Gesine Rode, Frauke Koops, Platonas N. Selemenakis, and Ioannis L. Aivaliotis for technical assistance, Zane Orinska for helpful discussion and Leo Zeef (University of Manchester) for the microarrays analysis.

### Funding

This work was supported in part by European Commission FP7 funding (INFLACARE agreement number 223151 and INSPIRE agreement number 284460) and Aristeia II from GSRT, Greece.

### Authors' Contributions

Conception and design by RB, VGG, SBP; Development of methodology by RB, ISP, OD, DAG, MP, PGF; Acquisition of data by RB, ISP, JP, OD, DAG, MP, PGF; Analysis and interpretation of data by RB, ISP, DAG, HS, KE, PGF, VGG, SBP; Writing, review, and/or revision of the manuscript by RB, MC, ISP, DAG, VGG, SBP; Study supervision by VGG, SBP.

### Supplemental Material

Supplemental data for this article can be accessed on the publisher's website.

### References

1. Jakobsiak M, Golab J, Lasek W. Interleukin 15 as a promising candidate for tumor immunotherapy. *Cytokine Growth Factor Rev* 2011; 22:99-108; PMID:21531164; <http://dx.doi.org/10.1016/j.cytogfr.2011.04.001>
2. van Heel DA. Interleukin 15: its role in intestinal inflammation. *Gut* 2006; 55:444-5; PMID:16531523; <http://dx.doi.org/10.1136/gut.2005.079335>
3. Doherty TM, Seder RA, Sher A. Induction and regulation of IL-15 expression in murine macrophages. *J Immunol* 1996; 156:735-41
4. Schulthess J, Meresse B, Ramiro-Puig E, Montcuquet N, Darche S, Begue B, Ruelmele F, Combadiere E, Gravelle C, Di Santo JP, Buzoni-Gatel D et al. Interleukin-15-dependent NKp46+ innate lymphoid cells control intestinal inflammation by recruiting inflammatory monocytes. *Immunity* 2012; 37:108-21; PMID:22705105; <http://dx.doi.org/10.1016/j.immuni.2012.05.013>
5. Giri JG, Ahdieh M, Eisenman J, Shanebeck K, Grabstein K, Kumaki S, Namen A, Park LS, Cosman D, Anderson D. Utilization of the beta and gamma chains of the IL-2 receptor by the novel cytokine IL-15. *EMBO J* 1994; 13:2822-30; PMID:8026467
6. Grabstein KH, Eisenman J, Shanebeck K, Rauch C, Srinivasan S, Fung V, Beers C, Richardson J, Schoenborn MA, Ahdieh M et al. Cloning of a T cell growth factor that interacts with the beta chain of the interleukin-2 receptor. *Science* 1994; 264:965-8; PMID:8178155; <http://dx.doi.org/10.1126/science.8178155>
7. Bamford RN, Grant AJ, Burton JD, Peters C, Kurys G, Goldman CK, Brennan J, Roessler E, Waldmann TA. The interleukin (IL) 2 receptor beta chain is shared by IL-2 and a cytokine, provisionally designated IL-T, that



- IL-10 subfamily, induces inflammatory responses in colonic subepithelial myofibroblasts. *Gastroenterology* 2005; 129:969-84; PMID:16143135; <http://dx.doi.org/10.1053/j.gastro.2005.06.071>
49. Kirchberger S, Royston DJ, Boulard O, Thornton E, Franchini F, Szabady RL, Harrison O, Powrie F. Innate lymphoid cells sustain colon cancer through production of interleukin-22 in a mouse model. *J Exp Med* 2013; 210:917-31; PMID:23589566; <http://dx.doi.org/10.1084/jem.20122308>
  50. Tosolini M, Kirilovsky A, Mlecnik B, Fredriksen T, Mauger S, Bindea G, Berger A, Bruneval P, Fridman WH, Pagès E, et al. Clinical impact of different classes of infiltrating T cytotoxic and helper cells (Th1, th2, treg, th17) in patients with colorectal cancer. *Cancer Res* 2011; 71:1263-71; PMID:21303976; <http://dx.doi.org/10.1158/0008-5472.CAN-10-2907>
  51. Wu S, Rhee KJ, Albesiano E, Rabizadeh S, Wu X, Yen HR, Huso DL, Brancati FL, Wick E, McAllister F et al. A human colonic commensal promotes colon tumorigenesis via activation of T helper type 17 T cell responses. *Nat Med* 2009; 15:1016-22; PMID:19701202; <http://dx.doi.org/10.1038/nm.2015>
  52. Perera LP, Goldman CK, Waldmann TA. IL-15 induces the expression of chemokines and their receptors in T lymphocytes. *J Immunol* 1999; 162:2606-12; PMID:10072502
  53. Chen JP, Liao NS, Lai SL, Hsu L, Mao WY, Ku MC, Liao F. Reduced 2,4-dinitro-1-fluorobenzene-induced contact hypersensitivity response in IL-15 receptor alpha-deficient mice correlates with diminished CCL5/RANTES and CXCL10/IP-10 expression. *Eur J Immunol* 2005; 35:690-8; PMID:15719370; <http://dx.doi.org/10.1002/eji.200425577>
  54. Chenoweth MJ, Mian MF, Barra NG, Alain T, Sonenberg N, Bramson J, Lichty BD, Richards CD, Ma A, Ashkar AA. IL-15 can signal via IL-15Ralpha, JNK, and NF-kappaB to drive RANTES production by myeloid cells. *J Immunol* 2012; 188:4149-57; PMID:22447977; <http://dx.doi.org/10.4049/jimmunol.1101883>
  55. Lavergne E, Combadiere C, Iga M, Boissonnas A, Bonduelle O, Maho M, Debr&Eacute; P, Combadiere B. Intratumoral CC chemokine ligand 5 overexpression delays tumor growth and increases tumor cell infiltration. *J Immunol* 2004; 173:3755-62; PMID:15356122; <http://dx.doi.org/10.4049/jimmunol.173.6.3755>
  56. Nesbeth Y, Scarlett U, Cubillos-Ruiz J, Martinez D, Engle X, Turk MJ, Conejo-Garcia JR. CCL5-mediated endogenous antitumor immunity elicited by adoptively transferred lymphocytes and dendritic cell depletion. *Cancer Res* 2009; 69:6331-8; PMID:19602595; <http://dx.doi.org/10.1158/0008-5472.CAN-08-4329>
  57. DePaolo RW, Abadie V, Tang F, Fehlner-Peach H, Hall JA, Wang W, Marietta EV, Kasarda DD, Waldmann TA, Murray JA et al. Co-adjutant effects of retinoic acid and IL-15 induce inflammatory immunity to dietary antigens. *Nature* 2011; 471:220-4; PMID:21307853; <http://dx.doi.org/10.1038/nature09849>
  58. Baier PK, Eggstein S, Wolff-Vorbeck G, Baumgartner U, Hopt UT. Chemokines in human colorectal carcinoma. *Anticancer Res* 2005; 25:3581-4; PMID:16101183
  59. Ullman TA, Itzkowitz SH. Intestinal inflammation and cancer. *Gastroenterology* 2011; 140:1807-16; PMID:21530747; <http://dx.doi.org/10.1053/j.gastro.2011.01.057>
  60. Gorgoulis VG, Vassiliou LV, Karakaidos P, Zacharatos P, Kotsinas A, Liloglou T, Venere M, Dittullo RA Jr, Kastrinakis NG, Levy B et al. Activation of the DNA damage checkpoint and genomic instability in human precancerous lesions. *Nature* 2005; 434:907-13; PMID:15829965; <http://dx.doi.org/10.1038/nature03485>
  61. Cooks T, Pateras IS, Tarcic O, Solomon H, Schetter AJ, Wilder S, Lozano G, Pikarsky E, Forsheo T, Rosenfeld N et al. Mutant p53 prolongs NF-kappaB activation and promotes chronic inflammation and inflammation-associated colorectal cancer. *Cancer Cell* 2013; 23:634-46; PMID:23680148; <http://dx.doi.org/10.1016/j.ccr.2013.03.022>
  62. Schramek D, Kotsinas A, Meixner A, Wada T, Elling U, Pospisilik JA, Neely GG, Zwick RH, Sigl V, Forni G et al. The stress kinase MKK7 couples oncogenic stress to p53 stability and tumor suppression. *Nat Genet* 2011; 43:212-9; PMID:21317887; <http://dx.doi.org/10.1038/ng.767>
  63. Kotsinas A, Evangelou K, Zacharatos P, Kittas C, Gorgoulis VG. Proliferation, but not apoptosis, is associated with distinct beta-catenin expression patterns in non-small-cell lung carcinomas: relationship with adenomatous polyposis coli and G(1)-to S-phase cell-cycle regulators. *Am J Pathol* 2002; 161:1619-34; PMID:12414510; [http://dx.doi.org/10.1016/S0002-9440\(10\)64440-9](http://dx.doi.org/10.1016/S0002-9440(10)64440-9)
  64. Gupta J, del Barco Barrantes I, Igea A, Sakellariou S, Pateras IS, Gorgoulis VG, Nebreda AR. Dual function of p38alpha MAPK in colon cancer: suppression of colitis-associated tumor initiation but requirement for cancer cell survival. *Cancer Cell* 2014; 25:484-500; PMID:24684847; <http://dx.doi.org/10.1016/j.ccr.2014.02.019>