

## INVITED REVIEW

# Genetics of human metabolism: an update

Gabi Kastenmüller<sup>1,4</sup>, Johannes Raffler<sup>1</sup>, Christian Gieger<sup>2,3</sup>  
and Karsten Suhre<sup>1,5,\*</sup>

<sup>1</sup>Institute of Bioinformatics and Systems Biology, <sup>2</sup>Research Unit of Molecular Epidemiology and <sup>3</sup>Institute of Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany, <sup>4</sup>German Center for Diabetes Research, Neuherberg, Germany and <sup>5</sup>Department of Physiology and Biophysics, Weill Cornell Medical College—Qatar, Doha, Qatar

\*To whom correspondence should be addressed at: Weill Cornell Medical College—Qatar, Qatar Foundation—Education City, PO Box 24144, Doha, Qatar. Tel: +974 4492 8482; Email: karsten@suhre.fr

## Abstract

Genome-wide association studies with metabolomics (mGWAS) identify genetically influenced metabolotypes (GIMs), their ensemble defining the heritable part of every human's metabolic individuality. Knowledge of genetic variation in metabolism has many applications of biomedical and pharmaceutical interests, including the functional understanding of genetic associations with clinical end points, design of strategies to correct dysregulations in metabolic disorders and the identification of genetic effect modifiers of metabolic disease biomarkers. Furthermore, it has been shown that GIMs provide testable hypotheses for functional genomics and metabolomics and for the identification of novel gene functions and metabolite identities. mGWAS with growing sample sizes and increasingly complex metabolic trait panels are being conducted, allowing for more comprehensive and systems-based downstream analyses. The generated large datasets of genetic associations can now be mined by the biomedical research community and provide valuable resources for hypothesis-driven studies. In this review, we provide a brief summary of the key aspects of mGWAS, followed by an update of recently published mGWAS. We then discuss new approaches of integrating and exploring mGWAS results and finish by presenting selected applications of GIMs in recent studies.

## Introduction

Homeostasis of blood metabolites and excretion of metabolites through urine are crucial for maintaining human health. Therefore, dysregulations of metabolite levels often indicate critical physiological states, a fact that is widely used in clinical chemistry for the diagnosis of common metabolic diseases, such as diabetes and chronic kidney disease, but also of rare metabolic disorders, including inborn errors of metabolism, such as phenylketonuria and medium-chain acyl-CoA dehydrogenase deficiency. The number and variety of metabolites that are detectable and quantifiable in biological samples in a single

experiment largely increased through the advent of modern metabolomics techniques, which are mainly based on nuclear magnetic resonance spectroscopy or mass spectrometry coupled to gas or liquid chromatography (1). As a consequence, blood and urine levels of a broad range of metabolites can be determined on an epidemiological scale from bio samples of thousands of individuals, opening new avenues to study human metabolism and its variation in health and disease based on large population studies (2).

Metabolite levels and their variation in the human body are influenced by various factors, specifically environmental conditions (day light, exposure to xenobiotics, etc.), lifestyle (nutrition

Received: May 31, 2015. Revised and Accepted: July 6, 2015

© The Author 2015. Published by Oxford University Press.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact [journals.permissions@oup.com](mailto:journals.permissions@oup.com)

habits, smoking and alcohol consumption, physical activity, etc.) and genetics. The genetic control of metabolite levels and its impact on human health are apparent in inborn errors of metabolism, where rare genetic variants disrupt a single gene that then leads to extreme, eventually toxic levels of related metabolites. Availability of population scale metabolomics and genotype data now allows to systematically investigate also the less apparent influences of more common and less deleterious genetic variants on human metabolism by conducting genome-wide association studies with metabolomics (mGWAS). This has been demonstrated in the first mGWAS by Gieger *et al.* (3), in which the authors performed a genome-wide scan for associations of common single-nucleotide polymorphisms (SNPs) with the levels of >350 metabolites and all pairwise ratios thereof. Despite the relatively small sample size of this first mGWAS ( $n = 284$ ), they identified four common SNPs that significantly alter functionally related metabolic traits. Moreover, they found that ratios between metabolites can serve as proxies for the biochemical conversion of metabolites and thereby significantly strengthen the associations, a concept that has been helpful in many subsequent studies (4).

SNP-metabolite trait associations [metabolite quantitative trait loci (mQTLs)] identified in mGWAS typically cluster in groups of SNPs in high linkage disequilibrium (LD) that are associated with the same and/or biochemically related metabolites. Each of these groups of mQTLs defines a distinct genetically influenced metabolite (GIM). Many GIMs identified so far share most of the following properties (2): (i) The variance explained by the common genetic variants in the observed metabolic traits is often large, exceeding 10%, and the effect sizes are strong, with minor allele homozygotes sometimes displaying 50% differences in their estimated enzymatic throughput compared with major allele homozygotes. (ii) For the majority of GIMs, the metabolic traits can be functionally linked to an enzyme, a transporter or a regulator of metabolism that is encoded at the genetic locus (e.g. the associated metabolites being substrates or products of the encoded enzyme). (iii) Associations found in GWAS with clinical end points are enriched in GIMs, and the related metabolites are candidate intermediate traits on the pathway to the disease [for details on the concept of the intermediate phenotype, see (5)]. For instance, Ried *et al.* (6) demonstrated the potential of mGWAS results to infer asthma-related metabolic markers by the identification of potentially deregulated phospholipids that associate with asthma and asthma risk alleles.

## Recent GWAS with Metabolic Traits

Recent mGWAS have extended previous studies in samples size, variety of metabolic traits and depth of genetic analysis (Table 1). As a result, the number of known GIMs largely increased. At the same time, most previously reported associations could be replicated in various cohorts, often using different metabolomics platforms and sometimes also different biofluids. Moreover, the functional interpretation of GIMs remarkably improved as recent studies started to systematically combine GIMs with additional data, such as expression quantitative trait loci (eQTLs) and metabolic pathway information. Thus, mGWAS published in recent years did not only lead to a more comprehensive and refined view on the inherited part of human metabolic individuality, but it also facilitated establishing potential links of this individuality to disease.

The largest leap in increasing the number of GIMs in the last 2 years has been achieved in the recent study by Shin *et al.* (23). The authors reported 145 GIMs (Fig. 1), 84 of which were new.

This mGWAS was based on relative quantification from liquid chromatography-mass spectrometry and gas chromatography-mass spectrometry (GC-MS) for 486 metabolites in blood samples of 7824 participants from two European cohorts. The metabolites and their associated genetic loci broadly cover representatives of all major metabolic pathways, providing the most comprehensive picture of how genetic variation affects homeostasis in blood metabolism to date. To facilitate further exploration and functional interpretation of the findings by the biomedical research community, all GIMs have been embedded into the metabolic network, which was reconstructed from the metabolomics data (Fig. 2). This network as well as a rich set of additional information (including mapped eQTLs, GWAS hits to clinical phenotypes, and drug-target information) is available at <http://gwas.eu/si>.

Recent mGWAS with similar or even larger sample sizes include the study by Draisma *et al.* (26), who analyzed 129 mostly lipid-related metabolites in 7478 individuals from seven European cohorts. They identified GIMs at 31 genomic loci, although only 4 of them were new. Nonetheless, the study refined patterns of associations of related lipid traits at already known loci and provided new insights into the complexity of lipid metabolism. Another example is the recent mGWAS by Rhee *et al.* (20), who identified specific patterns in the genetic associations of 46 triacylglycerols with different lengths of the fatty acid chain and degree of desaturation. Some of their loci have been previously linked to total triglyceride levels in blood, namely variants in glucokinase (hexokinase 4) Regulator (GCKR), FADS1-3 and APOA1/C3/A4/A5. While variants in GCKR showed stronger associations to triacylglycerols with shorter fatty acid chains, associations of variants in FADS1-3 were stronger with longer fatty acid chain lengths and a higher degree of desaturation, which is in line with the fatty acid desaturase function of the proteins encoded at the FADS1-3 locus.


While most early mGWAS were based on samples from individuals of European ancestry, Rueedi *et al.* (21) recently analyzed genetic associations of nuclear magnetic resonance (NMR)-derived metabolic traits in urine samples from 835 Europeans and replicated their findings in 601 samples from a Brazilian population with a diverse ethnic background. Another recent study by Yu *et al.* (22) performed an mGWAS in serum samples from almost 2000 individuals of African-American descent and thereby confirmed the robustness of various GIMs across different ethnicities, but also found novel independent variants at the same loci and even new GIMs, which have not been observed in Europeans before.

In addition to fully metabolome-wide mGWAS, many GWAS with smaller and more specialized sets of metabolic traits have been conducted. For instance, Stiles *et al.* (27) report genetic, anatomic and clinical determinants of human serum sterol and vitamin D levels; Ng *et al.* (28) analyzed genetic influences on blood levels of polychlorinated biphenyls and Xie *et al.* (29) conducted a GWAS focused on metabolites related to insulin sensitivity. The CHARGE Consortium (30) reported novel loci associated with plasma concentrations of four fatty acids in the *de novo* lipogenesis pathway. Further studies identified and refined associations with single metabolic traits, including disease-relevant markers, such as bilirubin (31–34), uric acid (35–38), dimethylarginine (39), homoarginine (40) and creatinine (41). These studies could identify new loci due to their much larger sample sizes, in some cases over 100 000 individuals, while using less expensive biochemistry-based methods.

Interestingly, most conclusions from mGWAS in humans also apply to animals and plants. Ghazalpour *et al.* (42) studied the genetic regulation of mouse liver metabolite levels. By analyzing 283 metabolites in 104 inbred and recombinant inbred mouse

**Table 1.** List of published mGWAS in humans and selected GWAS with multiple metabolic traits (in the order of publication date)

Biofluid	Metabolic trait(s)	Traits (N)	Platform	Study population	Cohort size (n)	Loci (n)	References
Serum	Targeted MS	363 + all ratios	MS	German	284	4	Gieger et al. (3)
Plasma and Serum	Sphingolipids	33 + 43 ratios	MS	European	4400	5	Hicks et al. (7)
Serum	Mainly phospholipids	163 + 26 406 ratios	MS	German, British	1809 + 422	9	Illig et al. (8)
Urine	NMR-derived metabolites	59 + 1661 ratios	NMR	German	862 + 992	5	Suhre et al. (9)
Serum	Non-targeted MS, knowns	276 + 37 179 ratios	MS	German, British	1768 + 1052	37	Suhre et al. (10)
Urine and Plasma	Urine: NMR peaks, plasma: mainly phospholipids	Urine: 512 peaks plasma: 163 + ratios	NMR + MS	British	211	3	Nicholson et al. (11)
Serum	Mainly lipid traits	117 + 99 ratios	NMR	Finnish	8330	31	Kettunen et al. (12)
Plasma	Phospholipids + sphingolipids	153	MS	European	4034	35	Demirkan et al. (13)
Serum	Mainly lipid traits	117 + 99 ratios	NMR	Finnish	8330	30	Tukiainen et al. (14)
Serum	Mainly lipid traits and low-weight metabolites	130	NMR	Finnish, British	1905 + 4703	34	Inouye et al. (15)
Serum	Non-targeted MS, unknowns	517	MS	German	1768	34	Krumsiek et al. (16)
Urine	NMR peaks	2425	NMR	Brazil	265	2	Montoliu et al. (17)
Plasma	NMR peaks	8600 + 124 750 ratios	NMR	German	1757	7	Raffler et al. (18)
Serum	MS peaks	6138	MS	Swedish	402 + 489	7	Hong et al. (19)
Plasma	Amino acids, amines, polar metabolites, lipids	217	MS	USA (European ancestry)	2076	31	Rhee et al. (20)
Urine	NMR peaks	1276	NMR	European, Brazil	835 + 601	11	Rueedi et al. (21)
Serum	Non-targeted MS, knowns and unknowns	308	MS	African American	1260	19	Yu et al. (22)
Serum	Non-targeted MS, knowns and unknowns	486 + 98 346 ratios	MS	European	7824	145	Shin et al. (23)
Serum	Targeted MS (mainly phospholipids) + non-targeted MS (knowns)	344 (151 + 193)	MS	European	1809 + 843	12 new	Ried et al. (24)
Serum	NMR-derived metabolites	42	NMR	European	2118	8	Demirkan et al. (25)
Serum	Mainly phospholipids	129	MS	European	7478 + 1182	31	Draisma et al. (26)

**Figure 1.** Manhattan plot of the mGWAS by Shin et al. (23). Upward pointing P-values: TwinsUK cohort, downward pointing P-values: KORA population study. Only SNPs with association to raw metabolites ( $P < 10^{-6}$ ) are displayed (no ratios). The green line indicates the genome-wide significance cutoff for the P-value ( $P < 10^{-10}$ ). Loci that reach genome-wide significance in either cohort are indicated by a short vertical black line. Loci with P-values  $< 10^{-30}$  are indicated with a red symbol. Loci that are further discussed in Figure 3 are highlighted and annotated [figure adapted from the Supplementary Material, Fig. S2 by Shin et al., *Nature Genetics*, 2014 (23)].


strains, they identified 240 loci, the majority of which accounted for 20–40% of total metabolite variation. Remarkably, more than one-third of the loci that regulate liver metabolites in mice also correspond to human GIMs, supporting the similarity in genetic regulation of metabolites between mice and humans. Chen et al. (43) conducted an mGWAS in rice, covering 840 metabolites and 6.4 million SNPs obtained from 529 diverse accessions of *Oryza sativa*. This study identified hundreds of common variants influencing numerous secondary metabolites with large effect sizes, and reported 36 candidate genes that modulate levels of metabolites of potential physiological and nutritional importance. The authors concluded that mGWAS provide a powerful tool

for large-scale interactive gene–metabolite annotation and identification, pathway elucidation and knowledge about crop improvement.

## New Concepts

### Identification of non-targeted metabolic traits

Many of the more recent mGWAS were based on non-targeted metabolomics approaches, which record all metabolite signals detectable by the specific NMR or MS method, including signals that could not be assigned to a specific biochemical molecule




**Figure 2.** Network integrating gene-metabolite associations and metabolite-metabolite correlations. Individual metabolites are lumped by pathway (colored circles) and colored by their general metabolic properties (see legend). Genetic loci (gray diamonds) are annotated by the gene that is most likely affected by the variant. Green edges between loci and metabolites represent significant genetic associations with metabolic traits. Gray edges between metabolites represent significant partial correlations between metabolic traits. The highlighted sub-network (shaded box) is further discussed in Figure 3. The full network is freely accessible in digital format at <http://gwas.eu/si> [figure adapted from Figure 2 by Shin et al., *Nature Genetics*, 2014 (23)].

(unknown metabolite). In contrast to targeted approaches that use optimized methods for the quantification of a set of predefined metabolites, non-targeted approaches usually provide less precise (relative) quantifications, but this for a biochemically broader range of metabolites. While less biological information can be gained from GIMs with signals of unknown identify, it has been shown that genetic association of data for unknown metabolites may allow metabolite identification in MS (16) and NMR (18,21). Often both known and unknown signals were considered in mGWAS to explore the entire breadth of the metabolite data from these non-targeted approaches. In 2013, Raffler et al. (18) reported an mGWAS with non-targeted NMR traits in blood plasma. While the concept of testing NMR signal intensities has already been introduced in previous mGWAS (11,17), this study was the first where the concept of testing ratios between NMR signals at different spectral positions was applied, resulting in the identification of four additional loci that displayed

genome-wide significant association signals. To elucidate the chemical identity of the metabolites that underlay the genetically associated NMR signals, the authors used pseudo-spectra to visualize either the strength of genetic associations of NMR signals ('association spectra') or the correlation between NMR signals and traits determined on complementary metabolomics platforms, such as MS and clinical biochemistry measurements ('correlation spectra'). In 2014, Rueedi et al. (21) presented an NMR-based mGWAS in urine where such pseudo-spectra were automatically annotated. There, the authors introduced the 'metabomatching' approach that compares association spectra with pure compound spectra derived from the Human Metabolome Database (HMDB) (44). Using this metabomatching approach, Rueedi et al. (21) could identify the metabolic nature of the signals in 6 out of 11 loci reported in their study. Hence, several methods are now available that allow to deduce the biochemical identity of non-targeted metabolic traits using genetic association data.





**Figure 3.** Example of the integration of mGWAS results in a biomedical context using data from different sources. This figure displays a 'Cardiovascular disease and hypertension metabolic sub-network', annotated based on correlations between molecular relationships and expert knowledge on blood pressure regulation, blood coagulation and known molecular risk factors for cardiovascular disease and hypertension. Metabolites (circles) and genes (diamonds) of the fibrinogen cleavage (left) and the kininogen/kinin system (right) and their interconnections were derived from the Shin *et al.* (see shaded box in Fig. 2) data. Gray nodes and edges display annotations of biochemical function based on expert knowledge (47). Colored nodes and edges correspond to reported associations based on genome-wide studies for blood pressure regulation (orange), blood coagulation (blue) and cholesterol levels (purple). This figure was first published as the Supplementary Material, Fig. S5 by Shin *et al.*, *Nature Genetics*, 2014 (23).

### Exome sequencing and imputation

Identification of causative variants in GWAS is always a major challenge. Intermediate phenotypes with large effect sizes, as obtained from metabolomics, may be useful to figure out the causative variants of associations with complex clinical end points when the genetic structure of the association with the intermediate trait and the disease end point are the same. For the first time, exome-sequencing-based metabolomics associations were reported by Demirkan *et al.* (25). The authors first performed an NMR-based mGWAS in serum using microarray genotype data of 2118 individuals. Following the mGWAS, they selected a subset of the GWAS cohort ( $n = 921$ ) and performed exome sequencing on candidate genes within eight candidate loci. They identified seven variants in or near four genes that modulate metabolite levels independently of the GWAS hits. For instance, the common SNP rs1047891 is a missense variant in CPS1 that was tagged by the genotyped SNP rs715. Interestingly, the association of rs1047891 to glycine was also previously reported by Shin *et al.* (23) using a fine-mapping approach of candidate loci based on 1000 Genomes Project data imputed genetic variants [see the Supplementary Material, Table S8 in Shin *et al.* (23)]. However, except for one variant, this fine-mapping approach did not significantly strengthen any association signal that was not already apparent using the genotyped SNPs. Fine mapping of association signals using imputed or exome-sequenced variants may thus help single out potentially causative SNPs; but for the moment, there is little evidence that it will help find truly novel association signals at the mGWAS level. This may change with larger sample numbers.

### Non-additive genetic models

In most mGWAS, association models between genotype and metabolite assume additive effects, which is intuitively supported

by the observation that many GIMs are related to rate limiting steps in enzymatic reactions or transport processes, which suggests a dose-dependent response to genetic variability. Tsepilov *et al.* (45) investigated systematically non-additive effects on a large panel of serum metabolites and all possible ratios ( $n = 22\,801$ ) in a population-based study ( $n = 1785$ ). They found that most genetic effects on metabolite concentrations and ratios were indeed additive, with a few notable exceptions that may allow to understand the genetic control of these loci more deeply. However, with larger sample sizes, rare variants with potentially larger effect sizes and recessive modes of inheritance may be detected.

### Epistasis and Mendelian randomization

Due to their large number and strong effect sizes, GIMs provide a test bed to identify and investigate more complex genetic and metabolic interactions. Shin *et al.* (23) provided an example of genetic interaction between two gene variants (epistasis), rs10469966 (NAT8) and rs4488133 (PYROXD2), and blood metabolite levels. The same study also reported an example of a Mendelian randomization analysis to establish causation, where expression of THEM4 was shown to mediate the association between rs6693388 and the ratio of linoleate (18:2n6) to 5,8-tetradecadien. However, it seems that as for now there were only few cases that display association signals that were strong enough to single out statistically significant cases of epistasis and Mendelian randomization.

### Multi-phenotype mGWAS

Strategies that combine several phenotypes in an mGWAS may potentially detect additional genetic loci and further their functional characterization. Inouye *et al.* (15) showed that association testing of multiple correlated phenotypes offers better power

than univariate analysis of single traits. Ried *et al.* (24) applied phenotype set enrichment analysis (PSEA), a method that tests sets of metabolites for association enrichment at genetic loci. In addition to confirming previously reported GIMs, the authors identified and validated 12 new loci.

### Augmenting GIMs with functional information

A crucial step in the interpretation of mGWAS results is to put the identified associations into the context of results from other association studies, including associations of genetic variants to traits at different phenotype layers. Most recently published mGWAS studies, therefore, go beyond the mere reporting of genetic associations by combining their results with additional -omics datasets (i.e. eQTLs) (23,25), by linking individual associations in a systems level approach (16,46), and by adding clinical association data to establish complex gene-to-disease networks (for an example, see Fig. 3). However, collecting and integrating publicly available association data still present a major bottleneck in the evaluation of mGWAS results, in large part due to the fact that data from different sources are reported on different, but highly correlated SNP sets. To facilitate this task, Arnold *et al.* (48) recently established the SNIIPA web service (<http://www.snipa.org>). SNIIPA contains LD information as well as functional annotations for almost all genetic variants of the latest 1000 Genomes Project dataset. The functional annotations include regulatory elements, eQTLs, associations to clinical traits and also associations to metabolic traits as provided by the NHGRI-EBI GWAS catalog (<http://www.ebi.ac.uk/gwas>). SNIIPA provides several user-friendly, interactive tools that are especially useful in the context of mGWAS, such as regional association plots and LD plots that are enriched with functional annotations. In particular, links to the Metabolomics GWAS server (<http://www.gwas.eu>) allow direct access to association data from the Suhre *et al.* (10) and Shin *et al.* (23) mGWAS, including many unpublished associations below the conservative level of genome-wide significance, that may prove to represent true positive associations when combined with additional evidence from user-provided studies (49).

### Application of GIMS for Hypothesis Generation

mGWAS results are more and more used as starting hypothesis for deeper functional research. The great potential of mGWAS as hypothesis generating tool is demonstrated when mGWAS 'rediscover'—sometimes decades-old—findings from biochemical experiments in the setting of modern genomic and metabolic studies (50). One interesting example is SNP rs37369, which is a non-synonymous variant (Val140Ile) in the coding region of *alanine-glyoxylate aminotransferase 2* (AGXT2). Several mGWAS linked this variant to changes in homeostasis of plasma beta-aminoisobutyrate (BAIB) (20), serum symmetric/asymmetric dimethylarginine (39) and serum homoarginine (40). Furthermore, SNP rs37369 was associated to elevated urinary excretion of BAIB (9,11,21) with BAIB concentrations >10 times higher in urine of homozygotes for the minor SNP allele. Since BAIB is one of AGXT2's substrates (51), it was hypothesized that rs37369 is causative for hyper-beta-aminoisobutyric aciduria (9,11), a heritable trait first described in the early 1950s (52,53). This mGWAS-generated hypothesis was recently validated by Kittel *et al.* (54) by *in vitro* studies where the authors demonstrated that the rs37369 polymorphism results in a significantly lower AGXT2 enzyme activity when compared with the wild type. These studies, inspired by a single mGWAS result, rejuvenated research interest in

AGXT2, indicating that altered AGXT2 activity may contribute to the pathogenesis of cardiovascular, renal, neurological and hematological diseases, and highlighted the unique role of AGXT2 at the intersection of key mitochondrial pathways, and even as a potential drug target (50). Inspired by an observed co-association of BAIB and certain triglyceride levels at the AGXT2 locus in their mGWAS, Rhee *et al.* (20) further set out to investigate the relationship between BAIB metabolism and lipid homeostasis. Using a morpholino knockdown of *agxt2* in zebrafish, the authors could indeed establish a functional link between BAIB, cholesterol ester and TAG metabolism, concluding that their data 'provides an example how the breadth of gene, metabolite, and phenotype data [ . . . ] can provide a springboard for research in metabolism'.

Another example where mGWAS data could have helped to formulate a hypothesis on a solute transporter's specificity has just been published by Nguyen *et al.* (55). Among the 12 novel loci reported by Ried *et al.* (24) was MFSD2A, a member of the major facilitator superfamily. The mouse homologue (*Mfsd2a*) was identified by Nguyen *et al.* (55) as a transporter for the essential omega-3 fatty acid docosahexaenoic acid. The authors showed that this previously orphan transporter transports lysolipids, writing: 'Unexpectedly, cell-based studies indicate that *Mfsd2a* transports DHA in the form of lysophosphatidylcholine (LPC), but not unesterified fatty acid, in a sodium-dependent manner. Notably, *Mfsd2a* transports common plasma LPCs carrying long-chain fatty acids such LPC oleate and LPC palmitate, but not LPCs with less than a 14-carbon acyl chain'. This finding was reflected in Ried *et al.*'s study data, which associated this locus with acyl-bound LPCs with side chain lengths C16:0, C17:0, C18:1, C20:4 and C18:0. This finding was further replicated by Draisma *et al.* (26), who found an association of SNP rs7529794 with these same LPCs, the top association at a *P*-value of  $2.8 \times 10^{-13}$ . Thus, had this information been available to the Nguyen *et al.*, they might actually have expected that *Mfsd2a* transports LPCs and have designed their experiments accordingly.

Pharmaceutical companies are exploring innovative ways of drug development from candidate selection to clinical proof of concept (56). One compelling application of GWAS for the validation of therapeutic targets through human genetics has been suggested by Plenge *et al.* (57). The authors describe the concept of using dose-response curves based on multiple functional variants in genes of pharmaceutical interest in order to prioritize molecular targets in drug development. However, for such an approach to be applicable, the target gene is required to harbor a causal variant that is unequivocally associated with a medical trait of interest, the biological function of the causal gene and causal variant needs to be known, and the gene has to harbor multiple causal variants of known biological function, thereby enabling the generation of genotype-phenotype dose-response curves. These criteria are unfortunately rarely met by genetic associations with clinical end points, especially in the case of complex disorders. However, when a GIM is overlapping such a locus, then, due to its generally much stronger effect size and its more direct link to the genetic variant, this GIM can provide intermediate and functionally relevant readouts (5) that allow for the identification of multiple causal variants in order to satisfy the criteria laid out by Plenge *et al.* (57).

### Conclusion and Outlook

With the era of GWAS with single clinical end points reaching maturation in meta-analyses that basically include all cohorts available worldwide for any given disease, what will the future bring

for GWAS with metabolic traits? Meta-analyses of mGWAS are certainly the next big step. However, merging data from different metabolomics platforms may require additional efforts in developing statistical methods that allow to combine phenotype information that match only partially (e.g. some platforms can differentiate between isoleucine and leucine, others only report their sum; some platforms resolve lipid traits to a high degree of detail, while other methods provide access to aggregated traits; MS or NMR peaks may be machine specific and hard to match between studies). Larger cohorts shall be phenotyped by multiple platforms in samples from the same donors. Moreover, samples from multiple biofluids are being collected in current population studies, allowing to study metabolic correlations between them, including possible genotype-dependent interactions with the human gut or saliva microbiome. A number of GWAS with other disease-relevant -omics phenotypes have also been conducted, including genome-wide gene expression (58), proteomics (59) and protein glycosylation (60), and further epigenome-wide association studies between DNA methylation and metabolomics (61) and studies on transcriptome-metabolome associations (62). We thus expect to see an exciting new series of GWAS in deeply omics-phenotyped cohorts, where the combination of multiple disease-relevant quantitative traits may allow the identification of functionally relevant genotype-pathway associations. In order to leverage the full potential of the multi-omics GWAS, new computational approaches for systems biology and network-based association tests need to be developed. We hope that the resulting growing number of GIMs will lead to a significant increase in our functional understanding of genetic variation in human metabolism and its pathologies, knowing that every single newly discovered GIM holds to potential to reveal an exciting and potentially health-relevant story.

*Conflict of Interest statement.* None declared.

## Funding

K.S. is supported by 'Biomedical Research Program' funds at Weill Cornell Medical College in Qatar, a program funded by the Qatar Foundation. The statements made herein are solely the responsibility of the author. C.G. is supported by a grant from the Helmholtz Association (Helmholtz-Russia Joint Research Group 310), the European Union's Seventh Framework Programme (FP7-Health-F5-2012) under Grant Agreement No. 305280 (MIMOmics) and BMBF e:Med project: e:AtheroSysMed—Systems medicine of myocardial infarction and stroke. Funding to pay the Open Access publication charges for this article was provided by Weill Cornell Medical College in Qatar.

## References

- Nicholson, J.K. and Lindon, J.C. (2008) Systems biology: metabolomics. *Nature*, **455**, 1054–1056.
- Suhre, K. and Gieger, C. (2012) Genetic variation in metabolic phenotypes: study designs and applications. *Nat. Rev. Genet.*, **13**, 759–769.
- Gieger, C., Geistlinger, L., Altmaier, E., Hrabce de Angelis, M., Kronenberg, F., Meitinger, T., Mewes, H.W., Wichmann, H.E., Weinberger, K.M., Adamski, J. et al. (2008) Genetics meets metabolomics: a genome-wide association study of metabolite profiles in human serum. *PLoS Genet.*, **4**, e1000282.
- Petersen, A.K., Krumsiek, J., Wagele, B., Theis, F.J., Wichmann, H.E., Gieger, C. and Suhre, K. (2012) On the hypothesis-free testing of metabolite ratios in genome-wide and metabolome-wide association studies. *BMC Bioinformatics*, **13**, 120.
- Kronenberg, F. (2012) Metabolic traits as intermediate phenotypes. In Suhre, K. (ed), *Genetics Meets Metabolomics*. Springer, New York, pp. 255–264.
- Ried, J.S., Baurecht, H., Stuckler, F., Krumsiek, J., Gieger, C., Heinrich, J., Kabesch, M., Prehn, C., Peters, A., Rodriguez, E. et al. (2013) Integrative genetic and metabolite profiling analysis suggests altered phosphatidylcholine metabolism in asthma. *Allergy*, **68**, 629–636.
- Hicks, A.A., Pramstaller, P.P., Johansson, A., Vitart, V., Rudan, I., Ugcocai, P., Aulchenko, Y., Franklin, C.S., Liebisch, G., Erdmann, J. et al. (2009) Genetic determinants of circulating sphingolipid concentrations in European populations. *PLoS Genet.*, **5**, e1000672.
- Illig, T., Gieger, C., Zhai, G., Romisch-Margl, W., Wang-Sattler, R., Prehn, C., Altmaier, E., Kastenmuller, G., Kato, B.S., Mewes, H.W. et al. (2010) A genome-wide perspective of genetic variation in human metabolism. *Nat. Genet.*, **42**, 137–141.
- Suhre, K., Wallaschofski, H., Raffler, J., Friedrich, N., Haring, R., Michael, K., Wasner, C., Krebs, A., Kronenberg, F., Chang, D. et al. (2011) A genome-wide association study of metabolic traits in human urine. *Nat. Genet.*, **43**, 565–569.
- Suhre, K., Shin, S.Y., Petersen, A.K., Mohnhey, R.P., Meredith, D., Wagele, B., Altmaier, E., CARDIoGRAM; Deloukas, P., Erdmann, J. et al. (2011) Human metabolic individuality in biomedical and pharmaceutical research. *Nature*, **477**, 54–60.
- Nicholson, G., Rantalainen, M., Li, J.V., Maher, A.D., Malmodin, D., Ahmadi, K.R., Faber, J.H., Barrett, A., Min, J.L., Rayner, N.W. et al. (2011) A genome-wide metabolic QTL analysis in Europeans implicates two loci shaped by recent positive selection. *PLoS Genet.*, **7**, e1002270.
- Kettunen, J., Tukiainen, T., Sarin, A.P., Ortega-Alonso, A., Tikkanen, E., Lyytikainen, L.P., Kangas, A.J., Soininen, P., Wurtz, P., Silander, K. et al. (2012) Genome-wide association study identifies multiple loci influencing human serum metabolite levels. *Nat. Genet.*, **44**, 269–276.
- Demirkan, A., van Duijn, C.M., Ugcocai, P., Isaacs, A., Pramstaller, P.P., Liebisch, G., Wilson, J.F., Johansson, A., Rudan, I., Aulchenko, Y.S. et al. (2012) Genome-wide association study identifies novel loci associated with circulating phospho- and sphingolipid concentrations. *PLoS Genet.*, **8**, e1002490.
- Tukiainen, T., Kettunen, J., Kangas, A.J., Lyytikainen, L.P., Soininen, P., Sarin, A.P., Tikkanen, E., O'Reilly, P.F., Savolainen, M.J., Kaski, K. et al. (2012) Detailed metabolic and genetic characterization reveals new associations for 30 known lipid loci. *Hum. Mol. Genet.*, **21**, 1444–1455.
- Inouye, M., Ripatti, S., Kettunen, J., Lyytikainen, L.P., Oksala, N., Laurila, P.P., Kangas, A.J., Soininen, P., Savolainen, M.J., Viikari, J. et al. (2012) Novel loci for metabolic networks and multi-tissue expression studies reveal genes for atherosclerosis. *PLoS Genet.*, **8**, e1002907.
- Krumsiek, J., Suhre, K., Evans, A.M., Mitchell, M.W., Mohnhey, R.P., Milburn, M.V., Wagele, B., Romisch-Margl, W., Illig, T., Adamski, J. et al. (2012) Mining the unknown: a systems approach to metabolite identification combining genetic and metabolite information. *PLoS Genet.*, **8**, e1003005.
- Montoliu, I., Genick, U., Ledda, M., Collino, S., Martin, F.P., le Coutre, J. and Rezzi, S. (2013) Current status on genome-metabolome-wide associations: an opportunity in nutrition research. *Genes Nutr.*, **8**, 19–27.
- Raffler, J., Romisch-Margl, W., Petersen, A.K., Pagel, P., Blochl, F., Hengstenberg, C., Illig, T., Meisinger, C., Stark, K., Wichmann, H.E. et al. (2013) Identification and MS-assisted interpretation of genetically influenced NMR signals in human plasma. *Genome. Med.*, **5**, 13.



19. Hong, M.G., Karlsson, R., Magnusson, P.K., Lewis, M.R., Isaacs, W., Zheng, L.S., Xu, J., Gronberg, H., Ingelsson, E., Pawitan, Y. et al. (2013) A genome-wide assessment of variability in human serum metabolism. *Hum. Mutat.*, **34**, 515–524.
20. Rhee, E.P., Ho, J.E., Chen, M.H., Shen, D., Cheng, S., Larson, M.G., Ghorbani, A., Shi, X., Helenius, I.T., O'Donnell, C.J. et al. (2013) A genome-wide association study of the human metabolome in a community-based cohort. *Cell Metab.*, **18**, 130–143.
21. Rueedi, R., Ledda, M., Nicholls, A.W., Salek, R.M., Marques-Vidal, P., Morya, E., Sameshima, K., Montoliu, I., Da Silva, L., Collino, S. et al. (2014) Genome-wide association study of metabolic traits reveals novel gene-metabolite-disease links. *PLoS Genet.*, **10**, e1004132.
22. Yu, B., Zheng, Y., Alexander, D., Morrison, A.C., Coresh, J. and Boerwinkle, E. (2014) Genetic determinants influencing human serum metabolome among African Americans. *PLoS Genet.*, **10**, e1004212.
23. Shin, S.-Y., Fauman, E.B., Petersen, A.-K., Krumsiek, J., Santos, R., Huang, J., Arnold, M., Erte, I., Forgetta, V., Yang, T.-P. et al. (2014) An atlas of genetic influences on human blood metabolites. *Nat. Genet.*, **46**, 543–550.
24. Ried, J.S., Shin, S.Y., Krumsiek, J., Illig, T., Theis, F.J., Spector, T.D., Adamski, J., Wichmann, H.E., Strauch, K., Soranzo, N. et al. (2014) Novel genetic associations with serum level metabolites identified by phenotype set enrichment analyses. *Hum. Mol. Genet.*, **23**, 5847–5857.
25. Demirkan, A., Henneman, P., Verhoeven, A., Dharuri, H., Amin, N., van Klinken, J.B., Karssen, L.C., de Vries, B., Meissner, A., Goraler, S. et al. (2015) Insight in genome-wide association of metabolite quantitative traits by exome sequence analyses. *PLoS Genet.*, **11**, e1004835.
26. Draisma, H.H., Pool, R., Kobl, M., Jansen, R., Petersen, A.K., Vaarhorst, A.A., Yet, I., Haller, T., Demirkan, A., Esko, T. et al. (2015) Genome wide association study identifies novel genetic variants contributing to variation in blood metabolite levels. *Nat. Commun.*, **6**, 7208.
27. Stiles, A.R., Kozlitina, J., Thompson, B.M., McDonald, J.G., King, K.S. and Russell, D.W. (2014) Genetic, anatomic, and clinical determinants of human serum sterol and vitamin D levels. *Proc. Natl. Acad. Sci. USA*, **111**, E4006–E4014.
28. Ng, E., Salihovic, S., Monica Lind, P., Mahajan, A., Syvanen, A.C., Axelsson, T., Ingelsson, E., Lindgren, C.M., van Bavel, B., Morris, A.P. et al. (2015) Genome-wide association study of plasma levels of polychlorinated biphenyls disclose an association with the CYP2B6 gene in a population-based sample. *Environ. Res.*, **140**, 95–101.
29. Xie, W., Wood, A.R., Lyssenko, V., Weedon, M.N., Knowles, J.W., Alkayyali, S., Assimes, T.L., Quertermous, T., Abbasi, F., Paananen, J. et al. (2013) Genetic variants associated with glycine metabolism and their role in insulin sensitivity and type 2 diabetes. *Diabetes*, **62**, 2141–2150.
30. Wu, J.H., Lemaitre, R.N., Manichaikul, A., Guan, W., Tanaka, T., Foy, M., Kabagambe, E.K., Djousse, L., Siscovick, D., Fretts, A.M. et al. (2013) Genome-wide association study identifies novel loci associated with concentrations of four plasma phospholipid fatty acids in the de novo lipogenesis pathway: results from the Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) consortium. *Circ. Cardiovasc. Genet.*, **6**, 171–183.
31. Iltis, J.N., Sebastiani, P., Solovieff, N., Hartley, S.W., Bhatnagar, P., Arking, D.E., Dworkis, D.A., Casella, J.F., Barron-Casella, E., Bean, C.J. et al. (2012) A genome-wide association study of total bilirubin and cholelithiasis risk in sickle cell anemia. *PLoS One*, **7**, e34741.
32. Chen, G., Ramos, E., Adeyemo, A., Shriner, D., Zhou, J., Doumatey, A.P., Huang, H., Erdos, M.R., Gerry, N.P., Herbert, A. et al. (2012) UGT1A1 is a major locus influencing bilirubin levels in African Americans. *Eur. J. Hum. Genet.*, **20**, 463–468.
33. Kang, T.W., Kim, H.J., Ju, H., Kim, J.H., Jeon, Y.J., Lee, H.C., Kim, K.K., Kim, J.W., Lee, S., Kim, J.Y. et al. (2010) Genome-wide association of serum bilirubin levels in Korean population. *Hum. Mol. Genet.*, **19**, 3672–3678.
34. Johnson, A.D., Kavousi, M., Smith, A.V., Chen, M.H., Dehghan, A., Aspelund, T., Lin, J.P., van Duijn, C.M., Harris, T.B., Cupples, L.A. et al. (2009) Genome-wide association meta-analysis for total serum bilirubin levels. *Hum. Mol. Genet.*, **18**, 2700–2710.
35. Kolz, M., Johnson, T., Sanna, S., Teumer, A., Vitart, V., Perola, M., Mangino, M., Albrecht, E., Wallace, C., Farrall, M. et al. (2009) Meta-analysis of 28,141 individuals identifies common variants within five new loci that influence uric acid concentrations. *PLoS Genet.*, **5**, e1000504.
36. Doring, A., Gieger, C., Mehta, D., Gohlke, H., Prokisch, H., Coassin, S., Fischer, G., Henke, K., Klopp, N., Kronenberg, F. et al. (2008) SLC2A9 influences uric acid concentrations with pronounced sex-specific effects. *Nat. Genet.*, **40**, 430–436.
37. Dehghan, A., Kottgen, A., Yang, Q., Hwang, S.J., Kao, W.L., Rivadeneira, F., Boerwinkle, E., Levy, D., Hofman, A., Astor, B.C. et al. (2008) Association of three genetic loci with uric acid concentration and risk of gout: a genome-wide association study. *Lancet*, **372**, 1953–1961.
38. Kottgen, A., Albrecht, E., Teumer, A., Vitart, V., Krumsiek, J., Hundertmark, C., Pistis, G., Ruggiero, D., O'Seaghdha, C.M., Haller, T. et al. (2013) Genome-wide association analyses identify 18 new loci associated with serum urate concentrations. *Nat. Genet.*, **45**, 145–154.
39. Seppala, I., Kleber, M.E., Lyytikainen, L.P., Hernesniemi, J.A., Makela, K.M., Oksala, N., Laaksonen, R., Pilz, S., Tomaschitz, A., Silbernagel, G. et al. (2014) Genome-wide association study on dimethylarginines reveals novel AGXT2 variants associated with heart rate variability but not with overall mortality. *Eur. Heart J.*, **35**, 524–531.
40. Kleber, M.E., Seppala, I., Pilz, S., Hoffmann, M.M., Tomaschitz, A., Oksala, N., Raitoharju, E., Lyytikainen, L.P., Makela, K.M., Laaksonen, R. et al. (2013) Genome-wide association study identifies 3 genomic loci significantly associated with serum levels of homoarginine: the AtheroRemo Consortium. *Circ. Cardiovasc. Genet.*, **6**, 505–513.
41. Chambers, J.C., Zhang, W., Lord, G.M., van der Harst, P., Lawlor, D.A., Sehmi, J.S., Gale, D.P., Wass, M.N., Ahmadi, K. R., Bakker, S.J. et al. (2010) Genetic loci influencing kidney function and chronic kidney disease. *Nat. Genet.*, **42**, 373–375.
42. Ghazalpour, A., Bennett, B.J., Shih, D., Che, N., Orozco, L., Pan, C., Hagopian, R., He, A., Kayne, P., Yang, W.P. et al. (2014) Genetic regulation of mouse liver metabolite levels. *Mol. Syst. Biol.*, **10**, 730.
43. Chen, W., Gao, Y., Xie, W., Gong, L., Lu, K., Wang, W., Li, Y., Liu, X., Zhang, H., Dong, H. et al. (2014) Genome-wide association analyses provide genetic and biochemical insights into natural variation in rice metabolism. *Nat. Genet.*, **46**, 714–721.
44. Wishart, D.S., Jewison, T., Guo, A.C., Wilson, M., Knox, C., Liu, Y., Djoumbou, Y., Mandal, R., Aziat, F., Dong, E. et al. (2013) HMDB 3.0—The Human Metabolome Database in 2013. *Nucleic Acids Res.*, **41**, D801–D807.
45. Tsepilov, Y.A., Shin, S.Y., Soranzo, N., Spector, T.D., Prehn, C., Adamski, J., Kastenmuller, G., Wang-Sattler, R., Strauch, K., Gieger, C. et al. (2015) Non-additive effects of genes in human metabolomics. *Genetics*, **200**, 707–718.



46. Krumsiek, J., Suhre, K., Illig, T., Adamski, J. and Theis, F.J. (2011) Gaussian graphical modeling reconstructs pathway reactions from high-throughput metabolomics data. *BMC Syst. Biol.*, **5**, 21.
47. Sainz, I.M., Pixley, R.A. and Colman, R.W. (2007) Fifty years of research on the plasma kallikrein-kinin system: from protein structure and function to cell biology and in-vivo pathophysiology. *Thromb. Haemost.*, **98**, 77–83.
48. Arnold, M., Raffler, J., Pfeufer, A., Suhre, K. and Kastenmuller, G. (2015) SNIIPA: an interactive, genetic variant-centered annotation browser. *Bioinformatics*, **31**, 1334–1336.
49. Ho, Y.Y., Baechler, E.C., Ortmann, W., Behrens, T.W., Graham, R.R., Bhangale, T.R. and Pan, W. (2014) Using gene expression to improve the power of genome-wide association analysis. *Hum. Hered.*, **78**, 94–103.
50. Rodionov, R.N., Jarzebska, N., Weiss, N. and Lentz, S.R. (2014) AGXT2: a promiscuous aminotransferase. *Trends Pharmacol. Sci.*, **35**, 575–582.
51. Kontani, Y., Kaneko, M., Kikugawa, M., Fujimoto, S. and Tamaki, N. (1993) Identity of D-3-aminoisobutyrate-pyruvate aminotransferase with alanine-glyoxylate aminotransferase 2. *Biochim. Biophys. Acta.*, **1156**, 161–166.
52. Harris, H. (1953) Family studies on the urinary excretion of beta-aminoisobutyric acid. *Ann. Eugen.*, **18**, 43–49.
53. Crumpler, H.R., Dent, C.E., Harris, H. and Westall, R.G. (1951) beta-Aminoisobutyric acid (alpha-methyl-beta-alanine); a new amino-acid obtained from human urine. *Nature*, **167**, 307–308.
54. Kittel, A., Muller, F., Konig, J., Mieth, M., Sticht, H., Zolk, O., Kralj, A., Heinrich, M.R., Fromm, M.F. and Maas, R. (2014) Alanine-glyoxylate aminotransferase 2 (AGXT2) polymorphisms have considerable impact on methylarginine and beta-aminoisobutyrate metabolism in healthy volunteers. *PLoS One*, **9**, e88544.
55. Nguyen, L.N., Ma, D., Shui, G., Wong, P., Cazenave-Gassiot, A., Zhang, X., Wenk, M.R., Goh, E.L. and Silver, D.L. (2014) Mfsd2a is a transporter for the essential omega-3 fatty acid docosahexaenoic acid. *Nature*, **509**, 503–506.
56. Owens, P.K., Raddad, E., Miller, J.W., Stille, J.R., Olovich, K.G., Smith, N.V., Jones, R.S. and Scherer, J.C. (2015) A decade of innovation in pharmaceutical R&D: the Chorus model. *Nat. Rev. Drug Discov.*, **14**, 17–28.
57. Plenge, R.M., Scolnick, E.M. and Altshuler, D. (2013) Validating therapeutic targets through human genetics. *Nat. Rev. Drug Discov.*, **12**, 581–594.
58. Westra, H.J., Peters, M.J., Esko, T., Yaghoobkar, H., Schurmann, C., Kettunen, J., Christiansen, M.W., Fairfax, B.P., Schramm, K., Powell, J.E. et al. (2013) Systematic identification of trans eQTLs as putative drivers of known disease associations. *Nat. Genet.*, **45**, 1238–1243.
59. Melzer, D., Perry, J.R., Hernandez, D., Corsi, A.M., Stevens, K., Rafferty, I., Lauretani, F., Murray, A., Gibbs, J.R., Paolisso, G. et al. (2008) A genome-wide association study identifies protein quantitative trait loci (pQTLs). *PLoS Genet.*, **4**, e1000072.
60. Lauc, G., Essafi, A., Huffman, J.E., Hayward, C., Knezevic, A., Kattla, J.J., Polasek, O., Gornik, O., Vitart, V., Abrahams, J.L. et al. (2010) Genomics meets glycomics—the first GWAS study of human N-glycome identifies HNF1alpha as a master regulator of plasma protein fucosylation. *PLoS Genet.*, **6**, e1001256.
61. Petersen, A.K., Zeilinger, S., Kastenmuller, G., Romisch-Margl, W., Brugger, M., Peters, A., Meisinger, C., Strauch, K., Hengstenberg, C., Pagel, P. et al. (2014) Epigenetics meets metabolomics: an epigenome-wide association study with blood serum metabolic traits. *Hum. Mol. Genet.*, **23**, 534–545.
62. Bartel, J., Krumsiek, J., Schramm, K., Adamski, J., Gieger, C., Herder, C., Carstensen, M., Peters, A., Rathmann, W., Roden, M. et al. (2015) The human blood metabolome-transcriptome interface. *PLoS Genet.*, **11**, e1005274.