



Published in final edited form as:

J Clin Immunol. 2015 August ; 35(6): 527–530. doi:10.1007/s10875-015-0177-x.

PLAID: A syndrome of complex patterns of disease and unique phenotypes

Joshua D. Milner

Genetics and Pathogenesis of Allergy Section, Laboratory of Allergic Diseases, NIAID, NIH, 10 Center Drive, NIH Building 10-CRC 5-3950, Bethesda, MD 20892, 3018273662

Abstract

PLCG2 associated antibody deficiency and immune dysregulation (PLAID) is a complex dominantly inherited disease characterized almost universally by cold urticaria, and variably by recurrent bacterial infection, autoimmunity and skin granuloma formation. Several striking phenotypes can emerge from this disease, and the pathophysiology leads to a complex mix of loss and gain of function in cellular signaling. This review discusses the key phenotypic characteristics and pathophysiologic observations seen in PLAID, and contrasts PLAID to several related disorders in order to best contextualize this fascinating disease.

Cold urticaria is a common physical urticaria typically characterized by classic-appearing hives elicited by contact with cold material. The response is rapid—with symptom onset within minutes, as is the resolution upon warming, although prolonged systemic cold exposure can lead to anaphylaxis. The onset of responsiveness to cold tends to happen in the second or third decade of life and can resolve after years of symptoms. The gold-standard for diagnosing this curious disorder is the elicitation of a hive after exposure to an ice cube. Little is known regarding etiology, except that serum transfer from affected individuals can lead to cold induced hives in unaffected recipients, and family history of atopy, although not cold-induced atopy, is frequently seen in some populations¹.

Familial cold urticaria was commonly used as a term to describe patients with FCAS due to NLRP3 mutations which usually lead to an excess of IL-1 production. NLRP3 mutations can also lead to other autoinflammatory conditions such as Schnitzler's syndrome, Muckle-Wells Syndrome, NOMID and others. The cold-induced “urticaria” observed in those patients is actually a neutrophilic infiltrate unrelated to mast cell degranulation, and is also associated with fever and inflammation. The disorder is inherited dominantly and symptoms were often triggered by cold exposures hours earlier². Delayed cold urticaria, which appears clinically more similar to typical cold urticaria except that symptom onset was hours after cold exposure, had also been described to be inherited in a dominant fashion³. Ghandi et al then described an immediate cold urticaria syndrome which was also inherited dominantly⁴. Mast cell degranulation was indeed seen in affected patients' cold-exposed skin. In contrast to typical cold urticaria, the syndrome seen in Ghandi et al was characterized by urticaria

from birth which did not resolve, a tendency to react to evaporative cooling more than contact with cold objects, and a negative ice cube test.

Subsequently, PLAID (PLCG2 associated antibody deficiency and immune dysregulation) was discovered after investigating an index patient who had diffuse granulomatous dermatitis which gradually worsened from birth.⁵

In addition to the granulomatous rash, further history showed that the index patient and other family members had other symptoms, inherited in an autosomal dominant fashion, the most common of which was an allergic reaction to cold. This index family was then compared to two additional families, some of members of whom were described in Ghandi et al, and a broad spectrum of disease was revealed^{5, 6}.

All patients with PLAID have an urticarial reaction to cold from infancy, including several patients who reported to having prolonged cyanosis when not placed rapidly under a warmer after birth. One patient was noted to first have symptoms when placed in a swing-- which generated a cool draft.

Evaporative cooling elicited the symptoms—breezes or exposure to air conditioning while perspiring were commonly described triggers, and patients had negative ice cube challenges. Cold swimming pools could trigger symptoms, and syncope was reported in situations of prolonged systemic cold exposure. Patients are more likely to have an erythematous, pruritic, localized rash in response to evaporative cooling which was not always raised, unlike typical hives.. Eating cold foods such as ice cream can trigger of burning sensations in the throat or retrosternal regions, however unlike food-induced anaphylaxis this reaction does not progress to throat closure or other systemic responses. ⁴

Two distinct cutaneous findings can occur in PLAID in addition to the acute urticarial responses. In a subset of patients, a blistering rash almost resembling a burn developed within the first few days of life at the tip of the nose, ears and fingers. In most of the patients in whom the rash appeared, there was spontaneous resolution. In a few, the rash actually worsened over time, and lead to soft tissue destruction of ear and nose cartilage, with sparing of axillary folds and other warmer areas of skin. In others who had the self-resolving neonatal rash, isolated granulomatous patches were developed later in life. Histopathological examination led to the diagnosis of sarcoidosis in at least one patient due to the presence of non-caseating granulomatous dermatitis, but there was little evidence suggest a clear etiology. ACE levels were all normal in PLAID patients, with and without granulomata⁶.

Patients with PLAID have a high frequency of positive anti-nuclear antibodies—up to 2/3 of the patients, while a number have clinically relevant autoimmunity in the form of autoimmune thyroiditis and vitiligo. Infectious phenotypes, when present include recurrent sinopulmonary infections varying from frequent colds and pharyngitis starting in childhood to the requirement of IVIG to treat recurrent pneumonias and bronchiectasis. Early-onset shingles and onychomycosis was noted in several patients as well⁵.

Aside from the urticaria, the other phenotypes were quite variable—multiple affected patients reported that cold-induced urticaria was their only symptom. Despite the

heterogeneity of symptoms, several laboratory measures were more consistent. In addition to the positive ANA, almost all patients had low serum IgM and circulating switched memory B-cells (IgM⁻, IgD⁻ CD27⁺). Many patients also had low IgA, and poor antibody responses to pneumococcal vaccines, as well as low or low-normal NK cells. CRP and ESR were elevated in the index patient with severe granulomata but this was not seen in any others⁵. In addition, cold agglutinins and cryoglobulins, seen in other states cold-induced inflammation, were universally negative.

Heterozygous genomic deletions in *PLCG2* are responsible for the majority of PLAID patients' disease. Phospholipase C gamma 2 catalyzes the hydrolysis of PIP₂ to IP₃ and DAG in response to receptor ligation in a variety of cell types. IP₃ then serves as a second messenger to release calcium from the endoplasmic reticulum and continue the cellular activation. While PLC1, a structural and functional homolog, is more ubiquitously expressed, PLC2 expression is largely limited to lymphoid and myeloid cells except for T-cells⁷. Deletion of *plcg2* in mice leads to poor calcium flux, poor long-term B-cell memory formation, the reduction of IgM, and several subclasses of IgG⁸. NK cell function is also impaired⁹, as is B-cell receptor editing¹⁰. Interestingly, two different mouse strains harboring *plcg2* gain of function mutations have been shown to develop dermatitis, autoinflammation and other phenotypes including, quite interestingly, skin inflammation which localizes to colder areas of the mouse body surface^{11, 12}. Patients with PLAID have phenotypes predicted to be due to both gain and loss of *PLCG2*, and as such, the mechanism by which PLAID-associated mutations lead to disease is complex. Most affected patients have a several kilobase genomic deletion spanning parts of an autoregulatory cSH2 domain in *PLCG2*. This cSH2 domain normally prevents enzymatic activity, and, upon receptor ligation, is thought to be displaced, revealing the active site of the enzyme⁷. Given the structural prediction, it was therefore not surprising that these deletions render the protein constitutively active in certain in vitro transfection conditions⁵. However, primary cells from the patients paradoxically have an anergic phenotype⁵ similar to that seen in *plcg2* null mice, when activated at body temperature. B-cell and NK cell activation were impaired, while T-cells, which do not express significant levels of *PLC2*, did not have a signaling defect. The loss of receptor-mediated responsiveness in *PLC2* expressing lymphocytes may be due to feedback inhibition secondary to the chronic activation, or to a direct role for the deleted regions in forming the signalosome necessary for normal receptor signal transduction.¹³

The cold urticaria in these patients likely results from the fact that mast cells expressing the mutant *PLCG2* spontaneously activate when exposed to lower temperatures in the absence of any antigen⁵. The temperatures which lead to activation in vitro are easily achieved in the surface of the skin with even mild cooling. It is not understood how the ice cube test could be negative in most PLAID patients, despite responses to evaporative or systemic cooling, nor is the molecular mechanism for how the cold immediately induces activation in *PLCG2* mutated patients. The complex intrinsic signaling abnormalities likely lead to temperature responses requiring a very narrow range, combining the inhibitory and excitatory states seen in cells from these patients.

Because the neonatal rash occurs in the coldest areas of the body surface, and the prolonged granulomatous rash seen in the severe cases spared warmer areas of the skin, it is quite

possible that other cell types may be activated by subtle drops in skin temperature in a more chronic fashion leading to the neonatal and chronic inflammation. Consistent with that clinical observation, neutrophils and monocytes from PLAID patients showed marked cold-induced activation. Furthermore it was not clear that neutrophils could produce superoxide in response to stimulation over and above the basal activation seen, making it possible that other neutrophil defects, in addition to cold-induced activation, could contribute to granuloma formation. It should be noted, however that the cold-induced activation of neutrophils and monocytes, as well as B-cells, was present in all patients, whether a granuloma was present or not^{5, 6}. Additional factors must modify the risk for granuloma formation, be they external temperature exposure variation, other intrinsic environmental exposures, or genetic modifiers.

Several families appear to have PLAID phenocopy without a detectable *PLCG2* deletion or mutation^{4, 6}. These patients' phenotypes suggest that the fascinating temperature-induced activation coupled with defective lymphocyte responses may be achievable through other genes in the signaling pathway, such that negative *PLCG2* sequencing can no longer be assumed to be definitive in ruling out a PLAID-like disease.

Treatment of PLAID patients centers around avoidance of evaporative or systemic cooling—this includes warming rapidly after showers, towel drying off sweat during and after exercise, avoiding drafts whenever possible and avoiding cold pools. Short and long-acting antihistamines can be effective as well. While the natural history of PLAID is limited to the 40–50 patients identified so far, it does not appear that the urticarial symptoms change much over time, rather affected individuals become more and more adept at avoiding the triggers. Otherwise healthy patients should be carefully monitored for changes in infection patterns, as CVID symptoms emerged only later in life in several patients. For those patients, the antibiotic prophylaxis or IVIG was determined by infectious history. The laboratory abnormalities are not as helpful, since immunoglobulin and specific antibody responses were so impaired in many otherwise asymptomatic patients. In the future, it may be possible to use *PLC2* inhibitors to restore normal function at body temperature and prevent spontaneous activation in the cold.

It should be mentioned that a heterozygous missense point mutation in *PLCG2* has been subsequently identified to cause APLAID (Autoinflammatory *PLCG2* associated antibody deficiency and immune dysregulation). APLAID was seen in a father and daughter early life onset of skin inflammation and granulomata, uveitis, colitis, lung inflammation (interstitial pneumonitis with respiratory bronchiolitis), and recurrent bacterial and viral sinopulmonary infections¹⁴. *Aspergillus* infection of the lung has been noted in one of the patients. Unlike PLAID patients, cold urticaria was not observed in either patient, and cellular signaling in B-cells and cells transfected with the mutant was enhanced at body temperature, with no response to cold. The skin lesions resembled sterile cellulitis, but granulomata of unknown etiology were also noted. Like other autoinflammatory diseases, symptoms have waxed and waned over the patients' lifetime. Similar to PLAID patients, both APLAID patients had low or low-normal serum IgA and IgM levels, poor responses to pneumococcal vaccine and nearly absent class-switched B-cells. However unlike in PLAID, the APLAID patients did not have substantial autoantibody formation. Inflammatory markers such as CRP, ESR and

others were normal. Why the point mutations in the same region lead to only minimal clinical overlap with the deletions seen in PLAID patients has yet to be understood.

Summary

Cold-induced allergic symptoms can be inherited in an autosomal dominant fashion and leads to distinct clinical phenotypes which can distinguish these patients from those with typical cold urticaria, and those with cold-induced autoinflammatory disorders. The mutations in PLAID lead to an intrinsic cold-induced activation which, depending on cell type, can lead to a variety of acute or chronic cold-induced phenotypes, including urticarial and granuloma. At the same time, a complex signaling defect exists, since at physiologic body temperature, signaling is impaired. This monogenic disease of atopy opens insight into mast cell and other leukocyte signaling and function, granuloma formation, and raises the notion that some mutations can lead to mendelian inheritance of acquired cold responsiveness in proteins not previously associated with cold responsiveness.

Acknowledgments

This research was supported by the Intramural Research Program of the NIH, NIAID

References

1. Abajian M, Mlynek A, Maurer M. Physical urticaria. *Current allergy and asthma reports*. 2012; 12(4):281–287. [PubMed: 22653630]
2. Mathur AN, Mathes EF. Urticaria mimickers in children. *Dermatologic therapy*. 2013; 26(6):467–475. [PubMed: 24552410]
3. Soter NA, Joshi NP, Twarog FJ, Zeiger RS, Rothman PM, Colten HR. Delayed cold-induced urticaria: a dominantly inherited disorder. *The Journal of allergy and clinical immunology*. 1977; 59(4):294–297. [PubMed: 66242]
4. Gandhi C, Healy C, Wanderer AA, Hoffman HM. Familial atypical cold urticaria: description of a new hereditary disease. *J Allergy Clin Immunol*. 2009; 124(6):1245–1250. [PubMed: 19910034]
5. Ombrello MJ, Remmers EF, Sun G, Freeman AF, Datta S, Torabi-Parizi P, et al. Cold urticaria, immunodeficiency, and autoimmunity related to PLCG2 deletions. *N Engl J Med*. 2012; 366(4):330–338. [PubMed: 22236196]
6. Aderibigbe OM, Priel DAL, Lee C-CR, Ombrello MJ, Prajapati VH, Liang MG, et al. Distinct cutaneous manifestations, autonomous and cold-induced leukocyte activation associated with PLCG2 mutations. *JAMA Dermatology*. 2015 in press.
7. Bunney TD, Esposito D, Mas-Droux C, Lamber E, Baxendale RW, Martins M, et al. Structural and functional integration of the PLCgamma interaction domains critical for regulatory mechanisms and signaling deregulation. *Structure*. 2012; 20(12):2062–2075. [PubMed: 23063561]
8. Wang D, Feng J, Wen R, Marine JC, Sangster MY, Parganas E, et al. Phospholipase Cgamma2 is essential in the functions of B cell and several Fc receptors. *Immunity*. 2000; 13(1):25–35. [PubMed: 10933392]
9. Tassi I, Presti R, Kim S, Yokoyama WM, Gilfillan S, Colonna M. Phospholipase C-gamma 2 is a critical signaling mediator for murine NK cell activating receptors. *J Immunol*. 2005; 175(2):749–754. [PubMed: 16002670]
10. Bai L, Chen Y, He Y, Dai X, Lin X, Wen R, et al. Phospholipase Cgamma2 contributes to light-chain gene activation and receptor editing. *Mol Cell Biol*. 2007; 27(17):5957–5967. [PubMed: 17591700]
11. Abe K, Fuchs H, Boersma A, Hans W, Yu P, Kalaydjiev S, et al. A novel ENU-induced mutation in Phospholipase C gamma 2 causes inflammatory arthritis, metabolic defects, and in vitro male infertility in the mouse. *Arthritis Rheum*. 2011

12. Yu P, Constien R, Dear N, Katan M, Hanke P, Bunney TD, et al. Autoimmunity and inflammation due to a gain-of-function mutation in phospholipase C gamma 2 that specifically increases external Ca²⁺ entry. *Immunity*. 2005; 22(4):451–465. [PubMed: 15845450]
13. Wang J, Sohn H, Sun G, Milner JD, Pierce SK. The autoinhibitory C-terminal SH2 domain of phospholipase C-gamma2 stabilizes B cell receptor signalosome assembly. *Science signaling*. 2014; 7(343):ra89. [PubMed: 25227611]
14. Zhou Q, Lee GS, Brady J, Datta S, Katan M, Sheikh A, et al. A Hypermorphic Missense Mutation in PLCG2, Encoding Phospholipase Cgamma2, Causes a Dominantly Inherited Autoinflammatory Disease with Immunodeficiency. *Am J Hum Genet*. 2012; 91(4):713–720. [PubMed: 23000145]

Appendix

Questions:

1. Which of the following statements regarding features in familial cold-induced autoinflammatory syndromes (also called familial cold urticaria) and PLAID are true:
 - A. Mutations in the causal gene for each disorder can lead to highly variable phenotypes
 - B. A subset of FCAS patients have typical mast-cell induced cold urticaria which resembles PLAID
 - C. PLAID patients do not respond to antihistamines, while FCAS patients do
 - D. FCAS patients' urticaria differs from PLAID patients' in that symptom onset and resolution are delayed, but the symptoms themselves are otherwise fairly similar
 - E. The histopathology of cold induced lesions is similar in FCAS and PLAID

Answer: A—NLRP3 mutations lead to multiple distinct syndromes, some of which include symptoms which are triggered by cold and some which are not, and organ system involvement can vary. PLCG2 mutations can lead to PLAID and APLAID which are quite distinct.

2. Features which distinguish PLAID from typical cold urticaria include
 - A. Both are usually characterized by neonatal onset, but patients with typical cold urticaria resolve at some point in time
 - B. Both are usually characterized by mendelian patterns of familial inheritance but PLAID tends to occur in a dominant fashion
 - C. Ice cube testing leads to raised hives in typical cold urticaria, and to pruritis and erythema, but not raised hives, in PLAID patients
 - D. Patients with PLAID can have multiple other co-morbidities, while patients with typical cold urticaria usually do not.
 - E. Both are characterized by cold-induced symptoms but only typical urticaria leads to anaphylaxis.

Answer D. PLAID patients can develop autoimmunity, granulomata and infection, while cold urticaria patients have few if any significant comorbidities.

3. Patients with mutations in PLAID have highly variable presentations with respect to which of the following phenotypes:
 - A. Cold urticaria
 - B. Immune deficiency
 - C. Laboratory measures of B-cell function
 - D. Reaction to ice cube test
 - E. Anatomical location of skin rashes

Answer: B The other symptoms are quite stable but only fraction have significant immune deficiency, and a larger fraction, but not all, have recurrent mild infections.

4. Enhanced antigen or Fc-receptor mediated signaling is seen in:
 - A. Many FCAS patients' cells
 - B. PLAID B-cells, but not mast cells, at body temperature
 - C. APLAID patient B-cells at physiologic body temperature
 - D. PLAID mast cells only after antigenic stimulation in the cold
 - E. APLAID lymphocytes exposed to cold

Answer: C. Only APLAID B-cells have been shown to exhibit increased BCR-mediated signal at physiologic temperatures.

5. 5. Laboratory abnormalities which can often be seen in PLAID include
 - A. Elevated ESR
 - B. Positive Cryoglobulins
 - C. Positive Cold Agglutinins
 - D. Low class-switched B-cells
 - E. Absent NK cells

Answer: D. Nearly all patients had marked diminution of class-switched B-cells.