

First Draft Genome Sequence of *Balamuthia mandrillaris*, the Causative Agent of Amoebic Encephalitis

H. Detering,^{a,b} T. Aebischer,^b P. W. Dabrowski,^{a,c} A. Radonić,^c A. Nitsche,^c B. Y. Renard,^a A. F. Kiderlen^b

Research Group Bioinformatics (NG4),^a Division for Mycotic, Parasitic and Mycobacterial Diseases (FG16),^b and Centre for Biological Threats and Special Pathogens (ZBS1),^c Robert Koch Institute, Berlin, Germany

The free-living amoeba *Balamuthia mandrillaris* is a rare but highly lethal agent of amoebic encephalitis in humans and many other mammalian species. Here, we announce the first draft genome sequence of the original 1990 isolate cultured from the brain of a deceased mandrill baboon.

Received 7 August 2015 Accepted 11 August 2015 Published 24 September 2015

Citation Detering H, Aebischer T, Dabrowski PW, Radonić A, Nitsche A, Renard BY, Kiderlen AF. 2015. First draft genome sequence of *Balamuthia mandrillaris*, the causative agent of amoebic encephalitis. *Genome Announc* 3(5):e01013-15. doi:10.1128/genomeA.01013-15.

Copyright © 2015 Detering et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to A. F. Kiderlen, kiderlena@rki.de.

Balamuthia mandrillaris is an opportunistic parasite that causes amoebic encephalitis (BAE) in humans and other mammals (1, 2), with a fatal outcome rate of >95%. The amoeba appears to be ubiquitous in soil (3–5) and freshwater habitats (6), and it invades its hosts via pulmonary (7), nasal/olfactory nerve (8), cutaneous (9), and possibly gastrointestinal (10) routes. BAE has been reported in immunocompromised (11) patients and in children and adults without clinical evidence for acquired or innate immunodeficiency (12, 13). Experimental infections in mice lacking lymphocyte subsets show that CD4⁺ T cells are important for resistance to intranasal *B. mandrillaris* infection and BAE (14).

Axenic cultivation of the CDC-V039 isolate was performed in modified Chang's special medium at 37°C in a humidified normal atmosphere supplemented with 5% CO₂ (15). Genomic DNA was isolated with the Qiagen DNeasy blood and tissue kit, according to the manufacturer's instructions. Genome sequencing was performed at the Robert Koch Institute on Roche 454 FLX, Illumina HiScan, and Illumina HiSeq 1500 machines and on a Pacific Biosciences RS II at the University of Lausanne (UNIL), Switzerland, with 20×, 1,600×, and 90× read coverage, respectively. Genome assembly of 756,805 PacBio reads with the HGAP3 (16) pipeline, followed by quality filtering of contigs, resulted in a genome of 68 Mbp in 1,605 contigs, with an *N*₅₀ value of 93,953 bp and an average G+C content of 46.8%. An integrated analysis using Illumina, 454, and PacBio data revealed a complex genome with indication of hyperploidy. A comparison of the mitochondrial genome structure and homology searches of *Acanthamoeba castellanii* protein sequences in the newly assembled genome suggest a more distant relationship of *B. mandrillaris* to its closest known relative than was previously assumed. Functional genome annotation is in progress and will help unravel the metabolic and pathogenic potential of this opportunistic parasite.

Nucleotide sequence accession numbers. The draft genome sequence of *B. mandrillaris* CDC-V039 has been deposited in GenBank under the accession no. [LFUI00000000](https://www.ncbi.nlm.nih.gov/nuclink/LFUI00000000). The version described in this paper is the first version, LFUI01000000.

ACKNOWLEDGMENTS

We thank Julia Hinzmann, Elke Radam, and Julia Tesch for excellent technical assistance and the staff of the Lausanne Genomic Technologies Facility Center for Integrative Genomics, especially Emmanuel Beaudoin, Mélanie Dupasquier, and Keith Harshman, for their swift and constructive support during PacBio sequencing and assembly.

We received no external funding in support of this work.

REFERENCES

1. Visvesvara GS, Martínez AJ, Schuster FL, Leitch GJ, Wallace SV, Sawyer TK, Anderson M. 1990. Leptomyxid amoeba. A new agent of amoebic meningoencephalitis in humans and animals. *J Clin Microbiol* 28:2750–2756.
2. Visvesvara GS, Schuster FL, Martínez AJ. 1993. *Balamuthia mandrillaris*, N. G., N. Sp., agent of amoebic meningoencephalitis in humans and other animals. *J Euk Microbiol* 40:504–514.
3. Dunnebacke TH, Schuster FL, Yagi S, Booton GC. 2004. *Balamuthia mandrillaris* from soil samples. *Microbiology* 150:2837–2842. <http://dx.doi.org/10.1099/mic.0.27218-0>.
4. Niyiyati M, Lorenzo-Morales J, Rezaeian M, Martin-Navarro CM, Haghi AM, Maciver SK, Valladares B. 2009. Isolation of *Balamuthia mandrillaris* from urban dust, free of known infectious involvement. *Parasitol Res* 106:279–281. <http://dx.doi.org/10.1007/s00436-009-1592-9>.
5. Schuster FL, Dunnebacke TH, Booton GC, Yagi S, Kohlmeier CK, Glaser C, Vugia D, Bakardjiev A, Azimi P, Maddux-Gonzalez M, Martínez AJ, Visvesvara GS. 2003. Environmental isolation of *Balamuthia mandrillaris* associated with a case of amoebic encephalitis. *J Clin Microbiol* 41:3175–3180. <http://dx.doi.org/10.1128/JCM.41.7.3175-3180.2003>.
6. Lares-Jiménez LF, Booton GC, Lares-Villa F, Velázquez-Contreras CA, Fuerst PA. 2014. Genetic analysis among environmental strains of *Balamuthia mandrillaris* recovered from an artificial lagoon and from soil in Sonora, Mexico. *Exp Parasitol* 145:S57–S61. <http://dx.doi.org/10.1016/j.exppara.2014.07.007>.
7. Martínez AJ, Visvesvara GS. 1997. Free-living, amphizoic and opportunistic amoebas. *Brain Pathol* 7:583–598. <http://dx.doi.org/10.1111/j.1750-3639.1997.tb01076.x>.
8. Kiderlen AF, Laube U. 2004. *Balamuthia mandrillaris*, an opportunistic agent of granulomatous amoebic encephalitis, infects the brain via the olfactory nerve pathway. *Parasitol Res* 94:49–53.
9. Bravo F, Sanchez MR. 2003. New and re-emerging cutaneous infectious diseases in Latin America and other geographic areas. *Dermatol Clin* 21:655–668. [http://dx.doi.org/10.1016/S0733-8635\(03\)00090-1](http://dx.doi.org/10.1016/S0733-8635(03)00090-1).
10. Kiderlen AF, Laube U, Radam E, Tata PS. 2007. Oral infection of immunocompetent and immunodeficient mice with *Balamuthia mandril-*

- laris amoebae*. Parasitol Res 100:775–782. <http://dx.doi.org/10.1007/s00436-006-0334-5>.
11. Anzil AP, Rao C, Wrzolek MA, Visvesvara GS, Sher JH, Kozlowski PB. 1991. Amebic meningoencephalitis in a patient with AIDS caused by a newly recognized opportunistic pathogen. Arch Pathol Lab Med 115:21–25.
 12. Bakardjiev A, Azimi PH, Ashouri N, Ascher DP, Janner D, Schuster FL, Visvesvara GS, Glaser C. 2003. Amebic encephalitis caused by *Balamuthia mandrillaris*: report of four cases. Pediatr Infect Dis J 22:447–452. <http://dx.doi.org/10.1097/01.inf.0000066540.18671.f8>.
 13. Yamasaki K, Sugimoto T, Futami M, Moriyama T, Uehara H, Takeshima H, Moriguchi S, Marutsuka K, Asada Y. 2011. Granulomatous amoebic encephalitis caused by *Balamuthia mandrillaris*. Neurol Med Chir (Tokyo) 51:667–670.
 14. Kiderlen AF, Radam E, Laube U, Martínez AJ. 2015. Resistance to intranasal infection with *Balamuthia mandrillaris* amoebae is T-cell dependent. J Eukaryot Microbiol 62:26–33. <http://dx.doi.org/10.1111/jeu.12189>.
 15. Kiderlen AF, Tata PS, Özel M, Laube U, Radam E, Schäfer H. 2006. Cytopathogenicity of *Balamuthia mandrillaris*, an opportunistic causative agent of granulomatous amoebic encephalitis. J Eukaryot Microbiol 53: 456–463. <http://dx.doi.org/10.1111/j.1550-7408.2006.00130.x>.
 16. Chin C, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. Nat Methods 10:563–569. <http://dx.doi.org/10.1038/nmeth.2474>.