

Review

Astaxanthin as a Potential Neuroprotective Agent for Neurological Diseases

Haijian Wu ¹, Huanjiang Niu ¹, Anwen Shao ², Cheng Wu ¹, Brandon J. Dixon ³, Jianmin Zhang ², Shuxu Yang ^{1,*} and Yirong Wang ^{1,*}

¹ Department of Neurosurgery, Sir Run Run Shaw Hospital, School of Medicine, Zhejiang University, Hangzhou 310016, China; E-Mails: haijwu@sina.com (H.W.); huanjiangniu@163.com (H.N.); chengwusy@sina.com (C.W.)

² Department of Neurosurgery, Second Affiliated Hospital, School of Medicine, Zhejiang University, Hangzhou 310009, China; E-Mails: awshao@sina.com (A.S.); zjmvip135@sina.com (J.Z.)

³ Department of Physiology and Pharmacology, School of Medicine, Loma Linda University, Loma Linda, CA, 92350, USA.; E-Mail: Bjdixon@llu.edu

* Authors to whom correspondence should be addressed; E-Mails: yangsxxy@163.com (S.Y.); srrsywang@sina.com (Y.W.); Tel./Fax: +86-571-8600-6166 (S.Y. and Y.W.).

Academic Editor: George Perry

Received: 6 June 2015 / Accepted: 7 September 2015 / Published: 11 September 2015

Abstract: Neurological diseases, which consist of acute injuries and chronic neurodegeneration, are the leading causes of human death and disability. However, the pathophysiology of these diseases have not been fully elucidated, and effective treatments are still lacking. Astaxanthin, a member of the xanthophyll group, is a red-orange carotenoid with unique cell membrane actions and diverse biological activities. More importantly, there is evidence demonstrating that astaxanthin confers neuroprotective effects in experimental models of acute injuries, chronic neurodegenerative disorders, and neurological diseases. The beneficial effects of astaxanthin are linked to its oxidative, anti-inflammatory, and anti-apoptotic characteristics. In this review, we will focus on the neuroprotective properties of astaxanthin and explore the underlying mechanisms in the setting of neurological diseases.

Keywords: astaxanthin; oxidative stress; inflammation; apoptosis; neuroprotection; neurological diseases

1. Introduction

Neurological diseases, exemplified by acute injuries (e.g., stroke and traumatic brain injury) and chronic neurodegeneration (e.g., Alzheimer's disease, Parkinson's disease, and Huntington's disease), are common causes of human death and disability [1,2]. Oxidative stress, inflammation, and apoptosis are some of the mechanisms involved in the pathogenesis of these diseases [3,4]. For example, highly insoluble amyloid beta peptide deposits and neurofibrillary tangles provide obvious stimuli for oxidative stress and inflammation in a brain with Alzheimer's disease (AD), which significantly contributes to neuronal death in this disease [5–7]. In addition, there is evidence demonstrating that mitochondrial deficits, oxidative and nitrosative stress, accumulation and aggregation of aberrant or misfolded proteins (*i.e.*, α -synuclein), and dysfunction of ubiquitin-proteasome system represents the principal molecular events that commonly underlie the pathogenesis of familial and sporadic forms of Parkinson's disease (PD) [8,9]. Additionally, highly polymorphic CAG tri-nucleotide repeat expansions in exon-1 of the *huntingtin* gene encodes an abnormally long poly-glutamine repeat, which is associated with Huntington's disease (HD)-related brain pathology [10]. Poly-glutamine expansion causes huntingtin to aggregate and accumulate in the nucleus. This leads to abnormal interactions with other proteins, which results in intra-nuclear accumulation of mutant huntingtin and the formation of neuropil aggregates that may ultimately lead to neuronal cell death [11]. Therefore, multi-targeted pharmacological agents may be effective for the treatment of these devastating diseases.

Astaxanthin, a unique member of the xanthophylls, is a deep red-colored phytonutrient that can be synthesized by a microalgae called *Haematococcus pluvialis* [12]. Distinct from other members of the xanthophylls, astaxanthin has two hydroxyl groups [13]. Astaxanthin spans the bi-lipid layer and is long enough that the two hydroxyl groups jut into the fluid phase near the membrane, and that when electrons are extracted from these hydroxyl groups by free radicals, the molecule is resonance stabilized. As a consequence, these properties allow astaxanthin to do a lot in the body. For instance, astaxanthin can dramatically decrease the risk of cardiovascular disease [14]. A diet supplemented with astaxanthin (75 or 200 mg/kg body weight per day) for 8 weeks has been shown to improve endothelium-dependent vasodilatation in resistance vessels, reduce systolic blood pressure, and improve cardiovascular remodeling in spontaneously hypertensive rats [15]. In addition, astaxanthin (100 and 500 mg/100 g) for 60 days protects against serum protein oxidation in hyper-cholesterolemic rabbits [16]. Studies have also demonstrated that astaxanthin can easily cross the BBB to protect the brain from acute injury and chronic neurodegeneration [17,18]. The neuroprotective properties of this molecule involves anti-oxidation, anti-inflammation, and anti-apoptosis [19–21]. Thus, this review article will focus on the beneficial effects of astaxanthin and explore the underlying mechanisms observed in experimental models of neurological diseases. We also propose that further studies involving astaxanthin are needed, in order to evaluate its potential application in the treatment of neurological disorders.

2. Astaxanthin: Source, Biochemistry, Bioavailability, and Safety

Xanthophyll is a class of oxygen-containing carotenoid pigments whose biosynthesis in plants derives from the lycopenene. Astaxanthin is a reddish pigment which belongs to the xanthophyll family [22].

This compound naturally exists in a wide variety of living organisms which includes microalgae, complex plants, and seafood [23]. The commercial form of astaxanthin is mainly synthesized from the algae *Haematococcus pluvialis* and the yeast *Phaffia rhodozyma*. As a member of the xanthophyll group, astaxanthin is closely related to other carotenoids such as β -carotene, lutein, and zeaxanthin. As a member of the xanthophyll group, astaxanthin is closely related to other carotenoids such as β -carotene, lutein, and zeaxanthin [24]. Similarly, they share many of the physiological and metabolic functions attributed to carotenoids [25]. Unlike β -carotene, astaxanthin does not have pro-vitamin A activity in the human body [26].

The molar mass of astaxanthin is 596.84 g/mol and the molecular formula is $C_{40}H_{52}O_4$. It is a symmetric molecule consisting of two terminal rings joined by a short polyene ring [22]. The hydroxyl group at the end of the molecule enables it to esterify fatty acids to form mono-esters or di-esters [13]. Natural astaxanthin mainly exists in an esterified form, while the synthetic form is produced in a free form [27]. Astaxanthin also contains conjugated double bonds, giving this molecule strong anti-oxidant properties by donating electrons and reacting with free radicals to terminate free radical chain reactions within cells [25,28].

Astaxanthin has both lipophilic and hydrophilic properties, since it is fat-soluble and can be carried by fat molecules directly to tissues and organs that need it the most, like the brain, retina, and skeletal muscle [22]. Astaxanthin is first absorbed into enterocytes through passive diffusion and undergoes facilitated diffusion in the presence of lipids [29]. The unesterified forms are incorporated into chylomicrons and are transported into the liver via the lymphatic system [30]. The liver does not biochemically convert these molecules into vitamin A [31]. Instead, it is incorporated into lipoproteins that are transported into organs and tissues via the circulation [32].

Astaxanthin is safe to consume with food and contains no reports of side effects [33,34]. One randomized clinical trial found that 6 mg/day of astaxanthin can be safely consumed by healthy adults [35]. In addition, numerous human clinical trials have shown that the astaxanthin rich extract, *Haematococcus pluvialis*, is safe as well [36,37]. Hoffman-La Roche confirmed the safety of astaxanthin with acute, mutagenicity, teratogenicity, embryotoxicity, and reproductive toxicity tests [38]. In addition, the United States Food and Drug Administration approved the use of astaxanthin as a dietary supplement in 1999 [25].

3. Neuroprotective Properties of Astaxanthin in Neurological Diseases

There have been numerous studies concerning the beneficial effects of astaxanthin. Astaxanthin-mediated neuroprotection in experimental models of neurological disorders involves anti-oxidation, anti-inflammation, and anti-apoptotic mechanisms [17,39]. The following sections will delve into these molecular mechanisms and their potential as treatments for neurological diseases.

3.1. Anti-Oxidant Effects

Oxidative stress is a key mediator in the pathology of neurological diseases [40–42]. Disturbance of the equilibrium status of pro-oxidant/anti-oxidant reactions in cells can lead to oxidative stress, which causes generation of reactive oxygen species (ROS) and free radicals [43]. When produced in excessive amounts, ROS like the superoxide anion radical (O_2^-) and its dismutation product, hydrogen peroxide

(H₂O₂), are detrimental to metabolic functions [44,45]. The O₂⁻ radical can oxidize the [4Fe-4S] clusters of dehydratases, such as aconitase, causing inactivation and release of Fe²⁺ [46,47]. Thereafter, Fe²⁺ reacts with H₂O₂ to yield the potent oxidizing free radical species hydroxyl radical (OH). These substances further react with key organic substrates, such as DNA, proteins, and lipids, to disturb cell function and cause cell death [48]. It is worth mentioning that astaxanthin can act as a safeguard against oxidative damage through various mechanisms, by quenching of singlet oxygen, scavenging of radicals, inhibiting lipid peroxidation, and regulating gene expression related to oxidative stress [49–52]. For example, astaxanthin exerts beneficial effects against HgCl₂-induced acute renal failure by preventing lipid and protein oxidation [53]. In an *in vivo* murine model, astaxanthin administration prevented *N*-Methyl-D-aspartate (NMDA)-triggered retinal damage, which is associated with decreasing lipid peroxidation and oxidative DNA damage [54]. Astaxanthin treatment ameliorates cyclophosphamide-induced oxidative stress and the subsequent DNA damage in rat hepatocytes [55]. The protective effect of this molecule is attributed to the activation of nuclear erythroid 2-related factor 2 (Nrf2) antioxidant response element (ARE) pathway, which eventually facilitates Nrf2-dependent gene expression of heme oxygenase-1 (HO-1) and NAD(P)H: quinone oxidoreductase-1 (NQO-1) [55]. In the human retinal pigment epithelial (RPE) cell line ARPE-19, astaxanthin inhibited the intracellular production of ROS and prevented H₂O₂-induced decrease in retinal pigment epithelial cell viability [56]. Astaxanthin also increased the nuclear translocation of Nrf2 and enhanced the expression of phase II anti-oxidant enzymes through the activation of the phosphoinositide 3-kinase (PI3K)/Akt pathway, which eventually provided protection against H₂O₂-induced oxidative stress in ARPE-19 cells [56].

The anti-oxidative effects of astaxanthin have also been investigated in experimental models of acute neurological conditions (Figure 1). Lee *et al.* reported that astaxanthin provides neuroprotective effects against oxidative stress induced by oxygen-glucose deprivation in SH-SY5Y cells and 10-min global cerebral ischemia in rats [57]. In a murine model of ischemic stroke, pre-treatment with astaxanthin decreased ROS production and alleviated lipid peroxidation in the ipsilateral brain of rats subjected to middle cerebral artery occlusion (MCAO) [17]. Simultaneously, astaxanthin reduced cerebral infarction and promoted locomotor function recovery following MCAO [17]. Zhang *et al.*, found that administration of astaxanthin had the potential of alleviating early brain injury (EBI) after subarachnoid hemorrhage (SAH) through its anti-oxidative properties [19]. Treatment with astaxanthin is believed to confer protective effects by restoring endogenous anti-oxidant enzymes of glutathione (GSH) and superoxide dismutase (SOD) following SAH [19]. Wu *et al.* reported that post-SAH treatment of astaxanthin facilitated the Nrf2-ARE pathway and ameliorated EBI in a prechiasmatic cistern model of SAH [58]. Astaxanthin activated the Nrf2-ARE signaling pathway to up-regulate the expression of Nrf2-regulated enzymes like HO-1, NQO-1 and glutathione-S-transferase- α 1 (GST- α 1) to resist oxidative stress [58].

Astaxanthin also plays a role in preventing the development of chronic neurodegeneration. It boosted the expression of HO-1 and protected neurons against A β -induced cytotoxicity [59,60]. Astaxanthin-stimulated activation of extracellular regulated protein kinase (ERK) signaling pathway facilitated the dissociation of Nrf2 from Keap1, promoting the nuclear translocation and DNA-binding activity of Nrf2 leading to up-regulation of HO-1 expression and protection against A β -induced neurotoxicity [59]. In a cellular model of PD, astaxanthin reduced the generation of intracellular ROS and provided cytoprotective effects against 1-methyl-4-phenylpyridinium (MPP⁺)-induced

cytotoxicity [61]. In addition, astaxanthin enhanced HO-1 expression and limited NADPH oxidase 2 (NOX2)-mediated oxidative damage in MPP⁺-treated PC12 cells [62]. Astaxanthin antagonized MPP⁺-induced oxidative stress through the regulation of specificity protein 1 (Sp1) and NMDA receptor subunit 1 (NR1) signaling pathway [63]. Pre-treatment with astaxanthin markedly inhibited the up-regulation and nuclear transfer of Sp1, thereby alleviated MPP⁺-induced production of intracellular ROS and cytotoxicity in PC12 cells [63]. Thus, astaxanthin provides protection against oxidative attacks in experimental neurological diseases.




Figure 1. The anti-oxidative effects of astaxanthin in neurological diseases. Astaxanthin facilitates the dissociation and nuclear translocation of nuclear erythroid 2-related factor (Nrf2), through activation of the PI3K/Akt and ERK signaling pathways, which contributes to increased expression of Nrf2-regulated enzymes like HO-1, NQO-1, and GST- α 1 that resist oxidative stress. In addition, astaxanthin negatively regulates Sp1/NR1 signaling pathway, alleviating the production of intracellular ROS and oxidative stress.

3.2. Anti-Inflammatory Effects

Inflammation is defined as series of complex immune responses that biologically occurs as a reaction to injuries of the body. It functions as a host defense mechanism to clear out damaged tissue from the original insult and initiates the tissue repair process [64]. However, excessive or uncontrolled inflammation is detrimental to the host and can cause damage to the host's cells and tissues [65]. In the central nervous system (CNS), inflammation has a critical role in both acute conditions (*i.e.*, stroke and traumatic injury) and chronic neurodegenerative conditions (e.g., AD, PD, and HD) [66]. Interestingly,

astaxanthin exhibits anti-inflammatory effects in lipopolysaccharide-induced uveitis by directly blocking the activity of inducible nitric oxide synthase (NOS) (Figure 2) [67]. In addition, astaxanthin suppressed gene expression of inflammatory mediators (*i.e.*, TNF- α and IL-1 β) and alleviated endotoxin-induced uveitis by blocking the NF- κ B-dependent signaling pathway [68]. Under normal conditions NF- κ B, a heterodimer composed of p50 and p65 subunits, interacts with inhibitor of NF- κ B (I κ B) and remains inactive in the cytosol [69]. Upon stimulation, I κ B undergoes phosphorylation by I κ B kinase β (IKK β) and is degraded via the ubiquitin proteasome pathway [70,71]. Dissociation of I κ B from the p50/p65 heterodimer exposes the nuclear localization signal on NF- κ B, which subsequently leads to the translocation of NF- κ B (p65) into the nucleus to regulate the transcription of inflammatory genes [72]. Astaxanthin treatment effectively alleviated NF- κ B-related inflammation in the liver of mice subjected to a high fructose and high fat diet by suppressing IKK β phosphorylation and nuclear translocation of NF- κ B (p65) subunit [73]. Astaxanthin also suppressed ROS-induced nuclear expression of NF- κ B (p65) and reduced the downstream production of pro-inflammatory cytokines (*i.e.*, IL-1 β , IL-6 and TNF- α) in U937 mononuclear cells by restoring the physiological levels of protein tyrosine phosphatase-1 (SHP-1) [74]. In a mouse model of experimental choroidal neovascularization, Izumi-Nagai demonstrated that astaxanthin treatment led to significant inhibition of macrophage infiltration into choroidal neovascularization [75]. Furthermore, astaxanthin suppressed I κ B- α degradation and NF- κ B nuclear translocation, resulting in subsequent down-regulation of inflammatory molecules (*i.e.*, IL-6, vascular endothelial growth factor (VEGF), intercellular adhesion molecule-1 (ICAM-1), and monocyte chemotactic protein 1 (MCP1) [75]. Astaxanthin also decreased gastric inflammation in mice infected with *Helicobacter pylori*, shifting the T-lymphocyte response from a Th1 response to a Th1/Th2 response [76]. Additionally, astaxanthin decreased nitric oxide (NO) production and inducible nitric oxide synthase (iNOS) activity in macrophages, resulting in inhibition of cyclooxygenase and down-regulation of prostaglandin E2 (PGE2) and TNF- α in mice [67]. Dietary administration of astaxanthin significantly suppressed aberrant NF- κ B activation in colonic mucosa, lowering gene expressions of IL-1 β , IL-6, and COX-2, which contributes to attenuation of dextran sulfate sodium (DSS)-induced colitis [77]. Lee and colleagues discovered that astaxanthin prevented inflammatory processes by suppressing the activation of NF- κ B signaling and the production of pro-inflammatory cytokines (e.g., TNF- α and IL-1 β) using both *in vitro* and *in vivo* models [78]. In human keratinocytes, Terazawa *et al.* demonstrated that astaxanthin interrupts the auto-phosphorylation and self-activation of mitogen- and stress-activated protein kinase-1 (MSK1), which results in decreased phosphorylation of NF- κ B (p65) and deficiency of NF- κ B DNA binding activity [79]. As a consequence, UVB-induced expression and secretion of PGE2 and IL-8 were down-regulated in these human keratinocytes [79].

In a prechiasmatic cistern SAH model, astaxanthin provides neuroprotection against EBI through suppression of cerebral inflammation [20]. Post-treatment with astaxanthin after SAH reduced neutrophil infiltration, suppressing the activity of NF- κ B, decreasing the protein and mRNA levels of inflammatory mediators IL-1 β , TNF- α , and ICAM-1, and dramatically reversed brain inflammation [20]. As a result secondary brain injury cascades, neuronal degeneration, BBB disruption, cerebral edema, and neurological dysfunction, were all alleviated after astaxanthin administration [20]. However, there is still a lack of research documenting the anti-inflammatory effects of astaxanthin on the treatment of neurological disorders. Several studies have reported that astaxanthin can enhance both humoral and

cell-mediated immune responses [26,80–83]. Dietary supplement of astaxanthin can stimulate T cell and B cell mitogen-induced lymphocyte proliferation, increase the cytotoxic activity of natural killer cell, and enhance IFN- γ and IL-6 production in young healthy adult female human subjects [84]. Additionally, Baliotti *et al.* showed gender-related differences in the anti-inflammatory effects of astaxanthin on the aging rat brain [85]. However, it is still unknown if this molecule exerts different anti-inflammatory effects in female and male brains under pathological conditions. Therefore, there is a need for future studies elucidating the inflammatory regulation mechanisms of astaxanthin.




Figure 2. The anti-inflammatory effects of astaxanthin in neurological diseases. Through suppression of I κ B- α degradation and NF- κ B nuclear translocation, astaxanthin inhibits the expression of inflammatory molecules IL-6, ICAM-1, and MCP1. Astaxanthin also suppresses nuclear expression of NF- κ B and reduces downstream production of pro-inflammatory cytokines by restoring physiological levels of SHP-1.

3.3. Anti-Apoptotic Effects

Apoptosis is a highly sophisticated energy-dependent process of programmed cell death [86]. Morphologically, it is characterized by shrinkage of the cell, membrane blebbing, nuclear fragmentation and chromatin condensation [87]. Under normal physiological conditions, apoptosis is vital for embryonic development and tissue homeostasis [88]. Under pathological conditions, uncontrolled apoptosis is harmful and contributes to the pathogenesis of a variety of human diseases including neurological disorders [89]. Kim *et al.* demonstrated that astaxanthin provided protection against H₂O₂-mediated apoptosis in a mouse neural progenitor cell culture model [90]. Astaxanthin is believed

to inhibit H₂O₂-mediated apoptotic cell death by maintaining mitochondria integrity, reducing cytochrome c release from the mitochondria, and inhibiting caspase activation in astaxanthin pre-treated cells through the modulation of p38 and mitogen-activated protein kinase kinase (MEK) signaling pathways in neural progenitor cells from mice [90]. Dong *et al.* reported that astaxanthin significantly reduced apoptotic death of retinal ganglion cells and alleviated diabetic retinopathy by oxidative stress inhibition [91]. In addition, astaxanthin administration increased Akt, enhanced Bad phosphorylation, and down-regulated the activation of downstream pro-apoptotic proteins (e.g., cytochrome c and caspase-3/9), leading to the amelioration of mitochondrial-related apoptosis and the attenuation of early acute kidney injury following severe burns [92].

Astaxanthin exerts a protective effect against neuronal apoptosis in the setting of neurological diseases as well (Figure 3). For example, astaxanthin mediated the activation of the PI3K/Akt survival pathway, promoted the phosphorylation-dependent inactivation of Bad, and decreased caspase-dependent neuronal apoptosis after SAH [21]. As a result, secondary brain injury in the early period of SAH, BBB disruption, cerebral edema, neurological deficits were all alleviated after treatment with astaxanthin [21]. Intra-cerebroventricular administration of astaxanthin antagonized ischemia/reperfusion-induced translocation of cytochrome c from the mitochondria to the cytoplasm, and prevented apoptosis in a transient MCAO model of ischemic stroke [17]. Lu *et al.* In addition, reported similar findings demonstrating that astaxanthin exhibits noticeable neuroprotection against cerebral ischemia-reperfusion insults through its anti-apoptotic actions [93]. In addition, pre-treatment with astaxanthin also significantly restored the mitochondrial membrane potential, prevented H₂O₂-induced neuronal apoptosis, decreased cerebral infarct volume, and improved neurological function after MCAO [93].

In an *in vitro* model of PD, Ikeda *et al.* demonstrated that astaxanthin attenuates 6-hydroxydopamine (6-OHDA)-induced apoptosis in human neuroblastoma SH-SY5Y cells [94]. Pre-treatment with astaxanthin significantly inhibits ROS generation and subsequent phosphorylation of p38 MAPK, ameliorates mitochondrial dysfunction, increases $\Delta\Psi_m$, reduces cytochrome c release, caspase activation, and rescues the cell from 6-OHDA-induced apoptosis [94,95]. Lee and coworkers found that astaxanthin treatment prevents MPP⁺-induced up-regulation of Bax and down-regulation of Bcl-2, alleviating $\Delta\Psi_m$ collapse in SH-SY5Y cells and protects the neuron against MPP⁺-induced mitochondrial damage and apoptosis [96]. Liu *et al.* demonstrated that astaxanthin has protective effects on 6-OHDA-induced cellular toxicity and apoptotic death of dopaminergic SH-SY5Y cells by inhibiting intracellular ROS generation, the decrease of mitochondrial membrane potential, the release of mitochondrial cytochrome c [97].

Intrestingly, it has been shown that astaxanthin induces cancer cell apoptosis through a mitochondrial-dependent pathway [98]. Astaxanthin mediates the inhibition of the Janus kinase 1 (JAK1)/STAT3 (signal transducer and activator of transcription 3) signaling pathway in hepatocellular carcinoma CBRH-7919 cells which down-regulates the anti-apoptotic gene expression of Bcl-2 and Bcl-xl, while also enhancing the pro-apoptotic gene expression of Bax resulting in apoptosis [99]. Another study also reported that astaxanthin induces caspase-mediated mitochondrial apoptosis by down-regulating the expression of anti-apoptotic Bcl-2 and survivin while up-regulating pro-apoptotic Bax and Bad [100]. It has also been reported that astaxanthin can induce the intrinsic apoptotic pathway in a hamster model of oral cancer through the inactivation of ERK/MAPK and PI3K/Akt

cascades which leads to the inhibition of NF- κ B and Wnt/ β -catenin [100]. Thus, depending on the pathological condition, astaxanthin may exert either anti-apoptotic or pro-apoptotic effects.




Figure 3. The anti-apoptotic effects of astaxanthin in neurological diseases. Astaxanthin induces the activation of PI3K/Akt survival pathway, promoting the phosphorylation-dependent inactivation of Bad, which leads to a decrease in caspase-dependent neuronal apoptosis. Astaxanthin also maintains mitochondria integrity through modulation of p38 and MEK signaling pathways, which reduces cytochrome c release and inhibits caspase-dependent apoptotic cell death.

4. Conclusions and Perspective

Astaxanthin confers multiple neuroprotective effects in various experimental models of neurological diseases, which includes both acute injuries and chronic neurodegenerative disorders. The protective effects of astaxanthin are associated with its anti-oxidative, anti-inflammatory, anti-apoptotic effects. Astaxanthin is a safe nutrient, with no toxic effects when it is consumed with food. Furthermore, as a fat-soluble compound, astaxanthin is able to effectively pass through the BBB. Therefore, astaxanthin is an excellent candidate for treating neurological diseases. It is essential that there continues to be further evaluations of the protective properties and underlying mechanisms of astaxanthin, which may eventually lead to astaxanthin becoming a novel neuroprotective agent.

Although the neuroprotective effects of astaxanthin have been examined in several experimental models of neurological disorders, there is a lack of research in some areas. Future studies should focus on the pharmaceutical potential and effects of astaxanthin esters in the treatment of neurological

disorders, especially since astaxanthin diesters can be easily absorbed into the metabolism and may increase biological activity more effectively than its free form [13]. Furthermore, it is important to note that a lot of the current data concerning astaxanthin-mediated neuroprotection mainly comes from ischemic stroke, SAH, AD, and PD. There is minimal evidence available regarding other neurological diseases such as traumatic brain injury, intracerebral hemorrhage, and HD. Therefore, future investigations should include these neurological disease models. The therapeutic time window, reliability of drug administration routes, and the optimal dosages of astaxanthin are other areas that need to be explored and determined. Most importantly, the development of clinical trials to assess astaxanthin as treatment of neurological diseases is warranted since there are a number of promising general safety results, neurological experimental model studies, and clinical trials in other diseases.

Acknowledgments

This study was supported by National Natural Science Foundation of China (Grant. 81171096 and 81371433), Public Technology Application Research Project of Zhejiang Province (2014C33G2010288).

Author Contributions

H.W. and H.N. collected the data and wrote the manuscript. A.S. and C.W. checked the data for accuracy. B.D. and J.Z. polished the language. Y.S. and Y.W. designed the review study.

Conflicts of Interest

The authors declare no conflicts of interest.

References

1. Corrigan, J.D.; Selassie, A.W.; Orman, J.A. The epidemiology of traumatic brain injury. *J. Head Trauma Rehabil.* **2010**, *25*, 72–80.
2. Mayeux, R. Epidemiology of neurodegeneration. *Annu. Rev. Neurosci.* **2003**, *26*, 81–104.
3. Bramlett, H.M.; Dietrich, W.D. Pathophysiology of cerebral ischemia and brain trauma: Similarities and differences. *J. Cereb. Blood Flow Metab.* **2004**, *24*, 133–150.
4. Bossy-Wetzel, E.; Schwarzenbacher, R.; Lipton, S.A. Molecular pathways to neurodegeneration. *Nat. Med.* **2004**, *10*, S2–S9.
5. Manczak, M.; Anekonda, T.S.; Henson, E.; Park, B.S.; Quinn, J.; Reddy, P.H. Mitochondria are a direct site of A beta accumulation in Alzheimer's disease neurons: Implications for free radical generation and oxidative damage in disease progression. *Hum. Mol. Genet.* **2006**, *15*, 1437–1449.
6. Cutler, R.G.; Kelly, J.; Storie, K.; Pedersen, W.A.; Tammara, A.; Hatanpaa, K.; Troncoso, J.C.; Mattson, M.P. Involvement of oxidative stress-induced abnormalities in ceramide and cholesterol metabolism in brain aging and Alzheimer's disease. *Proc. Natl. Acad. Sci. USA* **2004**, *101*, 2070–2075.
7. Hanzel, C.E.; Pichet-Binette, A.; Pimentel, L.S.; Iulita, M.F.; Allard, S.; Ducatzenzeiler, A.; do Carmo, S.; Cuello, A.C. Neuronal driven pre-plaque inflammation in a transgenic rat model of Alzheimer's disease. *Neurobiol. Aging* **2014**, *35*, 2249–2262.

8. Moore, D.J.; West, A.B.; Dawson, V.L.; Dawson, T.M. Molecular pathophysiology of Parkinson's disease. *Annu. Rev. Neurosci.* **2005**, *28*, 57–87.
9. Dawson, T.M.; Dawson, V.L. Molecular pathways of neurodegeneration in Parkinson's disease. *Science* **2003**, *302*, 819–822.
10. Ross, C.A.; Tabrizi, S.J. Huntington's disease: From molecular pathogenesis to clinical treatment. *Lancet Neurol.* **2011**, *10*, 83–98.
11. Bano, D.; Zanetti, F.; Mende, Y.; Nicotera, P. Neurodegenerative processes in Huntington's disease. *Cell Death Dis.* **2011**, *2*, e228.
12. Regnier, P.; Bastias, J.; Rodriguez-Ruiz, V.; Caballero-Casero, N.; Caballo, C.; Sicilia, D.; Fuentes, A.; Maire, M.; Crepin, M.; Letourneur, D.; *et al.* Astaxanthin from *Haematococcus Pluvialis* Prevents Oxidative Stress on Human Endothelial Cells without Toxicity. *Mar. Drugs* **2015**, *13*, 2857–2874.
13. Ambati, R.R.; Phang, S.M.; Ravi, S.; Aswathanarayana, R.G. Astaxanthin: Sources, extraction, stability, biological activities and its commercial applications—A review. *Mar. Drugs* **2014**, *12*, 128–152.
14. Fassett, R.G.; Coombes, J.S. Astaxanthin: A potential therapeutic agent in cardiovascular disease. *Mar. Drugs* **2011**, *9*, 447–465.
15. Monroy-Ruiz, J.; Sevilla, M.A.; Carron, R.; Montero, M.J. Astaxanthin-enriched-diet reduces blood pressure and improves cardiovascular parameters in spontaneously hypertensive rats. *Pharmacol. Res.* **2011**, *63*, 44–50.
16. Augusti, P.R.; Quatrin, A.; Somacal, S.; Conterato, G.M.; Sobieski, R.; Ruviano, A.R.; Maurer, L.H.; Duarte, M.M.; Roehrs, M.; Emanuelli, T. Astaxanthin prevents changes in the activities of thioredoxin reductase and paraoxonase in hypercholesterolemic rabbits. *J. Clin. Biochem. Nutr.* **2012**, *51*, 42–49.
17. Shen, H.; Kuo, C.C.; Chou, J.; Delvolve, A.; Jackson, S.N.; Post, J.; Woods, A.S.; Hoffer, B.J.; Wang, Y.; Harvey, B.K. Astaxanthin reduces ischemic brain injury in adult rats. *FASEB J.* **2009**, *23*, 1958–1968.
18. Ying, C.J.; Zhang, F.; Zhou, X.Y.; Hu, X.T.; Chen, J.; Wen, X.R.; Sun, Y.; Zheng, K.Y.; Tang, R.X.; Song, Y.J. Anti-inflammatory Effect of Astaxanthin on the Sickness Behavior Induced by Diabetes Mellitus. *Cell. Mol. Neurobiol.* **2015**, doi:10.1007/s10571-015-0197-3.
19. Zhang, X.S.; Zhang, X.; Zhou, M.L.; Zhou, X.M.; Li, N.; Li, W.; Cong, Z.X.; Sun, Q.; Zhuang, Z.; Wang, C.X.; *et al.* Amelioration of oxidative stress and protection against early brain injury by astaxanthin after experimental subarachnoid hemorrhage. *J. Neurosurg.* **2014**, *121*, 42–54.
20. Zhang, X.S.; Zhang, X.; Wu, Q.; Li, W.; Wang, C.X.; Xie, G.B.; Zhou, X.M.; Shi, J.X.; Zhou, M.L. Astaxanthin offers neuroprotection and reduces neuroinflammation in experimental subarachnoid hemorrhage. *J. Surg. Res.* **2014**, *192*, 206–213.
21. Zhang, X.S.; Zhang, X.; Wu, Q.; Li, W.; Zhang, Q.R.; Wang, C.X.; Zhou, X.M.; Li, H.; Shi, J.X.; Zhou, M.L. Astaxanthin alleviates early brain injury following subarachnoid hemorrhage in rats: Possible involvement of Akt/bad signaling. *Mar. Drugs* **2014**, *12*, 4291–4310.
22. Higuera-Ciapara, I.; Felix-Valenzuela, L.; Goycoolea, F.M. Astaxanthin: A review of its chemistry and applications. *Crit. Rev. Food Sci. Nutr.* **2006**, *46*, 185–196.

23. Hussein, G.; Nakamura, M.; Zhao, Q.; Iguchi, T.; Goto, H.; Sankawa, U.; Watanabe, H. Antihypertensive and neuroprotective effects of astaxanthin in experimental animals. *Biol. Pharm. Bull.* **2005**, *28*, 47–52.
24. Stahl, W.; Sies, H. Bioactivity and protective effects of natural carotenoids. *Biochim. Biophys. Acta* **2005**, *1740*, 101–107.
25. Guerin, M.; Huntley, M.E.; Olaizola, M. Haematococcus astaxanthin: Applications for human health and nutrition. *Trends Biotechnol.* **2003**, *21*, 210–216.
26. Jyonouchi, H.; Sun, S.; Gross, M. Effect of carotenoids on *in vitro* immunoglobulin production by human peripheral blood mononuclear cells: Astaxanthin, a carotenoid without vitamin A activity, enhances *in vitro* immunoglobulin production in response to a T-dependent stimulant and antigen. *Nutr. Cancer* **1995**, *23*, 171–183.
27. Yuan, J.P.; Chen, F. Hydrolysis kinetics of astaxanthin esters and stability of astaxanthin of Haematococcus pluvialis during saponification. *J. Agric. Food Chem.* **1999**, *47*, 31–35.
28. Yuan, J.P.; Peng, J.; Yin, K.; Wang, J.H. Potential health-promoting effects of astaxanthin: A high-value carotenoid mostly from microalgae. *Mol. Nutr. Food Res.* **2011**, *55*, 150–165.
29. Okada, Y.; Ishikura, M.; Maoka, T. Bioavailability of astaxanthin in Haematococcus algal extract: The effects of timing of diet and smoking habits. *Biosci. Biotechnol. Biochem.* **2009**, *73*, 1928–1932.
30. Coral-Hinostroza, G.N.; Ytrestoyl, T.; Ruyter, B.; Bjerkeng, B. Plasma appearance of unesterified astaxanthin geometrical *E/Z* and optical *R/S* isomers in men given single doses of a mixture of optical 3 and 3'*R/S* isomers of astaxanthin fatty acyl diesters. *Comp. Biochem. Physiol. C Toxicol. Pharmacol.* **2004**, *139*, 99–110.
31. Kistler, A.; Liechti, H.; Pichard, L.; Wolz, E.; Oesterheld, G.; Hayes, A.; Maurel, P. Metabolism and CYP-inducer properties of astaxanthin in man and primary human hepatocytes. *Arch. Toxicol.* **2002**, *75*, 665–675.
32. Parker, R.S. Absorption, metabolism, and transport of carotenoids. *FASEB J.* **1996**, *10*, 542–551.
33. Rao, A.R.; Reddy, R.L.R.; Baskaran, V.; Sarada, R.; Ravishankar, G.A. Characterization of microalgal carotenoids by mass spectrometry and their bioavailability and antioxidant properties elucidated in rat model. *J. Agric. Food Chem.* **2010**, *58*, 8553–8559.
34. Stewart, J.S.; Lignell, A.; Pettersson, A.; Elfving, E.; Soni, M.G. Safety assessment of astaxanthin-rich microalgae biomass: Acute and subchronic toxicity studies in rats. *Food. Chem. Toxicol.* **2008**, *46*, 3030–3036.
35. Spiller, G.A.; Dewell, A. Safety of an astaxanthin-rich Haematococcus pluvialis algal extract: A randomized clinical trial. *J. Med. Food* **2003**, *6*, 51–56.
36. Satoh, A.; Tsuji, S.; Okada, Y.; Murakami, N.; Urami, M.; Nakagawa, K.; Ishikura, M.; Katagiri, M.; Koga, Y.; Shirasawa, T. Preliminary Clinical Evaluation of Toxicity and Efficacy of A New Astaxanthin-Rich Haematococcus Pluvialis Extract. *J. Clin. Biochem. Nutr.* **2009**, *44*, 280–284.
37. Katagiri, M.; Satoh, A.; Tsuji, S.; Shirasawa, T. Effects of astaxanthin-rich Haematococcus pluvialis extract on cognitive function: A randomised, double-blind, placebo-controlled study. *J. Clin. Biochem. Nutr.* **2012**, *51*, 102–107.

38. Kidd, P. Astaxanthin, cell membrane nutrient with diverse clinical benefits and anti-aging potential. *Altern. Med. Rev.* **2011**, *16*, 355–364.
39. Yamagishi, R.; Aihara, M. Neuroprotective effect of astaxanthin against rat retinal ganglion cell death under various stresses that induce apoptosis and necrosis. *Mol. Vis.* **2014**, *20*, 1796–1805.
40. Gasche, Y.; Copin, J.C.; Sugawara, T.; Fujimura, M.; Chan, P.H. Matrix metalloproteinase inhibition prevents oxidative stress-associated blood-brain barrier disruption after transient focal cerebral ischemia. *J. Cereb. Blood Flow Metab.* **2001**, *21*, 1393–400.
41. Giasson, B.I.; Duda, J.E.; Murray, I.V.; Chen, Q.; Souza, J.M.; Hurtig, H.I.; Ischiropoulos, H.; Trojanowski, J.Q.; Lee, V.M. Oxidative damage linked to neurodegeneration by selective alpha-synuclein nitration in synucleinopathy lesions. *Science* **2000**, *290*, 985–989.
42. Wu, D.C.; Teismann, P.; Tieu, K.; Vila, M.; Jackson-Lewis, V.; Ischiropoulos, H.; Przedborski, S. NADPH oxidase mediates oxidative stress in the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine model of Parkinson's disease. *Proc. Natl. Acad. Sci. USA* **2003**, *100*, 6145–6150.
43. Valko, M.; Leibfritz, D.; Moncol, J.; Cronin, M.T.; Mazur, M.; Telser, J. Free radicals and antioxidants in normal physiological functions and human disease. *Int. J. Biochem. Cell Biol.* **2007**, *39*, 44–84.
44. Jilani, K.E.B.; Panee, J.; He, Q.; Berry, M.J.; Li, P.A. Overexpression of selenoprotein H reduces Ht22 neuronal cell death after UVB irradiation by preventing superoxide formation. *Int. J. Biol. Sci.* **2007**, *3*, 198–204.
45. Kamsler, A.; Segal, M. Hydrogen peroxide as a diffusible signal molecule in synaptic plasticity. *Mol. Neurobiol.* **2004**, *29*, 167–178.
46. Fridovich, I. Superoxide radical and superoxide dismutases. *Annu. Rev. Biochem.* **1995**, *64*, 97–112.
47. Sutton, V.R.; Mettert, E.L.; Beinert, H.; Kiley, P.J. Kinetic analysis of the oxidative conversion of the [4Fe-4S]²⁺ cluster of FNR to a [2Fe-2S]²⁺ Cluster. *J. Bacteriol.* **2004**, *186*, 8018–8025.
48. Tripathi, D.N.; Jena, G.B. Intervention of astaxanthin against cyclophosphamide-induced oxidative stress and DNA damage: A study in mice. *Chem. Biol. Interact.* **2009**, *180*, 398–406.
49. Stahl, W.; Sies, H. Antioxidant activity of carotenoids. *Mol. Asp. Med.* **2003**, *24*, 345–351.
50. Naito, Y.; Uchiyama, K.; Aoi, W.; Hasegawa, G.; Nakamura, N.; Yoshida, N.; Maoka, T.; Takahashi, J.; Yoshikawa, T. Prevention of diabetic nephropathy by treatment with astaxanthin in diabetic db/db mice. *Biofactors* **2004**, *20*, 49–59.
51. Camera, E.; Mastrofrancesco, A.; Fabbri, C.; Daubrawa, F.; Picardo, M.; Sies, H.; Stahl, W. Astaxanthin, canthaxanthin and beta-carotene differently affect UVA-induced oxidative damage and expression of oxidative stress-responsive enzymes. *Exp. Dermatol.* **2009**, *18*, 222–231.
52. Nishigaki, I.; Rajendran, P.; Venugopal, R.; Ekambaram, G.; Sakthisekaran, D.; Nishigaki, Y. Cytoprotective role of astaxanthin against glycated protein/iron chelate-induced toxicity in human umbilical vein endothelial cells. *Phytother. Res.* **2010**, *24*, 54–59.
53. Augusti, P.R.; Conterato, G.M.; Somacal, S.; Sobieski, R.; Spohr, P.R.; Torres, J.V.; Charao, M.F.; Moro, A.M.; Rocha, M.P.; Garcia, S.C.; *et al.* Effect of astaxanthin on kidney function impairment and oxidative stress induced by mercuric chloride in rats. *Food Chem. Toxicol.* **2008**, *46*, 212–219.

54. Nakajima, Y.; Inokuchi, Y.; Shimazawa, M.; Otsubo, K.; Ishibashi, T.; Hara, H. Astaxanthin, a dietary carotenoid, protects retinal cells against oxidative stress *in vitro* and in mice *in vivo*. *J. Pharm. Pharmacol.* **2008**, *60*, 1365–1374.
55. Tripathi, D.N.; Jena, G.B. Astaxanthin intervention ameliorates cyclophosphamide-induced oxidative stress, DNA damage and early hepatocarcinogenesis in rat: Role of Nrf2, p53, p38 and phase-II enzymes. *Mutat. Res.* **2010**, *696*, 69–80.
56. Li, Z.; Dong, X.; Liu, H.; Chen, X.; Shi, H.; Fan, Y.; Hou, D.; Zhang, X. Astaxanthin protects ARPE-19 cells from oxidative stress via upregulation of Nrf2-regulated phase II enzymes through activation of PI3K/Akt. *Mol. Vis.* **2013**, *19*, 1656–1666.
57. Lee, D.H.; Lee, Y.J.; Kwon, K.H. Neuroprotective Effects of Astaxanthin in Oxygen-Glucose Deprivation in SH-SY5Y Cells and Global Cerebral Ischemia in Rat. *J. Clin. Biochem. Nutr.* **2010**, *47*, 121–129.
58. Wu, Q.; Zhang, X.S.; Wang, H.D.; Zhang, X.; Yu, Q.; Li, W.; Zhou, M.L.; Wang, X.L. Astaxanthin activates nuclear factor erythroid-related factor 2 and the antioxidant responsive element (Nrf2-ARE) pathway in the brain after subarachnoid hemorrhage in rats and attenuates early brain injury. *Mar. Drugs* **2014**, *12*, 6125–6141.
59. Wang, H.Q.; Sun, X.B.; Xu, Y.X.; Zhao, H.; Zhu, Q.Y.; Zhu, C.Q. Astaxanthin upregulates heme oxygenase-1 expression through ERK1/2 pathway and its protective effect against beta-amyloid-induced cytotoxicity in SH-SY5Y cells. *Brain Res.* **2010**, *1360*, 159–167.
60. Motterlini, R.; Green, C.J.; Foresti, R. Regulation of heme oxygenase-1 by redox signals involving nitric oxide. *Antioxid. Redox Signal.* **2002**, *4*, 615–624.
61. Bae, J.W.; Kim, M.J.; Jang, C.G.; Lee, S.Y. Protective effects of heme oxygenase-1 against MPP(+)-induced cytotoxicity in PC-12 cells. *Neurol. Sci.* **2010**, *31*, 307–313.
62. Ye, Q.; Huang, B.; Zhang, X.; Zhu, Y.; Chen, X. Astaxanthin protects against MPP(+)-induced oxidative stress in PC12 cells via the HO-1/NOX2 axis. *BMC Neurosci.* **2012**, *13*, 156.
63. Ye, Q.; Zhang, X.; Huang, B.; Zhu, Y.; Chen, X. Astaxanthin suppresses MPP(+)-induced oxidative damage in PC12 cells through a Sp1/NR1 signaling pathway. *Mar. Drugs* **2013**, *11*, 1019–1034.
64. Turrin, N.P.; Rivest, S. Molecular and cellular immune mediators of neuroprotection. *Mol. Neurobiol.* **2006**, *34*, 221–242.
65. Brown, G.C.; Neher, J.J. Inflammatory neurodegeneration and mechanisms of microglial killing of neurons. *Mol. Neurobiol.* **2010**, *41*, 242–247.
66. Lucas, S.M.; Rothwell, N.J.; Gibson, R.M. The role of inflammation in CNS injury and disease. *Br. J. Pharmacol.* **2006**, *147*, S232–S240.
67. Ohgami, K.; Shiratori, K.; Kotake, S.; Nishida, T.; Mizuki, N.; Yazawa, K.; Ohno, S. Effects of astaxanthin on lipopolysaccharide-induced inflammation *in vitro* and *in vivo*. *Investig. Ophthalmol. Vis. Sci.* **2003**, *44*, 2694–2701.
68. Suzuki, Y.; Ohgami, K.; Shiratori, K.; Jin, X.H.; Ilieva, I.; Koyama, Y.; Yazawa, K.; Yoshida, K.; Kase, S.; Ohno, S. Suppressive effects of astaxanthin against rat endotoxin-induced uveitis by inhibiting the NF- κ B signaling pathway. *Exp. Eye Res.* **2006**, *82*, 275–281.
69. Ghosh, S.; May, M.J.; Kopp, E.B. NF- κ B and Rel proteins: Evolutionarily conserved mediators of immune responses. *Annu. Rev. Immunol.* **1998**, *16*, 225–260.

70. Yamamoto, Y.; Yin, M.J.; Gaynor, R.B. IkappaB kinase alpha (IKKalpha) regulation of IKKbeta kinase activity. *Mol. Cell. Biol.* **2000**, *20*, 3655–3666.
71. Poyet, J.L.; Srinivasula, S.M.; Lin, J.H.; Fernandes-Alnemri, T.; Yamaoka, S.; Tschlis, P.N.; Alnemri, E.S. Activation of the Ikappa B kinases by RIP via IKKgamma /NEMO-mediated oligomerization. *J. Biol. Chem.* **2000**, *275*, 37966–37977.
72. Shao, A.W.; Wu, H.J.; Chen, S.; Ammar, A.B.; Zhang, J.M.; Hong, Y. Resveratrol attenuates early brain injury after subarachnoid hemorrhage through inhibition of NF-kappaB-dependent inflammatory/MMP-9 pathway. *CNS Neurosci. Ther.* **2014**, *20*, 182–185.
73. Bhuvanewari, S.; Yogalakshmi, B.; Sreeja, S.; Anuradha, C.V. Astaxanthin reduces hepatic endoplasmic reticulum stress and nuclear factor-kappaB-mediated inflammation in high fructose and high fat diet-fed mice. *Cell Stress Chaperones* **2014**, *19*, 183–191.
74. Speranza, L.; Pesce, M.; Patruno, A.; Franceschelli, S.; de Lutiis, M.A.; Grilli, A.; Felaco, M. Astaxanthin treatment reduced oxidative induced pro-inflammatory cytokines secretion in U937: SHP-1 as a novel biological target. *Mar. Drugs* **2012**, *10*, 890–899.
75. Izumi-Nagai, K.; Nagai, N.; Ohgami, K.; Satofuka, S.; Ozawa, Y.; Tsubota, K.; Ohno, S.; Oike, Y.; Ishida, S. Inhibition of choroidal neovascularization with an anti-inflammatory carotenoid astaxanthin. *Investig. Ophthalmol. Vis. Sci.* **2008**, *49*, 1679–1685.
76. Bennedsen, M.; Wang, X.; Willen, R.; Wadstrom, T.; Andersen, L.P. Treatment of H. pylori infected mice with antioxidant astaxanthin reduces gastric inflammation, bacterial load and modulates cytokine release by splenocytes. *Immunol. Lett.* **1999**, *70*, 185–189.
77. Yasui, Y.; Hosokawa, M.; Mikami, N.; Miyashita, K.; Tanaka, T. Dietary astaxanthin inhibits colitis and colitis-associated colon carcinogenesis in mice via modulation of the inflammatory cytokines. *Chem. Biol. Interact.* **2011**, *193*, 79–87.
78. Lee, S.J.; Bai, S.K.; Lee, K.S.; Namkoong, S.; Na, H.J.; Ha, K.S.; Han, J.A.; Yim, S.V.; Chang, K.; Kwon, Y.G.; *et al.* Astaxanthin inhibits nitric oxide production and inflammatory gene expression by suppressing I(kappa)B kinase-dependent NF-kappaB activation. *Mol. Cells* **2003**, *16*, 97–105.
79. Terazawa, S.; Nakajima, H.; Shingo, M.; Niwano, T.; Imokawa, G. Astaxanthin attenuates the UVB-induced secretion of prostaglandin E2 and interleukin-8 in human keratinocytes by interrupting MSK1 phosphorylation in a ROS depletion-independent manner. *Exp. Dermatol.* **2012**, *21*, S11–S17.
80. Chew, B.P.; Wong, M.W.; Park, J.S.; Wong, T.S. Dietary beta-carotene and astaxanthin but not canthaxanthin stimulate splenocyte function in mice. *Anticancer Res.* **1999**, *19*, 5223–5227.
81. Chew, B.P.; Mathison, B.D.; Hayek, M.G.; Massimino, S.; Reinhart, G.A.; Park, J.S. Dietary astaxanthin enhances immune response in dogs. *Vet. Immunol. Immunopathol.* **2011**, *140*, 199–206.
82. Jyonouchi, H.; Sun, S.; Iijima, K.; Gross, M.D. Antitumor activity of astaxanthin and its mode of action. *Nutr. Cancer* **2000**, *36*, 59–65.
83. Kurihara, H.; Koda, H.; Asami, S.; Kiso, Y.; Tanaka, T. Contribution of the antioxidative property of astaxanthin to its protective effect on the promotion of cancer metastasis in mice treated with restraint stress. *Life Sci.* **2002**, *70*, 2509–2520.

84. Park, J.S.; Chyun, J.H.; Kim, Y.K.; Line, L.L.; Chew, B.P. Astaxanthin decreased oxidative stress and inflammation and enhanced immune response in humans. *Nutr. Metab. (Lond.)* **2010**, *7*, 18.
85. Baliotti, M.; Giannubilo, S.R.; Giorgetti, B.; Solazzi, M.; Turi, A.; Casoli, T.; Ciavattini, A.; Fattoretta, P. The effect of astaxanthin on the aging rat brain: Gender-related differences in modulating inflammation. *J. Sci. Food Agric.* **2015**, doi:10.1002/jsfa.7131.
86. Elmore, S. Apoptosis: A review of programmed cell death. *Toxicol. Pathol.* **2007**, *35*, 495–516.
87. Wu, H.J.; Pu, J.L.; Krafft, P.R.; Zhang, J.M.; Chen, S. The molecular mechanisms between autophagy and apoptosis: Potential role in central nervous system disorders. *Cell. Mol. Neurobiol.* **2015**, *35*, 85–99.
88. Nijhawan, D.; Honarpour, N.; Wang, X. Apoptosis in neural development and disease. *Annu. Rev. Neurosci.* **2000**, *23*, 73–87.
89. Shao, A.; Wang, Z.; Wu, H.; Dong, X.; Li, Y.; Tu, S.; Tang, J.; Zhao, M.; Zhang, J.; Hong, Y. Enhancement of Autophagy by Histone Deacetylase Inhibitor Trichostatin A Ameliorates Neuronal Apoptosis After Subarachnoid Hemorrhage in Rats. *Mol. Neurobiol.* **2014**, doi:10.1007/s12035-014-8986-0
90. Kim, J.H.; Choi, W.; Lee, J.H.; Jeon, S.J.; Choi, Y.H.; Kim, B.W.; Chang, H.I.; Nam, S.W. Astaxanthin inhibits H₂O₂-mediated apoptotic cell death in mouse neural progenitor cells via modulation of P38 and MEK signaling pathways. *J. Microbiol. Biotechnol.* **2009**, *19*, 1355–1363.
91. Dong, L.Y.; Jin, J.; Lu, G.; Kang, X.L. Astaxanthin attenuates the apoptosis of retinal ganglion cells in db/db mice by inhibition of oxidative stress. *Mar. Drugs* **2013**, *11*, 960–974.
92. Guo, S.X.; Zhou, H.L.; Huang, C.L.; You, C.G.; Fang, Q.; Wu, P.; Wang, X.G.; Han, C.M. Astaxanthin attenuates early acute kidney injury following severe burns in rats by ameliorating oxidative stress and mitochondrial-related apoptosis. *Mar. Drugs* **2015**, *13*, 2105–2123.
93. Lu, Y.P.; Liu, S.Y.; Sun, H.; Wu, X.M.; Li, J.J.; Zhu, L. Neuroprotective effect of astaxanthin on H₂O₂-induced neurotoxicity *in vitro* and on focal cerebral ischemia *in vivo*. *Brain Res.* **2010**, *1360*, 40–48.
94. Ikeda, Y.; Tsuji, S.; Satoh, A.; Ishikura, M.; Shirasawa, T.; Shimizu, T. Protective effects of astaxanthin on 6-hydroxydopamine-induced apoptosis in human neuroblastoma SH-SY5Y cells. *J. Neurochem.* **2008**, *107*, 1730–1740.
95. Gomez-Lazaro, M.; Galindo, M.F.; Concannon, C.G.; Segura, M.F.; Fernandez-Gomez, F.J.; Llecha, N.; Comella, J.X.; Prehn, J.H.; Jordan, J. 6-hydroxydopamine activates the mitochondrial apoptosis pathway through p38 MAPK-mediated, p53-independent activation of Bax and PUMA. *J. Neurochem.* **2008**, *104*, 1599–1612.
96. Lee, D.H.; Kim, C.S.; Lee, Y.J. Astaxanthin protects against MPTP/MPP⁺-induced mitochondrial dysfunction and ROS production *in vivo* and *in vitro*. *Food Chem. Toxicol.* **2011**, *49*, 271–280.
97. Liu, X.; Shibata, T.; Hisaka, S.; Osawa, T. Astaxanthin inhibits reactive oxygen species-mediated cellular toxicity in dopaminergic SH-SY5Y cells via mitochondria-targeted protective mechanism. *Brain Res.* **2009**, *1254*, 18–27.
98. Song, X.D.; Zhang, J.J.; Wang, M.R.; Liu, W.B.; Gu, X.B.; Lv, C.J. Astaxanthin induces mitochondria-mediated apoptosis in rat hepatocellular carcinoma CBRH-7919 cells. *Biol. Pharm. Bull.* **2011**, *34*, 839–844.

99. Song, X.; Wang, M.; Zhang, L.; Zhang, J.; Wang, X.; Liu, W.; Gu, X.; Lv, C. Changes in cell ultrastructure and inhibition of JAK1/STAT3 signaling pathway in CBRH-7919 cells with astaxanthin. *Toxicol. Mech. Methods* **2012**, *22*, 679–686.
100. Kavitha, K.; Kowshik, J.; Kishore, T.K.; Baba, A.B.; Nagini, S. Astaxanthin inhibits NF-kappaB and Wnt/beta-catenin signaling pathways via inactivation of Erk/MAPK and PI3K/Akt to induce intrinsic apoptosis in a hamster model of oral cancer. *Biochim. Biophys. Acta* **2013**, *1830*, 4433–4444.

© 2015 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>).