

Errata

Vol. 64, No. 1

In the report, “Incidence of Notifiable Diseases Among American Indian/Alaska Natives — United States, 2007–2011,” multiple errors occurred in the text and table.

In the third paragraph, the first sentence should read “For 26 notifiable diseases examined for 2007–2011, a total of **9,061,675** cases were recorded (Table).” The third sentence should read, “Missing data on race ranged from 0.8% for tuberculosis to 42% for giardiasis.”

In the fourth paragraph, the first sentence should read, “Of the 12 diseases with race information for >70% of records and

for which rates were higher among AI/ANs than among whites, the largest difference was for hantavirus pulmonary syndrome, which was reported **10** times more often among AI/ANs than among whites; however, only 20 cases were reported among AI/ANs of a total of 112 cases reported during 2007–2011.” The second sentence should read, “The second largest difference was for tularemia, which was reported 7.7 times as often among AI/ANs.”

On page 17, the Table should have read as follows:

TABLE. Number and incidence rate per 100,000 population for 26 selected notifiable diseases, by American Indian/Alaska Native (AI/AN), black, or white race — United States, 2007–2011

Disease	AI/ANs		Blacks		Whites		Total		Rate ratio: AI/ANs compared with whites	% with no race identified
	No.	Rate	No.	Rate	No.	Rate	No.	Rate		
Botulism, foodborne	26	0.12	28	0.01	339	0.03	672	0.04	4.38	35.42
Chickenpox (varicella)	503	2.45	7,086	3.41	78,776 78,776	6.45	110,634	7.22	0.38	17.82
<i>Chlamydia trachomatis</i>	77,072	374.83	2,189,748	1,052.68	1,841,172	150.74	6,283,761	409.90	2.49	29.84
Cryptosporidiosis	223	1.08	3,202	1.54	29,010	2.38	45,721	2.98	0.46	25.63
Ehrlichiosis, total	219	1.07	167	0.08	7,250	0.59	12,348	0.81	1.79	36.46
Gonorrhea	12,764	62.08	894,198	429.87	317,271	25.97	1,625,097	106.01	2.39	21.70
Giardiasis	385	1.87	6,875	3.31	38,506	3.15	93,164	6.08	0.59	41.55
<i>Haemophilus influenzae</i>	206	1.00	1,822	0.88	9,340	0.76	14,990	0.98	1.31	20.69
Hantavirus pulmonary syndrome	20	0.10	1	0.00	77	0.01	112	0.01	15.43	10.71
Hepatitis A, viral acute	66	0.32	677	0.33	5,607	0.46	10,544	0.69	0.70	28.15
Hepatitis B, viral acute	144	0.70	3,532	1.70	9,433	0.77	18,114	1.18	0.91	2.22
Hepatitis C, viral acute	88	0.43	261	0.13	3,220	0.26	4,553	0.30	1.62	19.33
Legionellosis	42	0.20	2,890	1.39	10,590	0.87	16,870	1.10	0.24	16.87
Lyme disease	476	2.31	1,649	0.79	85,721	7.02	160,209	10.45	0.33	38.68
Meningococcal disease	48	0.23	707	0.34	2,899	0.24	4,776	0.31	0.98	18.91
Pertussis	788	3.83	3,709	1.78	57,644	4.72	85,723	5.59	0.81	23.48
Salmonellosis	1,783	8.67	21,647	10.41	142,495	11.67	252,169	16.45	0.74	28.99
Shiga toxin–producing <i>Escherichia coli</i>	161	0.78	1,020	0.49	16,749	1.37	26,058	1.70	0.57	27.09
Shigellosis	1,115	5.42	17,822	8.57	37,309	3.05	85,172	5.56	1.78	28.71
Spotted fever rickettsiosis	519	2.52	434	0.21	7,325	0.60	11,108	0.72	4.21	23.17
<i>Streptococcus pneumoniae</i> , invasive (all ages)	575	2.80	8,652	4.26	28,766	2.36	49,548	3.23	1.19	20.76
<i>Streptococcus pneumoniae</i> , invasive (age <5 years)	297	15.92	2,249	13.38	9,214	12.04	16,102	15.95	1.32	21.96
Syphilis, primary and secondary	367	1.78	31,469	15.13	28,616	2.30	66,707	4.35	0.78	4.00
Tuberculosis	813	3.95	15,167	7.29	25,944	2.12	59,458	3.88	1.86	0.77
Tularemia	47	0.23	15	0.01	413	0.03	626	0.04	6.76	21.73
West Nile virus disease	184	0.89	348	0.17	5,142	0.42	7,439	0.49	2.13	21.86
Total	98,931	—	3,215,375	—	2,798,828	—	9,061,675	—	—	—

Errata

Vol. 64, No. 6

In the report, “Update on Progress in Selected Public Health Programs After the 2010 Earthquake and Cholera Epidemic — Haiti, 2014,” in the Figure on page 139, the data shown for “Eligible children receiving measles-rubella vaccination” pertain only to estimated coverage through routine immunization. Additional vaccinations were provided through special campaigns.