

Pgc-1 α and *Nr4a1* Are Target Genes of Circadian Melatonin and Dopamine Release in Murine Retina

Stefanie Kunst,^{1,2} Tanja Wolloscheck,¹ Debra K. Kelleher,¹ Uwe Wolfrum,² S. Anna Sargsyan,³ P. Michael Iuvone,³ Kenkichi Baba,⁴ Gianluca Tosini,⁴ and Rainer Spessert¹

¹Institute of Functional and Clinical Anatomy, University Medical Center of the Johannes Gutenberg University Mainz, Mainz, Germany

²Department of Cell and Matrix Biology, Institute of Zoology, Johannes Gutenberg University, Mainz, Germany

³Departments of Ophthalmology and Pharmacology, Emory University School of Medicine, Atlanta, Georgia, United States

⁴Neuroscience Institute and Department of Pharmacology and Toxicology, Morehouse School of Medicine, Atlanta, Georgia, United States

Correspondence: Rainer Spessert, Institute of Functional and Clinical Anatomy, University Medical Center of the Johannes Gutenberg University Mainz, Saarstraße 19-21, 55099 Mainz, Germany; spessert@uni-mainz.de.

Submitted: June 16, 2015

Accepted: August 9, 2015

Citation: Kunst S, Wolloscheck T, Kelleher DK, et al. *Pgc-1 α* and *Nr4a1* are target genes of circadian melatonin and dopamine release in murine retina. *Invest Ophthalmol Vis Sci*. 2015;56:6084-6094. DOI:10.1167/iov.15-17503

PURPOSE. The neurohormones melatonin and dopamine mediate clock-dependent/circadian regulation of inner retinal neurons and photoreceptor cells and in this way promote their functional adaptation to time of day and their survival. To fulfill this function they act on melatonin receptor type 1 (MT₁ receptors) and dopamine D₄ receptors (D₄ receptors), respectively. The aim of the present study was to screen transcriptional regulators important for retinal physiology and/or pathology (*Dbp*, *Egr-1*, *Fos*, *Nr1d1*, *Nr2e3*, *Nr4a1*, *Pgc-1 α* , *Rorb β*) for circadian regulation and dependence on melatonin signaling/MT₁ receptors or dopamine signaling/D₄ receptors.

METHODS. This was done by gene profiling using quantitative polymerase chain reaction in mice deficient in MT₁ or D₄ receptors.

RESULTS. The data obtained determined *Pgc-1 α* and *Nr4a1* as transcriptional targets of circadian melatonin and dopamine signaling, respectively.

CONCLUSIONS. The results suggest that *Pgc-1 α* and *Nr4a1* represent candidate genes for linking circadian neurohormone release with functional adaptation and healthiness of retina and photoreceptor cells.

Keywords: retina, circadian regulation, melatonin receptor type 1, dopamine D₄ receptor, *Nr4a1*, *Pgc-1 α*

The mammalian retina has the ability to adapt to the marked daily changes in the environment. This facilitates the retina, and in particular photoreceptor cells (PRCs), to comply with daily changes in metabolic and functional demands and on a long-term basis may contribute to their survival. Daily adaptation of retinal physiology¹⁻³ is driven by retinal clocks⁴ partly through the rhythmic release of the hormones melatonin from PRCs and dopamine from amacrine cells (for review, see Ref. 5). Melatonin and dopamine play opposing roles in the control of retinal adaptation (for reviews, see Refs. 6, 7). Melatonin release occurs at night and mediates dark adaptation of retina through G-protein receptors named melatonin type 1 (MT₁) and type 2 (MT₂) receptors (for review, see Ref. 8) that appear to function as MT₁/MT₂ heteromers.⁹ In this process it modulates visual processing and viability of PRCs.¹⁰⁻¹² On the other hand, dopamine release is higher during daytime^{13,14} and promotes adaptation of visual function to light (for reviews, see Refs. 6, 15, 16). Dopamine exerts its effects on various types of dopamine receptors widespread through the retina, but in particular on dopamine D₄ (D₄) receptors expressed primarily on PRCs to entrain light-adapted vision including contrast sensitivity function,^{12,16,17} the photoreceptor circadian clock in terms of protein phosphorylation,¹⁸ melatonin release,¹⁹ and adenylyl cyclase 1/cAMP signaling.²⁰

In the mammalian retina, transcriptional regulators are critical for visual function,^{21,22} cell fate determination (for reviews, see Refs. 23, 24), and homeostasis.²⁵⁻²⁸ They are also important for retinal cell health and viability. Thus, transcriptional regulators (1) underlie different forms of inherited retinal degeneration in humans, such as *Nr2e3*²⁹⁻³¹ and *Rorb β* ,³² and (2) are protective against retinal dystrophy, such as *Nr1d1*³³ and *Pgc-1 α* ,²⁷ or activate immune defense pathways in response to damage of the retina, such as *Egr-1* (*Krox24*; *Ngfi-a*; *Zif268*; *Zenk*^{28,34,35}; *Fos*^{34,36} and *Nr4a1* (*Ngfi-b*, *Nur77*³⁴). Remarkably, numerous transcriptional regulators display a daily rhythm of expression in the retina (*Dbp*³⁷; *Egr-1*³⁸; *Fos*³⁹; *Nr1d1*⁴⁰; *Nr2e3*⁴¹; *Nr4a1*⁴²; *Pgc-1 α* ⁴³; *Rorb β* ⁴⁴). Therefore, their positive role in retinal health may be based on their ability to promote daily adjustment of the retina to comply with ambient demands.

The aim of the present study was to determine transcriptional regulators that may mediate retinal responses to circadian neurotransmitter release in retina of melatonin-proficient mice not carrying *rd* mutations. For this, daily regulation of the transcriptional regulators *Dbp*, *Egr-1*, *Fos*, *Nr1d1*, *Nr2e3*, *Nr4a1*, *Pgc-1 α* , *Rorb β* was seen to be driven by a circadian clock, to take place in PRCs, and, most importantly, to depend on melatonin signaling/MT₁ receptors or dopamine signaling/D₄ receptors. As a result, *Pgc-1 α* and *Nr4a1* were identified as

TABLE 1. Primer Sequences Used for qPCR

Gene	Accession Number		Primer Sequence 5' to 3'	Length of PCR Product, bp
<i>Dbp</i>	NM_016974.3	Forward	GGAGGTGCTAATGACCTTTG	146
		Reverse	GGACTTTCCCTTGCCCTTCTTC	
<i>Egr-1</i>	NM_007913.5	Forward	CCCTTCCAGTGTCGAATC	127
		Reverse	CACCTCTGGCAAACCTTCC	
<i>Fos</i>	NM_010234	Forward	C'TTGAGCCAGTCAAGAG	141
		Reverse	TAGAAGGAACCGGACAGG	
<i>Gapdh</i>	BC082592	Forward	GTCATCCCAGAGCTGAAC	144
		Reverse	CTCAGATGCCTGCCTTCAC	
<i>Nr1d1</i>	NM_145434	Forward	CTGCATCATCGTCATCCTCTTC	150
		Reverse	ACAGTAGCACCATGCCATTC	
<i>Nr2e3</i>	NM_013708	Forward	GCCCTGGTCTCTTCAAAAC	112
		Reverse	GGTGGTGAGCCTTGCTATG	
<i>Nr4a1</i>	NM_010444	Forward	TGATGTTCCCGCCTTTGC	126
		Reverse	GCCATGTGCTCCTTCAGAC	
<i>Nrl</i>	NM_008736	Forward	GTGGAGGAACGGTCCAGATG	149
		Reverse	GAACTGGAGGGCTGGGTTAC	
<i>Pgc-1α</i>	NM_008904	Forward	GTGTTCCCGATCACCATATTTCC	101
		Reverse	GGTGTCTGTAGTGGCTTGATTC	
<i>Rorb</i>	NM_00146095.3	Forward	CCTGGCTGATCGAACCAAG	144
		Reverse	TGCAGACTGCCGTGATAG	
<i>Tb</i>	NM_009377.1	Forward	CAGCCCTACCAAGATCAAAC	129
		Reverse	GTACGGGTCAAACCTTCACAG	

putative transcriptional targets of the circadian melatonin/dopamine system in murine retina.

MATERIALS AND METHODS

Animals

Adult male and female mice with intact PRCs not carrying *rd* mutation were used in this study. As a rule the mice used were melatonin proficient (C3H/He (*rd*^{+/+}), C3H/*f*^{+/+}MT1^{+/+}, C3H/*f*^{+/+}MT1^{-/-}, C3H/*f*^{+/+}Drd4^{+/+}, and C3H/*f*^{+/+}Drd4^{-/-}) and where indicated melatonin deficient (C57BL/6Jb). Mice were genotyped by PCR analysis of genomic DNA. The mice were kept under standard laboratory conditions (illumination with fluorescent strip lights, 200 lux at cage level during the day and dim red light during the night; 20 \pm 1°C; water and food ad libitum) under light/dark 12:12 (LD 12:12) for 3 weeks. When indicated, the animals were then kept for one cycle under dim red light and killed during the next cycle. Animals (two for each time point) were killed at the indicated time points by decapitation following anesthesia with carbon dioxide. All dissections during the dark phase were carried out under dim red light. Retinas (four for each time point) were quickly removed, pooled, and immediately frozen or processed for laser microdissection and pressure catapulting (LMPC). Each experiment was carried out four times independently. Animal experimentation was carried out in accordance with the National Institutes of Health Guide on the Care and Use of Laboratory Animals and the ARVO Statement for the Use of Animals in Ophthalmic and Vision Research, and approved by the Institutional Animal Care and Use Committees of Morehouse School of Medicine, Emory University, and the European Communities Council Directive (86/609/EEC).

Laser Microdissection and Pressure Catapulting

To prepare the retinas for LMPC, the HOPE technique (HOPE, Hepes-glutamic acid buffer-mediated organic solvent protection effect; DCS, Hamburg, Germany) was applied for the

fixation.⁴⁵ Photoreceptor cells were isolated from the stained sections in a contact- and contamination-free manner by using the LMPC technique as described previously.⁴⁶ The purity grades of the preparations obtained were verified by using a specific gene marker of PRCs, namely, *Nrl* (as markers for rods⁴⁷), and of inner retinal neurons, namely, *Tb* (as a marker for amacrine cells⁴⁸). In comparison to whole-retina preparations, in PRCs collected by LMPC, the ratio of *Nrl* to *Tb* was increased 84-fold.

RNA Extraction, Reverse Transcription (RT), and Quantitative Polymerase Chain Reaction (qPCR)

RNA was isolated and reverse transcribed as described previously.⁴³ In brief, PCR amplification and quantification were performed in an i-Cycler (BioRad, Munich, Germany) according to the following protocol: denaturation for 3 minutes at 95°C, followed by 40 cycles of 30 seconds at 95°C, 20 seconds at 60°C, and 20 seconds at 72°C. All amplifications were carried out in duplicate. By using agarose gel electrophoresis, the generated amplicons for all genes under examination were shown to possess the predicted sizes (Table 1). The amount of RNA was calculated from the measured threshold cycles (C_t) using an internal standard curve with 10-fold serial dilutions (10¹-10⁸ copies/ μ L). The values were normalized with respect to the amount of glyceraldehyde 3-phosphate dehydrogenase (GAPDH) mRNA present.

Statistical Analysis

All PCR data are expressed as the mean \pm standard error of the mean (SEM) of four independent experiments including eight time points. Transcript levels were calculated relative to average expression of each dataset throughout 24 hours to plot temporal expression. Significance of daily regulation was defined by showing a $P < 0.05$ in ANOVA (one-way analysis of variance). Cosinor analysis was used to fit sine wave curves to the circadian data to mathematically estimate the time of peak gene expression (acrophase) and to assess the amplitude.^{49,50} The model can be expressed according to the equation $f(t) = A$

+ $B \cos [2\pi (t + C) / T]$ with the $f(t)$ indicating relative expression levels of target genes, t specifying the time of sampling (b), A representing the mean value of the cosine curve (MESOR; midline estimating statistic of rhythm), B indicating the amplitude of the curve (half of the sinusoid), and C indicating the acrophase (point of time when the function $f(t)$ is maximum). T gives the time of the period, which is 24 hours for this experimental setting.

RESULTS

Daily Profiling of the Transcriptional Regulators in Whole Retina of Melatonin-Proficient Mouse

With the exception of a recently published study,³⁷ daily profiling of gene expression in murine retina has so far been performed in mice strains that either are deficient for melatonin or carry *rd* mutation (resulting in damaged PRCs).^{51–55} The objective of this study—to investigate melatonin-dependent regulation of genes in the retina including PRCs—requires melatonin-proficient mice with intact PRCs (C3H/He (*rd*^{+/+})). It was seen that in these mice all transcriptional regulators under investigation display daily changes with peak expression during the day (*Dbp*, *Nr1d1*, *Nr2e3*, *Pgc-1 α*) or at night (*Egr-1*, *Fos*, *Nr4a1*, *Ror β*) (Fig. 1, blue lines; for statistical analysis, see Table 2).

Daily Profiling of the Transcriptional Regulators in PRCs of Melatonin-Proficient Mouse

In order to compare daily regulation of the transcriptional regulators between preparations of the whole retina and PRCs, the LMPC technique was applied.⁴⁶ The transcript levels of the genes were observed to exhibit daily rhythms in PRCs (Fig. 1, red lines; for statistical analysis, see Table 2) with profiles resembling those obtained from preparations of the whole retina (Fig. 1, blue lines; for statistical analysis, see Table 2). The findings obtained suggest that daily rhythmicity of the transcriptional regulators also occurs in PRCs. Interestingly, the relative strength of cycling of the genes (indicated in terms of the height of the amplitude) was different in PRCs (*Nr4a1* > *Fos* > *Dbp* > *Nr2e3* > *Pgc-1 α* > *Ror β*) and whole retina (*Nr4a1* > *Fos* > *Ror β* > *Dbp* > *Nr2e3* > *Pgc-1 α* > *Ror β*). This might reflect different circadian output in outer and inner retina.

Daily Profiling of the Transcriptional Regulators in Retina of Melatonin-Proficient Mice in Constant Darkness

To gain a view of transcriptional control of the retina by a circadian clock, light input, or both, mice (C3H/He (*rd*^{+/+})) adapted to LD 12:12 were kept in constant darkness (DD) for one cycle and monitored during the subsequent cycle (Fig. 1, black lines; for statistical analysis, see Table 2). Under these conditions, the daily rhythm in mRNA amount continued for all transcriptional regulators investigated (*Dbp*, *Egr-1*, *Fos*, *Nr1d1*, *Nr2e3*, *Nr4a1*, *Pgc-1 α* , *Ror β*). This observation suggests that daily rhythmicity of the transcriptional regulators investigated requires a circadian clock. As far as assessable ($P < 0.05$ in cosinor analysis), the retained rhythmicity of the transcriptional regulators occurred with reduced amplitude under DD conditions when compared with LD conditions with the exception of *Nr1d1*. Therefore clock-dependent rhythmicity of the transcriptional regulators appears to be enhanced by light/dark transitions.

Daily Profiling of the Transcriptional Regulators in Melatonin-Deficient Mice

The retinal clock system—probably in terms of a clock within PRCs^{40,46}—evokes a daily rhythm in melatonin release (for review, see Ref. 5). Therefore clock-driven rhythmicity of the transcriptional regulators may be conveyed by melatonin signaling. To test this assumption, daily profiling of the transcriptional regulators in melatonin-proficient mice C3H/He (*rd*^{+/+}) was compared with that of melatonin-deficient C57BL/6Jb mice (Fig. 2, blue versus red lines; for statistical analysis, see Table 2). This revealed that *Pgc-1 α* rhythmicity was prevented in melatonin-deficient mice. Conversely, the other transcriptional regulators under investigation (*Dbp*, *Egr-1*, *Fos*, *Nr1d1*, *Nr2e3*, *Nr4a1*, *Ror β*) were found to display daily patterns similar to those being observed in melatonin-proficient mice. These observations suggest that melatonin signaling is a prerequisite for the *Pgc-1 α* rhythmicity.

Daily Profiling of the Transcriptional Regulators in Melatonin-Proficient MT₁^{-/-} Mice

To verify and specify the response of the transcriptional regulators to melatonin signaling, their 24-hour profiles were compared in wild-type (WT) and MT₁ receptor-deficient retina (Fig. 3, blue versus red lines; for statistical analysis, see Table 3), with both genotypes deriving from a melatonin-proficient mice strain with intact PRCs (C3H/*f*^{+/+}). Consistent with the results obtained from melatonin-deficient mice, the loss of MT₁ receptors prevented the cycling of *Pgc-1 α* . As far as assessable ($P < 0.05$ in cosinor analysis), it was seen to slightly dampen the rhythmicity of the other transcriptional regulators under investigation (*Dbp*, *Nr1d1*, *Nr2e3*, *Nr4a1*, *Ror β*). This observation suggests that MT₁ receptors couple the pulsatile melatonin signal to the expression of *Pgc-1 α* and possibly play a general role in amplifying circadian regulation of retinal transcription.

Daily Profiling of the Transcriptional Regulators in Drd4^{-/-} Mice

Clock-dependent regulation of the transcriptional regulators may also involve dopamine signaling via dopamine D₄ receptors (for review, see Ref. 5). To investigate this possibility, 24-hour profiling of the transcriptional regulators was performed in *Drd4*-deficient mice (C3H/*f*^{+/+}, melatonin-proficient mice with intact PRCs) (Fig. 4, blue versus red lines; for statistical analysis, see Table 3). The rhythmicity of *Nr4a1* was seen to be dampened in *Drd4*^{-/-} mice, with a decrease in amplitude of 84%, whereas the other transcriptional regulators under investigation were seen to display similar daily profiles in both the genotypes. This finding suggests that *Nr4a1* links the clock-driven dopamine signaling with retinal gene expression.

DISCUSSION

The retinal clocks exert their influence on visual processing and survival of the retina through the neurohormones melatonin and dopamine and their respective action on MT₁ and D₄ receptors (for review, see Ref. 5). The main objective of this study was to identify candidate genes for linking clock-dependent neurohormone release with daily adaptation and healthiness of the retina. For this purpose, transcriptional regulators were screened for dependence on melatonin signaling/MT₁ receptors or dopamine signaling/D₄ receptors




FIGURE 1. Twenty-four-hour profiling in whole retina and photoreceptor cells. 24-hour profiling of the transcriptional regulators *Dbp*, *Egr-1*, *Fos*, *Nr1d1*, *Nr2e3*, *Nr4a1*, *Pgc-1α*, *Rorβ* in preparations of the whole retina under light/dark (LD) 12:12 (blue lines), in microdissected photoreceptor cells under light/dark (LD) 12:12 (red lines), and in preparations of the whole retina of mice under constant darkness (DD) (black lines). The mRNA levels are plotted as a function of Zeitgeber time (ZT), and the lines represent the periodic sinusoidal functions (solid and dotted lines for $P < 0.05$ and $P > 0.05$ in cosinor analysis). The solid bars indicate the dark period. Data represent a percentage of the average value of transcript amount during the 24-hour period. The value of ZT0 was plotted twice at both ZT0 and ZT24. Each value represents mean \pm SEM ($n = 4$). Statistical analysis of transcriptional profiling is provided in Table 2.

TABLE 2. Statistical Analysis of Transcriptional Profiling Illustrated in Figures 1 and 2

Gene	Retina of Mouse, C3H/He (<i>rd</i> ^{+/+})			PRCs of Mouse, C3H/He (<i>rd</i> ^{+/+})			Retina of Mouse, C3H/He (<i>rd</i> ^{+/+}); DD			WT C57BL/6Jb; LD		
	1-Way ANOVA		Cosinor Analysis	1-Way ANOVA		Cosinor Analysis	1-Way ANOVA		Cosinor Analysis	1-Way ANOVA		Cosinor Analysis
	P Value	Acrophase, h	Amplitude, %	P Value	Acrophase, h	Amplitude, %	P Value	Acrophase, h	Amplitude, %	P Value	Acrophase, h	Amplitude, %
<i>Dbbp</i>	< 0.001	7.9	32.4	< 0.001	8.6	42.1	= 0.039	< 0.05	5.0	25.7	< 0.001	< 0.05
<i>Egr1</i>	< 0.001	-	-	< 0.001	-	-	< 0.001	> 0.05	-	-	= 0.041	> 0.05
<i>Fos</i>	= 0.041	15.2	66.5	< 0.001	16.1	86.1	< 0.001	> 0.05	15.9	58.5	= 0.001	< 0.05
<i>Nr1d1</i>	= 0.041	5.5	19.6	= 0.032	-	-	= 0.043	< 0.05	2.8	24.8	= 0.047	< 0.05
<i>Nr2e3</i>	< 0.001	3.3	30.4	< 0.001	4.2	40.0	< 0.001	< 0.05	1.1	19.7	= 0.036	< 0.05
<i>Nr4a1</i>	< 0.001	16.4	77.6	< 0.001	17.4	105.3	< 0.001	< 0.05	16.6	62.7	< 0.001	< 0.05
<i>Pgc-1α</i>	= 0.005	10.8	30.0	< 0.001	6.5	35.1	= 0.047	< 0.05	5.9	18.7	= 0.477	> 0.05
<i>Rorb</i>	= 0.041	13.9	34.0	= 0.020	17.7	22.1	< 0.001	> 0.05	-	-	= 0.006	> 0.05

Significance of daily regulation was tested by ANOVA. The acrophase (time of peak gene expression) and the amplitude of daily profiles were obtained from cosinor analysis.

TABLE 3. Statistical Analysis of Transcriptional Profiling Illustrated in Figures 3 and 4

Gene	<i>C3H/f^{+/+}MT₁^{+/+}</i>			<i>C3H/f^{+/+}Drd₄^{+/+}</i>			<i>C3H/f^{+/+}Drd₄^{-/-}</i>					
	1-Way ANOVA		Cosinor Analysis	1-Way ANOVA		Cosinor Analysis	1-Way ANOVA		Cosinor Analysis			
	P Value	Acrophase, h	Amplitude, %	P Value	Acrophase, h	Amplitude, %	P Value	Acrophase, h	Amplitude, %			
<i>Dbbp</i>	= 0.014	6.7	32.1	< 0.001	6.7	28.8	= 0.013	< 0.05	4.4	43.8	= 0.010	> 0.05
<i>Egr1</i>	< 0.001	-	-	< 0.001	-	-	< 0.001	> 0.05	-	-	< 0.001	> 0.05
<i>Fos</i>	< 0.001	-	-	< 0.001	-	-	< 0.001	> 0.05	-	-	< 0.001	< 0.05
<i>Nr1d1</i>	< 0.001	7.0	25.0	< 0.001	5.1	21.9	< 0.001	< 0.05	4.6	22.5	< 0.001	< 0.05
<i>Nr2e3</i>	< 0.001	3.4	23.5	< 0.001	2.0	21.9	= 0.005	> 0.05	-	-	< 0.001	> 0.05
<i>Nr4a1</i>	< 0.001	14.1	61.8	< 0.001	14.3	48.7	= 0.040	< 0.05	15.3	46.9	= 0.040	< 0.05
<i>Pgc-1α</i>	= 0.001	7.3	25.8	= 0.066	-	-	= 0.001	< 0.05	6.7	16.4	= 0.003	< 0.05
<i>Rorb</i>	= 0.017	15.9	23.9	< 0.001	15.3	22.2	= 0.011	> 0.05	-	-	= 0.041	< 0.05

Significance of daily regulation was tested by ANOVA. The acrophase (the time of peak gene expression) and the amplitude of daily patterns were obtained from cosinor analysis.




FIGURE 2. Daily profiling in melatonin-proficient and -deficient mice. Daily profiling of the transcriptional regulators *Dbp*, *Egr-1*, *Fos*, *Nr1d1*, *Nr2e3*, *Nr4a1*, *Pgc-1α*, *Rorb* in retina of melatonin-proficient C3H/He (*rd*^{+/+}) mice (*blue lines*, corresponding to those in Fig. 1) and melatonin-deficient C57BL/6j mice (*red lines*). The mRNA levels are plotted as a function of Zeitgeber time (ZT), and the lines represent the periodic sinusoidal functions (*solid and dotted lines* for *P* < 0.05 and *P* > 0.05 in cosinor analysis). The *solid bars* indicate the dark period. Data represent a percentage of the average value of transcript amount during the 24-hour period. The value of ZT0 was plotted twice at both ZT0 and ZT24. Each value represents mean ± SEM (*n* = 4). Statistical analysis of transcriptional profiling is provided in Table 2.




FIGURE 3. Twenty-four-hour profiling in WT and $MT_1^{-/-}$ mice. 24-hour profiling of the transcriptional regulators *Dbp*, *Egr-1*, *Fos*, *Nr1d1*, *Nr2e3*, *Nr4a1*, *Pgc-1α*, *Rorb* in retina of WT (blue lines) and $MT_1^{-/-}$ mice (red lines). The mRNA levels are plotted as a function of Zeitgeber time (ZT), and the lines represent the periodic sinusoidal functions (solid and dotted lines for $P < 0.05$ and $P > 0.05$ in cosinor analysis). The solid bars indicate the dark period. Data represent a percentage of the average value of transcript amount during the 24-hour period. The value of ZT0 was plotted twice at both ZT0 and ZT24. Each value represents mean \pm SEM ($n = 4$). Statistical analysis of transcriptional profiling is provided in Table 3.




FIGURE 4. Twenty-four-hour profiling in WT and *Drd4*^{-/-} mice. 24-hour profiling of the transcriptional regulators *Dbp*, *Egr-1*, *Fos*, *Nr1d1*, *Nr2e3*, *Nr4a1*, *Pgc-1α*, *Rorb* in retina of WT (blue lines) and *Drd4*^{-/-} mice (red lines). The mRNA levels are plotted as a function of Zeitgeber time (ZT), and the lines represent the periodic sinusoidal functions (solid and dotted lines for $P < 0.05$ and $P > 0.05$ in cosinor analysis). The solid bars indicate the dark period. Data represent a percentage of the average value of transcript amount during the 24-hour period. The value of ZT0 was plotted twice at both ZT0 and ZT24. Each value represents mean ± SEM ($n = 4$). Statistical analysis of transcriptional profiling is provided in Table 3.

that fulfill the requirements to (1) cycle in PCRs (*Dbp*³⁷; Fig. 1; *Egr-1*; Fig. 1; *Fos*, *Nr1d1*⁴⁰; Fig. 1; *Nr2e3*; Fig. 1; *Nr4a1*; Fig. 1; *Pgc-1α*⁴³; Fig. 1; *Rorb*⁴⁰; Fig. 1), (2) be driven by a circadian clock (*Dbp*³⁷; Fig. 1; *Egr-1*³⁹; Fig. 1; *Fos*³⁹; Fig. 1; *Nr1d1*⁴⁰; Fig. 1; *Nr2e3*; Fig. 1; *Nr4a1*; Fig. 1; *Pgc-1α*; Fig. 1; *Rorb*³²; Fig. 1), and (3) influence retinal physiology or pathology (*Egr-1*, *Fos*^{34,36}, *Nr1d1*³³, *Nr2e3*²⁹⁻³¹, *Nr4a1*³⁴, *Nr1d1*; *Pgc-1α*²⁷; *Rorb*³²). As our experimental system we used mice with targeted deletion of MT₁ receptors and D₄ receptors, respectively, that we have crossed onto the C3H^{+/+} background.^{56,57} Thus the mouse model used in this study combines physiological melatonin formation and intact photoreceptors.

The data obtained determine the transcriptional coactivator *Pgc-1α* as a transcriptional target of circadian melatonin/MT₁ receptor signaling. This becomes evident from the finding that the daily rhythmicity of *Pgc-1α* mRNA amount was dampened in both melatonin-deficient (C57BL/6Jb) and MT₁^{-/-} mice.

Melatonin/MT₁ receptor-dependent control of *Pgc-1α* may take place in PRCs. This follows from the observation that PRCs combine a prominent rhythmicity of *Pgc-1α* (this study) with high density of MT₁ receptors.^{11,12} Of interest is that, in PCRs, melatonin appears to act as an autocrine signal.¹¹ Therefore PRCs may regulate their own *Pgc-1α* expression by melatonin release and its subsequent action on MT₁ receptors located at their cell membrane. Recent studies have shown that the action of melatonin in the PRCs is mediated by MT₁/MT₂ heteromers.⁹ Therefore, MT₁ receptor deficiency may result in nonfunctional MT₁/MT₂ heterodimers, and MT₂ knockout mice would have produced similar results from those obtained in MT₁.

Pgc-1α is a master regulator of glucose and energy metabolism in various tissues (for review, see Ref. 58) including retina.²⁷ Therefore the observed melatonin/MT₁ receptor-dependent regulation of *Pgc-1α* may contribute to daily changes in the energy metabolism of the retina and thus to its ability to comply with 24-hour changes in metabolic demands.⁵⁹

Remarkably, *Pgc-1α* knockout mice reportedly suffer from increased light damage susceptibility, overexpression of *Pgc-1α* has antiapoptotic effects, and *Pgc-1α* expression is reduced in different mouse models of retinitis pigmentosa.²⁷ This suggests that *Pgc-1α* provides protection against light damage and is therefore a promising candidate gene for transferring the positive effect of melatonin/MT₁ receptors on the viability of retinal PRCs and ganglion cells.¹¹ Accordingly, melatonin/MT₁ receptor-dependent control of *Pgc-1α* could contribute to the protection of retinal cells against light damage and on a long-term basis to the survival of retinal cells during aging.

Outside the retina, MT₁ receptor-dependent expression of *Pgc-1α* could play a role in the pathogenesis of type 2 diabetes. This becomes evident from the observations that the removal of MT₁ receptors⁶⁰ or *Pgc-1α* (for review, see Ref. 61) equally leads to increased insulin resistance in peripheral tissues.

As with *Pgc-1α* in the melatonergic system, the nuclear orphan receptor *Nr4a1* (Nur77, Ngfi-b) was identified to respond to the dopaminergic system. This is evident from the finding that the daily rhythmicity of *Nr4a1* is reduced in *Drd4*^{-/-} mice. Dopaminergic regulation of retinal *Nr4a1* probably occurs in PRC since this cell type combines circadian regulation of *Nr4a1* (this study) with the abundance of D₄ receptors.^{62,63} In the retina, clock-dependent dopamine release occurs from the unique population of cells in the inner nuclear layer that are either amacrine or interplexiform neurons.¹⁵ The dopamine neurons in retina express a full complement of core circadian clock genes.^{64,65} Therefore circadian/dopaminergic control of *Nr4a1* in PRCs appears to

be driven by these cell types. Furthermore, *Drd4*, the gene that encodes the D₄ receptor, is itself a circadian clock-controlled gene in retina,^{3,20,42,66} and this too may contribute to circadian regulation of *Nr4a1*.

Dopaminergic regulation of *Nr4a1* was until now believed to be confined to the motor/motivation system (for reviews, see Refs. 67, 68). Therefore its concurrent occurrence in the visual (this study) and the motor/motivation system suggests that it appears to be a more general feature of different functional brain systems affected by dopamine.

Furthermore, *Nr4a1* is a promising candidate gene for being responsible for dysfunction of the retinal dopaminergic system and its subsequent visual defects in type I diabetes mellitus.⁶⁹ This becomes evident from the observations that *Nr4a1* expression on one hand influences the functioning of the dopaminergic system (for reviews, see Refs. 67, 68) and on the other hand depends on glucose metabolism.⁷⁰⁻⁷²

Compelling experimental evidence indicates that *Nr4a1* not only contributes to dopaminergic responses but also is a decisive factor in dopamine-related neuroadaptation (for reviews, see Refs. 67, 68). If this is also valid in retina, circadian/dopaminergic regulation of *Nr4a1* expression (this study) may feed back to the dopaminergic system of the retina to promote its reported circadian plasticity.⁴²

It is important to note that—at variance with previous studies dealing with the role of melatonin in retina^{51,52}—the present investigation was performed in mouse strains/genotypes that combine melatonin proficiency with intact PRCs and thus in mouse models that reflect retinal physiology as closely as possible. On the other hand, a general limitation of the present study is that the observed phenotypes of the mouse models used may reflect not only a direct action of the neurotransmitters/receptors, but also the secondary effects downstream of them. Since melatonin signaling⁷³⁻⁷⁵ and dopamine signaling⁷⁶⁻⁷⁸ make a cellular feedback loop in which melatonin inhibits the release of dopamine through melatonin receptors,⁷³⁻⁷⁵ whereas dopamine downregulates melatonin formation, probably through dopamine D₄ receptors,^{76,77} this includes a possibility that genetic disturbance of melatonin signaling affects dopamine signaling and vice versa.

In conclusion, the data of the present investigation determined candidate transcriptional regulators for linking the circadian neurohormone systems to adaptation, function, and healthiness of the retina. They could provide a reliable basis for further investigations that may achieve a better understanding of how clock-dependent neurotransmitter release influences visual processing and survival of the retinal neurons.

Acknowledgments

The authors thank Ute Frederiksen and Kristina Schäfer for their excellent technical assistance, Susanne Rometsch and Bettina Wiechers-Schmied for secretarial help, Klaus Wolloscheck for statistical advice, and Russell G. Foster for providing us with C3H/He (*rd*^{+/+}) mice. The data contained in this study are included in the thesis of Stefanie Kunst as a partial fulfillment of her doctorate degree at the Johannes Gutenberg University, Mainz. There are no conflicts of interest, financial or otherwise, to be disclosed by the authors.

Supported by the BMBF “HOPE2” (01GM1108D to UW), the Faun-Stiftung Nurnberg (to UW), the European Community FP7/2009/241955 (SYSCILIA to UW), the Naturwissenschaftlich-medizinisches Forschungszentrum of the University Mainz (to UW and RS), the National Institutes of Health (Grants R01 EY004864 to PMI, P30 EY006360 to PMI, T32 EY007092 to PMI; EY022216 to GT) and an unrestricted grant from Research to Prevent Blindness,

Inc. (RPB). PMI was a recipient of a Senior Scientific Investigator Award from RPB.

Disclosure: **S. Kunst**, None; **T. Wolloscheck**, None; **D.K. Kelleher**, None; **U. Wolfrum**, None; **S.A. Sargsyan**, None; **P.M. Iuvone**, None; **K. Baba**, None; **G. Tosini**, None; **R. Spessert**, None

References

- Cameron MA, Barnard AR, Hut RA, et al. Electretinography of wild-type and Cry mutant mice reveals circadian tuning of photopic and mesopic retinal responses. *J Biol Rhythms*. 2008;23:489-501.
- Cameron MA, Barnard AR, Lucas RJ. The electroretinogram as a method for studying circadian rhythms in the mammalian retina. *J Genet*. 2008;87:459-466.
- Storch KF, Paz C, Signorovitch J, et al. Intrinsic circadian clock of the mammalian retina: importance for retinal processing of visual information. *Cell*. 2007;130:730-741.
- Tosini G, Menaker M. The clock in the mouse retina: melatonin synthesis and photoreceptor degeneration. *Brain Res*. 1998;789:221-228.
- McMahon DG, Iuvone PM, Tosini G. Circadian organization of the mammalian retina: from gene regulation to physiology and diseases. *Prog Retin Eye Res*. 2014;39:58-76.
- Iuvone PM, Tosini G, Pozdeyev N, et al. Circadian clocks, clock networks, arylalkylamine N-acetyltransferase, and melatonin in the retina. *Prog Retin Eye Res*. 2005;24:433-456.
- Tosini G, Pozdeyev N, Sakamoto K, Iuvone PM. The circadian clock system in the mammalian retina. *Bioessays*. 2008;30:624-633.
- Tosini G, Owino S, Guillaume JL, Jockers R. Understanding melatonin receptor pharmacology: latest insights from mouse models, and their relevance to human disease. *Bioessays*. 2014;36:778-787.
- Baba K, Benleulmi-Chaachoua A, Journe AS, et al. Heteromeric MT1/MT2 melatonin receptors modulate photoreceptor function. *Sci Signal*. 2013;6:ra89.
- Baba K, Mazzoni F, Owino S, et al. Age-related changes in the daily rhythm of photoreceptor functioning and circuitry in a melatonin-proficient mouse strain. *PLoS One*. 2012;7:e37799.
- Baba K, Pozdeyev N, Mazzoni F, et al. Melatonin modulates visual function and cell viability in the mouse retina via the MT1 melatonin receptor. *Proc Natl Acad Sci U S A*. 2009;106:15043-15048.
- Sengupta A, Baba K, Mazzoni F, et al. Localization of melatonin receptor 1 in mouse retina and its role in the circadian regulation of the electroretinogram and dopamine levels. *PLoS One*. 2011;6:e24483.
- Iuvone PM, Galli CL, Garrison-Gund CK, Neff NH. Light stimulates tyrosine hydroxylase activity and dopamine synthesis in retinal amacrine neurons. *Science*. 1978;202:901-902.
- Nir I, Haque R, Iuvone PM. Diurnal metabolism of dopamine in dystrophic retinas of homozygous and heterozygous retinal degeneration slow (rds) mice. *Brain Res*. 2000;884:13-22.
- Witkovsky P. Dopamine and retinal function. *Doc Ophthalmol*. 2004;108:17-40.
- Jackson CR, Ruan GX, Aseem F, et al. Retinal dopamine mediates multiple dimensions of light-adapted vision. *J Neurosci*. 2012;32:9359-9368.
- Hwang CK, Chaurasia SS, Jackson CR, et al. Circadian rhythm of contrast sensitivity is regulated by a dopamine-neuronal PAS-domain protein 2-adenylyl cyclase 1 signaling pathway in retinal ganglion cells. *J Neurosci*. 2013;33:14989-14997.
- Pozdeyev N, Tosini G, Li L, et al. Dopamine modulates diurnal and circadian rhythms of protein phosphorylation in photoreceptor cells of mouse retina. *Eur J Neurosci*. 2008;27:2691-2700.
- Cahill GM, Besharse JC. Resetting the circadian clock in cultured *Xenopus* eyecups: regulation of retinal melatonin rhythms by light and D2 dopamine receptors. *J Neurosci*. 1991;11:2959-2971.
- Jackson CR, Chaurasia SS, Hwang CK, Iuvone PM. Dopamine D(4) receptor activation controls circadian timing of the adenylyl cyclase 1/cyclic AMP signaling system in mouse retina. *Eur J Neurosci*. 2011;34:57-64.
- Yu RT, Chiang MY, Tanabe T, et al. The orphan nuclear receptor Tlx regulates Pax2 and is essential for vision. *Proc Natl Acad Sci U S A*. 2000;97:2621-2625.
- Haider NB, Mollema N, Gaule M, et al. Nr2e3-directed transcriptional regulation of genes involved in photoreceptor development and cell-type specific phototransduction. *Exp Eye Res*. 2009;89:365-372.
- Andreazzoli M. Molecular regulation of vertebrate retina cell fate. *Birth Defects Res C Embryo Today*. 2009;87:284-295.
- Bassett EA, Wallace VA. Cell fate determination in the vertebrate retina. *Trends Neurosci*. 2012;35:565-573.
- Zhang CL, Zou Y, Yu RT, Gage FH, Evans RM. Nuclear receptor TLX prevents retinal dystrophy and recruits the corepressor atrophin1. *Genes Dev*. 2006;20:1308-1320.
- Onishi A, Peng GH, Poth EM, et al. The orphan nuclear hormone receptor ERRbeta controls rod photoreceptor survival. *Proc Natl Acad Sci U S A*. 2010;107:11579-11584.
- Egger A, Samardzija M, Sothilingam V, et al. PGC-1alpha determines light damage susceptibility of the murine retina. *PLoS One*. 2012;7:e31272.
- Sharma YV, Cojocar RI, Ritter LM, et al. Protective gene expression changes elicited by an inherited defect in photoreceptor structure. *PLoS One*. 2012;7:e31371.
- Haider NB, Jacobson SG, Cideciyan AV, et al. Mutation of a nuclear receptor gene, NR2E3, causes enhanced S cone syndrome, a disorder of retinal cell fate. *Nat Genet*. 2000;24:127-131.
- Sharon D, Sandberg MA, Caruso RC, Berson EL, Dryja TP. Shared mutations in NR2E3 in enhanced S-cone syndrome, Goldmann-Favre syndrome, and many cases of clumped pigmentary retinal degeneration. *Arch Ophthalmol*. 2003;121:1316-1323.
- Wright AF, Reddick AC, Schwartz SB, et al. Mutation analysis of NR2E3 and NRL genes in Enhanced S Cone Syndrome. *Hum Mutat*. 2004;24:439.
- Andre E, Conquet F, Steinmayr M, et al. Disruption of retinoid-related orphan receptor beta changes circadian behavior, causes retinal degeneration and leads to vacillans phenotype in mice. *EMBO J*. 1998;17:3867-3877.
- Cruz NM, Yuan Y, Leehy BD, et al. Modifier genes as therapeutics: the nuclear hormone receptor Rev Erb alpha (Nr1d1) rescues Nr2e3 associated retinal disease. *PLoS One*. 2014;9:e87942.
- Kamphuis W, Dijk F, van Soest S, Bergen AA. Global gene expression profiling of ischemic preconditioning in the rat retina. *Mol Vis*. 2007;13:1020-1030.
- Bakalash S, Pham M, Koronyo Y, et al. Egr1 expression is induced following glatiramer acetate immunotherapy in rodent models of glaucoma and Alzheimer's disease. *Invest Ophthalmol Vis Sci*. 2011;52:9033-9046.
- Wurm A, Iandiev I, Uhlmann S, et al. Effects of ischemia-reperfusion on physiological properties of Muller glial cells in the porcine retina. *Invest Ophthalmol Vis Sci*. 2011;52:3360-3367.
- Hiragaki S, Baba K, Coulson E, et al. Melatonin signaling modulates clock genes expression in the mouse retina. *PLoS One*. 2014;9:e106819.

38. Agarwal N. Diurnal expression of NGF1-A mRNA in retinal degeneration slow (rds) mutant mouse retina. *FEBS Lett.* 1994; 339:253-257.
39. Humphries A, Carter DA. Circadian dependency of nocturnal immediate-early protein induction in rat retina. *Biochem Biophys Res Commun.* 2004;320:551-556.
40. Sandu C, Hicks D, Felder-Schmittbuhl MP. Rat photoreceptor circadian oscillator strongly relies on lighting conditions. *Eur J Neurosci.* 2011;34:507-516.
41. Mollema NJ, Yuan Y, Jelcick AS, et al. Nuclear receptor Rev-erb alpha (Nr1d1) functions in concert with Nr2e3 to regulate transcriptional networks in the retina. *PLoS One.* 2011;6:e17494.
42. Bai L, Zimmer S, Rickes O, et al. Daily oscillation of gene expression in the retina is phase-advanced with respect to the pineal gland. *Brain Res.* 2008;1203:89-96.
43. Kunst S, Wolloscheck T, Holter P, et al. Transcriptional analysis of rat photoreceptor cells reveals daily regulation of genes important for visual signaling and light damage susceptibility. *J Neurochem.* 2013;124:757-769.
44. Kamphuis W, Cailotto C, Dijk F, Bergen A, Buijs RM. Circadian expression of clock genes and clock-controlled genes in the rat retina. *Biochem Biophys Res Commun.* 2005;330:18-26.
45. Goldmann T, Burgemeister R, Sauer U, et al. Enhanced molecular analyses by combination of the HOPE-technique and laser microdissection. *Diagn Pathol.* 2006;1:2.
46. Schneider K, Tippmann S, Spiwoks-Becker I, et al. Unique clockwork in photoreceptor of rat. *J Neurochem.* 2010;115: 585-594.
47. Mears AJ, Kondo M, Swain PK, et al. Nrl is required for rod photoreceptor development. *Nat Genet.* 2001;29:447-452.
48. Osborne NN, Patel S, Vigny A. Dopaminergic neurones in various retinas and the postnatal development of tyrosine-hydroxylase immunoreactivity in the rabbit retina. *Histochemistry.* 1984;80:389-393.
49. Cornelissen G. Cosinor-based rhythmometry. *Theoret Biol Med Model.* 2014;11:16.
50. Refinetti R, Lissen GC, Halberg F. Procedures for numerical analysis of circadian rhythms. *Biol Rhythm Res.* 2007;38:275-325.
51. Dinet V, Ansari N, Torres-Farfan C, Korf HW. Clock gene expression in the retina of melatonin-proficient (C3H) and melatonin-deficient (C57BL) mice. *J Pineal Res.* 2007;42:83-91.
52. Dinet V, Korf HW. Impact of melatonin receptors on pCREB and clock-gene protein levels in the murine retina. *Cell Tissue Res.* 2007;330:29-34.
53. Liu X, Zhang Z, Ribelayga CP. Heterogeneous expression of the core circadian clock proteins among neuronal cell types in mouse retina. *PLoS One.* 2012;7:e50602.
54. Ait-Hmyed O, Felder-Schmittbuhl MP, Garcia-Garrido M, et al. Mice lacking Period 1 and Period 2 circadian clock genes exhibit blue cone photoreceptor defects. *Eur J Neurosci.* 2013;37:1048-1060.
55. Dkhissi-Benyahya O, Coutanson C, Knoblauch K, et al. The absence of melanopsin alters retinal clock function and dopamine regulation by light. *Cell Mol Life Sci.* 2013;70: 3435-3447.
56. Freedman MS, Lucas RJ, Soni B, et al. Regulation of mammalian circadian behavior by non-rod, non-cone, ocular photoreceptors. *Science.* 1999;284:502-504.
57. Lupi D, Cooper HM, Froehlich A, et al. Transgenic ablation of rod photoreceptors alters the circadian phenotype of mice. *Neuroscience.* 1999;89:363-374.
58. Liu C, Lin JD. PGC-1 coactivators in the control of energy metabolism. *Acta Biochim Biophys Sin (Shanghai).* 2011;43: 248-257.
59. Ames A III, Li YY. Energy requirements of glutamatergic pathways in rabbit retina. *J Neurosci.* 1992;12:4234-4242.
60. Contreras-Alcantara S, Baba K, Tosini G. Removal of melatonin receptor type 1 induces insulin resistance in the mouse. *Obesity (Silver Spring).* 2010;18:1861-1863.
61. Rowe GC, Arany Z. Genetic models of PGC-1 and glucose metabolism and homeostasis. *Rev Endocr Metab Disord.* 2014;15:21-29.
62. Cohen AI, Todd RD, Harmon S, O'Malley KL. Photoreceptors of mouse retinas possess D4 receptors coupled to adenylate cyclase. *Proc Natl Acad Sci U S A.* 1992;89:12093-12097.
63. Kay JN, De la Huerta I, Kim JJ, et al. Retinal ganglion cells with distinct directional preferences differ in molecular identity, structure, and central projections. *J Neurosci.* 2011;31:7753-7762.
64. Ruan GX, Zhang DQ, Zhou T, Yamazaki S, McMahon DG. Circadian organization of the mammalian retina. *Proc Natl Acad Sci U S A.* 2006;103:9703-9708.
65. Dorenbos R, Contini M, Hirasawa H, Gustincich S, Raviola E. Expression of circadian clock genes in retinal dopaminergic cells. *Vis Neurosci.* 2007;24:573-580.
66. Klitten LL, Rath MF, Coon SL, et al. Localization and regulation of dopamine receptor D4 expression in the adult and developing rat retina. *Exp Eye Res.* 2008;87:471-477.
67. Levesque D, Rouillard C. Nur77 and retinoid X receptors: crucial factors in dopamine-related neuroadaptation. *Trends Neurosci.* 2007;30:22-30.
68. Campos-Melo D, Galleguillos D, Sanchez N, Gysling K, Andres ME. Nur transcription factors in stress and addiction. *Front Mol Neurosci.* 2013;6:44.
69. Aung MH, Park HN, Han MK, et al. Dopamine deficiency contributes to early visual dysfunction in a rodent model of type 1 diabetes. *J Neurosci.* 2014;34:726-736.
70. Pei L, Waki H, Vaitheesvaran B, et al. NR4A orphan nuclear receptors are transcriptional regulators of hepatic glucose metabolism. *Nat Med.* 2006;12:1048-1055.
71. Fu Y, Luo L, Luo N, Zhu X, Garvey WT. NR4A orphan nuclear receptors modulate insulin action and the glucose transport system: potential role in insulin resistance. *J Biol Chem.* 2007; 282:31525-31533.
72. Kanzleiter T, Wilks D, Preston E, et al. Regulation of the nuclear hormone receptor nur77 in muscle: influence of exercise-activated pathways in vitro and obesity in vivo. *Biochim Biophys Acta.* 2009;1792:777-782.
73. Dubocovich ML. Melatonin is a potent modulator of dopamine release in the retina. *Nature.* 1983;306:782-784.
74. Boatright JH, Rubim NM, Iuvone PM. Regulation of endogenous dopamine release in amphibian retina by melatonin: the role of GABA. *Vis Neurosci.* 1994;11:1013-1018.
75. Ribelayga C, Wang Y, Mangel SC. A circadian clock in the fish retina regulates dopamine release via activation of melatonin receptors. *J Physiol.* 2004;554(pt 2):467-482.
76. Zawilska JB, Iuvone PM. Melatonin synthesis in chicken retina: effect of kainic acid-induced lesions on the diurnal rhythm and D2-dopamine receptor-mediated regulation of serotonin N-acetyltransferase activity. *Neurosci Lett.* 1992;135:71-74.
77. Nguyen-Legros J, Chanut E, Versaux-Botteri C, Simon A, Trouvin JH. Dopamine inhibits melatonin synthesis in photoreceptor cells through a D2-like receptor subtype in the rat retina: biochemical and histochemical evidence. *J Neurochem.* 1996;67:2514-2520.
78. Tosini G, Dirden JC. Dopamine inhibits melatonin release in the mammalian retina: in vitro evidence. *Neurosci Lett.* 2000; 286:119-122.