

SCIENTIFIC REPORTS



OPEN

A Casparian strip domain-like gene, *CASPL*, negatively alters growth and cold tolerance

Jinghua Yang^{1,2,*}, Changqing Ding^{1,2,*}, Baochen Xu^{1,2}, Cuiting Chen^{1,2}, Reena Narsai³, Jim Whelan³, Zhongyuan Hu^{1,2} & Mingfang Zhang^{1,2}

Received: 26 June 2015

Accepted: 24 August 2015

Published: 24 September 2015

A cold-induced transcript encoding a Casparian strip membrane domain (CASP)-like protein (*CICASPL*) was identified in watermelon (*Citrullus lanatus*). Fluorescence microscopy analysis showed that *CICASPL*-GFP is localized in the plasma membrane. The orthologous gene in *Arabidopsis thaliana* (*AtCASPL4C1*) was also found to play an important role in cold tolerance. Expression analysis using a β -glucuronidase (GUS) reporter reveals that *AtCASPL4C1* is widely expressed in a variety of organs and is cold inducible. Analysis of *AtCASPL4C1* T-DNA knock-out plants showed altered growth dynamics, faster growth, increased biomass (dry weight) and earlier flowering compared to wild type (Col-o) and *CICASPL* overexpressing plants. *AtCASPL4C1* knock-out plants showed elevated tolerance to cold stress, while overexpressing *CICASPL* resulted in increased sensitivity to cold stress in *Arabidopsis*. Interestingly, *AtCASPL4C1* knock-out plants did not display significant alterations in the Casparian strip formation in roots. Thus, the combination of these results suggests a role for *CICASPL* and *AtCASPL4C1* beyond Casparian strip formation in roots, possibly indicating a more fundamental role in vascular tissue.

Abiotic stresses are major environmental factors that adversely affect plant growth, development and yield. Plants have evolved complex signaling networks to adapt to abiotic stresses via modulating various physiological and biochemical processes^{1–3}. These stress signals are perceived by receptors, transduced and propagated by downstream effectors, ultimately altering the expression of a variety of genes that determine growth, tolerance and/or survival depending on the severity of the environmental conditions^{2,4}.

Transmembrane (TM) proteins located in the plasma membrane are known to have diverse physiological functions including signal perception and recognition, via ion and metabolite exchange. In *Arabidopsis*, approximately 6,500 proteins are predicted to be TM proteins⁵. Some integral membrane proteins are induced by stress, such as drought stress in *Triticum turgidum*⁶, or salt stress in *Xerophyta viscosa*⁷ as well as having a role during development, e.g. kernel development in *Hordeum vulgare*⁸.

The Casparian strip, first described by Robert Caspary in 1865⁹, is a ring-like cell wall structure in the root endodermis of vascular plants. The role of the Casparian strip is to block the passive flow of materials in vascular plants¹⁰. Recent reports showed that the Casparian strip is composed of a lignin polymer without suberin in *Arabidopsis*¹¹. This lignin polymer structure generates a para-cellular barrier, analogous to tight junctions in animals, that is thought to be crucial for selective nutrient uptake, exclusion of pathogens, and many other processes¹⁰. Casparian strip membrane domain proteins (CASP, CASP1/2/3/4/5) are crucial for Casparian strip formation in the endodermis in plants. CASP1/2/3/4/5 belong to ‘uncharacterized protein family’ UPF0497 (39 members) in *Arabidopsis*⁹. These CASPs are

¹Laboratory of Germplasm Innovation and Molecular Breeding, Institute of Vegetable Science, Zhejiang University, Hangzhou, 310058, P. R. China. ²Key laboratory of Horticultural Plant Growth, Development & Quality Improvement, Ministry of Agriculture, Hangzhou, 310058, P. R. China. ³Department of Animal, Plant and Soil Science, School of Life Science, Australian Research Council Centre of Excellence in Plant Energy Biology, LaTrobe University, Bundoora, Victoria 3086, Australian. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to M.Z. (email: mfzhang@zju.edu.cn)

proposed to form an extensive, transmembrane polymeric platform and were speculated to guide the assembly and activity of lignin biosynthetic enzymes⁹. Recently, evolutionary analysis of CASP family genes indicated that CASPL genes belong to the MARVEL (MAL and related proteins for vesicle trafficking and membrane link) protein family, which has only been experimentally described in metazoans, to date¹².

Previously, we identified a gene encoding a cold-induced integral membrane protein in watermelon¹³. The orthologue to this gene was identified to be *At3g55390* in *Arabidopsis thaliana*, where it is annotated as the CASP-LIKE PROTEIN 4C1. *At3g55390* belongs to the CASP protein family containing the five CASP genes (*CASP1/2/3/4/5*), which are known to mediate Casparian strip formation in plants⁹. Thereafter, these are referred to as *CICASPL* (from watermelon) and *AtCASPL4C1* (CASP-like) or *AtCASPL4C1* from *Arabidopsis*, respectively. In this study, we investigated the role of *CICASPL* and *AtCASPL4C1* in growth and cold tolerance in watermelon and *Arabidopsis*, respectively.

Materials and Methods

Plant Materials and treatment. An IVSM-9 inbred line of watermelon (*Citrullus lanatus*) and wild type (WT) of *Arabidopsis thaliana* (*Arabidopsis thaliana*) (Col-0) were used for amplification of *CICASPL* and *AtCASPL4C1* (*At3G55390*) genes. WT *Arabidopsis thaliana* (Col-0) was used for transformation to construct *OX-CICASPL* of *Arabidopsis*. The SALK_034800C line was used for screening of homozygous *Atcaspl4c1* knock-out mutant plants. 3-week-old plants in Jiffy seedling culture substrate or 2-week-old seedlings cultured in 1/2 MS medium of WT, *Atcaspl4c1* and *OX-CICASPL Arabidopsis* were used for analysis of cold stress and phenotypes evaluations. Tobacco (*Nicotianabens thamiana*) was used for transient expression of the *CICASPL* gene.

5 days old plants were transferred onto half-strength MS (Murashige-Skoog, sigma-Aldrich) medium and grew under 10°C, light/dark (16h/8h) conditions. For soil growth plants, 21 days old plants were used to cold treatments, under 10°C, light/dark (16h/8h) conditions. The pictures and data were collected at the indicated time. The values are means \pm SD (n = 20). Bar = 1 cm. Star signs indicate a significant difference (p < 0.05, student's *t* test).

Phylogenetic Tree Construction. The ClustalW program was used for alignment of *CICASPL* with the *AtCASPL4C1* protein sequence, which was obtained from the TAIR database. After alignment by ClustalW, a Neighbour-Joining tree was constructed by using MEGA 6.0, with 1000 as the number of bootstrap replications. The 39 members of CASP family genes were included from TAIR database. The TMHMM program was used for transmembrane region identification for *CICASPL* and *AtCASPL4C1*.

Transient Expression of *CICASPL* Gene in Tobacco. The amplification of *CICASPL* coding sequence without a termination codon was linked to pMDC83 binary expression vector to generate the *CICASPL*-pMDC83 by Gateway cloning¹⁴. The *CICASPL*-GFP plasmid was introduced into *Agrobacterium tumefaciens* strain (GV1301). Leaves from tobacco (*Nicotianabens thamiana*) were used for transformation and for checking transient expression according to the procedures¹⁵. Various organelle-targeted markers fused with RFP were used as controls to determine sub-cellular localization¹⁶. These tobacco leaves were observed using a confocal laser scanning microscope (ZEISS). The primers used in this study for GFP fusion are listed in Supplementary Table 1.

Expression of *CASPL*. *In silico* transcript abundance analysis for genes encoding proteins of the CASP family was carried out as outlined in Narsai *et al.*¹⁷. The promoter sequences were searched by A Database of Plant Promoter Sequences (<http://linux1.softberry.com/berry.phtml?topic=plant-prom&group=data&subgroup=plantprom>). The promoter region of *At3g55390* was amplified and inserted into a region upstream of the GUS gene of within the pMDC162 binary expression vector using the gateway system¹⁴. The resulting construct was transformed in the *Agrobacterium tumefaciens* strain GV3101. Transformation of *Arabidopsis* was conducted according to the floral-dip method¹⁸. Tissues from transgenic plants were collected in microcentrifuge tubes. Subsequently, the samples were stained as previously published procedures¹⁹.

Cold-induced expression of *CASPL* was analyzed using qPCR and *CASPL*-promoter analysis using *CICASPL* promoter-GUS activity after cold stress. GUS activity was measured as previously published procedures²⁰. The primers used for promoter amplification are listed in Supplementary Table 1.

***AtCASPL4C1* and *OX-CICASPL* Construction.** cDNA amplification of the *CICASPL* gene was inserted to pMDC32 binary expression vector to generate the *CICASPL*-pMDC32 by gateway approaches¹⁴. The *CICASPL*-pMDC32 plasmid was introduced into *Agrobacterium tumefaciens* strain (GV1301). The floral dipping method was used to generate *OX-CICASPL* of *Arabidopsis*¹⁸. For screening of transgenic plants, seeds harvested from transformed *Arabidopsis* plants were surface-sterilized and plated onto MS medium containing 50 mg/L hygromycin B. The plated seeds were vernalized at 4°C for 2–4 days in the dark to synchronize germination and then transferred to a growth chamber at 22°C (16/8h photo-period). 10-day-old *in vitro*-grown anti-hygromycin B seedlings were then transferred to soil mix. PCR and RT-PCR amplifications of the *CICASPL* gene from candidate transgenic *Arabidopsis* plants were

employed to confirm the successful transformation. WT and transgenic *Arabidopsis* T₃ lines were used in this study. The SALK_034800C line was used for the screening of homozygous *AtCASPL4C1* knock-out mutants of *Arabidopsis* using the three primers sets designed from the online service (<http://signal.salk.edu/tdnaprimers.2.html>). Primers used in this study for cDNA amplification of *CICASPL* gene and screening of *AtCASPL4C1* knock-out mutant of *Arabidopsis* are listed in Supplementary Table 1.

Casparian Strip Analysis. Propidium iodide (PI) staining was used to check Casparian strip formation in root as previously published procedures⁹. Roots from 5-day-old seedlings grown in MS medium were incubated in the dark for 10 min in 15 μM (10 μg/ml) PI (Invitrogen) and then were rinsed twice in water. The stained roots were observed using a fluorescence microscope (OLYMPUS BX51, Japan). Excitation and emission wavelengths are 488 nm and 500–550 nm.

Chlorophyll Fluorescence Analysis. 4-week-old seedlings of WT, *AtCASPL4C1* and *OX-CICASPL* of plants were cold treated at 10 °C and Fv/Fm measurements were taken. Individual plants were dark-adapted for 20 min prior to measurement. Chlorophyll fluorescence was measured in dark-adapted plants at 0, 6, 12, 24, 48 and 72 h under 10 °C cold stresses using a chlorophyll fluorescence system (M-Series Imaging-PAM, Germany).

Results

Characterization of CASPL. A CASP-like integral membrane protein gene, *CICASPL* (*Cl*a004012), was isolated in watermelon (*Citrullus lanatus*) using RT-PCR and genome blasting (Supplementary Figure 1-A). Bioinformatics analysis showed that *CICASPL* is orthologous to *At3g55390* from *Arabidopsis* (*AtCASPL4C1*, belonging to CASP family) (Fig. 1-A). Four transmembrane (TM) domains were predicted in *CICASPL* (amino acids 45–67, 87–109, 130–149 and 169–191) and *AtCASPL4C1* (*At3g55390*) proteins (amino acids 36–56, 78–98, 119–139 and 160–180 amino acid residues) by a variety of TM predication programs⁵ (Fig. 1-B). A total of 39 genes were identified that are defined as being part of the CASP family (UPF0497) in *Arabidopsis*, of which CASP1/2/3/4/5 has recently been identified to be associated with Casparian strip formation⁹. A phylogenetic tree for the CASP family in *Arabidopsis*²¹, defines 6 subfamilies using the Neighbor-Joining method and *CICASPL* branches in the same subfamily as *At3g55390* (*AtCASPL4C1*) of *Arabidopsis* (Fig. 1-C).

Subcellular Localization of CICASPL Protein. *CICASPL* is predicted to be targeted to the plasma membrane, according to the TargetP and SignalP subcellular localization prediction programs^{22,23}. The subcellular localization of *CICASPL* was investigated by the generation of a *CICASPL*-green fluorescent protein (GFP) fusion and transient expression in tobacco (*Nicotiana glauca*) leaves. Expression of the constructed *CICASPL-GFP* gene revealed a fluorescence signal exclusively in the plasma membrane, co-localizing with the plasma membrane marker¹⁶ that was tagged with red fluorescent protein (RFP) (Fig. 2).

Expression Analysis of CASPL. Transcriptomic analysis of public microarrays from the AtGenExpress developmental dataset⁵ and over seed germination¹⁷ indicated that *AtCASPL4C1* (*At3G55390*) is expressed in various tissues, but unlike some genes in this family it could not be defined as being predominantly expressed in root tissues (Fig. 3, black box). In contrast, a strong root expression profile is observed for several genes in this family, including CASP1/2/3/4/5 and other CASP-like genes (Fig. 3, yellow box). To determine spatial expression more precisely *AtCASPL4C1* expression patterns were determined using an *AtCASPL4C1*-promoter-GUS (for β-glucuronidase) reporter. Histochemical analysis expression in *planta*, with 12 independent lines all showed a very similar expression pattern, varied only slightly in the intensity of GUS staining. GUS-staining revealed that *AtCASPL4C1* expression is visible in roots and leaves (Fig. 4A). We can observe *AtCASPL4C1* expression in the vascular cylinder of roots, but no expression was detected in root tip (Fig. 4A,B). *AtCASPL4C1* expression was also detected in emerged lateral root (Fig. 4C). The filament, stigma and sepal of flowers also displayed intense staining (Fig. 4D), and signal could also be readily detected in siliques, but not in seeds (Fig. 4E). These results indicate that *AtCASPL4C1* is expressed in a wide range of variety of tissues and is not restricted to the vascular tissue of roots alone.

CASPL Negatively Regulates Growth Dynamics. To investigate the function of *CASPL* in plants, *AtCASPL4C1* knockout mutants and transgenic plants overexpressing *CICASPL* in *Arabidopsis* were generated (Supplementary Figure 1 and 2). Growth and development parameters were measured to quantify any changes in phenotype due to the absence or over-expression of *CICASPL*. The germination rate among wild type (Col-0), *AtCASPL4C1* knockout and *OX-CICASPL* plants did not significantly differ (Fig. 5A,C). The loss of *AtCASPL4C1*, however, did result in a small but significant increase in primary root length (Fig. 5D). In agreement with this, overexpressing *CICASPL* in *Arabidopsis* significantly decreased primary root length (Fig. 5D). *AtCASPL4C1* plants displayed significantly faster growth in several parameters, bigger plants (Fig. 5B,E), more biomass (Fig. 5H) and earlier rosette leaves development and flowering (Fig. 5F,G) compared to wild type (Col-0) and *OX-CICASPL* plants. Thus, the loss of *AtCASPL4C1* resulted in a positive growth effect, whereas over-expression gave the opposite effect. Thus,




Figure 1. Characterization of the watermelon (*Citrullus lanatus*) CICASPL and AtCASPL4C1 gene.

(A) Protein sequence alignment of CICASPL with AtCASPL4C1. (B) The four transmembrane (TM) regions of CICASPL and AtCASPL4C1 predicted by the TMHMM program. (C) Phylogenetic tree of CICASPL with AtCASPL4C1 (At3G55390) and CASP family (UPF0497, 39 members, CASP1/2/3/4/5 have been identified as being involved with Casparian strip formation) from *Arabidopsis*.

while only a single T-DNA insertion mutant is available for this gene in *Arabidopsis*, the complementary results of the knock-out and over-expressing plants strongly indicate that the phenotypes observed are due to the absence or increased abundance of CASPL.




Figure 2. Subcellular localization of CICASPL-GFP fusion protein in tobacco leaves. (A) Pattern of fluorescence for the CICASPL-GFP fusion protein. (B) Pattern of fluorescence of the plasma membrane RFP marker. (C) Merged images of (A,B) showing that these overlap. Bar = 50 μ m.




Figure 3. Transcript abundance profiles of 28 genes from the CASP protein family (UPF0497) that are represented on Affymetrix microarray chips. Transcript abundance is shown over germination and development. Expression levels (\log_2) are shown for the 28 genes (represented on microarrays) that encoding CASP or CASP-like genes. Hierarchical clustering shows strong co-expression of specific CASP like genes with known CASP genes (strongly co-expressed genes indicated by an asterisk*). The transcript abundance for *At3g55390* is shown in a black box, and genes that display a root enriched expression profile are shown in a yellow box. White boxes indicate no expression was detected.

Casparian Strip Formation Analysis. In the CASP family (UPF0497) of *Arabidopsis*, *CASP1/2/3/4/5* (Casparian strip membrane domain), were identified to mediate Casparian strip formation in the endodermis⁹. The Casparian strip is made up of a lignin polymer without suberin in *Arabidopsis*¹¹. Detection of Casparian strip formation by lignin staining in the root endodermis of *AtCASPL4C1*, wild type and *OX-CICASPL* clearly indicated lignin staining in root endodermis of *AtCASPL4C1*, WT and *OX-CICASPL* (Supplementary Figure 3A). *CASP1/2/3/4/5* transcript abundance was examined in root of wild type, *AtCASPL4C1* and *OX-CICASPL*. The transcript abundance of *CASP1* was significantly altered, with increased transcript abundance in *AtCASPL4C1* knock-outs and reduced abundance observed in *OX-CICASPL*. The transcript abundance of *CASP2*, *CASP3*, *CASP4* and *CASP5* were significantly increased transcript abundance in *AtCASPL4C1* knock-outs (Supplementary Figure 3B).

CASPL Negatively Mediated Plant Tolerance to Cold Stress. The transcript abundance of CASP protein family was examined under various kinds of stresses in *Arabidopsis* by transcriptomic analysis. These abiotic stresses result in significant up-regulation/down-regulation of the greatest number of CASP and CASP-like genes (Supplementary Figure 4).

The transcript abundance of *CICASPL* and *AtCASPL4C1* was examined under cold stress in watermelon and *Arabidopsis*. In watermelon, *CICASPL* was induced and peaked in expression 12 h after cold treatment, while in *Arabidopsis*, *AtCASPL4C1* was induced and peaked in expression 48 h after




Figure 4. Histochemical analysis of *CASPL* gene expression in *Arabidopsis*. (A) Seedling and root of 1-week-old plants. (B) Root tip. (C) Lateral root. (D) Flower. (E) Silique. Bars = 1 mm in (A,D,E).

cold treatment (Fig. 6A). Moreover, the *AtCASPLAC1* promoter-GUS activities after cold stress showed that expression of *AtCASPLAC1* was induced by cold stress in *Arabidopsis* (Fig. 6B). To study the role of *CASPL* in cold tolerance, different parameters were used to evaluate cold tolerance in wild type, *AtCASPLAC1* and *OX-CICASPL* plants. Growth was monitored for 5-day-old plants exposed to cold at 10°C for 7 days in medium. This examination showed that *AtCASPLAC1* grows better under cold stress, based on the significantly longer primary root length (Fig. 7A–C). Furthermore, 3-week-old *AtCASPLAC1* plants showed enhanced growth treated with cold stress, at 10°C for 10 days (Fig. 7D,E), keeping relatively higher chlorophyll fluorescence parameters compared to WT (Fig. 7F), as well as, more rosette leaves and greater biomass (Fig. 7G,H). These results displayed greater cold tolerance than wild type and *OX-CICASPL*.

Discussion

The Casparian strip in the root endodermis is a barrier that restricts the free diffusion of molecules between the inner cell layers of the root. Recent studies revealed that this endodermal barrier is not only




Figure 5. Phenotypes of wild type, *AtCASPL4C1* and *OX-CICASPL* under normal growth condition. (A) WT, *AtCASPL4C1* and *OX-CICASPL* growth for 5 days in MS medium. (B) Wild type, *AtCASPL4C1* and *OX-CICASPL* growth for 12 days. (C) Time course of germination for wild type, *AtCASPL4C1* and *OX-CICASPL* in medium. (D) Time course of primary root length for WT, *AtCASPL4C1* and *OX-CICASPL* in medium. (E) Wild type, *AtCASPL4C1* and *OX-CICASPL* growth for 28 days in Jiffy seedling culture substrate. (F) Wild type, *AtCASPL4C1* and *OX-CICASPL* growth for 42 days in Jiffy seedling culture substrate. (G) Time course of rosette leaves for wild type, *AtCASPL4C1* and *OX-CICASPL* in Jiffy seedling culture substrate. (H) Time course of dry weight for WT, *AtCASPL4C1* and *OX-CICASPL* in Jiffy seedling culture substrate. All growths are under growth condition of 22 °C and 16/8 h photoperiod in growth chamber. The star shows significance at 0.05 by Tukey test.

a regulator of water and nutrient uptake, but also probably acts as signal center for hormone-mediated control of growth^{24,25}. It may be considered as a signal communication hub in response to the external environment, facilitating the activation of hormone signaling pathways and the propagation of calcium waves^{24,25}. It also plays a central role in mediating cross talk between different cell layers during the




Figure 6. CASPL gene expression after cold stress. (A) Relative transcript abundance of *CICASPL* in watermelon (real line) and *AtCASPL4C1* in *Arabidopsis* (dash line) over the course of 72h of cold stress at 10°C. (B) GUS staining of CASPL-promoter after cold stress (10°C) for up to 48h in *Arabidopsis* seedlings.

development of new lateral organs, like lateral root emergence²⁶. The molecular mechanisms involved in the deposition of the Casparian strip in endodermal cells are beginning to be uncovered^{27,28}. The Casparian strip domain protein (CASP) anchors the plasma membrane to the cell wall and recruits the enzymes necessary for lignin biosynthesis^{27,28}.

Evolutionary analysis indicated plant CASP family is associated with the MARVEL protein family described only in metazoans with similarity in specific transmembrane domains, in the overall tetraspan protein structure and putative Cys bridge in the second extracellular loop¹². CASPs in this family are the earliest known proteins responsible for the CSD (Casparian strip membrane domain) formation⁹. The localizing at the CSD when expressed in the endodermis of CASPLs in this family prompts us to predict the role of yet-undiscovered CASPLs in mediating membrane subdomain formation and other plasma membrane domains in other cell types¹². Here, we identified the membrane-localizing *AtCASPL4C1* and *CICASPL* genes that encode proteins of the CASP family. Overall, the absence of *AtCASPL4C1* results in significantly increased growth under normal conditions. Based on analysis of *CASP1/2/3/4/5* transcript abundance and lignin staining in roots, it is suggested that mutation in *AtCASPL4C1* (*At3G55390*) did not dramatically affect the Casparian strip formation in roots. It has previously been shown in *Atcasp1* or *Atcasp3* single mutants, that Casparian strip formation was unaltered compared to WT, however, *casp* double mutants (*casp1/casp3*) displayed disorganized Casparian strip⁹. Thus, due to redundancy it is not possible to conclude if there is a direct role in Casparian strip formation for *AtCASPL4C1*. However, loss of function of *AtCASPL4C1* was observed to result in earlier vegetative and reproductive development and increased biomass (Fig. 5).

Plants respond to environmental changes by altering tissue differentiation and development, referred to as plant phenotypic plasticity. Here, we showed *CASPL* gene also regulates enhanced growth under cold stress conditions. While *AtCASPL4C1* belongs to this family based on predicted protein homology, the expression pattern based on *in silico* analyses and *GUS* reporter gene expression patterns suggest that the role of *AtCASPL4C1* may extend well beyond the Casparian strip in roots (Figs 3 and 4). The widespread expression pattern in vascular tissues also suggests that it plays a more fundamental role in plant growth and development, which is supported by the altered phenotype when it is inactivated or over-expressed.

In conclusion, we identified a CASP-like *CICASPL* gene in *Citrullus lanatus* and its ortholog *AtCASPL4C1* in *Arabidopsis* are associated with growth and cold tolerance. These findings offered us clues to further explore the distinguished role of the Casparian strip protein family genes in growth, development and environmental communications.




Figure 7. Phenotypes of wild type, *AtCASPL4C1* and OX-CICASPL under cold stress. (A) Wild type, *AtCASPL4C1* and OX-CICASPL seedling growth for 5 days at 22°C in MS medium. (B) 5-day-old seedlings of wild type, *AtCASPL4C1* and OX-CICASPL grown for 7 days under cold stress at 10°C in medium. (C) Time course of primary root length measurements for 5-day-old seedlings of wild type, *AtCASPL4C1* and OX-CICASPL grown for 7 days at 10°C in medium. (D) Wild type, *AtCASPL4C1* and OX-CICASPL grown for 21 days at 22°C in Jiffy seedling culture substrate. (E) 21-day-old seedlings of wild type, *AtCASPL4C1* and OX-CICASPL grown for 10 days at 10°C in Jiffy seedling culture substrate. (F) Time course of Fv/Fm fluorescence values after cold treatment at 10°C in 21-day-old seedlings of WT, *AtCASPL4C1* and OX-CICASPL plants. (G) Time course of rosette leaves for 21-day-old seedlings of WT, *AtCASPL4C1* and OX-CICASPL after cold stress at 10°C in Jiffy seedling culture substrate. (H) Time course of dry weight for 21-day-old seedlings of WT, *AtCASPL4C1* and OX-CICASPL after cold stress at 10°C in Jiffy seedling culture substrate. All growths are under growth condition of 16/8h photoperiod in growth chamber. The star shows significance at 0.05 by Tukey test. OX represent OX-CICASPL.

References

- Bohnert, H. J., Nelson, D. E. & Jensen, R. G. Adaptations to Environmental Stresses. *The Plant Cell* **7**, 1099–1111 (1995).
- Xiong, L. M., Schumaker, K. S. & Zhu, J. K. Cell signaling during cold, drought, and salt stress. *The Plant Cell* **14**, S165–S183 (2002).
- Zhu, J. H., Dong, C. H. & Zhu, J. K. Interplay between cold-responsive gene regulation, metabolism and RNA processing during plant cold acclimation. *Current Opinion in Plant Biology* **10**, 290–295 (2007).
- Mahajan, S. & Tuteja, N. Cold, salinity and drought stresses: an overview. *Archives of Biochemistry and Biophysics* **444**, 139–158 (2005).
- Schwacke, R. *et al.* ARAMEMNON, a novel database for *Arabidopsis* integral membrane proteins. *Plant Physiology* **131**, 16–26 (2003).
- Lucas, S., Dogan, E. & Budak, H. *TMP11* from wild emmer wheat: first characterisation of a stress-inducible integral membrane protein. *Gene* **483**, 22–28 (2011).
- Garwe, D., Thomson, J. A. & Mundree, S. G. Molecular characterization of *XVSAPI*, a stress-responsive gene from the resurrection plant *Xerophyta viscosa* Baker. *Journal of Experimental Botany* **54**, 191–201 (2003).
- Jang, C. S. *et al.* The molecular characterization of a cDNA encoding the putative integral membrane protein, *HvSec61* alpha, expressed during early stage of barley kernel development. *Plant Science* **168**, 233–239 (2005).
- Roppolo D. *et al.* A novel protein family mediates Casparian strip formation in the endodermis. *Nature* **473**, 380–383 (2011).
- Enstone, D. E., Peterson, C. A. & Ma, F. S. Root endodermis and exodermis: Structure, function, and responses to the environment. *Journal of Plant Growth Regulation* **21**, 335–351 (2002).
- Naseer, S. *et al.* Casparian strip diffusion barrier in *Arabidopsis* is made of a lignin polymer without suberin. *Proceedings of the National Academy of Science USA* **109**, 10101–10106 (2012).
- Roppolo, D. *et al.* Functional and evolutionary analysis of the CASPARIAN STRIP MEMBRANE DOMAIN PROTEIN family. *Plant Physiology* **165**, 1709–1722 (2014).
- Li, Y. M., Zhu, L., Xu, B. C., Yang, J. H. & Zhang, M. F. Identification of down-regulated genes modulated by an alternative oxidase pathway under cold stress conditions in watermelon plants. *Plant Molecular Biology Reporter* **30**, 214–224 (2012).
- Curtis, M. D. & Grossniklaus, U. A gateway cloning vector set for high-throughput functional analysis of genes in planta. *Plant Physiology* **133**, 462–469 (2003).
- Sparkes, I. A., Runions, J., Kearns, A. & Hawes, C. Rapid, transient expression of fluorescent fusion proteins in tobacco plants and generation of stably transformed plants. *Nature Protocol* **1**, 2019–2025 (2006).
- Nelson, B. K., Cai, X. & Nebenfuhr, A. A multicolored set of *in vivo* organelle markers for co-localization studies in *Arabidopsis* and other plants. *Plant Journal* **51**, 1126–1136 (2007).
- Narsai, R., Law, S. R., Carrie, C., Xu, L. & Whelan, J. In-depth temporal transcriptome profiling reveals a crucial developmental switch with roles for RNA processing and organelle metabolism that are essential for germination in *Arabidopsis*. *Plant Physiology* **157**, 1342–1362 (2011).
- Clough, S. J. & Bent, A. F. Floral dip: a simplified method for Agrobacterium-mediated transformation of *Arabidopsis thaliana*. *Plant Journal* **16**, 735–743 (1998).
- Weigel, D. & Glazebrook, J. *Arabidopsis*. A laboratory manual. Cold Spring Harbor, NY: Cold Spring Harbor Press (2002).
- Hong, J. K. & Hwang, B. K. The promoter of the pepper pathogen-induced membrane protein gene CaPIMP1 mediates environmental stress responses in plants. *Planta* **229**, 249–259 (2009).
- Tamura, K., Stecher, G., Peterson, D., Filipowski, A. & Kumar, S. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Molecular Biology Evolution* **30**, 2725–2729 (2013).
- Emanuelsson, O., Nielsen, H., Brunak, S. & von Heijne, G. Predicting subcellular localization of proteins based on their N-terminal amino acid sequence. *Journal of Molecular Biology* **300**, 1005–1016 (2000).
- Petersen, T. N., Brunak, S., von Heijne, G. & Nielsen, H. SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nature Methods* **8**, 785–786 (2011).
- Robbins II, N. E., Trontin, C., Duan, L., Dinneny, J. R. Beyond the barrier: communication in the root through the endodermis. *Plant Physiol.* **166**, 551–559 (2014).
- Dinneny, J. R. A gateway with a guard: how the endodermis regulated growth through hormone signaling. *Plant Science* **214**, 14019 (2014).
- Vermeer, J. E. M. *et al.* A spatial accommodation by neighboring cell is required for organ initiation in *Arabidopsis*. *Science* **343**, 178 (2014).
- Lee, Y., Rubio, M. C., Alassimone, J. & Geldner, N. A mechanism for localized lignin deposition in the endodermis. *Cell* **153**, 402–412 (2013).
- Hosmani, P. S. *et al.* Dirigent domain-containing proteins is part of the machinery required for formation of the lignin-based Casparian strip in the root. *Proceedings of the National Academy of Science USA* **110**, 14498–14503 (2013).

Acknowledgements

This work was supported by the earmarked fund for Modern Agro-Industry Technology Research System of China (CARS-26-17), key science and technology breeding program of agricultural new variety of Zhejiang Province (2012C129031-2-11) and Science and Technology Program of Zhejiang province (2011C12001). We thank Dr. J Ni to offer the RFP markers.

Author Contributions

J.Y. and M.Z. conceived and designed the study. C.D., B.X., C.C., R.N. and H.Z. performed the experiments and analyzed the data. J.Y., J.W., M.Z. wrote the paper. All authors read and approved the final manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Yang, J. *et al.* A Casparian strip domain-like gene, *CASPL*, negatively alters growth and cold tolerance. *Sci. Rep.* **5**, 14299; doi: 10.1038/srep14299 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>