

# The Calcium Ion Is a Second Messenger in the Nitrate Signaling Pathway of Arabidopsis<sup>1</sup>

Eleodoro Riveras, José M. Alvarez, Elena A. Vidal, Carolina Oses, Andrea Vega, and Rodrigo A. Gutiérrez\*

Fondo de Desarrollo de Areas Prioritarias Center for Genome Regulation, Millennium Nucleus Center for Plant Systems and Synthetic Biology, Departamento de Genética Molecular y Microbiología, Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile, Santiago, Chile 8331010 (E.R., J.M.A., E.A.V., C.O., A.V., R.A.G.); and Facultad de Agronomía e Ingeniería Forestal, Pontificia Universidad Católica de Chile, Santiago, Chile 7820436 (A.V.)

ORCID ID: 0000-0002-8208-7327 (E.A.V.).

Understanding how plants sense and respond to changes in nitrogen availability is the first step toward developing strategies for biotechnological applications, such as improvement of nitrogen use efficiency. However, components involved in nitrogen signaling pathways remain poorly characterized. Calcium is a second messenger in signal transduction pathways in plants, and it has been indirectly implicated in nitrate responses. Using aequorin reporter plants, we show that nitrate treatments transiently increase cytoplasmic  $\text{Ca}^{2+}$  concentration. We found that nitrate also induces cytoplasmic concentration of inositol 1,4,5-trisphosphate. Increases in inositol 1,4,5-trisphosphate and cytoplasmic  $\text{Ca}^{2+}$  levels in response to nitrate treatments were blocked by U73122, a pharmacological inhibitor of phospholipase C, but not by the nonfunctional phospholipase C inhibitor analog U73343. In addition, increase in cytoplasmic  $\text{Ca}^{2+}$  levels in response to nitrate treatments was abolished in mutants of the nitrate transporter NITRATE TRANSPORTER1.1/*Arabidopsis thaliana* NITRATE TRANSPORTER1 PEPTIDE TRANSPORTER FAMILY6.3. Gene expression of nitrate-responsive genes was severely affected by pretreatments with  $\text{Ca}^{2+}$  channel blockers or phospholipase C inhibitors. These results indicate that  $\text{Ca}^{2+}$  acts as a second messenger in the nitrate signaling pathway of Arabidopsis. Our results suggest a model where NRT1.1/AtNPF6.3 and a phospholipase C activity mediate the increase of  $\text{Ca}^{2+}$  in response to nitrate required for changes in expression of prototypical nitrate-responsive genes.

Plants are sessile organisms that evolved sophisticated sensing and response mechanisms to adapt to changing environmental conditions. Calcium, a ubiquitous second messenger in all eukaryotes, has been implicated in plant signaling pathways (Harper et al., 2004; Hetherington and Brownlee, 2004; Reddy and Reddy, 2004; Hepler, 2005). Multiple abiotic and biotic cues elicit specific and distinct spatiotemporal patterns of change in the

concentration of cytosolic  $\text{Ca}^{2+}$  ( $[\text{Ca}^{2+}]_{\text{cyt}}$ ) in plants (Sanders et al., 2002; Hetherington and Brownlee, 2004; Reddy and Reddy, 2004; Hepler, 2005). Abscisic acid and heat shock treatments cause a rapid intracellular  $\text{Ca}^{2+}$  increase that is preceded by a transient increase in the level of inositol 1,4,5-trisphosphate ( $\text{IP}_3$ ; Sanchez and Chua, 2001; Zheng et al., 2012).  $\text{Ca}^{2+}$  signatures are detected, decoded, and transmitted to downstream responses by a set of  $\text{Ca}^{2+}$  binding proteins that functions as  $\text{Ca}^{2+}$  sensors (White and Broadley, 2003; Dodd et al., 2010).

Nitrate is the main source of N in agriculture and a potent signal that regulates the expression of hundreds of genes (Wang et al., 2004; Vidal and Gutiérrez, 2008; Ho and Tsay, 2010). Despite progress in identifying genome-wide responses, only a handful of molecular components involved in nitrate signaling has been identified. Several pieces of evidence indicate that NITRATE TRANSPORTER1.1 (NRT1.1)/*Arabidopsis thaliana* NITRATE TRANSPORTER1 PEPTIDE TRANSPORTER FAMILY6.3 (AtNPF6.3) is a nitrate sensor in Arabidopsis (Ho et al., 2009; Gojon et al., 2011; Bouguyon et al., 2015). NRT1.1/AtNPF6.3 is required for normal expression of more than 100 genes in response to nitrate in Arabidopsis roots (Wang et al., 2009). Downstream of NRT1.1/AtNPF6.3, CALCINEURIN B-LIKE INTERACTING SER/THR-PROTEINE KINASE8 (CIPK8) is required for normal nitrate-induced expression

<sup>1</sup> This work was supported by the Howard Hughes Medical Institute, the Fondo de Desarrollo de Areas Prioritarias Center for Genome Regulation (grant no. 15090007), the Millennium Nucleus Center for Plant Systems and Synthetic Biology (grant no. NC130030), the Comisión Nacional de Investigación Científica y Tecnológica (PhD fellowship no. AT-24121649 to E.R., postdoctoral scholarship no. 3140336 to J.M.A., and PSD-74 Academy Insertion Fellowship to E.A.V.), and the Fondo Nacional de Desarrollo Científico y Tecnológico (grant nos. 11121225 to E.A.V., 11110095 to A.V., and 1141097 to R.A.G.).

\* Address correspondence to rgutierrez@bio.puc.cl.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors ([www.plantphysiol.org](http://www.plantphysiol.org)) is: Rodrigo A. Gutiérrez (rgutierrez@bio.puc.cl).

E.R. and R.A.G. designed the research; E.R., J.M.A., E.A.V., C.O., and A.V. performed the research; E.R., J.M.A., E.A.V., and R.A.G. wrote the article.

[www.plantphysiol.org/cgi/doi/10.1104/pp.15.00961](http://www.plantphysiol.org/cgi/doi/10.1104/pp.15.00961)

of primary nitrate response genes, and the CIPK23 kinase is able to control the switch from low to high affinity of NRT1.1/AtNPF6.3 (Ho et al., 2009; Hu et al., 2009; Ho and Tsay, 2010; Castaings et al., 2011). CIPKs act in concert with CALCINEURIN B-LIKE proteins, plant-specific calcium binding proteins that activate CIPKs to phosphorylate downstream targets (Albrecht et al., 2001). Early experiments using maize (*Zea mays*) and barley (*Hordeum vulgare*) detached leaves showed that nitrate induction of two nitrate primary response genes was altered by pre-treating leaves with the calcium chelator EGTA or the calcium channel blocker  $\text{LaCl}_3$  (Sakakibara et al., 1997; Sueyoshi et al., 1999), suggesting an interplay between nitrate response and calcium-related signaling pathways. However, the role of calcium as a second messenger in the nitrate signaling pathway has not been directly addressed.

We show that nitrate treatments cause a rapid increase of  $\text{IP}_3$  and  $[\text{Ca}^{2+}]_{\text{cyt}}$  levels and that blocking phospholipase C (PLC) activity inhibits both  $\text{IP}_3$  and  $[\text{Ca}^{2+}]_{\text{cyt}}$  increases after nitrate treatments. We provide evidence that NRT1.1/AtNPF6.3 is required for increasing both  $\text{IP}_3$  and  $[\text{Ca}^{2+}]_{\text{cyt}}$  in response to nitrate treatments. Altering  $[\text{Ca}^{2+}]_{\text{cyt}}$  or blocking PLC activity hinders regulation of gene expression of nitrate-responsive genes. Our results indicate that  $\text{Ca}^{2+}$  is a second messenger in the nitrate signaling pathway of Arabidopsis.

## RESULTS

### Nitrate Treatments Increase $[\text{Ca}^{2+}]_{\text{cyt}}$ Rapidly and Transiently in Arabidopsis Roots


Calcium is an essential second messenger in plant signaling processes (Bush, 1995; Trewavas and Malhó, 1998). Increase in  $[\text{Ca}^{2+}]_{\text{cyt}}$  has been recorded in cellular responses to several stimuli (Sanders et al., 1999). As a first step to determine whether calcium acts as a second messenger in the nitrate signaling pathway, we measured  $[\text{Ca}^{2+}]_{\text{cyt}}$  in roots of Arabidopsis, where the transcriptomic and phenotypic responses to nitrate have been well documented (Wang et al., 2004; Gifford et al., 2008; Gutiérrez et al., 2008; Vidal et al., 2010, 2013a,

2013b, 2014; Alvarez et al., 2014). Plants expressing cytoplasmic aequorin (WT-AQ; Gao et al., 2004) were grown hydroponically for 2 weeks with ammonium as the only N source. Plant roots were excised, and luminescence was recorded every 0.2 s after treating the roots with 5 mM  $\text{KNO}_3$  or KCl as control. As shown in Figure 1A, nitrate treatment elicited a rapid and transient increase in  $[\text{Ca}^{2+}]_{\text{cyt}}$  in roots. KCl treatment also generated a rapid and transient peak; however, this calcium peak was considerably lower than the one obtained after nitrate treatments (Fig. 1A). After reaching a maximum,  $[\text{Ca}^{2+}]_{\text{cyt}}$  decreased to near basal levels (Fig. 1A).

It is known that abiotic and biotic cues, such as sugar, salt, and drought stress, cause transient  $[\text{Ca}^{2+}]_{\text{cyt}}$  in roots and leaves (Furuichi et al., 2001; Choi et al., 2014; Michal Johnson et al., 2014). This increase in  $[\text{Ca}^{2+}]_{\text{cyt}}$  can be partially abolished by the use of  $\text{Ca}^{2+}$  channel blockers, such as lanthanum chloride (Knight et al., 1996; Choi et al., 2014). Pretreatment of WT-AQ root and seedlings with 5 mM  $\text{LaCl}_3$  for 1 h inhibited the  $[\text{Ca}^{2+}]_{\text{cyt}}$  increase observed in response to nitrate treatment (Fig. 1B). These results indicate that nitrate treatments cause a specific increase in  $[\text{Ca}^{2+}]_{\text{cyt}}$  in Arabidopsis.

### Phosphatidylinositol-PLC Activity Is Required for Changes in $[\text{Ca}^{2+}]_{\text{cyt}}$ Levels in Response to Nitrate Treatments in Arabidopsis Roots

To identify components of the signal transduction pathway mediating changes in cytoplasmic calcium levels in response to nitrate, we first determined whether a PLC-dependent pathway was implicated in this  $[\text{Ca}^{2+}]_{\text{cyt}}$  increase. We evaluated the effect of a PLC inhibitor (U73122) and a nonfunctional PLC inhibitor analog (U73343) in WT-AQ lines in response to  $\text{KNO}_3$  or KCl treatments. WT-AQ plants were pretreated for 1 h with 10  $\mu\text{M}$  U73122 or U73343, and luminescence of excised plant roots was recorded every 0.2 s after 5 mM  $\text{KNO}_3$  or KCl treatment. The presence of the PLC inhibitor (U73122) altered the  $[\text{Ca}^{2+}]_{\text{cyt}}$  increase in response to nitrate treatments (Fig. 2A). However, treatments with the nonfunctional analog



**Figure 1.** Nitrate treatments increase  $[\text{Ca}^{2+}]_{\text{cyt}}$  in Arabidopsis roots. Wild-type plants expressing WT-AQ were grown hydroponically for 2 weeks with 1 mM ammonium as the only N source. Aequorin was reconstituted by incubating plant roots in 2.5  $\mu\text{M}$  coelenterazine (CTZ) overnight in the dark.  $[\text{Ca}^{2+}]_{\text{cyt}}$  levels were monitored in excised roots in response to 5 mM  $\text{KNO}_3$  or KCl treatment (A) without pretreatment (B) or with pretreatment with the  $\text{Ca}^{2+}$  channel blocker lanthanum chloride ( $\text{LaCl}_3$ ; C). Values plotted correspond to the means of at least three independent biological replicates of five plants per treatment  $\pm$  SD.

(U73343) did not affect the  $[Ca^{2+}]_{cyt}$  increase in Arabidopsis roots (Fig. 2A). These results suggest that products of PLC enzyme activity or metabolites produced thereof trigger the  $[Ca^{2+}]_{cyt}$  increase in response to nitrate treatments. As an independent confirmation of PLC activity implicated in nitrate signaling, we measured IP<sub>3</sub> content after nitrate treatments in Arabidopsis roots. Wild-type plants were grown and treated with KNO<sub>3</sub> or KCl under the same experimental conditions described above, and roots were quickly collected and frozen in liquid nitrogen. Treatment with 5 mM KNO<sub>3</sub> resulted in a 3-fold increase in IP<sub>3</sub> levels compared with the KCl control 10 s after the treatment (Fig. 2B). Pretreatment of plants with U73122 (but not with U73343) completely blocked IP<sub>3</sub> increase in response to nitrate (Fig. 2B). In addition, LaCl<sub>3</sub> reduced IP<sub>3</sub> levels in all tested conditions, suggesting that a calcium-dependent PLC activity is implicated (Supplemental Fig. S1).

These results indicate that a PLC activity is required for IP<sub>3</sub> accumulation as well as increasing  $[Ca^{2+}]_{cyt}$  in response to nitrate treatments under our experimental conditions.

#### NRT1.1 Is a Positive Regulator of the $[Ca^{2+}]_{cyt}$ Increase in Response to Nitrate Treatments in Arabidopsis Roots

Several lines of evidence indicate that the nitrate transporter NRT1.1/AtNPF6.3 acts as a nitrate sensor in Arabidopsis plants (Ho et al., 2009; Wang et al., 2009; Gojon et al., 2011). To determine whether the increase in  $[Ca^{2+}]_{cyt}$  in response to nitrate requires a functional NRT1.1/AtNPF6.3, we generated a stable transgenic line that constitutively expresses aequorin in an *nrt1.1* mutant background. Aequorin-expressing *chlorate-resistant-1-5* (*chl1-5*) line (*chl1-5-AQ*) was generated by crossing *chl1-5* (Gao et al., 2004) with a transgenic line containing the 35S::aequorin construct (WT-AQ). We measured  $[Ca^{2+}]_{cyt}$  in *chl1-5-AQ* plant roots in response to nitrate using the same experimental strategy described


above. As shown in Figure 3A, the increase in  $[Ca^{2+}]_{cyt}$  elicited by nitrate was significantly reduced in the *chl1-5-AQ* line compared with in the wild-type plants.

We also evaluated  $[Ca^{2+}]_{cyt}$  in response to nitrate treatments in aequorin reporter lines in the *chl1-9* mutant background. *chl1-9* has a P492L point mutation that has been shown to reduce NRT1.1/AtNPF6.3 nitrate uptake without affecting the signaling function of NRT1.1 over the NRT2.1 nitrate transporter (Ho et al., 2009). It was recently shown that this point mutation causes abnormal NRT1.1/AtNPF6.3 localization (Bouguyon et al., 2015). As shown in Figure 3B,  $[Ca^{2+}]_{cyt}$  is lower in *chl1-9-AQ* roots compared with the wild type in response to nitrate treatments and comparable with the results obtained for the *chl1-5-AQ* line. These results indicate that the increase in  $[Ca^{2+}]_{cyt}$  by nitrate depends on NRT1.1/AtNPF6.3.


To evaluate whether NRT1.1/AtNPF6.3 was part of the nitrate-PLC-Ca<sup>2+</sup> pathway, we measured IP<sub>3</sub> content in *chl1-5* and *chl1-9* mutant plant roots after nitrate treatments. *chl1-5* and *chl1-9* plants were grown for 15 d and treated with 5 mM KNO<sub>3</sub> or KCl as control, and IP<sub>3</sub> content was measured. In contrast to the increase in IP<sub>3</sub> levels in wild-type roots, there was no significant increase in IP<sub>3</sub> content in *chl1-5* and *chl1-9* mutant roots after nitrate treatments (Fig. 3C). This result indicates that accumulation of IP<sub>3</sub> in Arabidopsis root in response to nitrate treatments also requires NRT1.1/AtNPF6.3 for activation of a PLC activity.

#### Nitrate-Induced Gene Expression Is Mediated by NRT1.1/NPF6.3, PLC, and Ca<sup>2+</sup>

To determine the impact of this signaling pathway on nitrate regulation of gene expression, we analyzed the expressions of nitrate-responsive genes, which have been shown to play important roles in nitrate-dependent root growth (Ho et al., 2009; Alvarez et al., 2014; Vidal et al., 2014) in wild-type, *chl1-5*, and *chl1-9*



**Figure 2.** A PLC inhibitor blocks the increase in  $[Ca^{2+}]_{cyt}$  and IP<sub>3</sub> levels in response to nitrate treatments in Arabidopsis roots. Wild-type plants expressing WT-AQ were grown hydroponically for 2 weeks with 1 mM ammonium as the only nitrogen source, and  $[Ca^{2+}]_{cyt}$  and IP<sub>3</sub> levels were assayed as described in the text.  $[Ca^{2+}]_{cyt}$  levels were monitored in excised roots pretreated with U73343 (nonfunctional analog; A) or U73122 (PLC inhibitor; B) after we treated with KNO<sub>3</sub> and KCl. C, Plants were pretreated with mock, U73122 (inhibitor of PLC), or U73343 (analogous no function), and we evaluated the IP<sub>3</sub> content in Arabidopsis roots in response to 5 mM KNO<sub>3</sub> or KCl. Values plotted correspond to the means of three independent biological replicates  $\pm$  sd. Gray bars represent time zero (before treatment), white bars represent KCl treatment, and black bars represent KNO<sub>3</sub> treatment. The letters indicate means that significantly differ between control and treatment conditions ( $P < 0.05$ ).




**Figure 3.** NRT1.1/AtNPF6.3 is required for increases in  $[Ca^{2+}]_{\text{cyt}}$  and  $IP_3$  levels in response to nitrate treatments in Arabidopsis roots. Wild-type, *chl1-5*, and *chl1-9* plants were grown hydroponically for 2 weeks with ammonium as the only nitrogen source, and  $[Ca^{2+}]_{\text{cyt}}$  and  $IP_3$  contents were assayed as described in the text. WT-AQ (A), *chl1-5*-AQ (B), and *chl1-9*-AQ (C) plants were reconstituted by incubating plant roots in 2.5  $\mu\text{M}$  CTZ overnight in the dark.  $[Ca^{2+}]_{\text{cyt}}$  levels were monitored over time. D, Wild-type (WT), *chl1-5*, and *chl1-9* plants were treated with 5 mM  $KNO_3$  and 5 mM KCl as control for 10 s, and then, we evaluated the  $IP_3$  content. Values plotted correspond to the means of at least three independent biological replicates  $\pm$  sd. Gray bars represent time zero (before treatment), white bars represent KCl treatment, and black bars represent  $KNO_3$  treatment. The letters indicate means that significantly differ between control and treatment conditions ( $P < 0.05$ ).

plants treated with the calcium channel blocker  $LaCl_3$  or the PLC inhibitor U73122. Total RNA was isolated from roots, and mRNA levels were measured for selected genes using quantitative real-time reverse transcription (qRT)-PCR. As shown in Figure 4, *NRT2.1*, *TGACG SEQUENCE-SPECIFIC BINDING PROTEIN1 (TGA1)*, and *AUXIN SIGNALING F-BOX3 (AFB3)* gene expressions are induced after  $KNO_3$  treatments. Consistent with previous reports (Ho et al., 2009; Alvarez et al., 2014; Vidal et al., 2014), nitrate regulation of gene expression of these genes was significantly altered in the *chl1-5* and *chl1-9* mutants under our experimental conditions. Similarly, nitrate inductions of *NRT2.1* and *TGA1* were significantly reduced in the presence of  $LaCl_3$  or U73122 but not mock or U73343 treatment (Fig. 4). Interestingly, induction of *AFB3* by nitrate was not significantly affected in the presence of U73122 or  $LaCl_3$  (Fig. 4). In addition, *NITRITE REDUCTASE (NIR)* and *NRT3.1* gene expressions behaved similarly to *TGA1*, with altered response to nitrate treatments in *chl1-5* or *chl1-9* mutant plants and in the presence of

U73122 or  $LaCl_3$  (Supplemental Fig. S2). This indicates that NRT1.1/AtNPF6.3, a PLC activity, and an increase in  $[Ca^{2+}]_{\text{cyt}}$  levels are required for changes in gene expression in response to nitrate treatments in Arabidopsis. Moreover, these results suggest the existence of  $Ca^{2+}$ -dependent and -independent pathways downstream of NRT1.1/AtNPF6.3 to control gene expression of nitrate-responsive genes (Fig. 5).

## DISCUSSION

Calcium is a second messenger implicated in various signaling pathways in plants (Sanders et al., 2002; Harper et al., 2004; Hetherington and Brownlee, 2004; Reddy and Reddy, 2004; Hepler, 2005; Dodd et al., 2010), and changes in  $[Ca^{2+}]_{\text{cyt}}$  are an important component of these calcium signaling networks. These changes can be induced by diverse environmental stimuli, including salt and oxidative stress, cold, light, hormones, and bacterial and fungal pathogens (Polisenky and Braam, 1996;



**Figure 4.** Regulation of gene expression in response to nitrate treatments is mediated by NRT1.1/AtNPF6.3, a PLC activity, and Ca<sup>2+</sup> in Arabidopsis roots. Col-0, *chl1-5*, and *chl1-9* plants were grown for 15 d. Plants were pretreated with 5 mM LaCl<sub>3</sub> (A) or 10 μM U73122 or U73343 (B) and then, treated with 5 mM KNO<sub>3</sub> or KCl as control. Values plotted correspond to the means of three independent biological replicates ± SD. White bars represent KCl treatment, and black bars represent KNO<sub>3</sub> treatment. The *ADAPTOR PROTEIN4 μ-ADAPTIN* gene (At4g24550) was used as a normalization reference (Aceituno et al., 2008). The letters indicate means that significantly differ between control and pharmacological treatment (*P* < 0.05). WT, Wild type.

Stoelzle et al., 2003; Chen and Kao, 2012; Choi et al., 2014; Gilroy et al., 2014). We found that nitrate is also able to trigger changes in [Ca<sup>2+</sup>]<sub>cyt</sub>. Moreover, we found that nitrate treatments increase IP<sub>3</sub> levels, which correlate with an increase in [Ca<sup>2+</sup>]<sub>cyt</sub>. In animals, IP<sub>3</sub> is generated by the cleavage of phosphatidylinositol (PI) 4,5-bisphosphate by PI-PLC enzymes (Alexandre et al., 1999; Hirose et al., 1999). This effect was abolished in *chl1-5* and *chl1-9* mutant plants, indicating that NRT1.1/AtNPF6.3 function is required for increased Ca<sup>2+</sup> and IP<sub>3</sub> in response to nitrate treatments. We found that gene expression in response to nitrate is affected by a PLC inhibitor and a Ca<sup>2+</sup> channel blocker, suggesting existence of a signaling pathway for nitrate sensing and signal transduction involving a perception event at or downstream of NRT1.1/AtNPF6.3, activation of a PLC activity, and calcium as a second messenger to regulate gene expression.

Arabidopsis has nine actively transcribed PI-PLC genes. AtPLC2 is expressed constitutively, but expressions of the remaining eight PI-PLC genes have been shown to be regulated by salt, cold and dehydration stress, abscisic acid, and other perturbations (Tasma et al., 2008). Interestingly, the expression of AtPLC4 and AtPLC5 genes is regulated by nitrate in Arabidopsis roots (Wang et al., 2003, 2004; Vidal et al., 2013a, 2013b;

Alvarez et al., 2014; Canales et al., 2014). Our results show that inhibition of PLC activity in plant roots blocks the increase in cytosolic IP<sub>3</sub> and Ca<sup>2+</sup> levels in response to nitrate treatments. In addition, LaCl<sub>3</sub> also blocked the increase in IP<sub>3</sub> and [Ca<sup>2+</sup>]<sub>cyt</sub> levels by nitrate treatments, suggesting a calcium-dependent PLC activity (Hunt et al., 2004). These results support the idea that one or more PLCs are implicated in Arabidopsis root nitrate signaling.

The mechanism by which PLC catalyzes the generation of diacylglycerol and IP<sub>3</sub> in animals is well understood (Alexandre et al., 1999; Hirose et al., 1999). However, although accumulation of IP<sub>3</sub> can be detected



**Figure 5.** A simplified model of the NRT1.1/AtNPF6.3 calcium-dependent and -independent nitrate signaling pathway. Nitrate is sensed by NRT1.1/AtNPF6.3 and activates a PLC activity that increases [Ca<sup>2+</sup>]<sub>cyt</sub>. Increase in [Ca<sup>2+</sup>]<sub>cyt</sub> activates gene expression of nitrate-responsive genes.

in plants in response to various stimuli and this increase in  $IP_3$  levels correlates with increases in cytoplasmic  $Ca^{2+}$  levels (Sanchez and Chua, 2001; Zheng et al., 2012), no homologs of animal  $IP_3$  receptors have been described in *Arabidopsis* (Nagata et al., 2004).  $IP_3$  can be further phosphorylated into inositol hexaphosphate ( $IP_6$ ; Laxalt and Munnik, 2002; Lemtiri-Chlieh et al., 2003; Meijer and Munnik, 2003; Munnik and Vermeer, 2010). Thus,  $IP_3$  levels may function directly or through its phosphorylated product  $IP_6$  in nitrate-mediated  $Ca^{2+}$  release. Similarly, DAG accumulation can lead to an increase in phosphatidic acid (PA), probably by action of a phospholipase D activity (Katagiri et al., 2001; Munnik, 2001; Sang et al., 2001). PA has been shown to act as second messenger in plant signaling pathways (Katagiri et al., 2001; Munnik, 2001; Sang et al., 2001), and previous work showed that phospholipase D $\epsilon$  and PA participate in N signaling during nitrogen deprivation in *Arabidopsis* (Hong et al., 2009). However, it is unclear whether PA has an effect over cytoplasmic calcium levels in *Arabidopsis*.

In *Arabidopsis* roots, the nitrate transporter NRT1.1/AtNPF6.3 is thought to be a nitrate sensor essential for regulation of gene expression in response to changes in external nitrate (Ho et al., 2009). Mutation of NRT1.1/AtNPF6.3 and U73122 treatments have a similar inhibitory effect over  $[Ca^{2+}]_{cyt}$  which suggests that NRT1.1/AtNPF6.3 and PLC belong to the same signal transduction pathway to control cytoplasmic calcium levels in response to nitrate. We found that normal response to nitrate of *NIR*, *NRT2.1*, *TGA1*, and *NRT3.1* depends on NRT1.1/AtNPF6.3, PLC activity, and  $Ca^{2+}$ . However, we did not observe an additional effect of U73122 or  $LaCl_3$  on nitrate regulation of gene expression in *chl1-5* or *chl1-9* mutant backgrounds. Our results indicate existence of a PLC-dependent signaling pathway downstream of NRT1.1/AtNPF6.3.

Treatment of detached maize and barley leaves with protein kinase inhibitors has been shown to alter the nitrate regulation of nitrate-responsive genes (Sakakibara et al., 1997; Sueyoshi et al., 1999). Furthermore, nitrate treatments induce changes in phosphorylation levels of proteins (Engelsberger and Schulze, 2012; Wang et al., 2012). Transcriptomics analysis of the nitrate response has shown that several protein kinases and phosphatases are regulated by nitrate availability (Canales et al., 2014), and the  $Ca^{2+}$ -dependent protein kinase CIPK8 controls the nitrate response of primary nitrate-responsive genes downstream of NRT1.1 (Hu et al., 2009). These studies are consistent with our results and suggest kinase targets of the nitrate-NRT1.1- $Ca^{2+}$  pathway described here to control gene expression.

We have previously shown that regulatory factors AFB3 and TGA1 are downstream of NRT1.1/AtNPF6.3 function in the *Arabidopsis* root nitrate response (Alvarez et al., 2014; Vidal et al., 2014). As our results indicate, TGA1 and its target *NRT2.1* would operate downstream of NRT1.1/AtNPF6.3 through a calcium-dependent signaling pathway, whereas AFB3 would operate downstream of NRT1.1/AtNPF6.3 through a calcium-independent signaling pathway. This observation is

consistent with previous results that indicate that AFB3- and TGA1-mediated responses act independently to control root system architecture in response to nitrate (Alvarez et al., 2014; Vidal et al., 2014). More recently, using transcriptomics and phenotypic analysis of NRT1.1/NPF6.3 mutants, Bouguyon et al. (2015) showed that multiple signaling pathways act downstream of NRT1.1/NPF6.3. Our results are also consistent with these observations and show that at least one signaling pathway downstream of NRT1.1/NPF6.3 depends on PLC,  $IP_3$ , and  $Ca^{2+}$ .

Our combined cell biology and molecular genetics approach allowed us to identify steps in the nitrate signaling pathway that involve  $Ca^{2+}$  as second messenger in the regulation of prototypical nitrate-responsive genes. Mapping components in the nitrate signaling pathway contributes to our understanding of how plants sense and respond to changes in N availability and provides unique targets for improving N use efficiency in crops.

## MATERIALS AND METHODS

### Plant Materials and Growth Conditions

*Arabidopsis* (*Arabidopsis thaliana*) ecotype Columbia-0 (Col-0) was used for all experiments. The *Arabidopsis* line expressing WT-AQ (Gao et al., 2004) was obtained from Christoph Plieth. The *chl1-5* and *chl1-9* mutants were donated by Yi-Fang Tsay. Plants were grown in hydroponic culture under long-day (16-h light/8-h-dark cycle) conditions at 22°C (in Percival incubators) using Murashige and Skoog salt medium without N (M531; Phytotechnology Laboratories) supplemented with 0.5 mM ammonium succinate and 0.1% (w/v) Suc. Plants were treated for the indicated periods of time at the beginning of the light cycle on day 15 with 5 mM  $KNO_3$  or KCl as a control.

### Chemical Treatment of Plants

U73122, U73343, and  $LaCl_3$  were purchased from Sigma-Aldrich. Before harvesting plant material for analysis of gene expression, Col-0 seedlings were pretreated in petri dishes for 1 h in the presence of 10  $\mu M$  U73122, 10  $\mu M$  U73343, or 5 mM  $LaCl_3$ . Plants were then treated for the indicated periods of time with 5 mM  $KNO_3$  or KCl. For aequorin measurements, plant pretreatment with all pharmacological agents was done 1 h before the addition of  $KNO_3$  or KCl to excised roots. U73122 and U73343 were dissolved in 0.1% (v/v) dimethyl sulfoxide, and  $LaCl_3$  was dissolved in water.

### In Vivo Reconstitution of Aequorin and $Ca^{2+}$ -Dependent Luminescence Measurements

Reconstitution of aequorin in vivo with CTZ was performed as described previously (Knight et al., 1996). Synthetic native CTZ was obtained from Sigma-Aldrich. Briefly, for each experiment, we incubated 14-d-old seedlings overnight in the dark with 2.5  $\mu M$  CTZ. Plant were washed with water, and roots were excised and placed in a cuvette to measure luminescence immediately after treatments. Luminescence was recorded for the duration of the experiment every 0.2 s. To convert luminescence into  $Ca^{2+}$  concentrations, 1 M  $CaCl_2$  and 10% (v/v) ethanol were added to discharge the remaining aequorin. Calculations of  $Ca^{2+}$  concentrations were performed as previously mentioned (Knight et al., 1996). Luminescence measurements were performed using a Sirius Single-Tube Luminometer (Berthold Detection Systems).

### $IP_3$ Assays

$IP_3$  was measured as described previously (Heilmann and Perera, 2013). Briefly, plants were treated with 5 mM  $KNO_3$  or KCl for 10 s, and roots were harvested and frozen immediately in liquid  $N_2$ . Frozen tissue (approximately 0.1 g) was grounded to powder and incubated with 200  $\mu L$  of 10% (v/v) perchloric

acid on ice for 20 min. Samples were centrifuged to remove the precipitate, the supernatant was transferred to a new tube, and the pH was adjusted to 7.5 using 1.5 M KOH and 60 mM HEPES. IP<sub>3</sub> was measured using the Inositol 1,4,5-Triphosphate (3H) Radioreceptor Assay Kit (Perkin Elmer) according to the instructions of the manufacturer.

## RNA Isolation and qRT-PCR

RNA was isolated from whole roots with the PureLink RNA Mini Kit (12183020; Life Technologies) according to the instructions of the manufacturer. Complementary DNA synthesis was carried out using the Improm-II Reverse Transcriptase according to the instructions of the manufacturer (Promega). qRT-PCR was carried out using the Brilliant SYBR Green QPCR Reagents on a Stratagene MX3000P qPCR System. The RNA levels were normalized relative to *ADAPTOR PROTEIN4*  $\mu$ -*ADAPTIN* (At4g24550; Aceituno et al., 2008).

## Supplemental Data

The following supplemental materials are available.

**Supplemental Figure S1.** LaCl<sub>3</sub> reduced the IP<sub>3</sub> levels in roots.

**Supplemental Figure S2.** NRT1.1/AtNPF6.3, PLC activity, and Ca<sup>2+</sup> are required for the nitrate-dependent up-regulation of NRT3.1 and NIR.

## ACKNOWLEDGMENTS

We thank Dr. Christoph Plieth for providing the Arabidopsis line expressing WT-AQ and Dr. Yi-Fang Tsay for donating *chl1-5* and *chl1-9* mutants.

Received June 23, 2015; accepted August 19, 2015; published August 24, 2015.

## LITERATURE CITED

- Aceituno FF, Moseyko N, Rhee SY, Gutiérrez RA (2008) The rules of gene expression in plants: organ identity and gene body methylation are key factors for regulation of gene expression in Arabidopsis thaliana. *BMC Genomics* 9: 438
- Albrecht V, Ritz O, Linder S, Harter K, Kudla J (2001) The NAF domain defines a novel protein-protein interaction module conserved in Ca<sup>2+</sup>-regulated kinases. *EMBO J* 20: 1051–1063
- Alexandra J, Lassalles JP, Kado RT (1999) Opening of Ca<sup>2+</sup> channels in isolated red beet root vacuole membrane by inositol 1,4,5-trisphosphate. *Nature* 343: 567–570
- Alvarez JM, Riveras E, Vidal EA, Gras DE, Contreras-López O, Tamayo KP, Aceituno F, Gómez I, Ruffel S, Lejay L, et al (2014) Systems approach identifies TGA1 and TGA4 transcription factors as important regulatory components of the nitrate response of *Arabidopsis thaliana* roots. *Plant J* 80: 1–13
- Bouguyon E, Brun F, Meynard D, Kubeš M, Pervent M, Leran S, Lacombe B, Krouk G, Guiderdoni E, Zajímalová E, et al (2015) Multiple mechanisms of nitrate sensing by Arabidopsis nitrate transceptor NRT1.1. *Nat Plants* 1: 1–8
- Bush DS (1995) Calcium regulation in plant cells and its role in signaling. *Annu Rev Plant Physiol Plant Mol Biol* 46: 95–122
- Canales J, Moyano TC, Villarreal E, Gutiérrez RA (2014) Systems analysis of transcriptome data provides new hypotheses about Arabidopsis root response to nitrate treatments. *Front Plant Sci* 5: 22
- Castaigns L, Marchive C, Meyer C, Krapp A (2011) Nitrogen signalling in Arabidopsis: how to obtain insights into a complex signalling network. *J Exp Bot* 62: 1391–1397
- Chen YH, Kao CH (2012) Calcium is involved in nitric oxide- and auxin-induced lateral root formation in rice. *Protoplasma* 249: 187–195
- Choi WG, Toyota M, Kim SH, Hilleary R, Gilroy S (2014) Salt stress-induced Ca<sup>2+</sup> waves are associated with rapid, long-distance root-to-shoot signaling in plants. *Proc Natl Acad Sci USA* 111: 6497–6502
- Dodd AN, Kudla J, Sanders D (2010) The language of calcium signaling. *Annu Rev Plant Biol* 61: 593–620
- Engelsberger WR, Schulze WX (2012) Nitrate and ammonium lead to distinct global dynamic phosphorylation patterns when resupplied to nitrogen-starved Arabidopsis seedlings. *Plant J* 69: 978–995
- Furuichi T, Mori IC, Takahashi K, Muto S (2001) Sugar-induced increase in cytosolic Ca<sup>2+</sup> in *Arabidopsis thaliana* whole plants. *Plant Cell Physiol* 42: 1149–1155
- Gao D, Knight MR, Trewavas AJ, Sattelmacher B, Plieth C (2004) Self-reporting Arabidopsis expressing pH and [Ca<sup>2+</sup>] indicators unveil ion dynamics in the cytoplasm and in the apoplast under abiotic stress. *Plant Physiol* 134: 898–908
- Gifford ML, Dean A, Gutierrez RA, Coruzzi GM, Birnbaum KD (2008) Cell-specific nitrogen responses mediate developmental plasticity. *Proc Natl Acad Sci USA* 105: 803–808
- Gilroy S, Suzuki N, Miller G, Choi WG, Toyota M, Devireddy AR, Mittler R (2014) A tidal wave of signals: calcium and ROS at the forefront of rapid systemic signaling. *Trends Plant Sci* 19: 623–630
- Gojon A, Krouk G, Perrine-Walker F, Laugier E (2011) Nitrate transceptor (s) in plants. *J Exp Bot* 62: 2299–2308
- Gutiérrez RA, Stokes TL, Thum K, Xu X, Obertello M, Katari MS, Tanurdzic M, Dean A, Nero DC, McClung CR, et al (2008) Systems approach identifies an organic nitrogen-responsive gene network that is regulated by the master clock control gene CCA1. *Proc Natl Acad Sci USA* 105: 4939–4944
- Harper JE, Breton G, Harmon A (2004) Decoding Ca<sup>2+</sup> signals through plant protein kinases. *Annu Rev Plant Biol* 55: 263–288
- Heilmann I, Perera IY (2013) Measurement of inositol (1,4,5) trisphosphate in plant tissues by a competitive receptor binding assay. *Methods Mol Biol* 1009: 33–41
- Hepler PK (2005) Calcium: a central regulator of plant growth and development. *Plant Cell* 17: 2142–2155
- Hetherington AM, Brownlee C (2004) The generation of Ca<sup>2+</sup> signals in plants. *Annu Rev Plant Biol* 55: 401–427
- Hirose K, Kadowaki S, Tanabe M, Takeshima H, Iino M (1999) Spatiotemporal dynamics of inositol 1,4,5-trisphosphate that underlies complex Ca<sup>2+</sup> mobilization patterns. *Science* 284: 1527–1530
- Ho CH, Lin SH, Hu HC, Tsay YF (2009) CHL1 functions as a nitrate sensor in plants. *Cell* 138: 1184–1194
- Ho CH, Tsay YF (2010) Nitrate, ammonium, and potassium sensing and signaling. *Curr Opin Plant Biol* 13: 604–610
- Hong Y, Devaiah SP, Bahn SC, Thamasandra BN, Li M, Welti R, Wang X (2009) Phospholipase D epsilon and phosphatidic acid enhance Arabidopsis nitrogen signaling and growth. *Plant J* 58: 376–387
- Hu HC, Wang YY, Tsay YF (2009) AtCIPK8, a CBL-interacting protein kinase, regulates the low-affinity phase of the primary nitrate response. *Plant J* 57: 264–278
- Hunt L, Otterhag L, Lee JC, Lasheen T, Hunt J, Seki M, Shinozaki K, Sommarin M, Gilmour DJ, Pical C, et al (2004) Gene-specific expression and calcium activation of *Arabidopsis thaliana* phospholipase C isoforms. *New Phytol* 162: 643–654
- Katagiri T, Takahashi S, Shinozaki K (2001) Involvement of a novel Arabidopsis phospholipase D, AtPLDdelta, in dehydration-inducible accumulation of phosphatidic acid in stress signalling. *Plant J* 26: 595–605
- Knight H, Trewavas AJ, Knight MR (1996) Cold calcium signaling in *Arabidopsis* involves two cellular pools and a change in calcium signature after acclimation. *Plant Cell* 8: 489–503
- Laxalt AM, Munnik T (2002) Phospholipid signalling in plant defence. *Curr Opin Plant Biol* 5: 332–338
- Lemtiri-Chlieh F, MacRobbie EAC, Webb AAR, Manison NF, Brownlee C, Skepper JN, Chen J, Prestwich GD, Brearley CA (2003) Inositol hexakisphosphate mobilizes an endomembrane store of calcium in guard cells. *Proc Natl Acad Sci USA* 100: 10091–10095
- Meijer HJG, Munnik T (2003) Phospholipid-based signaling in plants. *Annu Rev Plant Biol* 54: 265–306
- Michal Johnson J, Reichelt M, Vadassery J, Gershenzon J, Oelmüller R (2014) An Arabidopsis mutant impaired in intracellular calcium elevation is sensitive to biotic and abiotic stress. *BMC Plant Biol* 14: 162
- Munnik T (2001) Phosphatidic acid: an emerging plant lipid second messenger. *Trends Plant Sci* 6: 227–233
- Munnik T, Vermeer JEM (2010) Osmotic stress-induced phosphoinositide and inositol phosphate signalling in plants. *Plant Cell Environ* 33: 655–669
- Nagata T, Iizumi S, Satoh K, Ooka H, Kawai J, Carninci P, Hayashizaki Y, Otomo Y, Murakami K, Matsubara K, et al (2004) Comparative analysis of plant and animal calcium signal transduction element using plant full-length cDNA data. *Mol Biol Evol* 21: 1855–1870

- Polisensky DH, Braam J** (1996) Cold-shock regulation of the Arabidopsis *TCH* genes and the effects of modulating intracellular calcium levels. *Plant Physiol* **111**: 1271–1279
- Reddy VS, Reddy ASN** (2004) Proteomics of calcium-signaling components in plants. *Phytochemistry* **65**: 1745–1776
- Sakakibara H, Kobayashi K, Deji A, Sugiyama T** (1997) Partial characterization of the signaling pathway for the nitrate-dependent expression of genes for nitrogen-assimilatory enzymes using detached maize leaves. *Plant Cell Physiol* **38**: 837–843
- Sanchez JP, Chua NH** (2001) Arabidopsis PLC1 is required for secondary responses to abscisic acid signals. *Plant Cell* **13**: 1143–1154
- Sanders D, Brownlee C, Harper JF** (1999) Communicating with calcium. *Plant Cell* **11**: 691–706
- Sanders D, Pelloux J, Brownlee C, Harper JF** (2002) Calcium at the crossroads of signaling. *Plant Cell (Suppl)* **14**: S401–S417
- Sang Y, Cui D, Wang X** (2001) Phospholipase D and phosphatidic acid-mediated generation of superoxide in Arabidopsis. *Plant Physiol* **126**: 1449–1458
- Stoelzle S, Kagawa T, Wada M, Hedrich R, Dietrich P** (2003) Blue light activates calcium-permeable channels in Arabidopsis mesophyll cells via the phototropin signaling pathway. *Proc Natl Acad Sci USA* **100**: 1456–1461
- Sueyoshi K, Mitsuyama T, Sugimoto T, Kleinhofs A, Warner RL, Oji Y** (1999) Effects of inhibitors for signaling components on the expression of the genes for nitrate reductase and nitrite reductase in excised barley leaves. *Soil Sci Plant Nutr* **45**: 1015–1019
- Tasma IM, Brendel V, Whitham SA, Bhattacharyya MK** (2008) Expression and evolution of the phosphoinositide-specific phospholipase C gene family in *Arabidopsis thaliana*. *Plant Physiol Biochem* **46**: 627–637
- Trewavas AJ, Malhó R** (1998) Ca<sup>2+</sup> signalling in plant cells: the big network! *Curr Opin Plant Biol* **1**: 428–433
- Vidal EA, Álvarez JM, Gutiérrez RA** (2014) Nitrate regulation of AFB3 and NAC4 gene expression in Arabidopsis roots depends on NRT1.1 nitrate transport function. *Plant Signal Behav* **9**: e28501
- Vidal EA, Araus V, Lu C, Parry G, Green PJ, Coruzzi GM, Gutiérrez RA** (2010) Nitrate-responsive miR393/AFB3 regulatory module controls root system architecture in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* **107**: 4477–4482
- Vidal EA, Gutiérrez RA** (2008) A systems view of nitrogen nutrient and metabolite responses in Arabidopsis. *Curr Opin Plant Biol* **11**: 521–529
- Vidal EA, Moyano TC, Krouk G, Katari MS, Tanurdzic M, McCombie WR, Coruzzi GM, Gutiérrez RA** (2013a) Integrated RNA-seq and sRNA-seq analysis identifies novel nitrate-responsive genes in *Arabidopsis thaliana* roots. *BMC Genomics* **14**: 701
- Vidal EA, Moyano TC, Riveras E, Contreras-López O, Gutiérrez RA** (2013b) Systems approaches map regulatory networks downstream of the auxin receptor AFB3 in the nitrate response of Arabidopsis thaliana roots. *Proc Natl Acad Sci USA* **110**: 12840–12845
- Wang R, Okamoto M, Xing X, Crawford NM** (2003) Microarray analysis of the nitrate response in Arabidopsis roots and shoots reveals over 1,000 rapidly responding genes and new linkages to glucose, trehalose-6-phosphate, iron, and sulfate metabolism. *Plant Physiol* **132**: 556–567
- Wang R, Tischner R, Gutiérrez RA, Hoffman M, Xing X, Chen M, Coruzzi G, Crawford NM** (2004) Genomic analysis of the nitrate response using a nitrate reductase-null mutant of Arabidopsis. *Plant Physiol* **136**: 2512–2522
- Wang R, Xing X, Wang Y, Tran A, Crawford NM** (2009) A genetic screen for nitrate regulatory mutants captures the nitrate transporter gene *NRT1.1*. *Plant Physiol* **151**: 472–478
- Wang X, Bian Y, Cheng K, Zou H, Sun SSM, He JX** (2012) A comprehensive differential proteomic study of nitrate deprivation in Arabidopsis reveals complex regulatory networks of plant nitrogen responses. *J Proteome Res* **11**: 2301–2315
- White PJ, Broadley MR** (2003) Calcium in plants. *Ann Bot (Lond)* **92**: 487–511
- Zheng SZ, Liu YL, Li B, Shang ZL, Zhou RG, Sun DY** (2012) Phosphoinositide-specific phospholipase C9 is involved in the thermo-tolerance of Arabidopsis. *Plant J* **69**: 689–700