

Review Article

Nutritional Considerations for Performance in Young Athletes

JohnEric W. Smith, Megan E. Holmes, and Matthew J. McAllister

Department of Kinesiology, Mississippi State University, P.O. Box 6186, Mississippi State, MS 39762, USA

Correspondence should be addressed to JohnEric W. Smith; johneric.smith@msstate.edu

Received 31 May 2015; Accepted 2 August 2015

Academic Editor: Adrian W. Midgley

Copyright © 2015 JohnEric W. Smith et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Nutrition is an integral component to any athletes training and performance program. In adults the balance between energy intake and energy demands is crucial in training, recovery, and performance. In young athletes the demands for training and performance remain but should be a secondary focus behind the demands associated with maintaining the proper growth and maturation. Research interventions imposing significant physiological loads and diet manipulation are limited in youth due to the ethical considerations related to potential negative impacts on the growth and maturation processes associated with younger individuals. This necessary limitation results in practitioners providing nutritional guidance to young athletes to rely on exercise nutrition recommendations intended for adults. While many of the recommendations can appropriately be repurposed for the younger athlete attention needs to be taken towards the differences in metabolic needs and physiological differences.

1. Introduction

Current estimates suggest approximately 35 million youth between the ages of 5–18 years participate in organized sports each year [1]. While a majority of these young athletes are playing sports for the aspects of comradery and fun, a growing segment of young athletes train to enhance their opportunity to make a career of sport. While elite sport has long seen the presence of young athletes (Nadia Comaneci, 14 years of age (1976 Olympic Gold Medalist), Marjorie Gestring, 13 years of age (1936 Olympic Gold Medalist), and Dimitrios Loundras, 10 years of age (1896 Olympic Bronze Medalist)), the past few decades have experienced an expansion in the numbers of young athletes working to perform at higher levels as younger athletes. This expansion can be seen in the establishment of the many facilities open focusing specialized training for sports performance on not only elite athletes but also largely youth athletes.

While the increase in physical activity of youth is important we currently do not fully understand the effects such training has on the growth and development of youth. The American Academy of Pediatrics outlined potential risks associated with sports specialization in young athletes in a publication in 2000 [2]. Noted orthopedic surgeon, Dr. James Andrews, recently discussed the potential for negative

effects of specialized training on developing bodies and the rise in youth sport injuries he experienced since around the same 2000 timeframe [3]. It is not within the scope of this review to discuss the ethical considerations of having youth focus their training on a singular sport, nor to discuss the potential for injury as related to overuse injuries. However, with the continual trend in younger athletes training for high level performance it appears that our current options are to continue to underscore the potential risks while at the same time work with the participants providing as much assistance as possible to enhance safety.

Proper nutrition is a fundamental component of athletes' training and performance plan. Proper nutrition ensures that an individual is amassing the fuels necessary for the energy production needs related to activity and recovery. One of the areas needing to be addressed is the unique nutritional needs associated with intense exercise stress. However, our understanding of the effects of strenuous physiological training and nutritional variations in combination with exercise stress in youth athletes is greatly limited. This limited knowledge is most likely due to the ethical considerations of withholding nutrients and physiologically overstressing a vulnerable population such as children and adolescents still in the process of growth and development.

Our knowledge regarding the nutritional needs of youth is based on the needs related to proper growth and development in healthy children or those suffering from illness. Most of the knowledge we possess related to the physiological adaptations to training, exercise performance, and sports nutrition is based on research conducted in college aged, middle aged, and older adult populations. Therefore, most sports nutrition recommendations promoted in youth sport are actually based on findings in adult populations. While this is a starting point, research has shown that adolescent energy expenditure and metabolism can differ from those of their adult counterparts so many of these recommendations may not provide ideal insight into the nutritional needs of the youth athlete [4–6].

The goal of this review is to compile an overview of our understanding of the nutritional needs of the young athlete during training and competition. We will also identify the knowledge gaps that currently exist in our understanding of this vulnerable population's needs around physiologically stressing occasions. Due to the limited research on the young athlete population, in many instances the knowledge gained through research on adult populations is the only means to provide recommendations for the young athlete.

Nutrition for healthy growth and maturation is governed by a variety of parameters, each essential in the development from child to adult. This paper emphasizes the importance of adolescent nutrition by first examining gross total caloric intake to better understand the energy requirements of adolescents. Total caloric intake must be sufficient to meet the additional demands of growth, which vary at different stages of growth and maturation and between individual children. Likewise, the proportion of calories allocated to each macronutrient is heavily dependent on the situational constraints of the individual child, which is further complicated by the physiological constraints of a given level of development. This paper emphasizes the importance of each macronutrient with specific focus on the physiological nuances of adolescent metabolism specifically focused around the young athlete. Similarly, micronutrient needs are driven by demands of growth and maturation as well as activity levels. Unique demands of the growing adolescent have highlighted a few select micronutrients in the literature which will be reviewed here following a general overview of micronutrient needs.

2. Growth and Development

Growth, maturation, and development are three constructs paramount in any discussion regarding youth. While these terms often manifest concurrently in youth, they refer to three different parameters. Growth simply refers to the quantifiable increase in size, whereas maturation refers to timing and tempo of progress toward the mature state. Timing and tempo refer to the age at which specific maturational events occur and rate at which an individual progresses through these events. Both timing and tempo vary considerably between children [8]. Development is considered a social construct that typically focuses on behaviors and attitudes.

Behaviors and attitudes developed during childhood and adolescence provide the basis for adult behaviors and attitudes. Refinement of accepted behaviors in a society requires the development of competencies in an array of interrelated domains that ultimately shape a given behavior and attitude toward that behavior. Taken together, growth, maturation, and development synergistically influence an individual's general self-concept and self-esteem [8]. This holistic perspective is often overlooked when focused on specific pediatric topics, such as nutrition.

Much like business, calorie supply (i.e., energy intake) is dictated by demand (i.e., energy expenditure). Energy expenditure is represented by four major components in children and adolescents: basal/resting metabolic rate, thermic effect of food, thermic effect of activity, and the energy requirements of growth [8]. Basal and resting metabolic rates (BMR and RMR, resp.) vary chiefly on assessment methodology, but only marginally in amount of calories. For the purposes of this discussion, the term RMR will be used to represent both. In adults RMR increases proportionally with body mass, particularly lean body mass [9]. Similarly, RMR increases as children gain body mass. However, when RMR is examined per unit of body mass, RMR decreases as children progress to their adult size [10], which demonstrates the contribution of growth to RMR in children and adolescents. The thermic effect of food varies significantly by the proportion of macronutrients comprising the food consumed. On average, 6–8% of ingested calories are used in the digestive, absorptive, and storage processes of food. Thermic effect of activity is the most variable component of energy expenditure and refers to the calorie cost of movement. When estimating caloric requirements, activity levels are examined at three levels: light, moderate, and vigorous lifestyle physical activity. Given the significant participation in high energy demanding activities, vigorous lifestyle physical activity is exemplified in the youth athlete population. The energy cost of growth is examined in two parameters, the energy to synthesize tissue and the energy deposited in those tissues [7]. Growth varies according to the tempo of maturational development and is very rapid during infancy and early childhood and, thus, accounts for a greater proportion of caloric expenditure. Conversely, during late childhood and adolescence, growth accounts for 1–2%, which reflects a slower rate of growth [8]. With consideration to each of these four components, the FAO/WHO/UNU expert panel used typical weight gains per year to develop age specific and gender specific caloric recommendations [7]. Table 1 shows the caloric recommendations for boys and girls participating in vigorous lifestyles physical activity. Daily energy requirements increase with age and are similar between boys and girls until pubertal ages.

3. Protein

Protein is needed for normal cellular functioning as well as synthesis of various bodily tissues [11]. Athletes tend to have elevated demands for dietary protein intake compared to sedentary individuals [12]. As a general recommendation for maintaining health, current recommendations are between

TABLE 1: Age-specific energy requirements for boys and girls who participate in heavy physical activity levels.

Age (years)	Boys (kcal/day)	Girls (kcal/day)
6-7	1,800	1,650
7-8	1,950	1,775
8-9	2,100	1,950
9-10	2,275	2,125
10-11	2,475	2,300
11-12	2,700	2,475
12-13	2,925	2,625
13-14	3,175	2,725
14-15	3,450	2,855
15-16	3,650	2,875
16-17	3,825	2,875
17-18	3,925	2,875

Adapted from FAO/WHO/UNU, 2004 [7].

0.8 and 1.2 grams of protein per kg of body mass daily [13]. This recommendation is sufficient to meet the bodily demands of 97.5% of the population, which also accounts for variations in demographic BMI as well as gender [11]. A review by Nemet and Eliakim speculates these requirements are likely sufficient for children and youth athletes. The American College of Sports Medicine and American Dietetic Association recommend intakes between 1.2 and 1.8 g/kg of body mass for active adults [14, 15], which appears to be an adequate requirement for youth athletes [16, 17]. Protein synthesis is highest during infancy and, as such, during this time relative dietary protein intake is at an elevated demand [11]. The question of how much dietary protein is needed to maximize performance among athletes is a question that has been debated for more than 150 years [11, 18] and still remains a debate. Recent evidence suggests two to three times the RDA for protein intake may be optimal to enhance fat-free mass during periods of caloric restriction [19] which may be commonly practiced among athletic populations to achieve a body composition more favorable for performance. Investigation of dietary intakes for various youth age groups suggests that intakes this high are often achieved in normal dietary patterns [20], which indicates intake is sufficient to meet the elevated demands.

Many athletes make dietary modifications in attempt to maximize performance and meet body weight requirements for competitive classes [21]. Several studies have shown increased dietary protein intake accompanied by exercise intervention may aid in weight loss as well as preservation of lean body mass typically associated with reduced body weight [22–25]. Some suggest the mechanism may be partially attributed to increased thermogenesis and satiety associated with elevated protein intake. When compared to fat and carbohydrate, protein has a greater thermic effect that is likely only significant enough to result in weight loss when the high protein diet is maintained over the course of several months [26]. Additional research is needed to fully investigate this hypothesis. Several studies have demonstrated

satiating effects of high protein diets [27–29], which may better elucidate a mechanism of weight loss with this dietary intervention. Branched chain amino acids found in protein-rich foods are known to assist in preservation of lean body mass [30], which has significant implications for athletes particularly during periods of weight loss. Leucine specifically is one branched chain amino acid that is strongly associated with protein synthesis [31]. This amino acid can be ingested in supplement form; however, when determining the safe tolerable upper intake level for leucine intake, trials in youth are limited. One study suggests the upper level for young males aged 20–35 years is 500 mg/kg/day or 35 g/day based on plasma and urinary ammonia and leucine concentrations [32]. This recommendation has not been examined in youth and caution of leucine at these high levels is warranted. However, food sources such as egg whites and dairy sources contain multiple amino acids and, as such, protein-rich foods should be emphasized to a greater extent than single amino acids alone.

The most significant question is whether or not youth athletes are obtaining the amount of protein needed for their elevated demands. It has been documented that youth athletes in general are achieving protein intakes much greater than the RDA [20]. Given this evidence, it is unlikely beneficial to promote increased protein intake in youth. A thorough dietary evaluation is suggested before promoting increased protein intake in youth athletes. However, as a general recommendation, athletes should ingest balanced protein feedings throughout the day [19] and emphasize whole foods as opposed to protein based supplements due to the lack of scientific support for protein based supplements in comparison to protein-rich whole foods [33].

As previously mentioned, athletes require higher protein intake to maintain protein synthesis [33]. Additionally, research has shown the ingestion of 20 g of protein following exercise helps maintain positive protein balance following exercise [34]. Evidence suggests that protein based supplements are not required to meet this increased demand [2]. Nonetheless, protein supplements remain one of the most common dietary supplements [35] purchased by athletes who seek to increase markers of performance such as speed, strength, power, and hypertrophy [36]. Several reports have documented athletes' perception that protein supplements are necessary to build muscle [37, 38] and achieve peak performance [39]. This notion has been well investigated in adults but also appears true in the limited research regarding youth athletes, specifically high school football players [33]. This misconception among youth is partially driven by the lack of formal knowledge in nutrition [40]. Youth athletes gain a significant amount of nutrition information from magazines, family members, and coaches [41] and, thus, may not be able to make appropriate, evidence based decisions regarding the use of protein supplements [33]. Considering the lack of scientific support for protein based supplements being superior to natural protein containing foods, youth athletes should be advised to consume their protein from whole foods as opposed to supplements.

4. Fat

Dietary lipids are essential for the absorption of vitamins A, D, E, and K, as well as synthesis of cholesterol and other sex hormones [42]. In terms of caloric requirements, most sources recommend lipid intake should be limited to 25–30% of total caloric intake [43], which is relatively the same for both sedentary and active individuals. It is important to consider caloric demands are increased in athletic populations; therefore, *absolute* lipid intakes are likely to be higher. The average adolescent consumes roughly one-third of their dietary intake as lipids [44]. It is important to restrict lipid intake to avoid excessive caloric intake; however, there is no health benefit in diets with less than 15% of calories from lipids [2]. In terms of athletic requirements, an increase of dietary carbohydrate should account for a majority of the increased caloric demands, rather than an increase of dietary lipid. Adequate calorie consumption to support periods of rapid growth is of greatest concern when considering nutrition to maximize performance of adolescent athletes [45]. Roughly fifty percent of adult body mass as well as skeletal mass is achieved during puberty. Large increases in lean and adipose tissue are also seen in males and females during the transition from child to adult as well [44]. During this time, dietary fat is especially important to aid in the synthesis of hormones and assist in normal bodily functioning as well as healthy growth and maturation [46]. Dietary lipid intakes beyond 30% are not advised since this could contribute to excessive weight gain [47]. However, acutely, excessive lipid intake can also result in postprandial oxidative stress, which is associated with impaired vascular and metabolic functioning [48, 49]. Elevated lipid intake is also potentially associated with the pathogenesis of cardiovascular disease [42] which is particularly relevant for youth athletes, given that the origins of CVD begin at an early age and progress into adulthood [48, 50, 51]. The organized group setting provides an ideal platform for discussion of nutrition and physical activity habits among individuals who already acknowledge their value.

Adolescents are more efficient in terms of substrate utilization, which has been shown both at rest and during graded exercise tests since younger children derive a higher percentage of energy from lipids as indicated by lower RER values at submaximal intensities [52]. Improved aerobic efficiency is related to increased dependency on lipids for ATP production commonly noted in youth [5]. This could potentially be the result of an adaptive response since infants and toddlers (under the age of 2) require a higher percentage of energy from lipids to support their increased caloric and growth demands [53]. Alterations in dietary lipid intake could contribute to changes of enzymatic activity as well as elevated lipid metabolism [54]. A lack of glycolytic enzyme activity could be another reason for the aforementioned increased dependency on lipid metabolism [4]. During exercise, carbohydrates and lipids are the main sources of skeletal muscle ATP production, with lipids serving as an important source of energy during low and moderate intensity [45]. Chronic exercise training results in favorable mitochondrial adaptations in adults, which favor enhanced lipid metabolism as well [55].

Upon investigation of differences in lipid oxidation among different age and gender groups in children, a review by Aucouturier et al. [4] reported only miniscule differences between age groups among male and female adolescents. These miniscule differences are likely associated with a change in body size (e.g., acquisition of skeletal and muscle mass) during periods of growth and maturation among different age groups and are more significant in males compared to females. However, Aucouturier et al. reported all children (in general) depend more readily on lipids in comparison to adults. This metabolic characteristic could depend on pubertal status, since it has been shown that 12-year-old females demonstrate elevated lipid metabolism during exercise performed at 70% $\dot{V}O_{2\max}$ compared to 14-year-old females [56]. Similar findings have also been reported in boys aged 12 and 14 [57]. However, to our knowledge there is little to no evidence showing prepubescent males and females differ *significantly* in terms of relative fat and carbohydrate oxidation during submaximal exercise [4]. In comparison to adults, however, children lack the ability to sustain longer duration exercise, which may be related to a lack of the ability to store glycogen in children [58]. Generalizing substrate utilization during *prolonged* exercise is difficult given the paucity of experimental or quasi-experimental evidence which examines exercise testing in children for greater than one hour in duration [58], which is likely reflective of the general short and intermittent physical activity patterns and behaviors of that age group [59].

The composition of dietary fatty acids (e.g., chain length) can affect fat oxidation during submaximal exercise [54]. However, this response may vary among maturational levels of the young athlete since prepubescent males tend to have a higher percentage of fatty acid oxidation [60]. It is also important to consider the potential adaptations resulting from modifications of lipid intake. Short term elevations in lipid intake are likely to result in positive energy balance which may not be immediately matched with an increase in beta-oxidation [54]. However, trials in adult populations show that exercise can enhance lipid metabolism by stimulating mitochondrial biogenesis [61] as well as increasing activity of lipoprotein lipase [62] and carnitine palmitoyltransferase 1 [63]. The aforementioned adaptations can enhance lipid metabolism and contribute to an accommodated energy balance due to changes in fat metabolism [54]. This evidence is especially applicable to athletes since exercise training has been shown to accommodate the metabolic effects of short term high fat diets [64]. These adaptations, however, still do not constitute promotion of lipid intakes greater than 30% in youth athletes.

As mentioned earlier, factors such as the type of lipid ingested (i.e., composition of the hydrocarbon chain) can affect the subsequent metabolism [54] and potential storage [65, 66]. Piers et al. [67] demonstrated the substitution of saturated fat with monounsaturated fatty acids can potentially have a favorable effect on body composition. The contribution of saturated fatty acid intake to the development of CVD is important [42]; however, a meta-analysis by Siri-Tarino et al. [68] included twenty-one studies and reported no significant risk of coronary heart disease with elevated

saturated fatty acid intake. It is important to note coronary heart disease is a *chronic, progressive* disease, the origins of which begin early in life. Thus, youth require appropriate dietary advice that may potentially become healthy behaviors in adulthood as a means of preventing or reducing disease risk. Further, these findings [68] may not be applicable to all individuals since intensity of activity and lipid composition both affect lipid metabolism [54]. Considering the evidence to date, dietary unsaturated fatty acids should not serve as exclusively the sole source of lipids. However, as a general recommendation, and in the interest of promoting long-term healthy behaviors, unsaturated fatty acids should be emphasized to a greater extent.

Some athletes may believe certain lipid based supplements may allow for an ergogenic effect given the limited findings suggesting supplementation may enhance lipid metabolism by decreasing dependency on glycogen/glucose for energy metabolism [69]. Commercial marketing is based on the premise some lipid based dietary supplements can thereby increase aerobic capacity and performance, improve lipid metabolism, and reduce inflammatory damage [70]. Fish oil and conjugated linoleic acid (CLA) are two lipid based supplements that have been investigated in relation to potential ergogenic effects.

Fish oil contains two essential fatty acids (eicosapentaenoic acid [EPA] and docosahexaenoic acid [DHA]). Increased dietary intake of these essential fatty acids has been associated with decreased prevalence of cardiovascular diseases [71, 72], as well as reduced markers of inflammation [73, 74]. In regard to physical performance, a variety of trials failed to demonstrate an ergogenic effect of EPA and DHA [75–77]; Tartibian et al. [78] reported improved pulmonary functioning in young wrestlers [78]. In terms of the ability to improve athletic performance, the majority of data demonstrate a lack in ergogenic effect of fish oil ingestion on athletic performance in well trained athletes [76, 79, 80]. The overwhelming scientific support for fish oil supplementation highlights improvements in cardiovascular health and decreases markers of inflammation which *could* contribute to decreased recovery time between exercises as speculated by Macaluso et al. [69]. However, this has not yet been supported in literature. Furthermore, the majority of clinical trials utilize extremely high doses (>3 g/day) [69, 77, 81], which is difficult to achieve without dietary supplementation of fish oil.

Other benefits of fish oil consumption have been noted including improved cognitive function and reduced ADHD symptoms in children [82]. A meta-analysis by Yang et al. [83] indicated dietary fish and/or fish oil supplementation is also associated with a reduced prevalence of asthma in children [83]. However, additional trials in youth (specifically with athletic samples) are warranted as this population has not been investigated to our knowledge. Given the gap in the literature examining youth athletes, clear recommendations for fish oil consumption cannot be made and warrant further investigation.

CLA is another lipid based dietary supplement that has been proposed to improve athletic performance [69]. CLA is found naturally occurring in beef, lamb, and dairy products

such as milk and cheese [84] but is also available in supplement form. Animal studies utilizing CLA administration have demonstrated potentially favorable effects on body composition [85]. However, this may not be applicable to humans since Zambell et al. [86] failed to show a change in energy expenditure and lipid metabolism. A recent review by Macaluso et al. [69] reported fish oil and CLA supplementation can *potentially* have a favorable effect on anabolic effects of exercise which could be related to increased testosterone synthesis. However, given the strong relationship between growth, maturation, and anabolic hormone levels in youth, the demands for youth athletes to intentionally manipulate hormone levels are not advised.

5. Carbohydrate

Human metabolism relies primarily on the oxidation of fats and carbohydrates as its fuel sources. As physiological intensity increases from rest to vigorous there is a transition from fat functioning as the primary fuel source to carbohydrate supplying a majority of the body's fuel for energy. The sources of carbohydrate for metabolism are glycogen stores in the muscle, glycogen stores in the liver, and exogenous carbohydrate entering the blood stream through the ingestion of carbohydrate. Some confusion exists in athletes understanding of the specific carbohydrate needs and it has been suggested by Burke et al. to stem from the fact that many recommendations are based on percentage of total caloric intake which adds to the difficulty in understanding the dietary needs of carbohydrate in athletes that have caloric intakes which often exceed general recommendations [87].

General carbohydrate intake recommendations suggest adult athletes consume 5–12 grams of carbohydrate per kilogram per day dependent on their primary form of exercise/activity, activity intensity, sex, and environmental conditions [88]. The great variance that exists in the demands of sports, training, and level of play make it difficult to provide a single concise recommendation. As training duration and intensities increase carbohydrate requirements rise. Young athletes lack even the large range recommendations that are provided for adult athletes. The recommendations for young athletes suggest at least 50% of young athletes diet should be in the form of carbohydrate [89] or between 3 and 8 grams [90] of carbohydrate per kilogram of body mass dependent primarily on exercise intensity.

The role of carbohydrate ingestion around active occasions is an area of intense study. Early research into carbohydrate's role in exercise performance examined the effect of blood glucose levels and physiological state following prolonged exercise [91, 92]. Additional research investigated muscle glycogen's role in muscle fatigue [93, 94]. Subsequent research investigated the role of carbohydrate ingestion following exercise to restore muscle glycogen stores following the depletion related to exercise stress [95]. More recently the research focus has shifted to explore the role of carbohydrate ingestion during exercise stress in sustaining exercise intensity and improve performance [96–98].

It is commonly suggested that normal body stores of carbohydrate can be a significant fuel source for approximately 90–120 minutes of moderate to vigorous exercise. While this statement is accurate in many exercise settings where individuals are exercising at moderate-vigorous intensities typically associated with endurance exercise, more detailed investigation would demonstrate exercise intensity is a vital component in understanding glycogen depletion rates. Significant glycogen depletion can occur anywhere from ~10 minutes with supramaximal intensities to greater than 3 hours at low exercise intensities [99]. This oversimplification needs to be considered as we look at the findings from carbohydrate research. Many of the research investigations providing our understanding of carbohydrate needs around exercise are based on endurance type exercise. As we consider the sports typically involving young athletes some of our understanding may not fully translate.

As stated previously part of our limited understanding in the nutritional needs of young athletes is the result of proper research ethics. The research described above with adults involved muscle biopsies, exercise to failure, and exercise resulting in “poor” physiological states. This type of request would be inappropriate to make to children. Therefore most of our understanding of youth athletes comes from the utilization of less invasive techniques. Young athletes have been shown to have a lower respiratory exchange ratio (RER) during exercise at similar relative submaximal intensities ($\% \dot{V}O_{2 \max}$) as their adult counterparts [4]. Based on the RER changes resulting from the shift from fat as a primary fuel source to increasing carbohydrate utilization, this would suggest young athletes are better able to utilize fat as a fuel or are potentially limited in their maximal performance as a result of not being able to utilize carbohydrate readily enough at higher intensities.

Research has shown that increasing glycogen stores will enhance exercise performance [100] and reductions in muscle glycogen content correspond with increasing levels of fatigue. Unfortunately, young athletes have been shown to store less glycogen than adults [4]. During prolonged exercise and exercise at elevated intensities reduced glycogen levels will lead to early onsets of fatigue. Due to their lower glycogen stores, young athletes will likely experience accelerated rates of fatigue. This accelerated fatigue is a result of the inability of the body to maintain sufficient blood glucose levels to meet the young athletes elevated glucose needs of the brain as compared to adults [4].

With reduced muscle glycogen stores the need for exogenous sources of carbohydrate becomes increasingly more important in the maintenance of exercise intensity. Much of our understanding regarding the role of carbohydrate ingestion in exercising youth is the result of the work of the researchers in the Children's Exercise and Nutrition Centre at McMaster University. Riddell demonstrated adolescent athletes utilized lower absolute amounts of exogenous glucose as compared to values reported in adults [101]. However, exogenous carbohydrate utilization has been demonstrated as a greater relative contributor to total energy utilization in the young athlete even with lower absolute utilization rates [58, 101].

Current research foci have shifted more towards the improvement of health and wellness as opposed to performance. However recent research continues to demonstrate the ergogenic effects of carbohydrate ingestion on youth sport. Dougherty et al. demonstrated performance improvements with basketball skills test with carbohydrate compared to water ingestion [102]. Batatinha et al. demonstrated gymnasts experienced a reduction in the number of falls from a balance beam with carbohydrate ingestion during a session [103]. Smith et al. demonstrated performance was improved with carbohydrate ingestion during football skill performance [104].

While the need for carbohydrate is recognized as important, the differences between youth and adult are still not fully understood. Research has shown carbohydrate ingestion spares endogenous carbohydrate stores in exercising youth while at the same time youth seem to be unable to utilize carbohydrate at rates similar to those seen in adults [101]. Research in adults has led to the establishment of guidelines suggesting carbohydrate ingestion during activities lasting 45 minutes or longer provides an ergogenic effect with doses varying from “small amounts including mouth rinse” up to 90 g/h [88]. Currently, specific recommendations similar to what exists for adults are not available for the youth athlete.

For performance enhancement young athletes will benefit from the ingestion of carbohydrate during exercise. Without specific rates recommended for the young athletes, we must rely on the recommendations of adults and refine carbohydrate intakes during exercise based on trial and error methods. These recommendations suggest athletes should ingest simple sugars at a rate of 30–60 g/h for exercise lasting longer than 60 minutes. Additionally, athletes should ingest 1–1.5 g/kg of body mass in the 30 minutes following cessation of prolonged exercise [15]. The ingestion of carbohydrate during exercise should be considered an equally significant component of the training plan as the skill aspects of sport.

6. Water

As with their adult counterparts hydration status during sport is important to performance in young athletes. It could be argued that, due to their increased susceptibility to succumb to heat stressors, due to their greater body surface areas to body mass ratio, hydration is a more important consideration in young athletes [105]. In addition to an increased body surface area to body mass ratio, adolescents have been shown to have diminished sweat rates as compared to their adult counterparts [106]. Diminished sweat rates are advantageous as a result of their protection of body water status but are disadvantageous due to the reduced ability to dissipate heat. Added importance of hydration is due to the fact that, in addition to performance decrements, hypohydration has been shown to lead to increased physiological strain, increased risk of heat injury/illness, and increased perceived exertion at similar workloads [15, 107].

As a result of hypohydration, the body experience fluid shifts resulting in increased cardiovascular strain as plasma volume declines [108, 109]. Additionally, the impaired cardiovascular function also leads to diminished skin blood flow resulting in a decline in the ability to dissipate heat to the environment [110]. Hypohydration also leads to an increase in the perception of exertion required to maintain a steady work rate. Research investigating the role of hydration on exercise performance has shown fluid ingestion and the maintenance of proper hydration status improve performance [102, 111].

To add to the physiological strain associated with dehydration inherent with physical activity, most athletes have been found to arrive to practice in a hypohydrated state [106, 112]. Even with education emphasizing the need for proper hydration, preactivity hydration assessments of athletes involved in sport have demonstrated significant percentages of the population to be hypohydrated [113]. Kavouras et al. reported that educational interventions significantly improved preexercise hydration status, however following education regarding hydration 66% of the youth athletes reported to practice in a hypohydrated state [114].

While unlikely to be a broad risk in youth sport, attention must also be drawn to the potential to ingest fluids at excessive levels resulting in the risk of hypernatremia and potentially death. Since the first reports in the literature of hyponatremia in endurance events greater focus has continued to grow regarding the risks associated with over drinking [115]. Hyponatremia has been reported to be as high as 51% in ultramarathons performed under hotter ambient temperatures [115]. The incidence of hyponatremia increases as duration of activity increases along with fluid ingestion. A potential means to reduce the rate of plasma sodium concentration decline is through the ingestion of sodium containing beverages [116]. It should be noted that very few beverages actually contain sodium levels sufficient to maintain plasma sodium levels; however the ingestion of sodium containing beverages will reduce the rate of decline.

The American College of Sports Medicine's Position Stand on Nutrition and Athletic Performance recommends athletes to consume 5–7 mL/kg of body mass 4 hours prior to exercise, enough fluid to reduce body mass changes to less than 2% during activity, and 450–675 mL for every 0.5 kg of body mass lost during exercise [15]. These recommendations do not account for age but are likely a good place to begin for youth athletes since they are based on body mass rather than absolute volumes. Hydration goals should be to minimize body mass losses associated with dehydration while ensuring fluid ingestion does not exceed sweat losses. This is easily assessed through body weight measures immediately prior to and immediately following activity. Increases in body mass would inform the athlete fluid ingestion was too high and significant decreases in body mass would inform the athlete fluid ingestion was insufficient. Additionally, in situations where repeated days of exercise are performed in the heat sodium ingestion rates should be increased to maintain plasma sodium levels. Flavored fluids have been repeatedly shown to aid in the maintenance of fluid intake reducing voluntary dehydration.

Equation for determining sweat rate is as follows:

$$\frac{BW_0 + DF_0 - BW_E - DF_E}{\text{Time (hrs)}}, \quad (1)$$

where BW_0 is body mass before exercise, DF_0 is mass of exercise food and drink before exercise, BW_E is body mass after exercise, and DF_E is mass of exercise food and drink after exercise.

7. Micronutrients

Micronutrients categorically refer to vitamins and minerals used by the body during normal physiological functions. Generally, it is accepted that a well-balanced diet of sufficient caloric intake will provide the adequate micronutrients to support normal growth and maturation. The American Medical Association (AMA) and the American Dietetic Association (ADA) recommend nutrients be obtained from food sources rather than supplements in healthy children [117]. Likewise, the American Academy of Pediatrics (AAP) does not endorse regular supplementation of vitamins and minerals in healthy children (with the exception of fluoride in unfluoridated areas). However, AAP has noted that some children are at increased risk of nutrient deficiencies. Specifically, AAP suggests that children and adolescents with anorexia or poor appetites, chronic diseases, and food insecurity are at greater risk for nutrient deficiencies. Youth who do not consume adequate amounts of dairy or have sufficient sun exposure may also be at risk for deficiencies [118].

The daily time constraints of an elite young athlete can make achieving a balanced diet difficult, thus putting these individuals at a potential increased risk for micronutrient deficiencies as well. These deficiencies are most commonly observed in girls rather than boys and in mineral intake rather than vitamin intake. Athletic youth may actually be more likely to achieve the recommended intake of vitamins than nonathletic youth due to their increased total caloric intake. Adequate intake of minerals appears to be a bigger challenge for youth, particularly girls [119]. Specifically, iron and calcium are frequently noted as common nutritional concerns among children and adolescents.

Iron deficiency and subsequent anemia are common in adolescents [120]. Increases in hemoglobin production, blood volume, and muscle mass are normal characteristics of growth and maturation and account for the majority of increased iron needs in developing adolescents. However, iron requirements become even greater for girls at the onset of menses. Iron-deficient anemia has significant, negative implications on performance in adults [121, 122] and youth [123–125]. Diminished performance is most apparent in iron-deficient anemic athletes participating in activities with higher aerobic demands (i.e., endurance event athletes) [124]. The recommended intake of iron for boys and girls aged 9–13 years is 18 mg per day. Boys and girls aged 14–18 years should consume 11 mg and 15 mg per day, respectively [126]. In accordance with AMA, ADA, and AAP guidelines, youth should improve their iron status through consumption of iron-rich

foods at meals such as red meat, beans, and green vegetables. Athletes will likely find additional benefit by including other iron-rich foods such as peanuts and dried fruits and iron-fortified cereals as regular snacks. Furthermore, the inclusion of foods higher in ascorbic acid with these nonheme iron sources will improve iron absorption from these snacks [127].

Calcium requirements are greatest during adolescence, 1,300 mg per day for both boys and girls [128]. This higher requirement accommodates the prime opportunity for acquisition of bone mass that spans the pubertal ages. Availability of nutrients critical for bone development (e.g., calcium) and opportunity for increased bone loading (e.g., physical activity) prior to achieving skeletal maturation is critical in preventing osteoporosis later in life [129]. Despite the greater requirements and the clear benefits of consuming adequate amounts, United States children and adolescents' average intake falls below the minimum recommendations [130]. In the American diet, the majority of dietary calcium is obtained from milk and other dairy sources [131]. However, milk consumption has shown a general decline in this age group [132]. Given the perishability of most dairy products, young athletes face practicality issues scheduling regular consumption throughout the day. Breakfast consumption is associated with greater calcium intake among this age group [133] and should be strongly encouraged. Likewise, athletes should consume regular snacks that include rich sources of calcium (e.g., fortified orange juice, almonds, and broccoli), throughout the day. Furthermore, calcium absorption is dependent on adequate levels of vitamin D [128]. Thus, attention to sources of calcium fortified with vitamin D is warranted, particularly among individuals who are not likely to be exposed to sufficient sunlight to endogenously produce adequate vitamin D. While whole food sources are always preferred, the convenience of calcium supplements (often fortified with vitamin D) may make adequate intake a more viable possibility.

An additional consideration warranting attention in young athletes participating in large amounts of training and competition is the potential need to replenish sodium and potassium lost in sweat. The summation of the electrolyte loss resulting from sweat loss has been shown to be equivalent to daily intakes, even in young athletes [105]. Much of this additional loss in salt can be offset through the ingestion of sport drinks during practice and competition, which also partially addresses hydration concerns in the group.

8. Summary

Research regarding the nutritional needs of young competitive athletes is sparse and is primarily composed of investigations of youth-adult differences. In addition to the limited research, the majority of our current knowledge in the adult population is based on differences between typical adults compared to their more active counterparts. Research to date suggests similarities in the caloric and macronutrient needs of active adults and their younger counterparts; however, youth-adult differences in fuel utilization have also been clearly demonstrated. In addition to the energy needs of highly active

TABLE 2: General nutrition recommendations for maintaining health.

Protein	15–20% of total calories should come from protein; 0.8–1.2 g/kg/day derived from whole food sources
Fat	>15% and <30% of total calories should come from fat
Carbohydrate	>50% of total calories should come from carbohydrates, or 3–8 g/kg/day
Micronutrients	Regular supplementation is not recommended in healthy children and adolescents consuming a balanced diet

TABLE 3: Supplemental nutrition recommendations for athletes.

Protein	1.2–1.8 g/kg/day derived from whole food sources <i>After exercise:</i> 20 g of high quality protein shortly after exercise
Carbohydrate	<i>During exercise:</i> 30–60 g/hr for exercise lasting more than 1 hour <i>After exercise:</i> 1.0–1.5 g/kg of body mass within 30 minutes of exercise cessation
Fluid	<i>Before exercise:</i> 5–7 mL/kg 4 hrs prior to exercise <i>During exercise:</i> assess sweat rate and develop hydration plan to maintain body mass during exercise <i>After exercise:</i> 450–675 mL/0.5 kg and additional sodium consideration to account for loss through sweat
Micronutrients	<i>During exercise:</i> sodium to offset losses associated with sweat being lost in sweat

youth, nutritional intake plays a critical role in the growth and development of young athletes and should be a principal emphasis at this stage in their lives. General guidelines for nutrition for active youth are summarized in Table 2. These guidelines serve as recommendations that support healthy growth and development and also account for the additional caloric needs of active youth. As emphasis on performance outcome goals continues to increase in youth athletics, active youth quickly become young athletes. Likewise, nutritional considerations move beyond increased caloric needs and promotion of healthy growth and development to nutritional strategies that can optimize performance. A summary of available performance-based nutritional strategies can be found in Table 3. The paucity of data examining the unique needs of young athletes draws attention to increased need for further investigations on this subject which consider the distinct nutritional requirements of growth and maturation at all age and skill levels. Given the popularity of youth sports and the increasing demand for performance outcomes, increased attention on this topic is warranted by the scientific community.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Authors' Contribution

Megan E. Holmes and Matthew J. McAllister are coauthors.

References

- [1] V. Seefeldt and M. Ewing, "Youth sports in America, an overview," *President's Council on Physical Fitness and Sports Research Digest*, vol. 2, no. 11, pp. 1–10, 1997.
- [2] American College of Sports Medicine and American Dietetic Association & Dietitians of Canada, "Joint Position Statement: nutrition and athletic performance," *Medicine & Science in Sports & Exercise*, vol. 32, no. 12, pp. 2130–2145, 2000.
- [3] D. Manoloff, "Noted surgeon Dr. James Andrew wants your young athlete to stay healthy by playing less," 2013, http://www.cleveland.com/dman/index.ssf/2013/02/noted_surgeon_dr_james_andrews.html.
- [4] J. Aucouturier, J. S. Baker, and P. Duché, "Fat and carbohydrate metabolism during submaximal exercise in children," *Sports Medicine*, vol. 38, no. 3, pp. 213–238, 2008.
- [5] G. E. Duncan and E. T. Howley, "Substrate metabolism during exercise in children and the crossover concept," *Pediatric Exercise Science*, vol. 11, no. 1, pp. 12–21, 1999.
- [6] J. S. Harrell, R. G. McMurray, C. D. Baggett, M. L. Pennell, P. F. Pearce, and S. I. Bangdiwala, "Energy costs of physical activities in children and adolescents," *Medicine and Science in Sports and Exercise*, vol. 37, no. 2, pp. 329–336, 2005.
- [7] Food and Agriculture Organization of the United Nations, United Nations University, and World Health Organization, *Human Energy Requirements: Report of a Joint FAO/WHO/UNU Expert Consultation: Rome, 17–24 October 2001*, Food and Agricultural Organization of the United Nations, Rome, Italy, 2004.
- [8] R. M. Malina, C. Bouchard, and O. Bar-Or, *Growth, Maturation, and Physical Activity*, Human Kinetics, Champaign, Ill, USA, 2nd edition, 2004.
- [9] E. Ravussin and C. Bogardus, "Relationship of genetics, age, and physical fitness to daily energy expenditure and fuel utilization," *The American Journal of Clinical Nutrition*, vol. 49, supplement 5, pp. 968–975, 1989.
- [10] S. M. Garn and L. C. Clark Jr., "The sex difference in the basal metabolic rate," *Child Development*, vol. 24, no. 3–4, pp. 215–224, 1953.
- [11] D. Nemet and A. Eliakim, "Pediatric sports nutrition: an update," *Current Opinion in Clinical Nutrition and Metabolic Care*, vol. 12, no. 3, pp. 304–309, 2009.
- [12] K. D. Tipton, A. E. Jeukendrup, and P. Hespel, "Nutrition for the sprinter," *Journal of Sports Sciences*, vol. 25, supplement 1, pp. S5–S15, 2007.
- [13] E. Smit, J. Nieto, C. J. Crespo, and P. Mitchell, "Estimates of animal and plant protein intake in US adults: results from the Third National Health and Nutrition Examination Survey, 1988–1991," *Journal of the American Dietetic Association*, vol. 99, no. 7, pp. 813–820, 1999.
- [14] S. M. Phillips and L. J. C. Van Loon, "Dietary protein for athletes: from requirements to optimum adaptation," *Journal of Sports Sciences*, vol. 29, supplement 1, pp. S29–S38, 2011.
- [15] N. R. Rodriguez, N. M. Di Marco, and S. Langley, "American College of Sports Medicine position stand. Nutrition and athletic performance," *Medicine & Science in Sports & Exercise*, vol. 41, no. 3, pp. 709–731, 2009.
- [16] N. Boisseau, M. Vermorel, M. Rance, P. Duché, and P. Patureau-Mirand, "Protein requirements in male adolescent soccer players," *European Journal of Applied Physiology*, vol. 100, no. 1, pp. 27–33, 2007.
- [17] F. Meyer, H. O'Connor, and S. M. Shirreffs, "Nutrition for the young athlete," *Journal of Sports Sciences*, vol. 25, supplement 1, pp. S73–S82, 2007.
- [18] P. W. Lemon, "Beyond the zone: protein needs of active individuals," *Journal of the American College of Nutrition*, vol. 19, supplement 5, pp. 513S–521S, 2000.
- [19] C. H. Murphy, A. J. Hector, and S. M. Phillips, "Considerations for protein intake in managing weight loss in athletes," *European Journal of Sport Science*, vol. 15, no. 1, pp. 21–28, 2015.
- [20] A. Jeukendrup and L. Cronin, "Nutrition and elite young athletes," *Medicine and Sport Science*, vol. 56, pp. 47–58, 2011.
- [21] S. Mettler, N. Mitchell, and K. D. Tipton, "Increased protein intake reduces lean body mass loss during weight loss in athletes," *Medicine and Science in Sports and Exercise*, vol. 42, no. 2, pp. 326–337, 2010.
- [22] D. K. Layman, R. A. Boileau, D. J. Erickson et al., "A reduced ratio of dietary carbohydrate to protein improves body composition and blood lipid profiles during weight loss in adult women," *Journal of Nutrition*, vol. 133, no. 2, pp. 411–417, 2003.
- [23] D. K. Layman, E. Evans, J. I. Baum, J. Seyler, D. J. Erickson, and R. A. Boileau, "Dietary protein and exercise have additive effects on body composition during weight loss in adult women," *Journal of Nutrition*, vol. 135, no. 8, pp. 1903–1910, 2005.
- [24] H. J. Leidy, N. S. Carnell, R. D. Mattes, and W. W. Campbell, "Higher protein intake preserves lean mass and satiety with weight loss in pre-obese and obese women," *Obesity*, vol. 15, no. 2, pp. 421–429, 2007.
- [25] K. A. Meckling and R. Sherfey, "A randomized trial of a hypocaloric high-protein diet, with and without exercise, on weight loss, fitness, and markers of the metabolic syndrome in overweight and obese women," *Applied Physiology, Nutrition, and Metabolism*, vol. 32, no. 4, pp. 743–752, 2007.
- [26] T. L. Halton and F. B. Hu, "The effects of high protein diets on thermogenesis, satiety and weight loss: a critical review," *The Journal of the American College of Nutrition*, vol. 23, no. 5, pp. 373–385, 2004.
- [27] J. D. Latner and M. Schwartz, "The effects of a high-carbohydrate, high-protein or balanced lunch upon later food intake and hunger ratings," *Appetite*, vol. 33, no. 1, pp. 119–128, 1999.
- [28] S. D. Poppitt, D. McCorMacK, and R. Buffenstein, "Short-term effects of macronutrient preloads on appetite and energy intake in lean women," *Physiology and Behavior*, vol. 64, no. 3, pp. 279–285, 1998.
- [29] A. R. Skov, S. Toubro, B. Rønn, L. Holm, and A. Astrup, "Randomized trial on protein vs carbohydrate in *ad libitum* fat reduced diet for the treatment of obesity," *International Journal of Obesity*, vol. 23, no. 5, pp. 528–536, 1999.
- [30] D. K. Layman, "Protein quantity and quality at levels above the RDA improves adult weight loss," *Journal of the American College of Nutrition*, vol. 23, no. 6, supplement, pp. 631s–636s, 2004.

- [31] J. C. Anthony, T. G. Anthony, S. R. Kimball, and L. S. Jefferson, "Signaling pathways involved in translational control of protein synthesis in skeletal muscle by leucine," *Journal of Nutrition*, vol. 131, no. 3, pp. 856s–860s, 2001.
- [32] R. Elango, K. Chapman, M. Rafii, R. O. Ball, and P. B. Pencharz, "Determination of the tolerable upper intake level of leucine in acute dietary studies in young men," *The American Journal of Clinical Nutrition*, vol. 96, no. 4, pp. 759–767, 2012.
- [33] M. C. Duellman, J. M. Lukaszuk, A. D. Prawitz, and J. P. Brandenburg, "Protein supplement users among high school athletes have misconceptions about effectiveness," *Journal of Strength and Conditioning Research*, vol. 22, no. 4, pp. 1124–1129, 2008.
- [34] M. Beelen, L. M. Burke, M. J. Gibala, and L. J. C. van Loon, "Nutritional strategies to promote postexercise recovery," *International Journal of Sport Nutrition and Exercise Metabolism*, vol. 20, no. 6, pp. 515–532, 2010.
- [35] D. Nemet, B. Wolach, and A. Eliakim, "Proteins and amino acid supplementation in sports: are they truly necessary?" *Israel Medical Association Journal*, vol. 7, no. 5, pp. 328–332, 2005.
- [36] K. Froiland, W. Koszewski, J. Hingst, and L. Kopecky, "Nutritional supplement use among college athletes and their sources of information," *International Journal of Sport Nutrition and Exercise Metabolism*, vol. 14, no. 1, pp. 104–120, 2004.
- [37] B. H. Jacobson and S. G. Aldana, "Current nutrition practice and knowledge of varsity athletes," *The Journal of Strength & Conditioning Research*, vol. 6, no. 4, pp. 232–238, 1992.
- [38] B. G. Wiita and I. A. Stombaugh, "Nutrition knowledge, eating practices, and health of adolescent female runners: a 3-year longitudinal study," *International Journal of Sport Nutrition and Exercise Metabolism*, vol. 6, no. 4, pp. 414–425, 1996.
- [39] S. S. Jonnalagadda, C. A. Rosenbloom, and R. Skinner, "Dietary practices, attitudes, and physiological status of collegiate freshman football players," *Journal of Strength and Conditioning Research*, vol. 15, no. 4, pp. 507–513, 2001.
- [40] C. A. Rosenbloom, S. S. Jonnalagadda, and R. Skinner, "Nutrition knowledge of collegiate athletes in a Division I National Collegiate Athletic Association institution," *Journal of the American Dietetic Association*, vol. 102, no. 3, pp. 418–420, 2002.
- [41] B. H. Jacobson, C. Sobonya, and J. Ransone, "Nutrition practices and knowledge of college varsity athletes: a follow-up," *Journal of Strength and Conditioning Research*, vol. 15, no. 1, pp. 63–68, 2001.
- [42] A. H. Lichtenstein, E. Kennedy, P. Barrier et al., "Dietary fat consumption and health," *Nutrition Reviews*, vol. 56, no. 5, part 2, pp. S3–S28, 1998.
- [43] J. W. Smith and A. Jeukendrup, "Performance nutrition for young athletes," in *Nutrition and Enhanced Sports Performance: Muscle Building, Endurance, and Strength*, S. D. N. Bagchi and C. K. Sen, Eds., pp. 523–529, Elsevier, 2013.
- [44] R. Wahl, "Nutrition in the adolescent," *Pediatric Annals*, vol. 28, no. 2, pp. 107–111, 1999.
- [45] P. Bjorntorp, "Importance of fat as a support nutrient for energy: metabolism of athletes," *Journal of Sports Sciences*, vol. 9, pp. 71–76, 1991.
- [46] B. A. Bushman, "Calorie requirements for young competitive female athletes," *ACSM's Health & Fitness Journal*, vol. 16, no. 5, pp. 4–8, 2012.
- [47] E. Jequier, "Is fat intake a risk factor for fat gain in children?" *The Journal of Clinical Endocrinology & Metabolism*, vol. 86, no. 3, pp. 980–983, 2001.
- [48] R. J. Bloomer, M. M. Kabir, K. E. Marshall, R. E. Canale, and T. M. Farney, "Postprandial oxidative stress in response to dextrose and lipid meals of differing size," *Lipids in Health and Disease*, vol. 9, article 79, 2010.
- [49] K. H. Fisher-Wellman and R. J. Bloomer, "Exacerbated postprandial oxidative stress induced by the acute intake of a lipid meal compared to isoenergetically administered carbohydrate, protein, and mixed meals in young, healthy men," *Journal of the American College of Nutrition*, vol. 29, no. 4, pp. 373–381, 2010.
- [50] C. Bouziotas, Y. Koutedakis, A. Nevill et al., "Greek adolescents, fitness, fatness, fat intake, activity, and coronary heart disease risk," *Archives of Disease in Childhood*, vol. 89, no. 1, pp. 41–44, 2004.
- [51] L. T. Mahoney, T. L. Burns, W. Stanford et al., "Coronary risk factors measured in childhood and young adult life are associated with coronary artery calcification in young adults: the Muscatine study," *Journal of the American College of Cardiology*, vol. 27, no. 2, pp. 277–284, 1996.
- [52] M. Máček and J. Vávra, "Prolonged exercise in 14-year-old girls," *International Journal of Sports Medicine*, vol. 2, no. 4, pp. 228–230, 1981.
- [53] American Academy of Pediatrics and National Cholesterol Education Program, "Report of the expert panel on blood cholesterol levels in children and adolescents," *Pediatrics*, vol. 89, no. 3, part 2, pp. 525–584, 1992.
- [54] J. A. Cooper, A. C. Watras, T. Shriver, A. K. Adams, and D. A. Schoeller, "Influence of dietary fatty acid composition and exercise on changes in fat oxidation from a high-fat diet," *Journal of Applied Physiology*, vol. 109, no. 4, pp. 1011–1018, 2010.
- [55] R. J. Tunstall, K. A. Mehan, G. D. Wadley et al., "Exercise training increases lipid metabolism gene expression in human skeletal muscle," *The American Journal of Physiology—Endocrinology and Metabolism*, vol. 283, no. 1, pp. E66–E72, 2002.
- [56] B. W. Timmons, O. Bar-Or, and M. C. Riddell, "Energy substrate utilization during prolonged exercise with and without carbohydrate intake in preadolescent and adolescent girls," *Journal of Applied Physiology*, vol. 103, no. 3, pp. 995–1000, 2007.
- [57] B. W. Timmons, O. Bar-Or, and M. C. Riddell, "Influence of age and pubertal status on substrate utilization during exercise with and without carbohydrate intake in healthy boys," *Applied Physiology, Nutrition and Metabolism*, vol. 32, no. 3, pp. 416–425, 2007.
- [58] B. W. Timmons, O. Bar-Or, and M. C. Riddell, "Oxidation rate of exogenous carbohydrate during exercise is higher in boys than in men," *Journal of Applied Physiology*, vol. 94, no. 1, pp. 278–284, 2003.
- [59] G. J. Welk, C. B. Corbin, and D. Dale, "Measurement issues in the assessment of physical activity in children," *Research Quarterly for Exercise and Sport*, vol. 71, no. 2, pp. S59–S73, 2000.
- [60] M. C. Riddell, V. K. Jamnik, K. E. Iscoe, B. W. Timmons, and N. Gledhill, "Fat oxidation rate and the exercise intensity that elicits maximal fat oxidation decreases with pubertal status in young male subjects," *Journal of Applied Physiology*, vol. 105, no. 2, pp. 742–748, 2008.
- [61] J. O. Holloszy and E. F. Coyle, "Adaptations of skeletal muscle to endurance exercise and their metabolic consequences," *Journal of Applied Physiology Respiratory Environmental and Exercise Physiology*, vol. 56, no. 4, pp. 831–838, 1984.
- [62] A. Raben, E. Mygind, and A. Astrup, "Lower activity of oxidative key enzymes and smaller fiber areas in skeletal muscle

- of postobese women," *The American Journal of Physiology—Endocrinology and Metabolism*, vol. 275, no. 3, part 1, pp. E487–E494, 1998.
- [63] P. M. Berthon, R. A. Howlett, G. J. F. Heigenhauser, and L. L. Spriet, "Human skeletal muscle carnitine palmitoyltransferase I activity determined in isolated intact mitochondria," *Journal of Applied Physiology*, vol. 85, no. 1, pp. 148–153, 1998.
- [64] K. C. Hansen, Z. Zhang, T. Gomez, A. K. Adams, and D. A. Schoeller, "Exercise increases the proportion of fat utilization during short-term consumption of a high-fat diet," *The American Journal of Clinical Nutrition*, vol. 85, no. 1, pp. 109–116, 2007.
- [65] P. J. H. Jones, J. E. Ridgen, P. T. Phang, and C. L. Birmingham, "Influence of dietary fat polyunsaturated to saturated ratio on energy substrate utilization in obesity," *Metabolism*, vol. 41, no. 4, pp. 396–401, 1992.
- [66] J. C. Lovejoy, S. R. Smith, C. M. Champagne et al., "Effects of diets enriched in saturated (palmitic), monounsaturated (oleic), or trans (elaidic) fatty acids on insulin sensitivity and substrate oxidation in healthy adults," *Diabetes Care*, vol. 25, no. 8, pp. 1283–1288, 2002.
- [67] L. S. Piers, K. Z. Walker, R. M. Stoney, M. J. Soares, and K. O'Dea, "Substitution of saturated with monounsaturated fat in a 4-week diet affects body weight and composition of overweight and obese men," *British Journal of Nutrition*, vol. 90, no. 3, pp. 717–727, 2003.
- [68] P. W. Siri-Tarino, Q. Sun, F. B. Hu, and R. M. Krauss, "Meta-analysis of prospective cohort studies evaluating the association of saturated fat with cardiovascular disease," *The American Journal of Clinical Nutrition*, vol. 91, no. 3, pp. 535–546, 2010.
- [69] F. Macaluso, R. Barone, P. Catanese et al., "Do fat supplements increase physical performance?" *Nutrients*, vol. 5, no. 2, pp. 509–524, 2013.
- [70] A. E. Jeukendrup and S. Aldred, "Fat supplementation, health, and endurance performance," *Nutrition*, vol. 20, no. 7–8, pp. 678–688, 2004.
- [71] É. E. Dewailly, C. Blanchet, S. Gingras et al., "Relations between n-3 fatty acid status and cardiovascular disease risk factors among Quebecers," *The American Journal of Clinical Nutrition*, vol. 74, no. 5, pp. 603–611, 2001.
- [72] A. M. Hill, J. D. Buckley, K. J. Murphy, and P. R. C. Howe, "Combining fish-oil supplements with regular aerobic exercise improves body composition and cardiovascular disease risk factors," *The American Journal of Clinical Nutrition*, vol. 85, no. 5, pp. 1267–1274, 2007.
- [73] P. M. M. Andrade, B. G. Ribeiro, M. T. Bozza, L. F. B. C. Rosa, and M. G. T. do Carmo, "Effects of the fish-oil supplementation on the immune and inflammatory responses in elite swimmers," *Prostaglandins, Leukotrienes & Essential Fatty Acids*, vol. 77, no. 3–4, pp. 139–145, 2007.
- [74] T. H. Lee, R. L. Hoover, J. D. Williams et al., "Effect of dietary enrichment with eicosapentaenoic and docosahexaenoic acids on in vitro neutrophil and monocyte leukotriene generation and neutrophil function," *The New England Journal of Medicine*, vol. 312, no. 19, pp. 1217–1224, 1985.
- [75] M. Bortolotti, L. Tappy, and P. Schneiter, "Fish oil supplementation does not alter energy efficiency in healthy males," *Clinical Nutrition*, vol. 26, no. 2, pp. 225–230, 2007.
- [76] J. D. Buckley, S. Burgess, K. J. Murphy, and P. R. C. Howe, "DHA-rich fish oil lowers heart rate during submaximal exercise in elite Australian Rules footballers," *Journal of Science and Medicine in Sport*, vol. 12, no. 4, pp. 503–507, 2009.
- [77] G. E. Peoples, P. L. McLennan, P. R. C. Howe, and H. Groeller, "Fish oil reduces heart rate and oxygen consumption during exercise," *Journal of Cardiovascular Pharmacology*, vol. 52, no. 6, pp. 540–547, 2008.
- [78] B. Tartibian, B. H. Maleki, and A. Abbasi, "The effects of omega-3 supplementation on pulmonary function of young wrestlers during intensive training," *Journal of Science and Medicine in Sport*, vol. 13, no. 2, pp. 281–286, 2010.
- [79] G. S. Oostenbrug, R. P. Mensink, M. R. Hardeman, T. De Vries, F. Brouns, and G. Hornstra, "Exercise performance, red blood cell deformability, and lipid peroxidation: effects of fish oil and vitamin E," *Journal of Applied Physiology*, vol. 83, no. 3, pp. 746–752, 1997.
- [80] T. Raastad, A. T. Hostmark, and S. B. Stramme, "Omega-3 fatty acid supplementation does not improve maximal aerobic power, anaerobic threshold and running performance in well-trained soccer players," *Scandinavian Journal of Medicine and Science in Sports*, vol. 7, no. 1, pp. 25–31, 1997.
- [81] C. Y. Guezennec, J. F. Nadaud, P. Satabin, F. Leger, and P. Lafargue, "Influence of polyunsaturated fatty acid diet on the hemorrheological response to physical exercise in hypoxia," *International Journal of Sports Medicine*, vol. 10, no. 4, pp. 286–291, 1989.
- [82] M. J. Heilskov Rytter, L. B. B. Andersen, T. Houmann et al., "Diet in the treatment of ADHD in children—a systematic review of the literature," *Nordic Journal of Psychiatry*, vol. 69, no. 1, pp. 1–18, 2014.
- [83] H. Yang, P. Xun, and K. He, "Fish and fish oil intake in relation to risk of asthma: a systematic review and meta-analysis," *PLoS ONE*, vol. 8, no. 11, Article ID e80048, 2013.
- [84] H. B. MacDonald, "Conjugated linoleic acid and disease prevention: a review of current knowledge," *Journal of the American College of Nutrition*, vol. 19, no. 2, supplement, pp. 111s–118s, 2000.
- [85] Y. W. Wang and P. J. Jones, "Conjugated linoleic acid and obesity control: efficacy and mechanisms," *International Journal of Obesity and Related Metabolic Disorders*, vol. 28, no. 8, pp. 941–955, 2004.
- [86] K. L. Zambell, N. L. Keim, M. D. Van Loan et al., "Conjugated linoleic acid supplementation in humans: effects on body composition and energy expenditure," *Lipids*, vol. 35, no. 7, pp. 777–782, 2000.
- [87] L. M. Burke, B. Kiens, and J. L. Ivy, "Carbohydrates and fat for training and recovery," *Journal of Sports Sciences*, vol. 22, no. 1, pp. 15–30, 2004.
- [88] L. M. Burke, J. A. Hawley, S. H. S. Wong, and A. E. Jeukendrup, "Carbohydrates for training and competition," *Journal of Sports Sciences*, vol. 29, no. 1, pp. S17–S27, 2011.
- [89] L. Bonci, "Sports nutrition for young athletes," *Pediatric Annals*, vol. 39, no. 5, pp. 300–306, 2010.
- [90] P. M. Nisevich, "Sports nutrition for young athletes," *IDEA Fitness Journal*, pp. 65–67, 2008.
- [91] B. Gordon, L. A. Kohn, S. A. Levine, M. Matton, W. M. Scriver, and W. B. Whiting, "Sugar content of the blood following a marathon race with especial reference to the prevention of hypoglycemia: further observations," *The Journal of the American Medical Association*, vol. 85, no. 7, pp. 508–509, 1925.
- [92] S. A. Levine, B. Gordon, and C. L. Derick, "Some changes in the chemical constituents of the blood following a marathon race with special reference to the development of hypoglycemia," *The Journal of the American Medical Association*, vol. 82, no. 22, pp. 1778–1779, 1924.

- [93] J. Bergstrom and E. Hultman, "A study of the glycogen metabolism during exercise in man," *Scandinavian Journal of Clinical & Laboratory Investigation*, vol. 19, no. 3, pp. 218–228, 1967.
- [94] L. Hermansen, E. Hultman, and B. Saltin, "Muscle glycogen during prolonged severe exercise," *Acta Physiologica Scandinavica*, vol. 71, no. 2, pp. 129–139, 1967.
- [95] D. L. Costill, W. M. Sherman, W. J. Fink, C. Maresh, M. Witten, and J. M. Miller, "The role of dietary carbohydrates in muscle glycogen resynthesis after strenuous running," *American Journal of Clinical Nutrition*, vol. 34, no. 9, pp. 1831–1836, 1981.
- [96] E. F. Coyle and A. R. Coggan, "Effectiveness of carbohydrate feeding in delaying fatigue during prolonged exercise," *Sports Medicine*, vol. 1, no. 6, pp. 446–458, 1984.
- [97] J. W. Smith, D. D. Pascoe, D. H. Passe et al., "Curvilinear dose-response relationship of carbohydrate (0–120 g.h⁻¹) and performance," *Medicine & Science in Sports & Exercise*, vol. 45, no. 2, pp. 336–341, 2013.
- [98] J. W. Smith, J. J. Zachwieja, F. Péronnet et al., "Fuel selection and cycling endurance performance with ingestion of [¹³C]glucose: evidence for a carbohydrate dose response," *Journal of Applied Physiology*, vol. 108, no. 6, pp. 1520–1529, 2010.
- [99] P. D. Gollnick, K. Piehl, and B. Saltin, "Selective glycogen depletion pattern in human muscle fibres after exercise of varying intensity and at varying pedalling rates," *The Journal of Physiology*, vol. 241, no. 1, pp. 45–57, 1974.
- [100] W. M. Sherman, D. L. Costill, W. J. Fink, and J. M. Miller, "Effect of exercise-diet manipulation on muscle glycogen and its subsequent utilization during performance," *International Journal of Sports Medicine*, vol. 2, no. 2, pp. 114–118, 1981.
- [101] M. C. Riddell, O. Bar-Or, H. P. Schwarcz, and G. J. F. Heigenhauser, "Substrate utilization in boys during exercise with [¹³C]glucose ingestion," *European Journal of Applied Physiology*, vol. 83, no. 4–5, pp. 441–448, 2000.
- [102] K. A. Dougherty, L. B. Baker, M. Chow, and W. L. Kenney, "Two percent dehydration impairs and six percent carbohydrate drink improves boys basketball skills," *Medicine & Science in Sports & Exercise*, vol. 38, no. 9, pp. 1650–1658, 2006.
- [103] H. A. P. Batatinha, C. E. da Costa, E. de França et al., "Carbohydrate use and reduction in number of balance beam falls: implications for mental and physical fatigue," *Journal of the International Society of Sports Nutrition*, vol. 10, article 32, 2013.
- [104] J. W. Smith, K. A. Lee, J. P. Dobson, T. J. Roberts, and A. E. Jeukendrup, "Fluid and carbohydrate ingestion improve performance in American football players," *Medicine & Science in Sports & Exercise*, vol. 45, no. 5, supplement, p. 549, 2013.
- [105] H. J. Petrie, E. A. Stover, and C. A. Horswill, "Nutritional concerns for the child and adolescent competitor," *Nutrition*, vol. 20, no. 7–8, pp. 620–631, 2004.
- [106] E. A. Stover, J. Zachwieja, J. Stofan, R. Murray, and C. A. Horswill, "Consistently high urine specific gravity in adolescent American football players and the impact of an acute drinking strategy," *International Journal of Sports Medicine*, vol. 27, no. 4, pp. 330–335, 2006.
- [107] D. J. Casa, L. E. Armstrong, S. K. Hillman et al., "National athletic trainers' association position statement: fluid replacement for athletes," *Journal of Athletic Training*, vol. 35, no. 2, pp. 212–224, 2000.
- [108] J. González-Alonso, R. Mora-Rodríguez, P. R. Below, and E. F. Coyle, "Dehydration markedly impairs cardiovascular function in hyperthermic endurance athletes during exercise," *Journal of Applied Physiology*, vol. 82, no. 4, pp. 1229–1236, 1997.
- [109] S. J. Montain and E. F. Coyle, "Influence of graded dehydration on hyperthermia and cardiovascular drift during exercise," *Journal of Applied Physiology*, vol. 73, no. 4, pp. 1340–1350, 1992.
- [110] R. Murray, "Dehydration, hyperthermia, and athletes: science and practice," *Journal of Athletic Training*, vol. 31, no. 3, pp. 248–252, 1996.
- [111] L. B. Baker, D. E. Conroy, and W. L. Kenney, "Dehydration impairs vigilance-related attention in male basketball players," *Medicine and Science in Sports and Exercise*, vol. 39, no. 6, pp. 976–983, 2007.
- [112] G. Arnaoutis, S. A. Kavouras, Y. P. Kotsis, Y. E. Tsekouras, M. Makrillos, and C. N. Bardis, "Ad libitum fluid intake does not prevent dehydration in suboptimally hydrated young soccer players during a training session of a summer camp," *International Journal of Sport Nutrition and Exercise Metabolism*, vol. 23, no. 3, pp. 245–251, 2013.
- [113] N. R. Decher, D. J. Casa, S. W. Yeargin et al., "Hydration status, knowledge, and behavior in youths at summer sports camps," *International Journal of Sports Physiology and Performance*, vol. 3, no. 3, pp. 262–278, 2008.
- [114] S. A. Kavouras, G. Arnaoutis, M. Makrillos et al., "Educational intervention on water intake improves hydration status and enhances exercise performance in athletic youth," *Scandinavian Journal of Medicine and Science in Sports*, vol. 22, no. 5, pp. 684–689, 2012.
- [115] M. D. Hoffman, T. Hew-Butler, and K. J. Stuempfle, "Exercise-associated hyponatremia and hydration status in 161-km ultramarathoners," *Medicine and Science in Sports and Exercise*, vol. 45, no. 4, pp. 784–791, 2013.
- [116] D. M. J. Vrijens and N. J. Rehrer, "Sodium-free fluid ingestion decreases plasma sodium during exercise in the heat," *Journal of Applied Physiology*, vol. 86, no. 6, pp. 1847–1851, 1999.
- [117] American Dietetic Association, "Position of the American Dietetic Association: food fortification and dietary supplements," *Journal of the American Dietetic Association*, vol. 101, no. 1, pp. 115–125, 2001.
- [118] American Academy of Pediatrics, Committee on Nutrition, and L. A. Barnes, *Pediatric Nutrition Handbook*, American Academy of Pediatrics, Elk Grove Village, Ill, USA, 6th edition, 2009.
- [119] T. Rankinen, M. Fogelholm, U. Kujala, R. Rauramaa, and M. Uusitupa, "Dietary intake and nutritional status of athletic and nonathletic children in early puberty," *International Journal of Sport Nutrition*, vol. 5, no. 2, pp. 136–150, 1995.
- [120] World Health Organization, *Nutrition in Adolescence—Issues and Challenges for the Health Sector*, WHO, Geneva, Switzerland, 2005.
- [121] B. Ekblom, A. N. Goldberg, and B. Gullbring, "Response to exercise after blood loss and reinfusion," *Journal of Applied Physiology*, vol. 33, no. 2, pp. 175–180, 1972.
- [122] P. S. Hinton, "Iron and the endurance athlete," *Applied Physiology, Nutrition, and Metabolism*, vol. 39, no. 9, pp. 1012–1018, 2014.
- [123] J. Malczewska, G. Raczyński, and R. Stupnicki, "Iron status in female endurance athletes and in non-athletes," *International Journal of Sport Nutrition and Exercise Metabolism*, vol. 10, no. 3, pp. 260–276, 2000.
- [124] R. A. Raunikaar and H. Sabio, "Anemia in the adolescent athlete," *American Journal of Diseases of Children*, vol. 146, no. 10, pp. 1201–1205, 1992.

- [125] T. W. Rowland, S. A. Black, and J. F. Kelleher, "Iron deficiency in adolescent endurance athletes," *Journal of Adolescent Health Care*, vol. 8, no. 4, pp. 322–326, 1987.
- [126] Institute of Medicine (U.S.) Panel on Micronutrients, *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc*, National Academy Press, Washington, DC, USA, 2001.
- [127] J. D. Cook and M. B. Reddy, "Effect of ascorbic acid intake on nonheme-iron absorption from a complete diet," *The American Journal of Clinical Nutrition*, vol. 73, no. 1, pp. 93–98, 2001.
- [128] A. C. Ross, J. E. Manson, S. A. Abrams et al., "The 2011 report on dietary reference intakes for calcium and vitamin D from the institute of medicine: what clinicians need to know," *Journal of Clinical Endocrinology and Metabolism*, vol. 96, no. 1, pp. 53–58, 2011.
- [129] J.-P. Bonjour and T. Chevalley, "Pubertal timing, bone acquisition, and risk of fracture throughout life," *Endocrine Reviews*, vol. 35, no. 5, pp. 820–847, 2014.
- [130] A. C. Looker, C. M. Loria, M. D. Carroll, M. A. McDowell, and C. L. Johnson, "Calcium intakes of Mexican Americans, Cubans, Puerto Ricans, non-Hispanic whites, and non-Hispanic blacks in the United States," *Journal of the American Dietetic Association*, vol. 93, no. 11, pp. 1274–1279, 1993.
- [131] L. Gueguen and A. Pointillart, "The bioavailability of dietary calcium," *Journal of the American College of Nutrition*, vol. 19, no. 2, pp. 119S–136S, 2000.
- [132] C. Cavadini, A. M. Siega-Riz, and B. M. Popkin, "U.S. adolescent food intake trends from 1965 to 1996," *Archives of Disease in Childhood*, vol. 83, no. 1, pp. 18–24, 2000.
- [133] B. A. Barton, A. L. Eldridge, D. Thompson et al., "The relationship of breakfast and cereal consumption to nutrient intake and body mass index: the National Heart, Lung, and Blood Institute Growth and Health Study," *Journal of the American Dietetic Association*, vol. 105, no. 9, pp. 1383–1389, 2005.